

Alternative reproductive strategies in black-winged territorial males of *Paraphlebia zoe* (Odonata, Thaumatoneturidae)

Anais Rivas-Torres ^{Corresp., 1}, Rosa Ana Sánchez-Guillén ², Adolfo Cordero-Rivera ¹

¹ ECOEVO Lab, Departamento de Ecología e Biología Animal, Universidade de Vigo, Pontevedra, Galiza, Spain

² Department of Ecology and Animal Biology, INECOL, Xalapa, Veracruz, Mexico

Corresponding Author: Anais Rivas-Torres
Email address: anarivas@uvigo.es

Alternative reproductive strategies are commonly associated with male dimorphism. In *Paraphlebia zoe*, a species of damselfly whose males are dimorphic in wing coloration, black-and-white-winged (BW) males defend territories, while hyaline-winged (HW) males usually play the role of satellites. We found that several BW males can sometimes share a territory, and we hypothesized that within this morph there are two alternative tactics: submissive and dominant. We conducted an experiment to test whether dominant and submissive roles are plastic or stable and fixed on each individual. To this end, we manipulated black and white spots of BW males in four treatments: (i) painting over white and black spots without changing their size, (ii) erasing the white spot using black painting, (iii) increasing the black spot and moving the white spot maintaining its size and (iv) control males. Additionally, we investigated the correlation between some phenotypic variables (wing asymmetry, survival and recapture probabilities) and male behaviour (in terms of quality of the territory). We found that the two behavioural roles (submissive and dominant) were not affected by the manipulative experiments, therefore suggesting that they are stable and fixed. Additionally, we found a positive correlation between body size and survival in both sexes, and a positive effect of territory quality and lifespan on mating success. Moreover, the largest and youngest BW males were the most symmetrical. We conclude that *Paraphlebia zoe* holds high behavioural diversity, with two types of strategies in BW males, dominant and submissive. The occurrence of this intra-morph behavioural diversity might depend on demographic factors such as population density and/or the relative frequency of the different morphs.

1 **Alternative reproductive strategies in black-winged territorial males of *Paraphlebia zoe* (Odonata,**
2 **Thaumatoneuridae)**

3 Anais Rivas-Torres¹, Rosa Ana Sánchez-Guillén² & Adolfo Cordero-Rivera¹

4 ¹ECOEVO Lab, Departamento de Ecoloxía e Bioloxía Animal, Universidade de Vigo, Pontevedra, Galiza,
5 Spain

6 ²Department of Ecology and Animal Biology, INECOL, Xalapa, Veracruz, Mexico

7 **Author for correspondence:** Anais Rivas-Torres, **address:** ECOEVO Lab, Universidade de Vigo, Escola de
8 Enxeñaría Forestal, Campus A Xunqueira, 36005 Pontevedra, Galiza, Spain, **telephone number:**
9 0034657885598 **email:** arivasto@gmail.com

10 **Running title:** Alternative reproductive tactics in a dimorphic damselfly.

11 ABSTRACT

12 Alternative reproductive strategies are commonly associated with male dimorphism. In *Paraphlebia zoe*,
 13 a species of damselfly whose males are dimorphic in wing coloration, black-and-white-winged (BW) males
 14 defend territories, while hyaline-winged (HW) males usually play the role of satellites. We found that
 15 several BW males can sometimes share a territory, and we hypothesized that within this morph there are
 16 two alternative tactics: submissive and dominant. We conducted an experiment to test whether dominant
 17 and submissive roles are plastic or stable and fixed on each individual. To this end, we manipulated black
 18 and white spots of BW males in four treatments: (i) painting over white and black spots without changing
 19 their size, (ii) erasing the white spot using black painting, (iii) increasing the black spot and moving the
 20 white spot maintaining its size and (iv) control males. Additionally, we investigated the correlation
 21 between some phenotypic variables (wing asymmetry, survival and recapture probabilities) and male
 22 behaviour (in terms of quality of the territory). We found that the two behavioural roles (submissive and
 23 dominant) were not affected by the manipulative experiments, therefore suggesting that they are stable
 24 and fixed. Additionally, we found a positive correlation between body size and survival in both sexes, and
 25 a positive effect of territory quality and lifespan on mating success. Moreover, the largest and youngest
 26 BW males were the most symmetrical. We conclude that *Paraphlebia zoe* holds high behavioural diversity,
 27 with two types of strategies in BW males, dominant and submissive. The occurrence of this intra-morph
 28 behavioural diversity might depend on demographic factors such as population density and/or the relative
 29 frequency of the different morphs.

30

INTRODUCTION

The existence of alternative reproductive behaviours in individuals of the same sex is widespread in nature. Most of examples of alternative reproductive phenotypes are readily interpreted as alternative tactics within a conditional strategy, *i.e.* monomorphic individuals adopting one strategy or another, based on a condition-dependent trait, such as body size, fat reserves and so on (Gross, 1996). In contrast, alternative strategies linked to genetic polymorphisms are uncommon, and they are usually associated with male polymorphism in insects and other invertebrates. Examples include wing dimorphism in odonates, and size in horned beetles and cuttlefish (Nomakuchi, 1992; Emlen, 1997; Norman, Finn & Tregenza, 1999). In polymorphic species, each morph is usually associated with different behaviours, like two morphs of newts which differed in boldness (Winandy & Denoël, 2015). In harvestmen, a case of trimorphism in chelicerae size has been recently reported, associated with alternative reproductive tactics (Painting et al., 2015).

In odonates, colour polymorphism is common in females (Fincke et al., 2005; Van Gossum, Sherratt & Cordero-Rivera, 2008; Sánchez-Guillén et al., 2017). Female-limited colour polymorphism (thorax coloration) is associated with alternative reproductive strategies, normally one female has a male-like coloration and behaviour (androchrome; behaving more aggressively than the gynochrome morphs). However, male-limited colour polymorphism (wing coloration) is a rare phenomenon in damselflies and only found in some families like Calopterygidae (genus *Mnais*; Siva-Jothy and Tsubaki 1989; Plaistow and Tsubaki 2000), Platystictidae (Gonzalez-Soriano, Novelo-Gutierrez & Verdugo-Garza, 1982), Synlestidae (Samways, 2006) and in the genera *Thaumatoneura* and *Paraphlebia*, (Garrison, von Ellenrieder, & Louton, 2010; Paulson, 2016). In *Diastatops obscura* (Libellulidae), there are three types of males: territorial males near the riverbank (the best territories), territorial males in the middle of the river, and satellites in the riverbank vegetation, who do not defend territories (Bañuelos Irusta & Araújo, 2007). Territories can be even shared by several males, like in *Libellula luctuosa*, and are surrounded by non-territorial satellite males (Moore, 1987).

The damselfly *Paraphlebia zoe* Selys in Hagen, 1861, is a polymorphic species, with two male colour morphs, one with coloured wings, with a black spot and a white stripe (BW), while the other morph shows hyaline wings (HW), similar to females. In *Megaloprepus caerulatus*, a monomorphic species whose males have a similar wing pattern to BW *P. zoe* males, it has been found that white wing bands of males are correlated with territory residence and they might be a signal of male condition, because they are based on wax (Schultz & Fincke, 2009; Xu & Fincke, 2015). In *P. zoe*, each morph has been associated with a specific reproductive strategy. BW males are more territorial and aggressive than HW males, secure more matings and copulate for shorter times (Romo-Beltrán, Macías-Ordóñez & Córdoba-Aguilar, 2009). This species is therefore a typical example of genetic polymorphism associated with alternative reproductive strategies, presumably maintained by frequency-dependent selection (Gross, 1996). Some BW males that do not defend a territory have been previously found in *P. zoe* (Romo-Beltrán, Macías-Ordóñez & Córdoba-Aguilar, 2009), and are considered as satellite males. Our preliminary observations suggested that BW territorial males can behave as *dominants*, particularly when their territory is close to the water or as *submissive*, when they occupy a territory away from the water, in a similar way as the behaviour described for *D. obscura* (Bañuelos Irusta & Araújo, 2007). Dominant BW males are highly territorial and highly aggressive against submissive males, probably giving them resource advantages (Baker, 1983), and submissive males are less aggressive, usually avoiding confrontation with dominants, but nevertheless defend their territory against satellites. *Paraphlebia zoe* does not show precopulatory courtship (Wong-Muñoz et al., 2013), consequently, females could be choosing territory quality instead of male quality.

Previous studies have indicated that longevity and body size are among the main correlates of male reproductive success in damselflies (e.g. Banks & Thompson, 1987; Cordero, 1995). The main aim of our study was to determine whether dominant and submissive roles of BW males are related to phenotypic characters, in particular body size, wing asymmetry and size of the wing pigmentation (black spot and a white stripe), and how this behavioural polymorphism relates to reproductive success. To this end, we first, using mark-recapture methods, estimated survivorship rates and the effect of body size on survival of BW males (dominant and submissive). Second, fluctuating asymmetry (Forbes, Leung & Schalk, 1997; Beck & Pruett-Jones, 2002) has been widely used as a correlate of genetic quality, because it is presumed that, for traits with bilateral expression, small deviations between sides are due to the inability of the organism to buffer environmental stressors (Leung & Forbes, 1997a). In odonates, some contrasting results about the effect of asymmetry on reproductive success have been found. Harvey & Walsh (1993) found a negative effect, while others did not find any effect (Leung & Forbes, 1997b; Koga et al., 1998; Carchini et al., 2001). We therefore analysed the effect of male wing spot asymmetry on mating success of BW males (taking into account territory quality and male strategy). Third, some studies have suggested that male mating success might be more related to the ability to be in the right place at the right time, which is expected to be related with the capacity to maintain the territory (Switzer, 2002). For this reason, we studied mating behaviour and evaluated which factors affect copulation duration, including male phenotype, time of day and temperature (e.g. Cordero, 1990), because these variables might affect territory tenancy and hence reproductive success. Finally, we designed a manipulative experiment to test whether dominant and submissive BW male roles are plastic or fixed over time. Based on previous studies with odonates that have found that more pigmented males have higher success (e.g. Alex Córdoba-Aguilar, 2002) we hypothesized that increasing the black-wing spot of the BW submissive males would change their status towards dominance. This prediction is based on the idea that males adapt their aggressiveness to the result of previous interactions, so that other males would tend to become submissive in confrontations with a male with an enlarged black spot, and the manipulated male would increase its aggressiveness (Senar & Camerino, 1998). However, increasing the black spot size of the BW dominant males would not affect their status, because they are already dominant. For the white spot, there is no information about its function in this species. Therefore, if we conceal the white spot, we could either expect that BW dominant males change their status to submissive if this spot is related to aggressiveness (Schultz & Fincke, 2009; Xu & Fincke, 2015), or that they maintain the same dominance, if the spot is not related to dominance. In the case of submissive BW males, concealing their white spot would not affect their behaviour under the assumption that the white spot is not related to dominance.

METHODS

Morphometrics and asymmetry

Paraphlebia zoe was studied in Teocelo, Veracruz, Mexico (19°40'42" N, 96°98'28" W) in a creek depression in a runoff water habitat, from 11 June to 25 July, 2017. Fieldwork was carried out from 9:00 to 16:30, every day if the weather allowed it (*i.e.* if there were no major showers), by two observers who recorded date and time of capture, male morph and body size, measured to the nearest 0.1 mm using a caliper. Males and females were captured with an entomological net and marked with a small number on the left anterior wing, using a black *Staedtler* permanent lumocolor pen, a method that does not affect the survival of the damselflies (Munguía-Steyer, Córdoba-Aguilar & Romo-Beltrán, 2010). The number was situated far from the wing spots (see Figure 1), to minimize the possible effect of this mark on behaviour.

During our fieldwork period, HW males were extremely rare or absent, and were therefore not included in the study. For BW males, behavioural status was divided into two categories: territorial males, defined

as those individuals who stayed in the same place for more than two days and returned to the same perch site after participating in a fight in flight with a conspecific male, while individuals who did not meet this criterion were considered as non-territorial BW (satellite) males (Romo-Beltrán, Macías-Ordóñez & Córdoba-Aguilar, 2009). Within the BW territorial males, we found two subcategories: the dominant territorial males and the submissive males. The subcategories of the BW males were assigned observing them during 3 days, using 10-min focal observations each day. The male which won more than 60% of the interactions in which he was involved, was labelled as the dominant, whereas the submissive male was the one that won less than 50% of the interactions (Figure 2). Moreover, the histogram of figure 2 shows that there is no overlap between dominant and submissive males, and therefore they can be unambiguously identified. When a focal individual was observed perching outside the area of territories, it was considered to have switched to submissive. However, for focal males not observed during a day, we maintained their originally assigned status because they could be feeding, or making a break in their reproductive activities.

Body size was measured as length from the head to the end of the abdomen (including cerci) to the nearest 0.1 mm with a caliper. After that, each BW male was photographed with the wings spread for asymmetry measurements. The software Image J (Schneider, Rasband & Eliceiri, 2012) was used to measure area of the black and white spots from BW males as proxies of spot size. To control for trait size, asymmetry was calculated as the absolute value of the difference between right and left sides, divided by the average value of both sides (Cuervo, 2000). Four estimates of asymmetry were obtained for each BW male (one for each black and white spot), and we calculated the total asymmetry by adding up the four partial estimates. To distinguish measurement error from actual asymmetry, each individual was measured twice and the variance due to replicate and wing size was estimated following Crawley (2007). A sample of ten males, marked and photographed as young individuals and recaptured after 15 to 37 days, were photographed again to study ontogenetic changes in spot size. If spot size is related to genetic quality, and is not condition-dependent, we expect spot size to remain invariable over time. This could not be compared among dominant and submissive roles because the number of young individuals that were recaptured was too low.

Estimation of survival rates

To estimate survival (Φ) and recapture (p) probabilities we analysed the recapture histories of marked animals using the software MARK 8.1 (White & Burnham, 1999). Sex, time and their interaction were included as factors in the models (Lebreton et al., 1992). First, we tested the fit of the full time-dependent Cormack-Jolly-Seber model by groups (model $\{\Phi(g*t) p(g*t)\}$, where g is sex and t is time) using program RELEASE. The results indicated highly significant lack of fit, due to heterogeneity in males (TEST2+TEST3, males $\chi^2_{87}=204.25$, $p<0.001$; females $\chi^2_{31}=8.31$, $p=1$). We therefore increased the number of groups to account for BW and HW males. This model with three groups was again not appropriate due to heterogeneity in the group of BW males (TEST2+TEST3, BW males $\chi^2_{75}=176.59$, $p<0.001$; HW males $\chi^2_{15}=2.81$, $p=1$; females $\chi^2_{31}=9.72$, $p=1$). Finally, we subdivided BW males into two behavioural categories (dominant and submissive), and found that the model fitted all the groups, except for dominant BW males, which still showed lack of fit (TEST2+TEST3, dominant BW males $\chi^2_{43}=70.85$, $p=0.005$). Nevertheless, the overall fit was acceptable (TEST2+TEST3, four groups $\chi^2_{146}=145.46$, $p=0.497$), and we started with this model, given that no other phenotypic variables could be used to subdivide the group of dominant BW males.

Then, we estimated the variance inflation factor (c -hat; needed to correct for overdispersion) using two procedures: by dividing the c -hat obtained from model $\{\Phi(g*t) p(g*t)\}$ by the mean c -hat of the bootstrap

simulations in MARK, and by dividing the mean deviance of the saturated model by the mean deviance of the bootstrap simulations. The first method yielded an estimation of $\hat{c}=1$, and the second $\hat{c}=1.007$. We used the second value (more conservative) to adjust parameter estimates and standard errors. In a first step, we ran models without individual covariates, to select the most supported model, as the one that minimizes the quasi-likelihood version of the Akaike's Information Criterion adjusted for small sample sizes (QAICc). In a second step, body length was included as an individual covariate.

Reproductive behaviour

During the first 10 days of fieldwork, we conducted focal observations to estimate the limits of the territories and the alternative tactics (dominant and submissive; see Results) that BW territorial males showed. In territorial damselflies, males tend to occupy first high quality areas, and therefore the amount of fighting for territories increases with the number of males (Switzer, 2002). In *P. zoe*, little is known about which areas offer the best resources, but males concentrate in the areas closer to the water, where most matings are observed. We assume that males choose the best territories first (those close to the water), and territory quality could be estimated as the average number of males observed in a territory over the three days of focal observations. Copulations were timed to the nearest second. We studied 16 territories, whose limits were defined by the flight path that the territorial male used for patrolling (following Romo-Beltrán, Macías-Ordóñez, & Córdoba-Aguilar, 2009). Two observers, from 9:30 to 16:00, carried out focal observations. The duration of each focal observation was 10 min per territory. In these observations, we registered the identity of males, BW alternative tactics and reproductive success (number of copulations and copulation duration).

Dominant and submissive roles: plastic or fixed?

We conducted a manipulative experiment to test whether wing spot size is related to the dominant/submissive roles. To this end, we manipulated black and white spots in four treatments: (i) painting over white and black spots without changing their size (control 1), (ii) unmanipulated males (control 2), (iii) erasing the white spot using black painting (EWS), (iv) increasing the black spot and moving the white spot maintaining its size (IBS) (Figure 1). We used a black Staedtler® permanent Lumocolor pen and a white Edding® 751 paint marker. Each male was followed for three days after manipulation to establish the territorial male subcategory (*i.e.* dominant or submissive) as described above (Romo-Beltrán, Macías-Ordóñez, & Córdoba-Aguilar, 2009). The response variable was change of status, coded as 0 if the male maintained the previous state (dominant or submissive) and as 1 when he changed status or was found far from the territory. This variable was analysed with a GLM with Bernoulli distribution and logit link, entering as explanatory variables the treatment and the initial status. We compared the change of status between day 0 and the following three days, in separate GLMs.

The number of males that copulated once, twice, and so on, followed a negative binomial distribution (Goodness of fit test, $\chi^2_2=0.004$, $p=0.998$). Therefore, we used a GLM with a negative binomial distribution and a log link function (with the dispersion parameter estimated from the residual deviance) to test the effect of longevity, body length, territory quality, spot size (mean of black and white spots of fore- and hindwings entered separately), asymmetry and male strategy (dominant or submissive) on the number of matings obtained by each BW male.

Air temperature was obtained from a datalogger situated in the shade, which took a measure every 5 min, to consider this variable in further analyses, as it is known to influence copula duration in odonates (Michiels, 1992). We found that copulation duration did not follow a normal distribution, but was

normalized using a Box-Cox transformation with $\lambda = -0.6$. Using a GLM with normal errors, we tested the effect of time of start, temperature, asymmetry, body length, territory quality, age and status (dominant/submissive) on copulation duration.

Statistical tests were carried out using XLSTAT 2018 software (www.xlstat.com) and Genstat 18.1 software (GenStat, 2015). Mean values are presented with their standard error and sample size.

RESULTS

Overall, we observed 931 interactions between BW males. On average BW males considered dominant won 11.05 ± 0.77 disputes (mean \pm SE) and lost 1.17 ± 0.15 , whereas BW males assigned as submissive won 0.81 ± 0.14 and lost 9.7 ± 0.67 interactions. Figure 2 shows that there were clearly two groups of males (*i.e.* submissives and dominants) without any intermediate category.

Morphometrics and asymmetry

Over the period of study, we marked 212 males and 111 females. Table 1 shows the number of BW and HW males, females and their recapture rates. The average size (area) of black spots varied between 0.44 and 1.14 cm², and white spots between 0.08 and 0.40 cm² (Table 2). The analysis of the size of the spots in young males that were recaptured after 15-37 days, indicates that spot area diminished with age (Figure S1). The slope of the regression between initial spot size and final spot size was compared to 1, the expectation if the spots do not change over ontogeny. With the exception of the white spot of the fore wings, the slope was significantly lower than 1 (Figure S1), indicating that their size diminished with age.

The mean asymmetry was 0.341 ± 0.015 (N=168). There were no significant differences between dominant and submissive males in asymmetry or size of the spots (Table 2). There was a significant negative relationship between body length and total asymmetry (Pearson $r = -0.278$, $p < 0.0001$; Figure 3).

Estimation of survival rates

Among BW males, dominant males were resighted a mean of six times in contrast to three times for the submissive males. HW males and females were resighted 1-2 times (Table 1). The analysis of daily survival and recapture probabilities using MARK indicated that the most supported model was $\{\Phi(g) p(g^*t)\}$, implying an effect of the group (three types of males and females) on survival and a variable recapture probability due to the effect of the group, the time and their interaction (Table S1). Introducing body length as an individual covariate improved the statistical support for the model (Table S1). We fitted models with common and different intercepts for the covariate, and also a model with a quadratic term of body length, to test for stabilizing selection. Results indicated that the model with a common intercept and without the quadratic term is the most supported (model $\{\Phi(g^*length) p(g^*t)\}$ -common intercept, table S1). The best three models, all included daily variation of recapture rates and the effect of the group, and their interaction. We estimated survival probabilities using the model averaging option of Mark (Table 1, and present the recapture rates of the first model with the term $p(g)$, to allow a comparison between groups.

The effect of the individual covariate, body length, was positive on survival rate in all types of individuals (Figure 4). Hence, larger individuals had increased survivorship, even among non-territorial HW males and females.

243 ***Reproductive behaviour***

244 BW males (dominant and submissive) were identified and their reproductive activity in terms of number
245 of matings, sperm translocation and copulation duration was registered.

246 Sperm translocation occurred in precopulatory tandem, and had a duration of 19.3 ± 0.4 (53), with a range
247 from 12 to 25 sec. Copulation duration oscillated between 3 and 16 minutes, with an average of 6.6 ± 0.5
248 min (N=49 matings, all BW males), and only occasionally copulation was interrupted by short breaks.
249 Copulation duration was not correlated with any of the variables measured using bi-variate correlations
250 (results not shown; Spearman r ; variables tested: asymmetry, body length, quality of territory (number of
251 individuals in the territory), temperature and age). Using a GLM with normal errors (see Methods), we
252 found that copulation duration was not different between dominant and submissive males, and the only
253 marginally significant variable was temperature, which had a negative effect (estimate = -0.039 ± 0.020 ,
254 $p = 0.066$; Table S2).

255 We analysed lifetime mating success of marked males. This analysis suggests that longevity had a positive
256 effect on mating success (estimate: 0.042 ± 0.011 , $t_{158} = 3.90$, $p < 0.001$; Figure 5). Territory quality also had
257 a positive effect (estimate: 0.232 ± 0.057 , $t_{158} = 4.07$, $p < 0.001$; Figure 6) and BW-submissive strategy had a
258 negative effect compared to BW-dominant strategy (estimate: -1.040 ± 0.225 , $t_{158} = -4.62$, $p < 0.001$). The
259 other variables were not significant (see Supplementary Table S3).

260 ***Dominant and submissive roles: plastic or fixed?***

261 The sample of 79 BW males, whose spots were manipulated (Figure 1) to test whether spot size (black
262 and white) had an effect on their mating strategy, were observed during three days after manipulation.
263 Comparing day 0 to day 3, the experimental treatments had no significant effects, suggesting that the
264 strategy was not affected by spot size manipulation (Table 3). However, in two comparisons (day 0 to day
265 1, and day 0 to day 2) the initial status had a negative significant effect, i.e. animals with dominant status
266 were more likely to become submissive over time than the opposite (Table 3).

267

268 **DISCUSSION**

269 Our results indicate that BW males can behave as dominant or submissive. We found evidence for a
270 positive effect of territory quality (measured as the number of males occupying the area over time) and
271 lifespan on mating success, and that male morph strategy and sex affected the probability of recapture,
272 and larger individuals had higher survival. However, black wing spots asymmetry was not correlated with
273 the reproductive strategy of BW males, and the experimental manipulation of the black and white wing
274 spots did not affect male dominance. We will discuss these findings in relation to what is known in this
275 species and other territorial animals.

276 BW dominant males were resighted more often than BW submissive males, HW males and females. This
277 is easily explained since BW dominant males are more prone to stay on a territory for long periods,
278 because they are expected to be the best males, the only ones able to defend the territories close to the
279 water (see also Bañuelos Irusta & Araújo, 2007). In many animals, territorial males are usually larger, and
280 defend their territory from other males continuously (e.g. Baker, 1983), a situation commonly found in
281 other territorial damselflies (reviewed by Córdoba-Aguilar & Cordero-Rivera, 2005 and Suhonen, Rantala,

& Honkavaara, 2008) and other taxa (e.g. Mathis, 1991). Territory holders behave aggressively and this is expected to be costly. In fact, experimentally increasing the level of aggressiveness with methoprene (an analog of the juvenile hormone) impaired fat reserves and reduced survival in *Calopteryx* damselflies (Contreras-Garduño et al., 2011) and reduced survival in *Argia* (Córdoba-Aguilar & Munguía-Steyer, 2015). Being large and being territorial can be therefore costly, and the expectation is that only males of high quality are able to defend their territory for long periods of time (Tsubaki & Ono, 1987). In our case, larger individuals had higher survival rates not only in the two BW male morph strategies, but also in HW males and females (Figure 3). This contrasts with previous results on the same population, that only found a positive effect of size on survival in females but not in males (Munguía-Steyer, Córdoba-Aguilar & Romo-Beltrán, 2010). The general trade-off between territory defence and survival, under the assumption that territory holding is costly (Blanckenhorn, 2000), did not hold in our experiment, suggesting again that BW dominant males were of high quality. In other damselfly species, larger males presented higher survivorship (e.g. *Heteragrion cooki*: Rivas-Torres, et.al, 2017; *Polythore mutata*; Cordero-Rivera et al, unpublished), and similar results were also found in other taxa, such as the mosquito *Aedes aegypti* (Maciel-De Freitas, et.al, 2007). Large size is favoured in adult odonates, but probably not in larvae, contributing to the stasis in body size observed over evolutionary time (Waller & Svensson, 2017).

We did not find differences in the asymmetry of the black wing spots between BW dominant and BW submissive males. Nonetheless, our results indicated that larger males (independently of their reproductive strategy) were more symmetric. Therefore, asymmetry *per se* does not seem to affect the ability of males to behave in a dominant way, but perhaps could affect indirectly *via* the effect of the body size. Furthermore, those males marked when young and recaptured as mature, indicated that spot size diminished with age, a result in agreement with previous findings in other wing-pigmented damselflies (*Mnais*: Sanmartín-Villar, 2017; *Polythore*: Sanmartín-Villar & Cordero-Rivera, 2016). Whether these changes are adaptive, as apparently occurs with changes in coloration with age in other taxa (Dreiss & Roulin, 2010; Lopez-Idiaquez et al., 2016) remains to be studied. It seems, therefore, that black wing spots in *P. zoe* are important for agonistic interactions, and that over their life, males lose part of their signals of quality. The expectation is that old males will be unable to hold territories when confronted by young freshly coloured males (reviewed by Suhonen et al., 2008).

Our observations on mating behaviour and copula duration are consistent with other studies on the genus *Paraphlebia*, with the exception that in *P. zoe* copulation was usually not interrupted, a fact commonly observed in *P. quinta*, where it was interpreted as evidence for sperm ejection by females (González-Soriano & Córdoba-Aguilar, 2003). We found that copulation is brief for odonates, with an average of 7 minutes, a value almost identical to previous studies of the same species (Wong-Muñoz et al., 2013), and of similar duration for dominant and submissive males. During our fieldwork period, HW males were very rare, but previous studies showed that HW males copulate for longer (Wong-Muñoz et al., 2013). Figure 5 shows that mating success was positively related to longevity, a result that is general in many odonates (Fincke, Waage & Koenig, 1997; Koenig, 2008) and other animals (e.g. Madsen & Shine, 1992). Given the unpredictable nature of weather, particularly in some regions, animals able to live for long periods are likely to be more successful (Cordero, 1995), and in our species this correlates with a large body size. The fact that previous studies have also found a positive effect of body size on male success in *P. zoe* (Bello-Bedoy et al., 2015) is an argument for the necessity to demonstrate these effects in more than one study. Contrasting results between studies could be due to variable selection regimes among years, which is a reason to replicate research (Kelly, 2006) and increase our confidence on the generality of results.

Another factor that correlates with mating success is territory quality (Figure 6). In our case, the territory quality was estimated as the number of males that occupied the area over time, although only one of

them was the dominant. Note that we assumed that males are able to identify the best territories (but see below). Moreover, the higher the number of BW submissive males that surrounded a dominant BW male, the more successful the dominant BW male was, suggesting that submissive BW males do not represent a real threat for the dominant BW male, or at least the presence of submissive BW males does not completely counterbalance whatever factors are affecting mating success of the BW dominant male. Submissive BW males probably behave more as sneakers, which would make this system particularly interesting as HW males may also behave as territorials, despite its more common satellite strategy, as found by Romo-Beltrán et al (2009). These authors found that some territories were consistently more attractive for females than others, which is similar to our findings (Figure 6). They suggested that males did not recognize the best territories because in their study male turnover did not correlate with the number of matings observed in each area (Romo-Beltrán, Macías-Ordóñez & Córdoba-Aguilar, 2009). Nevertheless, this could depend on the season: when density is low, some territories would be vacant, and males could in these cases perform better by occupying a vacant site than by trying to fight for an already occupied territory. Being able to find and defend the best territories, i.e., those that attract a higher number of females, seems to be the key for success in *P. zoe* males, and territorial defence is apparently related to fat reserves (Romo-Beltrán, Macías-Ordóñez & Córdoba-Aguilar, 2009).

Contrarily to our predictions, we found that manipulating black and white wing spot size did not change behavioural status in BW males; submissive males remained submissive, even after an enlargement of their black spot. In agreement with this, we did not find differences in the black and white wing spot size between dominant and submissive males (Table 2), and the size of the black and white wing spots did not correlate with mating success. In birds, increasing the size of the spots, changed the aggressive behaviour of manipulated males (Senar, 2006). Our reasoning was that wing spot size would be relevant in intra-sexual selection (Romo-Beltrán, Macías-Ordóñez & Córdoba-Aguilar, 2009), but apparently the effect, if any, was negligible. The only change detected was a tendency for dominant males to lose their status over time (Table 3), but without relation to the experiment. It is possible that in species whose males do not court, as *P. zoe*, females have to rely on cues that are not related with the attractiveness of the males (like wing spots) and follow rules related with the resources offered by the territories, which in this species are basically the best sites to oviposit.

CONCLUSIONS

Our findings add further arguments to the recognition of ethodiversity as a basic element of biodiversity (Cordero-Rivera, 2017). We have found that *Paraphlebia zoe*, a species with male dimorphism (BW and HW morphs) associated with alternative reproductive tactics, holds high behavioural diversity, because at least three types of strategies, dominant, submissive and satellite, are concealed in males. This intra-morph behavioural diversity might be related to male phenotype and environmental factors, such as population density or different proportions of BW:HW males in different periods of the reproductive season (Wong-Muñoz et al., 2013), which may affect male reproductive behaviour in damselflies (Cordero-Rivera & Andrés, 2002). Performing similar manipulative experiments with the HW males when they are in sufficient density would be of great relevance to understand the relationship between male morph and dominance. Furthermore, future studies should take into account the UV wavelength of the black and white bands, which in other species have been found to relate to territoriality (Xu & Fincke, 2015).

ACKNOWLEDGMENTS

We are grateful to Pedro Démanos, Janet Nolasco, Daniel Tokman, Fernanda Baena and Ramsés Chávez for field assistance, and to Rogelio Macías Ordóñez for his suggestions and advice during all the study period. Thanks to M. Olalla Lorenzo-Carballea for the help on manuscript editing, and to two anonymous referees, who greatly improved the readability of our text.

REFERENCES

- Baker RR. 1983. Insect territoriality. *Annual Review of Entomology* 28:65–89.
- Banks MJ., Thompson DJ. 1987. Lifetime reproductive success of females of the damselfly *Coenagrion puella*. *Journal of Animal Ecology* 56:815–832.
- Bañuelos Irusta J., Araújo A. 2007. Reproductive tactics of sexes and fitness in the dragonfly, *Diastatops obscura*. *Journal of Insect Science* 7:24.
- Beck ML., Pruett-Jones S. 2002. Fluctuating asymmetry, sexual selection, and survivorship in male Dark-Winged damselflies. *Ethology* 108:779–791.
- Bello-Bedoy R., González-Santoyo I., Serrano-Meneses MA., Vrech D., Rivas M., Munguía-Steyer R., Córdoba-Aguilar A. 2015. Is allometry of sexual traits adaptive? A field test with territorial damselflies. *Biological Journal of the Linnean Society* 114:327–334.
- Blanckenhorn WU. 2000. The evolution of body size: what keeps organisms small? *Quarterly Review of Biology* 75:385–407.
- Carchini G., Chiarotti F., Di Domenico M., Mattoccia M., Paganotti G. 2001. Fluctuating asymmetry, mating success, body size and heterozygosity in *Coenagrion scitulum* (Rambur) (Odonata: Coenagrionidae). *Animal Behaviour* 61:661–669. DOI: 10.1006/anbe.2000.1621.
- Contreras-Garduño J., Córdoba-Aguilar A., Azpilicueta Amorín M., Cordero-Rivera A. 2011. Juvenile hormone favors sexually-selected traits but impairs fat reserves and abdomen mass in males and females. *Evolutionary Ecology* 25:845–856.
- Cordero-Rivera A. 2017. Behavioral diversity (ethodiversity): a neglected level in the study of biodiversity. *Frontiers in Ecology and Evolution* 5:1–7. DOI: 10.3389/fevo.2017.00007.
- Cordero-Rivera A., Andrés JA. 2002. Male coercion and convenience polyandry in a Calopterygid damselfly (Odonata). *Journal of Insect Science* 2:14–Available online: insectscience.org/2.14. DOI: 10.1673/031.002.1401.
- Cordero A. 1990. The adaptive significance of the prolonged copulations of the damselfly, *Ischnura graellsii* (Odonata: Coenagrionidae). *Animal Behaviour* 40:43–48.
- Cordero A. 1995. Correlates of male mating success in two natural populations of the damselfly *Ischnura graellsii* (Odonata: Coenagrionidae). *Ecological Entomology* 20:213–222.
- Córdoba-Aguilar A. 2002. Wing pigmentation in territorial male damselflies, *Calopteryx haemorrhoidalis*: a possible relation to sexual selection. *Animal Behaviour* 63:759–766. DOI: 10.1006/anbe.2001.1974.
- Córdoba-Aguilar A., Cordero-Rivera A. 2005. Evolution and ecology of Calopterygidae (Zygoptera: Odonata): status of knowledge and research perspectives. *Neotropical Entomology* 34:861–879. DOI: 10.1590/S1519-566X2005000600001.
- Córdoba-Aguilar A., Munguía-Steyer R. 2015. To be or not to be? Mating success and survival trade-offs when switching between alternative reproductive tactics. *Journal of Evolutionary Biology* 28:2119–

2124. DOI: 10.1111/jeb.12721.
- Crawley MJ. 2007. *The R Book*. Chichester: John Wiley & Sons. DOI: 10.1002/9780470515075.
- Cuervo JJ. 2000. El estudio de la asimetría fluctuante. *Etología* 18:27–60.
- Dreiss AN., Roulin A. 2010. Age-related change in melanin-based coloration of Barn owls (*Tyto alba*): Females that become more female-like and males that become more male-like perform better. *Biological Journal of the Linnean Society* 101:689–704. DOI: 10.1111/j.1095-8312.2010.01503.x.
- Emlen DJ. 1997. Alternative reproductive tactics and male-dimorphism in the horned beetle *Onthophagus acuminatus* (Coleoptera: Scarabaeidae). *Behavioral Ecology and Sociobiology* 41:335–341. DOI: 10.1007/s002650050393.
- Fincke OM., Jödicke R., Dennis R., Schultz TD., Paulson DR. 2005. The evolution and frequency of female color morphs in Holarctic Odonata: why are male-like females typically the minority? *International Journal of Odonatology* 8:183–202. DOI: 10.1080/13887890.2005.9748252.
- Fincke OM., Waage JK., Koenig WD. 1997. Natural and sexual selection components of odonate mating patterns. In: Choe JC, Crespi BJ eds. *The evolution of mating systems in insects and arachnids*. Cambridge: Cambridge University Press, 58–74.
- Forbes MRL., Leung B., Schalk G. 1997. Fluctuating asymmetry in *Coenagrion resolutum* (Hagen) in relation to age and male pairing success (Zygoptera: Coenagrionidae). *Odonatologica* 26:9–16.
- Garrison RW., von Ellenrieder N., Louton JA. 2010. *Damselfly genera of the New World*. Baltimore: Johns Hopkins University Press.
- GenStat. 2015. *GenStat for Windows*. Oxford: VSN International Ltd.
- González-Soriano E., Córdoba-Aguilar A. 2003. Sexual behaviour in *Paraphlebia quinta* calvert: Male diamorphism and a possible example of female control (Zygoptera: Megapodagrionidae). *Odonatologica* 32:345–353.
- González Soriano E., Novelo Gutiérrez R., Verdugo Garza M. 1982. Reproductive behavior of *Palaemnema desiderata* Selys (Odonata: Platystictidae). *Adv. Odonatol.* 1:55–62.
- Van Gossum H., Sherratt TN., Cordero-Rivera A. 2008. The evolution of sex-limited colour polymorphisms. In: Córdoba-Aguilar A ed. *Dragonflies: model organisms for ecological and evolutionary studies*. Oxford: Oxford University Press, 219–229.
- Gross MR. 1996. Alternative reproductive strategies and tactics: diversity within sexes. *Trends in Ecology and Evolution* 11:92–98.
- Harvey IF., Walsh KJ. 1993. Fluctuating asymmetry and lifetime mating success are correlated in males of the damselfly *Coenagrion puella* (Odonata: Coenagrionidae). *Ecological Entomology* 18:198–202.
- Kelly CD. 2006. Replicating empirical research in Behavioral Ecology: how and why it should be done but rarely ever is. *The Quarterly Review of Biology* 81:221–236. DOI: 10.1086/506236.
- Koenig WD. 2008. Lifetime reproductive success and sexual selection theory. In: Córdoba-Aguilar A ed. *Dragonflies and damselflies. Model organisms for ecological and evolutionary research*. Oxford: Oxford University Press, 153–166.
- Koga T., Backwell-Patricia RY., Jennions MD., Christy JH. 1998. Elevated predation risk changes mating behaviour and courtship in a fiddler crab. *Proceedings of the Royal Society B: Biological Sciences* 265:1385–1390.
- Lebreton JD., Burnham KP., Clobert J., Anderson DR. 1992. Modeling survival and testing biological

- hypotheses using marked animals: a unified approach with case studies. *Ecological Monographs* 62:67–118.
- Leung B., Forbes MR. 1997a. Modelling fluctuating asymmetry in relation to stress and fitness. *Oikos* 78:397–405.
- Leung B., Forbes MR. 1997b. Fluctuating asymmetry in relation to indices of quality and fitness in the damselfly, *Enallagma ebrium* (Hagen). *Oecologia* 110:472–477.
- Lopez-Idiaquez D., Vergara P., Fargallo JA., Martinez-Padilla J. 2016. Old males reduce melanin-pigmented traits and increase reproductive outcome under worse environmental conditions in common kestrels. *Ecology and Evolution* 6:1224–1235. DOI: 10.1002/ece3.1910.
- Madsen T., Shine R. 1992. Determinants of reproductive success in female adders, *Vipera berus*. *Oecologia* 92:40–47.
- Mathis A. 1991. Large male advantage for access to females: evidence of male-male competition and female discrimination in a territorial salamander. *Behavioral Ecology and Sociobiology* 29:133–138. DOI: 10.1007/BF00166488.
- Michiels NK. 1992. Consequences and adaptive significance of variation in copulation duration in the dragonfly *Sympetrum danae*. *Behavioral Ecology and Sociobiology* 29:429–435.
- Moore AJ. 1987. The behavioral ecology of *Libellula luctuosa* (Burmeister) (Anisoptera: Libellulidae): I. Temporal changes in the population density and the effects of male territorial behaviour. *Ethology* 75:246–254.
- Munguía-Steyer R., Córdoba-Aguilar A., Romo-Beltrán A. 2010. Do individuals in better condition survive for longer? Field survival estimates according to male alternative reproductive tactics and sex. *Journal of Evolutionary Biology* 23:175–184.
- Nomakuchi S. 1992. Male reproductive polymorphism and form-specific habitat utilization of the damselfly *Mnais pruinosa* (Zygoptera: Calopterygidae). *Ecological Research* 7:87–96.
- Norman MD., Finn J., Tregenza T. 1999. Female impersonation as an alternative reproductive strategy in giant cuttlefish. *Proceedings of the Royal Society B: Biological Sciences* 266:1347–1349.
- Painting CJ., Probert AF., Townsend DJ., Holwell GI. 2015. Multiple exaggerated weapon morphs: A novel form of male polymorphism in harvestmen. *Scientific Reports* 5:16368. DOI: 10.1038/srep16368.
- Paulson DR. 2004. Critical species of Odonata in the Neotropics. *International Journal of Odonatology* 7:163–188.
- Plaistow SJ., Tsubaki Y. 2000. A selective trade-off for territoriality and non-territoriality in the polymorphic damsel y *Mnais costalis*.
- Romo-Beltrán A., Macías-Ordóñez R., Córdoba-Aguilar A. 2009. Male dimorphism, territoriality and mating success in the tropical damselfly, *Paraphlebia zoe Selys* (Odonata: Megapodagrionidae). *Evolutionary Ecology* 23:699–709. DOI: 10.1007/s10682-008-9265-1.
- Samways MJ. 2006. Open and banded wings: Hypotheses on damselfly wing position (Zygoptera). *Odonatologica* 35:67–73.
- Sánchez-Guillén RA., Wellenreuther M., Chávez-Ríos JR., Beatty CD., Rivas-Torres A., Velasquez-Velez M., Cordero-Rivera A. 2017. Alternative reproductive strategies and the maintenance of female color polymorphism in damselflies. *Ecology and Evolution* 7:5592–5602. DOI: 10.1002/ece3.3083.
- Sanmartín-Villar I. 2017. Eco-etología de cabaliños do demo con polimorfismo de cor. University of Vigo.

491 Sanmartín-Villar I., Cordero-Rivera A. 2016. Female colour polymorphism and unique reproductive
492 behaviour in Polythore damselflies (Zygoptera: Polythoridae). *Neotropical Entomology* 45:658–664.
493 DOI: 10.1007/s13744-016-0417-7.

494 Schneider CA., Rasband WS., Eliceiri KW. 2012. NIH Image to ImageJ: 25 years of image analysis. *Nature*
495 *Methods* 9:671–675. DOI: 10.1038/nmeth.2089.

496 Schultz TD., Fincke OM. 2009. Structural colours create a flashing cue for sexual recognition and male
497 quality in a Neotropical giant damselfly. *Functional Ecology* 23:724–732. DOI:
498 10.1109/ICICI.2017.8365278.

499 Senar JC. 2006. Color displays as intrasexual signals of aggression and dominance. *Bird coloration* 2:87–
500 136.

501 Senar JC., Camerino M. 1998. Status signalling and the ability to recognize dominants: An experiment
502 with siskins (*Carduelis spinus*). *Proceedings of the Royal Society B: Biological Sciences* 265:1515–
503 1520. DOI: 10.1098/rspb.1998.0466.

504 Siva-Jothy MT., Tsubaki Y. 1989. Variation in copulation duration in *Mnais pruinosa pruinosa* Selys
505 (Odonata: Calopterygidae) 2. Causal factors. *Behavioral Ecology and Sociobiology* 25:261–267.

506 Suhonen J., Rantala MJ., Honkavaara J. 2008. Territoriality in odonates. In: Córdoba-Aguilar A ed.
507 *Dragonflies and damselflies. Model organisms for ecological and evolutionary research*. Oxford:
508 Oxford University Press, 203–218.

509 Switzer P V. 2002. Territory quality, habitat selection, and competition in the amberwing dragonfly,
510 *Perithemis tenera* (Say) (Odonata : Libellulidae): Population patterns as a consequence of
511 individual behavior. *Journal of the Kansas Entomological Society* 75:145–157.

512 Tsubaki Y., Ono T. 1987. Effects of age and body size on the male territorial system of the dragonfly,
513 *Nannophya pygmaea* Rambur (Odonata: Libellulidae). *Animal Behaviour* 35:518–525.

514 Waller JT., Svensson EI. 2017. Body size evolution in an old insect order: no evidence for Cope’s Rule in
515 spite of fitness benefits of large size. *Evolution* 71:2178–2193. DOI: 10.1111/evo.13302.

516 White GC., Burnham KP. 1999. Program MARK: Survival estimation from populations of marked animals.
517 *Bird Study* 46 Supplem:120–138.

518 Winandy L., Denoël M. 2015. The aggressive personality of an introduced fish affects foraging behavior
519 in a polymorphic newt. *Behavioral Ecology* 26:1528–1536. DOI: 10.1093/beheco/arv101.

520 Wong-Muñoz J., Anderson CN., Munguía-Steyer R., Córdoba-Aguilar A. 2013. Body size and morph as
521 drivers of copulation duration in a male dimorphic damselfly. *Ethology* 119:407–416.

522 Xu M., Fincke OM. 2015. Ultraviolet wing signal affects territorial contest outcome in a sexually
523 dimorphic damselfly. *Animal Behaviour* 101:67–74. DOI: 10.1016/j.anbehav.2014.12.018.

524

Table 1(on next page)

The number of individuals marked and recaptured in each category, with their body length.

The number of individuals marked and recaptured in each category, with their body length. Survival rate was estimated by model averaging over the three most supported models (Table S1). Recapture rate was different for each day and morph. Here we present the estimates obtained from model $\{\Phi(g, \text{length}) p(g)\}$ (where g is group and t refers to time; see Table S1), to allow a comparison between groups. BW= black-winged male; HW= hyaline winged male. Mean \pm SE

1

2 The number of individuals marked and recaptured in each category, with their body length. Survival rate
3 was estimated by model averaging over the three most supported models (Table S1). Recapture rate was
4 different for each day and morph. Here we present the estimates obtained from model $\{\Phi(g, \text{length})$
5 $p(g)\}$ (where g is group and t refers to time; see Table S1), to allow a comparison between groups. BW=
6 black-winged male; HW= hyaline winged male. Mean \pm SE

7

Variable	BW dominant	BW submissive	HW	Females
Number marked	54	133	25	111
Recaptured (%)	44 (81.5)	61 (45.9)	7 (28.0)	22 (19.8)
Number of recaptures	6.3 \pm 0.7	2.8 \pm 0.3	1.9 \pm 0.4	1.3 \pm 0.1
Body length (cm)	48.9 \pm 0.2	48.1 \pm 0.2	43.7 \pm 0.3	41.8 \pm 0.2
Daily survival rate	0.926 \pm 0.019	0.825 \pm 0.022	0.917 \pm 0.029	0.888 \pm 0.038
Daily recapture rate	0.558 \pm 0.024	0.408 \pm 0.024	0.197 \pm 0.046	0.062 \pm 0.015

8

Table 2 (on next page)

Descriptive statistics of asymmetry and spot size (cm²) for BW males of *Paraphlebia zoe*, in function of their mating strategy (D= dominant, S= submissive).

Descriptive statistics of asymmetry and spot size (cm²) for BW males of *Paraphlebia zoe*, in function of their mating strategy (D= dominant, S= submissive). P-value refers to the comparison between dominant and submissive males using an ANOVA.

1 Descriptive statistics of asymmetry and spot size (cm²) for BW males of *Paraphlebia zoe*, in function of
 2 their mating strategy (D= dominant, S= submissive). P-value refers to the comparison between dominant
 3 and submissive males using an ANOVA.

4

Variable	Sample size	Minimum	Maximum	Mean	Std. deviation	p-value
Asymmetry D	55	0.035	0.722	0.316	0.159	0.257
Asymmetry S	113	0.066	1.066	0.353	0.211	
Fore-black right D	55	0.556	0.901	0.730	0.077	0.974
Fore-black right S	113	0.525	1.013	0.730	0.086	
Fore-black left D	55	0.547	0.909	0.741	0.087	0.859
Fore-black left S	113	0.472	1.144	0.744	0.098	
Back-black right D	55	0.509	0.905	0.685	0.087	0.664
Back-black right S	113	0.452	0.926	0.679	0.088	
Back-black left D	55	0.497	0.878	0.693	0.089	0.562
Back-black left S	113	0.436	0.900	0.684	0.090	
Fore-white right D	55	0.136	0.309	0.215	0.041	0.651
Fore-white right S	113	0.135	0.395	0.212	0.042	
Fore-white left D	55	0.132	0.298	0.216	0.040	0.957
Fore-white left S	113	0.097	0.357	0.216	0.044	
Back-white right D	55	0.086	0.289	0.157	0.037	0.245
Back-white right S	113	0.083	0.266	0.151	0.033	
Back-white left D	55	0.088	0.227	0.153	0.035	0.851
Back-white left S	113	0.082	0.270	0.152	0.035	

5

Table 3 (on next page)

Results of the experiment of manipulation of black and white spot size of BW males of *Paraphlebia zoe*.

Results of the experiment of manipulation of black and white spot size of BW males of *Paraphlebia zoe*. Values are parameter estimates (\pm SE) after a GLM with Bernoulli distribution and logit link, where the response variate is “change of status”, coded as 1 if changed or 0 if not. Treatments include (i) painting over white and black spots without changing their size (control 1), (ii) unmanipulated control males (control 2), (iii) erasing the white spot using black painting (EWS), (iv) increasing the black spot (IBS) and moving the white spot maintaining its size (reference level for the comparisons). Strategy refers to the initial reproductive strategy of the males, dominant (D) or submissive (S). Spot size manipulation significantly affected only whether males were dominant or submissive, and only for Day 0 versus Day 1 and Day 0 versus Day 2, as indicated by values of $p < 0.05$ in **bold**.

1 Results of the experiment of manipulation of black and white spot size of BW males of *Paraphlebia zoe*. Values are parameter estimates (\pm SE) after
2 a GLM with Bernoulli distribution and logit link, where the response variate is “change of status”, coded as 1 if changed or 0 if not. Treatments
3 include (i) painting over white and black spots without changing their size (control 1), (ii) unmanipulated control males (control 2), (iii) erasing the
4 white spot using black painting (EWS), (iv) increasing the black spot (IBS) and moving the white spot maintaining its size (reference level for the
5 comparisons). Strategy refers to the initial reproductive strategy of the males, dominant (D) or submissive (S). Spot size manipulation significantly
6 affected only whether males were dominant or submissive, and only for Day 0 versus Day 1 and Day 0 versus Day 2, as indicated by values of
7 $p < 0.05$ in **bold**.

8

Factor	Comparison between days					
	0 to 1	0 to 2	0 to 3	1 to 2	1 to 3	2 to 3
control 1 (compared to IBS)	-1.07 \pm 1.00	-0.67 \pm 1.00	-0.51 \pm 0.68	-0.10 \pm 1.45	-0.13 \pm 0.70	-0.06 \pm 0.79
control 2 (compared to IBS)	-0.99 \pm 1.00	0.39 \pm 0.88	-0.63 \pm 0.71	1.59 \pm 1.17	0.00 \pm 0.71	-1.57 \pm 1.17
EWS (compared to IBS)	-0.39 \pm 0.89	0.71 \pm 0.86	-0.93 \pm 0.74	0.76 \pm 1.27	-0.92 \pm 0.81	-1.57 \pm 1.17
Strategy (S compared to D)	-1.80\pm0.83^a	-1.69\pm0.71^b	-0.89 \pm 0.52	-0.68 \pm 0.79	-0.71 \pm 0.53	0.02 \pm 0.70

9 ^a $p = 0.030$, ^b $p = 0.01$

Figure 1(on next page)

The four manipulative experiments.

(A) Control 1; painting over white and black spots. (B) Control 2; unmanipulated males. (C) IBS; increasing the black spot and moving the white spot maintaining its size. (D) EWS; erasing the white spot using black painting.







Figure 2 (on next page)

Histogram of the proportion of interactions won in BW males of *P. zoe*.

We considered dominant males, these that won more than 60% of the interactions, and submissive males these that won less than 50% of the interactions. In black, dominant BW males. In grey, submissive BW males.




Figure 3(on next page)

The relationship between body length and total asymmetry for BW males of *P. zoe*.

The line represents the regression equation and the ellipse the 95% confidence interval (Pearson $r = -0.167$, $p = 0.283$).




Figure 4(on next page)

The relationship between body length and survival rate for territorial (BW), non-territorial (HW) and females of *Paraphlebia zoe*.

The black line represents the estimate and the grey lines the confidence intervals. In all cases, larger individuals survived better.




Figure 5(on next page)

The relationship between longevity and reproductive success (number of matings) for BW males of *P. zoe*.

The line represents the regression equation and the ellipse the 95% confidence interval (Pearson $r = 0.39$, $p < 0.001$).




Figure 6(on next page)

The relationship between the quality of territory (number of individuals in the territory) and reproductive success (number of matings) for BW males of *P. zoe*.

The line represents the regression equation and the ellipse the 95% confidence interval (Pearson $r = 0.29$, $p < 0.001$).

