

# Validation of reference genes for gene expression studies in tartary buckwheat (*Fagopyrum tataricum* Gaertn.) using quantitative real-time PCR

Chenglei Li<sup>1</sup>, Haixia Zhao<sup>1</sup>, Maofei Li<sup>1</sup>, Panfeng Yao<sup>1</sup>, Qingqing Li<sup>1</sup>, Xuerong Zhao<sup>1</sup>, Anhu Wang<sup>2</sup>, Hui Chen<sup>1</sup>, Zizhong Tang<sup>1</sup>, Tongliang Bu<sup>1</sup>, Qi Wu<sup>Corresp. 1</sup>

<sup>1</sup> College of life science, Sichuan Agricultural University, Ya'an, Sichuan, China

<sup>2</sup> Xichang College, Xichang, Sichuan, China

Corresponding Author: Qi Wu  
Email address: wuqi@sicau.edu.cn

Quantitative real-time reverse transcriptase polymerase chain reaction (qRT-PCR) is a sensitive technique for quantifying gene expression levels. By implementing three distinct algorithms (geNorm, normFinder and BestKeeper), we have validated the stability of the expression of seven candidate reference genes in tartary buckwheat, including *FtSAND*, *FtCACS*, *FtExpressed1*, *FtGAPDH*, *FtActin*, *FtEF-1a* and *FtH3*. In this study, the results indicated that *FtCACS* and *FtSAND* were the best reference genes for 'abiotic cotyledons', and *FtExpressed1* and *FtEF-1a* were the best reference genes for aluminium treatment; *FtCACS* and *FtExpressed1* performed the best for the immature seed stage; *FtCACS* was best for the abiotic treatment, and *FtH3* appeared to be the most suitable reference gene for the abiotic treatment in hypocotyls and all samples in this study. In contrast, *FtActin* and *FtGAPDH* are unsuitable genes. Our findings offer additional stable reference genes for gene expression research on tartary buckwheat at the immature seed stage and under abiotic treatment.

# Validation of reference genes for gene expression studies in tartary buckwheat (*Fagopyrum tataricum* Gaertn.) using quantitative real-time PCR

Chenglei Li<sup>1</sup>, Haixia Zhao<sup>1</sup>, Maofei Li<sup>1</sup>, Panfeng Yao<sup>1</sup>, Qingqing Li<sup>1</sup>, Xuerong Zhao<sup>1</sup>, Anhu Wang<sup>2</sup>, Hui Chen<sup>1</sup>, Zizhong Tang<sup>1</sup>, Tongliang Bu<sup>1</sup>, Qi Wu<sup>1\*</sup>

<sup>1</sup>College of life science, Sichuan Agricultural university, Ya'an 625014, Sichuan, China

<sup>2</sup>Xichang college, Xichang 615013, Sichuan, China

\*Corresponding Author: Qi Wu

Sichuan Agricultural university, Ya'an 625014, Sichuan, China

Email addresses: wuqi@sicau.edu.cn

## Abstract

Quantitative real-time reverse transcriptase polymerase chain reaction (qRT-PCR) is a sensitive technique for quantifying gene expression levels. By implementing three distinct algorithms (geNorm, normFinder and BestKeeper), we have validated the stability of the expression of seven candidate reference genes in tartary buckwheat, including *FtSAND*, *FtCACS*, *FtExpressed1*, *FtGAPDH*, *FtActin*, *FtEF-1a* and *FtH3*. In this study, the results indicated that *FtCACS* and *FtSAND* were the best reference genes for 'abiotic cotyledons', and *FtExpressed1* and *FtEF-1a* were the best reference genes for aluminium treatment; *FtCACS* and *FtExpressed1* performed the best for the immature seed stage; *FtCACS* was best for the abiotic treatment, and *FtH3* appeared to be the most suitable reference gene for the abiotic treatment in hypocotyls and all samples in this study. In contrast, *FtActin* and *FtGAPDH* are unsuitable genes. Our findings offer additional stable reference genes for gene expression research on tartary buckwheat at the immature seed stage and under abiotic treatment.

## 28 Introduction

29 Quantitative real-time reverse transcription polymerase chain reaction (qRT-PCR) has become  
 30 the most prevalent quantification method used in assays of gene expression on account of its  
 31 specificity, accuracy, efficiency and high sensitivity (Jiang et al. 2014). The data from qRT-PCR  
 32 can be analysed using absolute or relative quantification. Absolute quantification yields an exact  
 33 gene copy number by transforming quantification cycles (Hruz et al.) into a standard curve.  
 34 Relative quantification relies on internal control genes as references to present the qRT-PCR data  
 35 of the target genes (Ginzinger 2002). Relative quantification has become a more widely used  
 36 method in gene expression assays because most researchers mainly focus on the analysis of  
 37 differences in gene expression (Wu et al. 2016). To normalize qRT-PCR data, a reference gene is  
 38 needed, and it should be universally valid, with a similar expression level across all feasible  
 39 cells, tissue samples and experimental treatments (Huang et al. 2014). However, ideal reference  
 40 genes do not actually exist (Remans et al. 2014). The selection of the appropriate reference gene  
 41 is a critical step in controlling the variability of samples when using a sensitive qRT-PCR  
 42 technique (Tong et al. 2009). If the reference gene is appropriate, the discrepancies that may  
 43 exist in terms of the initial sample amount, RNA integrity, RNA recovery, and the efficiency of  
 44 cDNA synthesis will be eliminated. Statistical algorithms, such as geNorm (Vandesompele et al.  
 45 2002), normFinder (Andersen et al. 2004) and BestKeeper (Pfaffl et al. 2004), have been  
 46 developed to help to select appropriate reference genes.

47 Most studies about gene expression in bacteria (Wu et al. 2017), yeast (Cankorur-Cetinkaya et  
 48 al. 2012) and mammals (Terzi et al. 2010) now include reference gene validation (Chapman &  
 49 Waldenstrom 2015). However, the validation of reference genes in plants has received very little  
 50 attention, and housekeeping genes tend to be used as references without any appropriate  
 51 validation (Gutierrez et al. 2008). In 2005, Czechowski et al were the first to present a list of  
 52 stably expressed Arabidopsis genes under a large range of experimental conditions, and the  
 53 evidence clearly showed that several genes are expressed more stably than traditional reference  
 54 genes (Czechowski et al. 2005). Since then, suitable reference genes for gene expression studies  
 55 have been reported in higher plants such as flax (Huis et al. 2010), soybean (Hu et al. 2009),  
 56 tomato (Lovdal & Lillo 2009), *Pisum sativum* (Die et al. 2010), *Brachypodium distachyon* (Hong  
 57 et al. 2008) and carrot (Tian et al. 2015). However, reference gene stability is not consistent

across experimental conditions and plant species. Consequently, it is necessary to find additional reference genes for different conditions and species.

Tartary buckwheat (*Fagopyrum tataricum* Gaertn.) belongs to the Polygonaceae family (Kim et al. 2013), and as an important functional food material, it has a relatively high flavonoid content. Tartary buckwheat is capable of thriving in regions with poor soil or harsh climates (Kim et al. 2009). The results of genome sequencing also show that tartary buckwheat has a remarkable ability to cope with highly variable environmental stress, including drought, salinity, UV-B and cold (Zhang et al. 2017). The stress resistance of tartary buckwheat is mainly due to its abundant flavonoids (Suzuki et al. 2005). In particular, tartary buckwheat is a naturally aluminium (Al) tolerant species (Wang et al. 2015). Abiotic stresses and flavonoid metabolism regulate the expression of these genes in plants at both transcriptional and post-transcriptional levels. There are many reports on tartary buckwheat under various experimental conditions in which gene expression is normalized to a reference gene (*H3*) for semi-quantitative RT-PCR or qRT-PCR (Bai et al. 2014). However, to date, there is no systematic strategy to analyse tartary buckwheat reference genes at the immature seed stage or under abiotic stress.

In this work, we aimed to evaluate the potential use of different reference genes for internal normalization to more accurately measure the expression level of genes of interest in tartary buckwheat. Seven candidate reference genes were selected, and the stability of their expression was assessed in tartary buckwheat at the immature seed stage and under different abiotic stress treatments. Evaluating the stability of the expression of candidate reference genes depends on statistical analysis. Three different statistical software programs (geNorm, normFinder, and BestKeeper) were used to calculate the variability of the expression of the candidate genes and determine which were the most suitable.

## Materials & Methods

### Plant materials and treatments

Tartary buckwheat ("Xiqiao No. 2") seeds were grown in the field on a farm (XTBG; 29°59'N, 102°59'E; 800 m elevation) at Sichuan Agricultural University, Ya'an, Sichuan, China (Li et al. 2012). Tissues, including roots, stems, leaves, flowers, immature seed 1 (seed formation started) and immature seed 2 (seeds in the milk) were collected at the immature seed stage (Gupta et al. 2011). The 7-day-old seedlings were stressed with saline or drought by adding 100 mM NaCl or

20% PEG 2000, respectively, to the medium. The 7-day-old seedlings were exposed in a chamber at 4°C with a 16 h photoperiod for cold treatment (Gao et al. 2016). UV-B treatment was conducted under UV-B (302 nm, 0.1 mW/cm<sup>2</sup>) in a chamber. After 0, 2, 4, 6, 12 and 24 h of treatment, all stressed seedlings were collected and separated into cotyledons and hypocotyls. For the Al treatment, the samples were processed according to a previous report (Zhu et al. 2015). Root tips (0-2 cm) and basal roots (2-4 cm) were sampled under both -Al and +Al conditions. All samples were collected in two biological replicates, and RNA was extracted immediately.

#### **Total RNA isolation and cDNA synthesis**

Total RNA was isolated from various samples with an RNAout 2.0 kit (Tiandz, China) according to the manufacturer's instructions. To remove trace DNA from samples, total RNA extractions were treated with RNase free DNase I. The RNA integrity was detected using 2% agarose gels. A Bio-RAD smart spec<sup>TM</sup>plus spectrophotometer was used to determine the RNA purity. cDNA was synthesized with a PrimeScript<sup>TM</sup> RT reagent kit and gDNA Eraser (Perfect Real Time) (TaKaRa, Dalian, China).

#### **Selection of candidate reference genes and design of qRT-PCR primers**

Potential homologues of the seven candidate genes were identified from the transcriptome sequencing data of tartary buckwheat ('Xiqiao No. 2') (Yao et al. 2017). Primers were designed using Primer Premier 5.0. All primers used in this research are listed in Table 1. The specificity of the amplification was assessed based on the presence of a single band of the expected size in a 1.5% agarose gel following electrophoresis and a single peak in the qRT-PCR melting curve.

#### **Quantitative real-time PCR**

The qRT-PCR procedure was designed in accordance with MIQE guidelines (Bustin et al. 2009). qRT-PCR was executed in a CFX96 Real Time PCR system (Bio Rad, Singapore) with a SYBR Premix EX *Taq* kit (TaKaRa, Japan) in a total reaction volume of 15 µL that included 10 µL of SYBR Green mix, primers at 0.5 µM each and 1 µL of cDNA. The amplification conditions were as follows: 95°C for 30 s and 40 cycles of 95°C for 5 s and 60°C for 20 s. A melting curve from 60 to 95°C was used to verify the specificity of the PCR amplification. All genes were amplified from cDNA. Ten-fold serial dilutions of cDNA samples from young leaves were used to establish the standard curves to calculate the amplification efficiency of each primer pair.

## Statistical analysis

Three software programs (geNorm v3.5, the Excel add in of normFinder v0.953 and BestKeeper v1) were used to analysis the stability of reference gene expression across all experimental sets (Andersen et al. 2004; Pfaffl et al. 2004; Vandesompele et al. 2002). SPSS v17.0 was used to calculate the span of Cq values for each gene by drawing a box-whisker plot, and the expression levels of *FtSTAR* and *FtDFR* were showed by mean  $\pm$  standard deviation (SD).

## Results

### RNA solution and quality

A series of 58 samples from tartary buckwheat were divided into six different groups. ‘Abiotic cotyledons’ was composed of cotyledons from all stress-treated samples. ‘Abiotic hypocotyls’ included hypocotyls from all stress-treated samples. ‘Abiotic total’ was composed of ‘abiotic cotyledons’ and ‘abiotic hypocotyls’. ‘Al treatment’ comprised all samples that were treated with Al. ‘Immature seed stage’ consisted of six tissues (roots, stems, leaves, flowers, immature seed 1 and immature seed 2) at the immature seed stage based on the state of the seed. Finally, ‘total’ included all the samples in this study. Protein and organic pollutants were isolated and removed from all samples via RNA extraction. The total RNAs, with  $A_{260}/A_{280}$  ratios of 1.8-2.0, were reverse transcribed into cDNA as templates for qRT-PCR detection.

### Expression profiles of candidate reference genes

We selected the best reference genes (among the 7 candidate genes) from the seven candidate genes in six different groups (sequencing data was shown in Supplemental Fig. S1). The amplification efficiencies (E) of all reactions ranged from 95.5% to 107.5% and were calculated from standard curves with good linear relationships ( $R^2 > 0.99$ ). Single-peak melting curves were obtained for all qRT-PCR amplifications (Supplemental Fig. S2). A simple and commonly used method to identify stably expressed genes is to compare the span of quantification cycle (Hruz et al.) values for each gene in the qRT-PCR reactions. The results showed that the candidate reference genes spanned a wide range of Cq values, ranging from 14.98 (*FtGAPDH*) to 29.65 (*FtActin*), with the median Cq values of the genes ranging from 17.59 (*FtGADPH*) to 25.5 (*FtSAND*). The reference gene expression levels, in descending order, were *FtGAPDH*, *FtEF-1*, *FtH3*, *FtCACS*, *FtActin*, *FtExpressed1* and *FtSAND*. *FtActin* (median, 23.55; Cq variation,

11.38) showed the most variability, and the *FtH3* gene (median, 20.19; Cq variation, 3.67) showed the least variability (Fig. 1).

# **Software: geNorm, NormFinder and BestKeeper**

After a simple comparison of the raw Cq values, the three software programs were used to further analyse the stability of the expression of the seven candidate genes. The genes were ranked in descending order in terms of the stability of their expression in each of six groups (Table 2), and the details of the cotyledons and hypocotyls under abiotic stress are provided in Supplemental Table S1. GeNorm performed a stepwise exclusion of the most unstable gene and then recalculated M until only two genes remained, and these two genes had the most stable expression (Vandesompele et al. 2002). The gene with the lower M value was considered to have the most stable expression. An M value limit of <1.5 was suggested by geNorm. All the results from geNorm were lower than 1.5 in ‘abiotic cotyledons’ and ‘total’; *FtCACS* and *FtH3* (M = 0.33, 0.60) were chosen as the most stable genes, while *FtH3* and *FtSAND* (M = 0.65, 0.55) were identified as the most stable genes in the ‘abiotic hypocotyls’ and ‘abiotic total’ groups, respectively. In ‘immature seed stage’, *FtCACS* and *FtExpressed1* (M = 0.16) were recognized as the most stable genes. Finally, *FtEF-1α* and *FtExpressed1* (M = 0.16) were the most stable genes in ‘Al treatment’.

To determine the optimal number of reference genes required for accurate normalization, geNorm was used to calculate the pairwise variation ( $V_n/V_{n+1}$ ) between the sequential normalization factors (NFs) ( $NF_n$  and  $NF_{n+1}$ ). As suggested, a threshold value of 0.15 was adopted. As depicted in Fig. 2, pairwise variation analysis indicated that the ideal number of reference genes may be different for the different groups. For instance, only two genes are necessary for normalization for ‘immature seed stage’, ‘abiotic cotyledons’ and ‘Al treatment’, but the pairwise variation of the other three groups was above the threshold of 0.15. The pairwise variations for cotyledons and hypocotyls under different abiotic treatments are provided in Supplemental Fig. S3.

Unlike geNorm, normFinder (Andersen et al. 2004) depends on a variance estimation approach, which allows the comparison of inter/intra-group variation. Genes with the lowest average expression stability values are the most stable. In the normFinder analysis, *FtH3* and *FtCACS* were the most stable genes in ‘abiotic cotyledons’, and *FtSAND* and *FtExpressed1* were the most stable genes under ‘Al treatment’. In addition, the most stable reference genes in

‘immature seed stage’ were *FtSAND* and *FtGAPDH*, while the other three groups had the same two most stable reference genes (*FtSAND* and *FtCACS*). The least stable gene in all groups was *FtActin*.

BestKeeper (Pfaffl et al. 2004) determines the most stable genes by taking the coefficient of variance (CV) and SD of the Cq values. The more stably expressed genes are indicated by the lower SD and CV values. The results showed that the most stably expressed genes were *FtSAND* (CV  $\pm$  SD = 1.32  $\pm$  0.33) and *FtCACS* (CV  $\pm$  SD = 1.47  $\pm$  0.31) for ‘abiotic cotyledons’. In ‘abiotic hypocotyls’, *FtH3* (CV  $\pm$  SD = 2.47  $\pm$  0.51) and *FtSAND* (CV  $\pm$  SD = 2.68  $\pm$  0.69) were the most stable genes. *FtH3* (CV  $\pm$  SD = 2.55  $\pm$  0.52, CV  $\pm$  SD = 2.69  $\pm$  0.54) and *FtCACS* (CV  $\pm$  SD = 2.70  $\pm$  0.61, CV  $\pm$  SD = 2.33  $\pm$  0.53) were the most stable genes in the ‘abiotic total’ and ‘total’ groups, respectively. *FtExpressed1* (CV  $\pm$  SD = 0.73  $\pm$  0.17) and *FtEF-1 $\alpha$*  (CV  $\pm$  SD = 1.55  $\pm$  0.25) showed the most stable expression in ‘Al treatment’, and *FtExpressed1* (CV  $\pm$  SD = 2.00  $\pm$  0.50) and *FtCACS* (CV  $\pm$  SD = 2.43  $\pm$  0.57) showed the highest expression stabilities in ‘immature seed stage’. The least stable gene was *FtActin*, but the *FtGAPDH* gene, with a CV  $\pm$  SD of 7.59  $\pm$  1.39 in ‘abiotic hypocotyls’, was considered the least acceptable for gene expression normalization.

### Reference gene validation

To validate the availability of a reference gene, the expression levels of *FtSTAR* under Al treatment and of the *FtDFR* gene under UV treatment were determined using the seven candidate reference genes for normalization. For Al treatment, *STAR* has a conserved response in plants, and this has been shown in previous reports, such as reports on rice *STAR2* (Huang et al. 2009), *Arabidopsis ALS* (Larsen et al. 2005) and tartary buckwheat *FtSTAR2* (Zhu et al. 2015), whose expression increased after exposure to Al. The expression of *FtSTAR2* was greatly increased (Fig. 3), which was reinforced by the Al-induced expression of *STAR2*. The use of the most favourable reference gene (*FtExpressed1*) resulted in the greatest variation, resulting in increases of approximately 1.67-fold in root tips and 6.24-fold in basal roots, and the use of the other most stable gene (*FtEF-1 $\alpha$* ) resulted in increases of approximately 1.42-fold and 6.19-fold, respectively. Finally, the use of the least stable gene (*FtActin*) resulted in increases of approximately 1.90-fold in root tips and 2.48-fold in basal roots. These genes also showed a pattern of increased expression, regardless of whether stable or unstable reference genes were used for normalization. However, the relative expression using the most stable genes showed the



higher fold increases compared with the unstable genes. Therefore, *FtExpressed1* and *FtEF-1a* are the most suitable genes under AI treatment.

DFR catalyses the reduction of dihydroflavonols to leuco-anthocyanins and is a key enzyme in the biosynthesis of anthocyanins (Yuan et al. 2007). The contents of anthocyanin increased under UV treatment in tartary buckwheat hypocotyls (Eguchi & Sato 2009). In hypocotyls under UV treatment, the relative quantification of *FtDFR* using the most stable genes (*FtCACS*, *FtH3* and *FtActin*) for normalization exhibited similar expression patterns but different expression levels. When other less stable genes were used for normalization, the results obviously differed from those with the most stable genes (Fig. 4). The expression levels of the *FtDFR* gene under UV treatment showed similar trends after normalization to stable genes in hypocotyls. Thus, we selected the most stable reference genes for gene expression normalization in tartary buckwheat.

## Discussion

SYBR Green I is a fluorescent reporter dye used in qRT-PCR; when it binds double-stranded DNA, its fluorescence increases nearly 1000-fold (Morrison et al. 1998). The problems in qRT-PCR caused by the variability of RNA templates and inappropriate data normalization are obvious and widely known but disregarded (Nolan et al. 2006). To address these questions, proper reference genes should be used. Thus, validating candidate reference genes and selecting stable reference genes are becoming general interests for researchers.

Candidate reference genes can be selected from reference gene validation papers that use the same experimental conditions or closely related species (orthologies) or by mining transcriptomic data for stably expressed genes or traditional housekeeping genes (Hruz et al. 2011). The seven candidate reference genes tested in tartary buckwheat include three genes that are stably expressed in common buckwheat (*SAND*, *CACS* and *Expressed1*), three commonly used reference genes (*GAPDH*, *Actin* and *EF-1a*) and one unique gene (*H3*). Neither high (>30) nor low (<15) Cq values are recommended based on general guidelines (Wan et al. 2010). In this study, the Cq values of the reference genes ranged from 14.98 (*FtGAPDH*) to 29.65 (*FtActin*), suggesting variable expression. Although the raw Cq value comparison can provide a rough estimate of the stability of gene expression, it is not sufficient to accurately evaluate the expression patterns of reference genes.

Scientists have developed several approaches to select stable reference genes for normalization. However, to date, there is no consistent algorithm that can be used to test gene stability. These software programs have their own merits and drawbacks: NormFinder can avoid the misinterpretations caused by the artificial selection of co-regulated genes (Andersen et al. 2004); geNorm is based on the assumption that none of the analysed genes are co-regulated (Vandesompele et al. 2002); and geNorm can estimate the fewest number of reference genes needed for accurate normalization (Stamova et al. 2009). It seems possible that more reliable controls can be obtained using a combination of different algorithms (de Almeida et al. 2010). We used three software programs that were based on different statistical approaches to assess seven candidate genes in tartary buckwheat. The distinct statistical algorithms are likely to generate inconsistent rankings of stability. After integrating the outcomes of the three programs above in this study, we recommend the reference genes in tartary buckwheat as follows: *FtCACS* + *FtSAND* for ‘abiotic cotyledons’; *FtExpressed1* + *FtEF-1α* for ‘Al treatment’; *FtCACS* + *FtExpressed1* for ‘immature seed stage’; *FtCACS* for ‘abiotic total’; and *FtH3* for ‘abiotic hypocotyls’ and ‘total’. The *H3* gene is the most commonly used internal gene for normalization in tartary buckwheat (Luo et al. 2016). Our results indicate that *FtH3* is a stable gene under abiotic treatment. However, it is not stable across different organs or after Al treatment. In common buckwheat (Demidenko et al. 2011), *CACS* and *Expressed1* were validated as the most stable genes in the development and fruit stages. *FtCACS* and *FtExpressed1* being the most stable reference genes in the immature seed stage of tartary buckwheat supports the statement that the orthologues of identified reference genes could serve the same purpose in other species. In tomato, similar results were obtained in a study on reference gene selection (Exposito-Rodriguez et al. 2008).

*GAPDH* and *Actin* are the most generally used reference genes for the analysis of gene expression in various plant species (Kumar et al. 2011). However, our analysis indicates that *GAPDH* and *Actin* are not reliable genes for comparative expression analysis. The cause of these fluctuations at the gene expression level is probably that *GAPDH* and *Actin* have several biological functions, such as participating in the glycolytic pathway and other processes (Stürzenbaum & Kille 2001). *Actin* supports the cell and determines its shape, and it also takes part in other cellular functions (Kravets et al. 2017). Although *Actin* is a stable reference gene in developmental studies, it is not stable under various conditions.

# Conclusions

As far as we know, our studies take advantage of three software (geNorm, NormFinder and BestKeeper) to analysis the stability of seven candidate reference genes for the first time in tartary buckwheat using qRT-PCR. Three software identified sightly differet genes as most suited for normalization prompted us to merge the data. The result showed that the expression of *GAPDH* or *Actin* is unstable across all samples. We also provide a list with the stable reference genes in six group and under certain conditions.

# References

- Andersen CL, Jensen JL, and Orntoft TF. 2004. Normalization of real-time quantitative reverse transcription-PCR data: a model-based variance estimation approach to identify genes suited for normalization, applied to bladder and colon cancer data sets. *Cancer Res* 64:5245-5250. 10.1158/0008-5472.CAN-04-0496
- Bai YC, Li CL, Zhang JW, Li SJ, Luo XP, Yao HP, Chen H, Zhao HX, Park SU, and Wu Q. 2014. Characterization of two tartary buckwheat R2R3-MYB transcription factors and their regulation of proanthocyanidin biosynthesis. *Physiol Plant* 152:431-440. 10.1111/ppl.12199
- Bustin SA, Benes V, Garson JA, Hellemans J, Huggett J, Kubista M, Mueller R, Nolan T, Pfaffl MW, Shipley GL, Vandesompele J, and Wittwer CT. 2009. The MIQE guidelines: minimum information for publication of quantitative real-time PCR experiments. *Clin Chem* 55:611-622. 10.1373/clinchem.2008.112797
- Cankorur-Cetinkaya A, Dereli E, Eraslan S, Karabekmez E, Dikicioglu D, and Kirdar B. 2012. A novel strategy for selection and validation of reference genes in dynamic multidimensional experimental design in yeast. *PLoS One* 7:e38351. 10.1371/journal.pone.0038351
- Chapman JR, and Waldenstrom J. 2015. With Reference to Reference Genes: A Systematic Review of Endogenous Controls in Gene Expression Studies. *PLoS One* 10:e0141853. 10.1371/journal.pone.0141853
- Czechowski T, Stitt M, Altmann T, Udvardi MK, and Scheible WR. 2005. Genome-wide identification and testing of superior reference genes for transcript normalization in *Arabidopsis*. *Plant Physiol* 139:5-17. 10.1104/pp.105.063743
- de Almeida MR, Ruedell CM, Ricachenevsky FK, Sperotto RA, Pasquali G, and Fett-Neto AG. 2010. Reference gene selection for quantitative reverse transcription-polymerase chain reaction normalization during *in vitro* adventitious rooting in *Eucalyptus globulus* Labill. *BMC Mol Biol* 11:73. 10.1186/1471-2199-11-73
- Demidenko NV, Logacheva MD, and Penin AA. 2011. Selection and validation of reference genes for quantitative real-time PCR in buckwheat (*Fagopyrum esculentum*) based on transcriptome sequence data. *PLoS One* 6:e19434. 10.1371/journal.pone.0019434
- Die JV, Roman B, Nadal S, and Gonzalez-Verdejo CI. 2010. Evaluation of candidate reference genes for expression studies in *Pisum sativum* under different experimental conditions. *Planta* 232:145-153. 10.1007/s00425-010-1158-1
- Eguchi K, and Sato T. 2009. Differences in the Ratios of Cyanidin-3-O-glucoside and Cyanidin-3-O-rutinocide to Total Anthocyanin under UV and Non-UV Conditions in

- 318 Tartary Buckwheat (*Fagopyrum tataricum* Garten). *Plant Production Science*  
319 12:150-155. Doi 10.1626/Pps.12.150
- 320 Exposito-Rodriguez M, Borges AA, Borges-Perez A, and Perez JA. 2008. Selection of  
321 internal control genes for quantitative real-time RT-PCR studies during tomato  
322 development process. *BMC Plant Biol* 8:131. 10.1186/1471-2229-8-131
- 323 Gao F, Zhao HX, Yao HP, Li CL, Chen H, Wang AH, Park SU, and Wu Q. 2016.  
324 Identification, isolation and expression analysis of eight stress-related R2R3-MYB  
325 genes in tartary buckwheat (*Fagopyrum tataricum*). *Plant Cell Rep* 35:1385-1396.  
326 10.1007/s00299-016-1971-5
- 327 Ginzinger DG. 2002. Gene quantification using real-time quantitative PCR. *Experimental*  
328 *Hematology* 30:503-512. 10.1016/s0301-472x(02)00806-8
- 329 Gupta N, Sharma SK, Rana JC, and Chauhan RS. 2011. Expression of flavonoid  
330 biosynthesis genes vis-a-vis rutin content variation in different growth stages of  
331 *Fagopyrum* species. *Journal of Plant Physiology* 168:2117-2123.  
332 10.1016/j.jplph.2011.06.018
- 333 Gutierrez L, Mauriat M, Guenin S, Pelloux J, Lefebvre JF, Louvet R, Rusterucci C, Moritz  
334 T, Guerineau F, Bellini C, and Van Wuytswinkel O. 2008. The lack of a systematic  
335 validation of reference genes: a serious pitfall undervalued in reverse  
336 transcription-polymerase chain reaction (RT-PCR) analysis in plants. *Plant*  
337 *Biotechnol J* 6:609-618. 10.1111/j.1467-7652.2008.00346.x
- 338 Hong SY, Seo PJ, Yang MS, Xiang F, and Park CM. 2008. Exploring valid reference genes  
339 for gene expression studies in *Brachypodium distachyon* by real-time PCR. *BMC*  
340 *Plant Biol* 8:112. 10.1186/1471-2229-8-112
- 341 Hruz T, Wyss M, Docquier M, Pfaffl MW, Masanetz S, Borghi L, Verbrugghe P, Kalaydjieva  
342 L, Bleuler S, Laule O, Descombes P, Gruissem W, and Zimmermann P. 2011.  
343 RefGenes: identification of reliable and condition specific reference genes for RT-  
344 qPCR data normalization. *BMC Genomics* 12:156. 10.1186/1471-2164-12-156
- 345 Hu R, Fan C, Li H, Zhang Q, and Fu YF. 2009. Evaluation of putative reference genes for  
346 gene expression normalization in soybean by quantitative real-time RT-PCR. *BMC*  
347 *Mol Biol* 10:93. 10.1186/1471-2199-10-93
- 348 Huang CF, Yamaji N, Mitani N, Yano M, Nagamura Y, and Ma JF. 2009. A bacterial-type  
349 ABC transporter is involved in aluminum tolerance in rice. *Plant Cell* 21:655-667.  
350 10.1105/tpc.108.064543
- 351 Huang L, Yan H, Jiang X, Yin G, Zhang X, Qi X, Zhang Y, Yan Y, Ma X, and Peng Y. 2014.  
352 Identification of candidate reference genes in perennial ryegrass for quantitative  
353 RT-PCR under various abiotic stress conditions. *PLoS One* 9:e93724.  
354 10.1371/journal.pone.0093724
- 355 Huis R, Hawkins S, and Neutelings G. 2010. Selection of reference genes for quantitative  
356 gene expression normalization in flax (*Linum usitatissimum* L.). *BMC Plant Biol*  
357 10:71. 10.1186/1471-2229-10-71
- 358 Jiang Q, Wang F, Li MY, Ma J, Tan GF, and Xiong AS. 2014. Selection of suitable  
359 reference genes for qPCR normalization under abiotic stresses in *Oenanth*  
360 *javanica* (Bl.) DC. *PLoS One* 9:e92262. 10.1371/journal.pone.0092262
- 361 Kim YB, Park SY, Thwe AA, Seo JM, Suzuki T, Kim SJ, Kim JK, and Park SU. 2013.  
362 Metabolomic analysis and differential expression of anthocyanin biosynthetic  
363 genes in white- and red-flowered buckwheat cultivars (*Fagopyrum esculentum*). *J*  
364 *Agric Food Chem* 61:10525-10533. 10.1021/jf402258f
- 365 Kim YK, Li XH, Xu H, Il Park N, Uddin MR, Pyon JY, and Park SU. 2009. Production of  
366 phenolic compounds in hairy root culture of tartary buckwheat (*Fagopyrum*  
367 *tataricum* Gaertn). *Journal of Crop Science and Biotechnology* 12:53-57.  
368 10.1007/s12892-009-0075-y


- 369 Kravets EA, Yemets AI, and Blume YB. 2017. Cytoskeleton and nucleoskeleton  
370 involvement in processes of cytomixis in plants. *Cell Biol Int*. 10.1002/cbin.10842
- 371 Kumar V, Sharma R, Trivedi PC, Vyas GK, and Khandelwal V. 2011. Traditional and novel  
372 references towards systematic normalization of qRT-PCR data in plants.  
373 *Australian Journal of Crop Science* 5:1455-1468. 10.1016/j.fcr.2011.04.016
- 374 Larsen PB, Geisler MJ, Jones CA, Williams KM, and Cancel JD. 2005. ALS3 encodes a  
375 phloem-localized ABC transporter-like protein that is required for aluminum  
376 tolerance in *Arabidopsis*. *Plant J* 41:353-363. 10.1111/j.1365-313X.2004.02306.x
- 377 Li C, Bai Y, Li S, Chen H, Han X, Zhao H, Shao J, Park SU, and Wu Q. 2012. Cloning,  
378 characterization, and activity analysis of a flavonol synthase gene FtFLS1 and its  
379 association with flavonoid content in tartary buckwheat. *J Agric Food Chem*  
380 60:5161-5168. 10.1021/jf205192q
- 381 Lovdal T, and Lillo C. 2009. Reference gene selection for quantitative real-time PCR  
382 normalization in tomato subjected to nitrogen, cold, and light stress. *Anal*  
383 *Biochem* 387:238-242. 10.1016/j.ab.2009.01.024
- 384 Luo XP, Zhao HX, Xue J, Li CL, Chen H, Park SU, and Wu Q. 2016. Cloning of two basic  
385 helix-loop-helix transcription factor genes from Tartary buckwheat (*Fagopyrum*  
386 *tataricum*) and their expression under abiotic stress. *Turkish Journal of Biology*  
387 40:1192-1201. 10.3906/biy-1511-36
- 388 Morrison TB, Weis JJ, and Wittwer CT. 1998. Quantification of low-copy transcripts by  
389 continuous SYBR Green I monitoring during amplification. *Biotechniques* 24:954-  
390 958, 960, 962. 10.1271/bbb.62.1271
- 391 Nolan T, Hands RE, and Bustin SA. 2006. Quantification of mRNA using real-time RT-  
392 PCR. *Nat Protoc* 1:1559-1582. 10.1038/nprot.2006.236
- 393 Pfaffl MW, Tichopad A, Prgomet C, and Neuvians TP. 2004. Determination of stable  
394 housekeeping genes, differentially regulated target genes and sample integrity:  
395 BestKeeper--Excel-based tool using pair-wise correlations. *Biotechnol Lett*  
396 26:509-515. 10.1023/B:BILE.0000019559.84305.47
- 397 Remans T, Keunen E, Bex GJ, Smeets K, Vangronsveld J, and Cuypers A. 2014. Reliable  
398 gene expression analysis by reverse transcription-quantitative PCR: reporting and  
399 minimizing the uncertainty in data accuracy. *Plant Cell* 26:3829-3837.  
400 10.1105/tpc.114.130641
- 401 Stürzenbaum SR, and Kille P. 2001. Control genes in quantitative molecular biological  
402 techniques: the variability of invariance. *Comparative Biochemistry and*  
403 *Physiology Part B: Biochemistry and Molecular Biology* 130:281-289.  
404 10.1016/s1096-4959(01)00440-7
- 405 Stamova BS, Apperson M, Walker WL, Tian Y, Xu H, Adamczyk P, Zhan X, Liu DZ, Ander  
406 BP, Liao IH, Gregg JP, Turner RJ, Jickling G, Lit L, and Sharp FR. 2009.  
407 Identification and validation of suitable endogenous reference genes for gene  
408 expression studies in human peripheral blood. *BMC Med Genomics* 2:49.  
409 10.1186/1755-8794-2-49
- 410 Suzuki T, Honda Y, and Mukasa Y. 2005. Effects of UV-B radiation, cold and desiccation  
411 stress on rutin concentration and rutin glucosidase activity in tartary buckwheat  
412 (*Fagopyrum tataricum*) leaves. *Plant Science* 168:1303-1307.  
413 10.1016/j.plantsci.2005.01.007
- 414 Terzi V, Morcia C, Spini M, Tudisco R, Cutrignelli MI, Infascelli F, Stanca AM, and Faccioli  
415 P. 2010. Identification and validation of reference genes for gene expression  
416 studies in water buffalo. *Animal* 4:853-860. 10.1017/S1751731110000042
- 417 Tian C, Jiang Q, Wang F, Wang GL, Xu ZS, and Xiong AS. 2015. Selection of suitable  
418 reference genes for qPCR normalization under abiotic stresses and hormone  
419 stimuli in carrot leaves. *PLoS One* 10:e0117569. 10.1371/journal.pone.0117569

- 420 Tong Z, Gao Z, Wang F, Zhou J, and Zhang Z. 2009. Selection of reliable reference genes  
421 for gene expression studies in peach using real-time PCR. *BMC Mol Biol* 10:71.  
422 10.1186/1471-2199-10-71
- 423 Vandesompele J, De Preter K, Pattyn F, Poppe B, Van Roy N, De Paepe A, and Speleman  
424 F. 2002. Accurate normalization of real-time quantitative RT-PCR data by  
425 geometric averaging of multiple internal control genes. *Genome Biol*  
426 3:RESEARCH0034. 10.1186/gb-2002-3-7-research0034
- 427 Wan H, Zhao Z, Qian C, Sui Y, Malik AA, and Chen J. 2010. Selection of appropriate  
428 reference genes for gene expression studies by quantitative real-time polymerase  
429 chain reaction in cucumber. *Anal Biochem* 399:257-261. 10.1016/j.ab.2009.12.008
- 430 Wang H, Chen RF, Iwashita T, Shen RF, and Ma JF. 2015. Physiological characterization  
431 of aluminum tolerance and accumulation in tartary and wild buckwheat. *New*  
432 *Phytol* 205:273-279. 10.1111/nph.13011
- 433 Wu BY, Ye JR, Huang L, He LM, and Li DW. 2017. Validation of reference genes for RT-  
434 qPCR analysis in Burkholderia pyrrocinia JK-SH007. *J Microbiol Methods* 132:95-  
435 98. 10.1016/j.mimet.2016.10.004
- 436 Wu ZJ, Tian C, Jiang Q, Li XH, and Zhuang J. 2016. Selection of suitable reference genes  
437 for qRT-PCR normalization during leaf development and hormonal stimuli in tea  
438 plant (*Camellia sinensis*). *Sci Rep* 6:19748. 10.1038/srep19748
- 439 Yao H, Li C, Zhao H, Zhao J, Chen H, Bu T, Anhu W, and Wu Q. 2017. Deep sequencing of  
440 the transcriptome reveals distinct flavonoid metabolism features of black tartary  
441 buckwheat (*Fagopyrum tataricum* Garetn.). *Prog Biophys Mol Biol* 124:49-60.  
442 10.1016/j.pbiomolbio.2016.11.003
- 443 Yuan T, Fujioka S, Takatsuto S, Matsumoto S, Gou XP, He K, Russell SD, and Li J. 2007.  
444 *BEN1*, a gene encoding a dihydroflavonol 4-reductase (DFR)-like protein,  
445 regulates the levels of brassinosteroids in *Arabidopsis thaliana*. *The Plant Journal*  
446 51:220-233. 10.1111/j.1365-313X.2007.03129.x
- 447 Zhang L, Li X, Ma B, Gao Q, Du H, Han Y, Li Y, Cao Y, Qi M, Zhu Y, Lu H, Ma M, Liu L,  
448 Zhou J, Nan C, Qin Y, Wang J, Cui L, Liu H, Liang C, and Qiao Z. 2017. The Tartary  
449 Buckwheat Genome Provides Insights into Rutin Biosynthesis and Abiotic Stress  
450 Tolerance. *Mol Plant* 10:1224-1237. 10.1016/j.molp.2017.08.013
- 451 Zhu H, Wang H, Zhu Y, Zou J, Zhao FJ, and Huang CF. 2015. Genome-wide transcriptomic  
452 and phylogenetic analyses reveal distinct aluminum-tolerance mechanisms in the  
453 aluminum-accumulating species buckwheat (*Fagopyrum tataricum*). *BMC Plant*  
454 *Biol* 15:16. 10.1186/s12870-014-0395-z

# Figure 1


Cq values of seven candidate reference genes across all experimental sets in tartary buckwheat.

A line across the box depicts the median. The box indicates the 25% and 75% percentiles. Whiskers represent the maximum and minimum values, and asterisks indicate extremes.



# Figure 2

Optimal number of reference genes required for accurate normalization entile.






# Figure 3

Relative quantification of *FtSTAR2* gene expression under AI treatment in tartary buckwheat.


Error bars represent standard deviation of the mean, data shown are means  $\pm$  SD (n = 3).



# Figure 4

Relative quantification of *FtDFR* gene expression under UV treatment in tartary buckwheat hypocotyls.

Error bars represent standard deviation of the mean, data shown are means  $\pm$  SD (n = 3).



**Table 1**(on next page)

Genes, primers and different features derived from qRT-PCR analysis

1 **Table 1** Genes, primers and different features derived from qRT-PCR analysis

Gene symbol	Gene name	Amplification length (bp)	Primer sequences	E%	T <sub>M</sub> °C
<i>FtSAND</i>	SAND family protein gene	79	GACCCCCTTGCAGACAAAGCATTGGCA TCTCGTTCTCAACGTCTTTTACCCACTGG	98.9	81
<i>FtCACS</i>	Clathrin adapter complex subunit family protein	125	AAGACAGTCAGTTTCGTGCCACCTGA TCCATGCGTGTCTACCCAACCTCCTT	90.3	82.5
<i>FtExpressed1</i>	Expressed protein of unknown function	127	AGGCCAGTTCCTGCTGAATGTAATGC TAGCCTGATCCAAACAAGCCTGGCAA	90.9	83
<i>FtH<sub>3</sub></i>	Histones 3	158	GAAATTCGCAAGTACCAGAAGAG CCAACAAGGTATGCCTCAGC	109.3	85
<i>FtGAPDH</i>	Glyceradehyde-3-phosphate dehydrogenase gene	155	TGGAGCTGCTAAGGCTGTCTG TGATAGCACTCTTGATGTCCTCGTA	91.9	83
<i>FtActin</i>	Actin 2 genes	118	GGAAGTATAGCGTCTGGATTGGC CACTTGCGGTGAACGATTGC	93.1	82.5
<i>FtEF-1α</i>	Elongation factor-1α gene	108	GCTGCTGAGATGAACAAGAGGTC CTCAAACCTCCACAACGCGAT	91.2	82.5

2

# **Table 2**(on next page)

The stability of the expression of seven candidate reference genes in six groups as analysed by geNorm, NormFinder, BestKeeper

**Table 2** The stability of the expression of seven candidate reference genes in six groups as analysed by geNorm, NormFinder, BestKeeper

Group	Rank	geNorm		NormFinder		BestKeeper		
		Gene	Stability	Gene	Stability	Gene	SD	CV
Abiotic Cotyledons	1	<i>FtCACS</i>	0.33	<i>FtH<sub>3</sub></i>	0.058	<i>FtSAND</i>	0.33	1.32
	2	<i>FtH<sub>3</sub></i>	0.33	<i>FtCACS</i>	0.080	<i>FtCACS</i>	0.31	1.47
	3	<i>FtSAND</i>	0.41	<i>FtSAND</i>	0.129	<i>FtH<sub>3</sub></i>	0.33	1.64
	4	<i>FtExpressed1</i>	0.54	<i>FtExpressed1</i>	0.280	<i>FtExpressed1</i>	0.59	2.48
	5	<i>FtEF-1α</i>	0.73	<i>FtEF-1α</i>	0.683	<i>FtEF-1α</i>	1.04	5.70
	6	<i>FtGAPDH</i>	0.84	<i>FtGAPDH</i>	0.751	<i>FtGAPDH</i>	1.07	6.26
	7	<i>FtActin</i>	1.31	<i>FtActin</i>	1.678	<i>FtActin</i>	1.46	6.36
Abiotic Hypocotyls	1	<i>FtH<sub>3</sub></i>	0.65	<i>FtSAND</i>	0.416	<i>FtH<sub>3</sub></i>	0.51	2.47
	2	<i>FtSAND</i>	0.65	<i>FtCACS</i>	0.437	<i>FtSAND</i>	0.69	2.68
	3	<i>FtCACS</i>	0.72	<i>FtH<sub>3</sub></i>	0.449	<i>FtActin</i>	0.79	3.38
	4	<i>FtExpressed1</i>	0.84	<i>FtEF-1α</i>	0.469	<i>FtExpressed1</i>	0.81	3.34
	5	<i>FtEF-1α</i>	0.91	<i>FtExpressed1</i>	0.475	<i>FtCACS</i>	0.86	3.75
	6	<i>FtGAPDH</i>	1.05	<i>FtGAPDH</i>	0.811	<i>FtEF-1α</i>	0.92	5.32
	7	<i>FtActin</i>	1.17	<i>FtActin</i>	0.895	<i>FtGAPDH</i>	1.35	7.59
Abiotic total	1	<i>FtH<sub>3</sub></i>	0.55	<i>FtCACS</i>	0.304	<i>FtH<sub>3</sub></i>	0.52	2.55
	2	<i>FtSAND</i>	0.55	<i>FtSAND</i>	0.327	<i>FtCACS</i>	0.61	2.70
	3	<i>FtCACS</i>	0.61	<i>FtH<sub>3</sub></i>	0.349	<i>FtSAND</i>	0.63	2.46
	4	<i>FtExpressed1</i>	0.72	<i>FtExpressed1</i>	0.380	<i>FtExpressed1</i>	0.73	3.02
	5	<i>FtEF-1α</i>	0.89	<i>FtEF-1α</i>	0.694	<i>FtEF-1α</i>	1.02	5.71
	6	<i>FtGAPDH</i>	1.01	<i>FtGAPDH</i>	0.782	<i>FtActin</i>	1.17	5.07
	7	<i>FtActin</i>	1.30	<i>FtActin</i>	1.329	<i>FtGAPDH</i>	1.20	6.89
AI treatment	1	<i>FtEF-1α</i>	0.16	<i>FtSAND</i>	0.091	<i>FtExpressed1</i>	0.17	0.73
	2	<i>FtExpressed1</i>	0.16	<i>FtExpressed1</i>	0.100	<i>FtEF-1α</i>	0.25	1.55
	3	<i>FtSAND</i>	0.21	<i>FtH<sub>3</sub></i>	0.105	<i>FtH<sub>3</sub></i>	0.31	1.56
	4	<i>FtGAPDH</i>	0.23	<i>FtEF-1α</i>	0.106	<i>FtGAPDH</i>	0.34	1.88
	5	<i>FtH<sub>3</sub></i>	0.27	<i>FtGAPDH</i>	0.156	<i>FtCACS</i>	0.35	1.57
	6	<i>FtCACS</i>	0.30	<i>FtCACS</i>	0.282	<i>FtSAND</i>	0.37	1.43
	7	<i>FtActin</i>	0.49	<i>FtActin</i>	0.652	<i>FtActin</i>	1.05	4.16
Immature	1	<i>FtExpressed1</i>	0.16	<i>FtSAND</i>	0.097	<i>FtExpressed1</i>	0.50	2.00

e seed stage	2	<i>FtCACS</i>	0.16	<i>FtGAPDH</i>	0.097	<i>FtCACS</i>	0.57	2.43
	3	<i>FtSAND</i>	0.30	<i>FtH<sub>3</sub></i>	0.167	<i>FtSAND</i>	0.82	3.03
	4	<i>FtGAPDH</i>	0.36	<i>FtCACS</i>	0.292	<i>FtEF-1α</i>	0.83	4.32
	5	<i>FtH<sub>3</sub></i>	0.45	<i>FtExpressed1</i>	0.422	<i>FtGAPDH</i>	0.86	4.51
	6	<i>FtEF-1α</i>	0.62	<i>FtEF-1α</i>	0.675	<i>FtH<sub>3</sub></i>	1.11	5.29
	7	<i>FtActin</i>	1.01	<i>FtActin</i>	1.354	<i>FtActin</i>	2.16	9.96
Total	1	<i>FtH<sub>3</sub></i>	0.60	<i>FtCACS</i>	0.245	<i>FtCACS</i>	0.53	2.33
	2	<i>FtCACS</i>	0.60	<i>FtSAND</i>	0.304	<i>FtH<sub>3</sub></i>	0.54	2.69
	3	<i>FtSAND</i>	0.66	<i>FtH<sub>3</sub></i>	0.313	<i>FtSAND</i>	0.57	2.24
	4	<i>FtExpressed1</i>	0.71	<i>FtExpressed1</i>	0.377	<i>FtExpressed1</i>	0.68	2.85
	5	<i>FtGAPDH</i>	0.87	<i>FtGAPDH</i>	0.650	<i>FtGAPDH</i>	0.97	5.53
	6	<i>FtEF-1α</i>	0.97	<i>FtEF-1α</i>	0.780	<i>FtEF-1α</i>	1.03	5.91
	7	<i>FtActin</i>	1.33	<i>FtActin</i>	1.480	<i>FtActin</i>	1.36	5.70

3