

Placental vascular pathology and increased thrombin generation as mechanisms of disease in obstetrical syndromes

Obstetrical complications including preeclampsia, fetal growth restriction, preterm labor, preterm prelabor rupture of membranes and fetal demise are all the clinical endpoint of several underlying mechanisms (i.e. infection inflammation, thrombosis, endocrine disorder, immunologic rejection, genetic, and environmental), therefore, they may be regarded as syndromes. Placental vascular pathology and increased thrombin generation were reported in all of these obstetrical syndromes. Moreover, elevated concentrations of thrombin-anti-thrombin III complexes and changes in the coagulation as well as anticoagulation factors can be detected in the maternal circulation prior to the clinical development of the disease in some of these syndromes. In this review, we will assess the changes in the hemostatic system during normal and complicated pregnancy in maternal blood, maternal-fetal interface and amniotic fluid, and describe the contribution of thrombosis and vascular pathology to the development of the great obstetrical syndromes.

1 Placental Vascular Pathology and Increased Thrombin Generation as Mechanisms of 2 Disease In Obstetrical Syndromes

3 Salvatore Andrea Mastrolia^{1,2}, Moshe Mazor², Giuseppe Loverro¹, Vered Klaitman², and Offer
4 Erez²

5 ¹ Department of Obstetrics and Gynecology, Azienda Ospedaliero-Universitaria Policlinico di
6 Bari, School of Medicine, University of Bari “Aldo Moro”, Bari, Italy
7 ²Obstetrics and Gynecology, Soroka University Medical Center, School of Medicine, Ben Gurion
8 University of the Negev, Beer Sheva, Israel

9 Corresponding author

10 Offer Erez M.D.

11 Acting director Maternal Fetal Medicine Unit

12 Department of Obstetrics and Gynecology

13 Soroka University Medical Center

14 School of Medicine, Faculty of Health Sciences

15 Ben Gurion University of the Negev

16 P.O.Box 151,

17 Beer Sheva 84101

18 Israel

19 erezof@bgu.ac.il

20 1. Introduction

21 Obstetrical complications including preeclampsia, fetal growth restriction, preterm labor, preterm
22 prelabor rupture of membranes and fetal demise are all the clinical endpoint of several underlying
23 mechanisms (i.e. infection inflammation, thrombosis, endocrine disorder, immunologic rejection,
24 genetic, and environmental), therefore, they may be regarded as syndromes. In this review, we
25 will assess the changes in the hemostatic system during normal and complicated pregnancy in
26 maternal blood, maternal-fetal interface and amniotic fluid, and describe the contribution of
27 thrombosis and vascular pathology to the development of the great obstetrical syndromes.

28 2. What are the great obstetrical syndromes?

The major obstetrical complications including preeclampsia, intrauterine growth restriction (IUGR), preterm labor (PTL), preterm prelabor rupture of membranes (PROM), fetal demise, and recurrent abortions are all syndromes, also defined as "great obstetrical syndromes". As reported in The Oxford Medical Dictionary a syndrome is 'a combination of symptoms and/or signs that form a distinct clinical picture indicative of a particular disorder'. Hence, they represent the clinical manifestation of many possible underlying mechanisms of disease¹. Key features of these syndromes are²: multiple etiologies; long preclinical stage; frequent fetal involvement; clinical manifestations which are often adaptive in nature; and predisposition to a particular syndrome is influenced by gene–environment interaction and/or complex gene-gene interactions involving maternal and/or fetal genotypes. These mechanisms of disease were identified and reported in all the obstetrical complications listed above. This review is focused on the role of thrombosis and vascular pathology of the placenta in these syndromes.

3. What are the changes in the coagulation system during normal pregnancy?

In term of the coagulation and hemostatic systems there are several major compartments: the maternal circulation, the fetal maternal interface (the placenta, and membranes), amniotic fluid and the fetus that each has a specific behavior during gestation. The changes in the coagulation system during gestation are considered to be adaptive mechanisms and are aimed to: 1) the prevention of bleeding at the time of trophoblast implantation and the delivery of the fetus; 2) allow the laminar flow and the intervillous space; and 3) seal amniotic fluid leak and reduce obstetrical bleeding³⁻⁷. Of interest, the fetus is somewhat less involved and its coagulation system develops during gestation, and this subject is beyond the scope of this review. Indeed, normal pregnancy has been associated with excessive maternal thrombin generation^{3, 8} and a tendency for platelets to aggregate in response to agonists^{9, 10}. Pregnancy is accompanied by 2 to 3-fold increase in fibrinogen concentrations and 20% to 1000% increase in factors VII, VIII, IX, X, and XII, all of which peak at term¹¹. The concentrations of vWF increase up to 400% by term¹¹. By contrast, those of pro-thrombin and factor V remain unchanged while the concentrations of factors XIII and XI decline modestly¹². Indeed there is evidence of chronic low-level thrombin and fibrin generation throughout normal pregnancy as indicated by enhanced concentrations of pro-thrombin fragment 1.2, thrombin-antithrombin (TAT) III complexes, and soluble fibrin polymers¹³. Free protein S concentration declines significantly (up to 55%) during pregnancy due to increased circulating complement 4B-binding protein its molecular carrier. Protein S nadir at delivery and this reduction is exacerbated by cesarean delivery and infection^{11, 12}. As a consequence, pregnancy is associated with an increase in resistance to activated protein

C^{12, 13}. The concentrations of PAI-1 increase by 3 to 4-folds during pregnancy while plasma PAI-2 values, which are negligible before pregnancy reach concentrations of 160 mg/L at delivery¹¹. Thus, pregnancy is associated with increased clotting potential, as well as decreased anticoagulant properties, and fibrinolysis¹⁴. Therefore, it can be defined as a prothrombotic state. One of the most important mediators of the hypercoagulable state of normal pregnancy is tissue factor (TF). Indeed, there is a substantial increase in TF concentrations in the decidua and myometrium¹⁵⁻¹⁸, preventing placental abruption since this leads to an increase in the efficiency of clotting function¹⁴. The placenta is a source of TF, since trophoblast cells constitutively express it, behaving as activated endothelium, and leading to a condition of procoagulant state that, if not controlled by anticoagulant mechanisms, predisposes to thrombotic complications¹⁵. The principal anticoagulant mechanism inhibiting TF activation pathway is tissue factor pathway inhibitor (TFPI), which mRNA is highly expressed in the macrophages in the villi in term placenta¹⁹. Similarly, high TF concentrations have been detected in the fetal membranes (mainly the amnion) and amniotic fluid^{7, 20-23}. TFPI has been found in amniotic fluid as well²⁰, but it is not clear if the presence of TF and its natural inhibitor is related to coagulation per se or is somehow connected with embryonic development²⁴. In contrast to the changes detected in the amniotic fluid and the decidua, the median maternal plasma immunoreactive TF concentration of normal pregnant women do not differ significantly from that of non-pregnant patients^{3, 25}. However, labor at term increases significantly the maternal plasma immunoreactive TF concentration in comparison to the non-pregnant state²⁰. In addition to the changes in TF, normal pregnancy is associated with increased thrombin generation^{3, 8}, as determined by the elevation of maternal concentrations of fibrinopeptide A, prothrombin fragments (PF) 1 and 2, and thrombin–antithrombin (TAT) III complexes^{7, 26-28}. The concentration of these complexes further increases during and after normal parturition^{27, 29}, and subsequently decreases during the puerperium^{27, 29}.

4. What are the changes in the hemostatic system associated with the great obstetrical syndromes?

The great obstetrical syndromes are associated with changes in the hemostatic and vascular systems in the compartments mentioned above: 1) the maternal circulation; 2) the fetomaternal interface of placenta and membranes; 3) and the amniotic fluid.

4.1 Changes in the hemostatic system of women with obstetrical syndromes.

The involvement of the hemostatic system in the pathophysiology of these obstetrical syndromes is becoming more and more apparent. Indeed, increased thrombin generation is reported in the maternal circulation of women with preeclampsia³⁰⁻³⁴, IUGR^{30-32, 35, 36}, fetal demise³⁷, PTL^{8, 37, 38} and preterm PROM^{8, 37, 39}.

There are several possible explanations for the increased thrombin generation in these patients: 1) increased activation of coagulation cascade in the maternal circulation due to pathological processes including bleeding or inflammation; and 2) depletion of anticoagulation proteins that subsequently leads to increased thrombin generation (Table 1).

4.1.1 Increased activation of the coagulation cascade and thrombin generation in the maternal circulation in patients with pregnancy complications

All the obstetrical syndromes including preeclampsia^{30-34, 40, 41}, IUGR^{31, 32, 35, 36}, fetal demise³⁷, PTL^{8, 38} and preterm PROM^{8, 37, 39} are associated with a higher maternal thrombin generation than a normal pregnancy. These may be of clinical implication since in women with preterm labor, elevated maternal plasma TAT III complexes concentration was associated with a higher chance to deliver within <7 days from admission³⁷ (Fig. 1). To further understand how does thrombin affect the duration of pregnancy and the clinical phenotype of patients with the obstetrical syndromes we need to consider what are the mechanisms leading to thrombin generation and how it affects the fetomaternal unit.

Increased thrombin generation can result from the following underlying mechanisms: 1) decidual hemorrhage that leads to a retro-placental clot formation⁴²; 2) intra-amniotic infection/inflammation which can induce decidual bleeding and sub-clinical abruption⁴³, as well as increased intra-amniotic TAT complexes³⁷; and 3) an increased maternal systemic inflammatory response⁴⁴ that may activate the extrinsic pathway of coagulation due to the expression and release of TF by activated monocytes⁴⁵.

Thrombin affects many systems including also the following: 1) stimulation of decidual cell secretion of matrix metalloproteinase (MMP) (i.e. MMP-1 and MMP-3) that can degrade the extracellular matrix of the chorioamniotic membranes^{46, 47} (as in preterm PROM); 2) myometrial activation and uterine contractions generation that may lead to preterm labor with or without rupture of membranes and a subsequent preterm delivery^{38, 48, 49}; and 3) thrombin has an inhibitory effect on the production of TFPI by endothelial cells⁵⁰, and the increased thrombin generation observed in patients with PTL may be associated with a concomitant reduction in TFPI production by the maternal vascular endothelium (the depletion of anticoagulant proteins will be discussed in the following section of this review).

There is evidence to support that the extrinsic pathway of coagulation is activated in many of these pregnancy complications and it is the source of the increased thrombin generation⁵¹. Indeed, increased immunoreactive TF concentrations were reported in women with preeclampsia and those with preterm PROM⁵². Moreover, the contribution of preeclampsia to elevated maternal immunoreactive TF persisted also among patients with fetal demise, while those with fetal death who were normotensive did not have higher median TF concentration than normal pregnant women. Moreover, the median TF concentration of patients with preeclampsia was also higher than in patients with fetal demise without hypertension. These findings are consistent with previous studies^{3, 53}, suggesting that elevated TF immunoreactivity and activity may be associated with the pathophysiologic process leading to preeclampsia, rather than being a consequence of the fetal death.

In some of the obstetrical syndromes there was elevated TF activity in the maternal circulation without a concomitant increase in the plasma concentration of this factor. This was the case among patients with a small for gestational age (SGA) neonate and those with preterm labor^{31, 54} (Table 1). This suggests that the increased TF activity among patients with PTL as well as those with an SGA neonate, contributes to a higher generation of factor Xa that, along with the physiologic increase in the maternal plasma concentrations of factor VII and factor X during gestation^{11, 55-57}, may be the underlying mechanism leading to the increased thrombin generation reported these syndromes.

The differences between PTL and preterm PROM in term of maternal plasma TF concentration and activity may derive from the specific component of the common pathway of parturition, which is activated in each obstetrical syndrome⁵⁸. While preterm PROM is associated with the activation of the decidua and the membranes, myometrial activation is the major component of preterm labor with intact membranes⁵⁸. This is relevant because the decidua and the membranes have a high TF concentration^{17, 18, 59}.

In summary, the evidence brought herein suggests that increased thrombin generation in patients with the great obstetrical syndromes may reflect the activation of the coagulation cascade mainly through the extrinsic arm. This activation can be attributed to various underlying mechanisms.

4.1.2 Depleted or insufficient anticoagulant proteins concentration

In the normal state there is a delicate balance between the proteins activating/participating the coagulation cascade and their inhibitors. Increased thrombin generation may result, as we presented above, from activation of the coagulation cascade due to higher concentrations or activities of the proteins included in the coagulation cascade. However, thrombin generation can also result from insufficient concentration or activity of anticoagulation proteins.

Tissue factor pathway inhibitor (TFPI), a glycoprotein comprising of three Kunitz domain⁶⁰ that are specific inhibitors of trypsin-like proteinases⁶¹, is the main inhibitor of the extrinsic pathway of coagulation. TFPI inhibits thrombin generation through the inactivation of activated factor X and the factor VIIa/TF complex^{60, 62}. The mean maternal plasma concentrations of total TFPI increases during the first half of pregnancy, remains relatively constant in the second half⁶³ and decreases during labor²⁰. There are two types of TFPI: 1) TFPI-1 is the more prevalent form in the non-pregnant state in the maternal circulation and can also be found in the fetal blood, platelets, endothelial cells and other organs^{19, 64}; and 2) TFPI-2- the major form of TFPI in the placenta⁶⁵⁻⁶⁸, also known as Placental Protein 5 (PP5)^{69, 70}. During pregnancy, the maternal plasma concentration of TFPI-2 increases gradually, reaches a plateau at 36 weeks and subsides after delivery⁷¹⁻⁷⁴.

The overall balance between the concentration and activity of the coagulation factors and the anti-coagulation proteins is one of the determining factors of thrombin generation. In the normal state, the immunoreactive concentrations of TFPI in the plasma are 500 to 1000 times higher than that of TF⁷⁵, suggesting that an excess of anti-coagulant proteins closely controls the coagulation cascade activity. The median maternal plasma TFPI concentration increases during preeclampsia^{53, 76}, which is associated with an exaggerated maternal systemic inflammatory response. However, the increase in the median maternal TF plasma concentration is such that the overall balance between TF and its inhibitor is affected leading to increased thrombin generation in this syndrome. In contrast to preeclampsia, maternal plasma TFPI concentration decreases in patients with PTL⁵² and preterm PROM⁷⁷ regardless to the presence of intra-amniotic infection/inflammation, as well as in women with fetal demise⁵⁴, and does not change in mothers with SGA fetuses⁵³. Overall these findings suggest that the increased thrombin generation observed among these patients may derive not only from an increased activation of the hemostatic system, but also from insufficient anti-coagulation, as reflected by the lower TFPI concentrations (Fig. 2).

A possible explanation of the lower maternal plasma concentration observed in some of the obstetrical syndromes may be that during these syndromes there is a reduction in the placental production of TFPI^{65, 66, 69, 76} (mainly TFPI-2), contributing to the low maternal plasma concentrations detected in patients with PTL, in addition to the thrombin inhibitory effect to TFPI expression on endothelial cells, as above mentioned. Indeed, patients with vascular complications of pregnancy (preeclampsia, eclampsia, placental abruption, fetal growth restriction, and fetal

demise) have a lower placental concentration of total TFPI, and TFPI mRNA expression than in women with normal pregnancies^{78, 79}. Other proteins implicated in the inhibitory control of the coagulation cascade are protein S, protein C and protein Z. Protein S is a cofactor to protein C in the inactivation of factors Va and VIIIa. This protein exists in two forms: a free form and a complex form bound to complement protein C4b-binding protein (C4BP). Only the free form is active⁸⁰. Protein S also acts as a TFPI cofactor, in the presence of weak pro-coagulant stimuli, by enhancing the interaction of TFPI with factor Xa while using Ca²⁺ and phospholipids in the process⁸¹ without increasing inhibition of factor VIIa-TF by TFPI⁸². During pregnancy there is a physiologic change in the relationship between the bound and the free forms of protein S in the maternal plasma. The increase in C4BP during gestation reduces free protein S concentration in up to 55% of its value out of pregnant state, reaching its nadir at delivery. Of interest, cesarean delivery and infection exacerbate the reduction in free protein S concentrations^{11, 83}. Moreover, a functional protein S deficiency can explain a poor response to activated protein C⁸⁴. The association between the alteration of concentration and function of protein S and protein C in the great obstetrical syndromes is not completely clear. The evidence regarding the association of protein S and protein C deficiency and preeclampsia is controversial^{85, 86}. While some reported an association between protein S deficiency and an increased risk for this syndrome (especially for early onset preeclampsia)⁸⁵ others could not demonstrate this effect⁸⁶. There is some evidence regarding the relation of protein S deficiency and increased risk of stillbirth⁸⁷ and mid-trimester IUGR⁸⁸. An increased risk of stillbirth has been reported in patients with protein S deficiency while the risk was not significantly increased in cases of protein C deficiency⁸⁷, and Kupferminc et al⁸⁸ found that protein S, but not protein C deficiency, was significantly associated with severe mid-trimester IUGR. Protein Z, in complex with protein Z-dependent protease inhibitor (ZPI) (Fig. 3)⁸⁹⁻⁹¹, acts as a physiologic inhibitor of activation of prothrombin by factor Xa. Protein Z is a vitamin K-dependent plasma glycoprotein⁹² that is an essential cofactor for ZPI activity. In the absence of protein Z, the activity of ZPI is reduced by more than 1000-fold⁹¹. Normal pregnancy is characterized by an increased plasma concentration of protein Z⁹³, probably as a compensation for the increase of factor X concentration. Women with preterm labor without intra-amniotic infection or inflammation and those with vaginal bleeding who delivered preterm had a lower median maternal plasma protein Z concentration than women with a normal pregnancy and those with vaginal bleeding who delivered at term⁹⁴. The changes of protein Z concentrations in other pregnancy complications are controversial. Some demonstrated that the median plasma

227 concentration of protein Z in patients with preeclampsia, IUGR, and late fetal death were not
 228 significantly different than that of patients with a normal pregnancy⁹⁵. Others reported lower
 229 median maternal plasma protein Z concentrations in women with preeclampsia or pyelonephritis
 230 and higher proportion of protein Z deficiency (defined as protein Z plasma concentration below
 231 the 5th percentile) in patients with preeclampsia or fetal demise than in those with a normal
 232 pregnancy⁹⁶. Moreover, increased maternal plasma anti-protein Z antibodies concentrations were
 233 associated with SGA neonates, fetal demise and preeclampsia.
 234 The information presented above suggest that it is not only the concentration of one coagulation
 235 factor or anticoagulation protein, but rather the overall balance between the coagulation factors
 236 and their inhibitors that increases thrombin generation in the great obstetrical syndromes. Indeed,
 237 although preterm labor was not associated with a significant change in the median maternal
 238 plasma TF concentration, the TFPI/TF ratio of these patients was lower than that of normal
 239 pregnant women, mainly due to decreased TFPI concentrations.
 240 This observation was also reported in patients with preterm PROM⁷⁷, and those with
 241 preeclampsia⁵³. The lower TFPI/TF ratio in patients with preeclampsia occurs despite the increase
 242 in the median maternal plasma TFPI concentration observed in these patients. This suggests that
 243 the balance between TF and its natural inhibitor may better reflect the overall activity of the TF
 244 pathway of coagulation, than the individual concentrations of TF or TFPI.
 245 Collectively, these observations suggest that our attention should be focused not only on the
 246 coagulation protein but also on their inhibitors since an imbalance between them may contribute
 247 to increased thrombin generation leading to the onset of the great obstetrical syndromes.

4.2 Changes in the feto-maternal interface

Normal placental development and the establishment of an adequate feto-maternal circulation are key points for a successful pregnancy. The networks of the placental vascular tree either on the maternal or fetal side are dynamic structures that can be substantially altered in cases of abnormal placentation and trophoblast invasion. The human trophoblast has properties of endothelial cells and can regulate the degree of activation of the coagulation cascade in the intervillous space^{97, 98}. The villous trophoblasts express heparin sulfate, protein C and protein Z on their surface that serve as anticoagulant that sustain laminar blood flow through the intervillous space. On the other hand, unlike the endothelium of other organs, the trophoblast constantly presents the active placental isoform of TF on its surface⁹⁸⁻¹⁰¹. This isoform has a higher affinity for factor VIIa¹⁰², which may lead to increased activation of the coagulation cascade. One of the leading pathological processes observed in all these syndromes is thrombosis and vascular abnormality of the placenta at the maternal-fetal interface. The incidence of these pathological processes varies among the different syndromes being more prevalent in preeclampsia, IUGR, and fetal demise than in PTL and preterm PROM^{30, 31, 37, 38}.

4.2.1 Placental pathology in the Great Obstetrical Syndromes

There is a range of placental vascular and thrombotic lesions that are being observed in placentas of patients with pregnancy complications. Thrombotic events of placental vessels can cause an impairment of placental perfusion, leading to fetal growth restriction (FGR), preeclampsia and fetal death as well as in some extents to PTL and preterm PROM^{103, 104}. The frequency of the specific vascular placental lesions varies among these obstetrical syndromes¹⁰⁵. Placental vascular lesions are divided into maternal or fetal vascular origin (figure 4-5)^{106, 107}. Lesions of the maternal vascular compartment include placental marginal and retro-placental hemorrhages, lesions related to maternal under perfusion (acute atherosclerosis and mural hypertrophy, increased syncytial knots, villous agglutination, increased intervillous fibrin deposition, villous infarcts)¹⁰⁶. Placental fetal vascular obstructive lesions are the result of stasis, hypercoagulability and vascular damage within the fetal circulation of the placenta. Placental fetal vascular abnormalities include: cord-related abnormalities (as torsion of cord, over-coiling, strictures and tight knots¹⁰⁸) and vascular lesions consistent with fetal thrombo-occlusive disease (thrombosis of the chorionic plate and stem villous vessels, fibrotic, hypo-vascular and avascular villi¹⁰⁶). In addition, villitis of unknown etiology or chronic villitis, defined as lymphohistiocytic

inflammation localized to the stroma of terminal villi but often extending to the small vessels of upstream villi is also associated with obliterative fetal vasculopathy¹⁰⁶ (Fig. 4-5).

Preeclampsia: The classical example for an association between obstetrical syndrome and vascular placental lesions is preeclampsia. Women who develop preeclampsia have an increased rate of abnormalities of the maternal side of the placental circulation and maternal underperfusion^{109, 110}. The frequency of these lesions is inversely related to the gestational age in which the hypertensive disorder was diagnosed. The earlier the development of hypertension/preeclampsia the more severe are the vascular lesions^{111, 112}. Moreover, Kovo et al¹¹³ reported that the presence of fetal growth restriction in women with preeclampsia increases also the frequency of fetal vascular lesions. Indeed, patients with early-onset preeclampsia complicated by FGR had a higher rate of fetal-vascular supply lesions consistent with fetal thrombo-occlusive disease than women with early-onset disease without FGR¹¹³. An assessment of the pathologic changes in placental hemostatic system has been performed in patients with preeclampsia. Teng et al¹¹⁴ studied TF and TFPI placental levels in pregnant patients with preeclampsia, compared to normal pregnancies. They found increased TF placental expression and a reduced expression of TFPI-1 and TFPI-2, with a significant correlation between the levels of TF and TFPI-2 between maternal plasma and placenta.

Fetal growth restriction: Placentas from pregnancies complicated by FGR are smaller and have significantly increased maternal and fetal vascular lesions compared to placentas from normal pregnancies with appropriate for gestational age neonates (AGA)^{115, 116}. Maternal vascular lesions were detected in about 50% of placentas from pregnancies complicated with FGR at term, compared to only 20% in normal pregnancies, while fetal vascular lesions were observed in 11% of FGR pregnancies compared to only 4% in placentas from normal pregnancies¹¹³. Placentas from normotensive pregnancies complicated by early-onset FGR (<34 weeks of gestation) had a higher rate of low placental weight (<10th percentile) and maternal underperfusion, as compared to placentas of women who delivered AGA neonates \leq 34 weeks of gestation¹¹⁵. Of interest, placentas from the late onset FGR group (after 34 weeks of gestation), in addition to the high maternal vascular abnormalities, show also more fetal vascular abnormalities, compared with AGA controls who delivered >34 weeks¹¹⁷.

Fetal demise: Placental disease has been recognized as an important contributor to unexplained fetal demise. Fetal vascular abnormalities¹⁰⁵ are extensively involved in early and late fetal death rather than maternal vascular lesions. In fetal death occurring prior to 34 weeks, an earlier and

extended insult in the placental development occurs. On the other hand, late fetal demise is an unpredicted event that is mostly characterized by non-thrombotic cord related lesions and less placental vascular compromise¹⁰⁷.

Preterm labor and preterm PROM: Placental studies in PTL demonstrated a combination of inflammatory and vascular lesions. PTL is generally attributed to an inflammatory response involving the bacterial induction of cytokine and prostanoid production¹¹⁸. Finding of histological chorioamnionitis in PTL¹¹⁹ has established infection and inflammation as a causative factor of preterm birth, moreover, noninfectious trigger may also contribute to the development of preterm labor and in some instances may be evident by placental sterile inflammatory response¹²⁰. In addition, isolated placental vascular lesions, mostly of maternal supply, were reported in 20% of cases of PTL and an additional 20% had combined inflammatory and vascular lesions. Moreover, there are consistent reports describing increased rate of failure of transformation of the spiral arteries in women with preterm labor without intrauterine infection/inflammation and in those with preterm PROM than in women with normal pregnancies¹²¹. Such findings imply that an inadequate uteroplacental blood flow due to abnormal placentation plays an important role in pathogenesis of preterm parturition^{121, 122}.

Collectively, placental vascular lesions were reported in all the great obstetrical syndromes. The severity of these lesions is associated with the timing of diagnosis of the disease. The more severe the vascular injury, the more likely these complications will become clinically evident prior to 34 weeks of gestation. Of interest, vascular lesions often come along with evidence of acute inflammation or lesions associated with chronic inflammatory processes, suggesting that sometimes more than one mechanism is involved in development of a specific obstetrical syndrome.

4.3. Hemostatic changes in the amniotic fluid of women with obstetrical syndromes

During normal pregnancy, there is an increase in the amniotic fluid TF concentration^{7, 20-23}. In order to demonstrate the association of hemostatic changes and the development of obstetrical complications, Erez et al⁵⁴ studied the changes in the intra-amniotic concentration of TAT III complexes, as well as TF concentration and activity, in cases of fetal demise and in normal pregnancies. Patients with a fetal demise had higher median amniotic fluid–TF concentration and activity than those with normal pregnancies. Moreover, among patients with a FD there was a significant correlation (Fig. 6) between the amniotic fluid–TF concentrations and activity ($r = 0.88$, P

<0.0001). The median amniotic fluid– TAT III complexes concentration did not differ significantly between the groups (normal pregnancy: median: 66.3 mg/l, range 11.4–2265.4 vs. FD: median: 59.3 mg/l, range: 13.6–15,425.3; $P=0.7$). In their study, the median amniotic fluid–TF concentration in normal pregnant women was 10 fold higher than in maternal plasma. The changes in amniotic fluid thrombin generation were reported also in women with preterm parturition. Indeed, intra-amniotic infection and/or inflammation is associated with an increased amniotic fluid TAT III complexes (Fig. 7). This is important since it represents an increased thrombin generation in the amniotic cavity during infection and/or inflammation that may contribute to uterine contractility and the development of preterm birth¹²³. Of interest, elevated intra-amniotic TAT III concentrations were associated with a shorter amniocentesis to delivery interval and an earlier gestational age at delivery only in patients with preterm labor without intra-amniotic infection or inflammation¹²³. This observation suggests that in a subset of patients with preterm labor, activation of the coagulation system can generate preterm parturition and delivery; while in those with intra-amniotic infection and/or inflammation the activation of the coagulation and thrombin generation is a byproduct of the inflammatory process leading to preterm birth. This represents evidence of the activation and propagation of coagulation cascade, being thrombin generation the witness of the former mechanisms and the inhibitor of the initiation step⁵⁴.

5. Conclusion

The evidence presented herein suggests a role for increased thrombin generation and vascular placental lesions in the pathogenesis of the great obstetrical syndromes. This process can be the result of the contribution of procoagulant and vascular abnormalities as well as inflammatory and infectious mechanisms, representing the starting point for pregnancy complications based on vascular disease. As presented, these changes affect the mother, the placenta, membranes and amniotic fluid. Moreover, preliminary evidence suggest that some of the changes in the hemostatic system in the mother and in the amniotic fluid predate the clinical presentation of the disease. Suggesting that better understanding of the vascular and coagulation changes associated with the great obstetrical syndromes may assist us in earlier detection and the development or introduction of therapeutic modalities for these syndromes.

- 377 1. *Concise Medical Dictionary*: Oxford University Press, 2010.
- 378 2. Romero R. Prenatal medicine: the child is the father of the man. 1996. J Matern Fetal
379 Neonatal Med 2009;22:636-9.
- 380 3. Bellart J, Gilabert R, Miralles RM, Monasterio J, Cabero L. Endothelial cell markers and
381 fibrinopeptide A to D-dimer ratio as a measure of coagulation and fibrinolysis balance in
382 normal pregnancy. Gynecol Obstet Invest 1998;46:17-21.
- 383 4. Walker MC, Garner PR, Keely EJ, Rock GA, Reis MD. Changes in activated protein C
384 resistance during normal pregnancy. Am J Obstet Gynecol 1997;177:162-9.
- 385 5. Sørensen JD, Secher NJ, Jespersen J. Perturbed (procoagulant) endothelium and
386 deviations within the fibrinolytic system during the third trimester of normal pregnancy. A
387 possible link to placental function. Acta Obstet Gynecol Scand 1995;74:257-61.
- 388 6. Yuen PM, Yin JA, Lao TT. Fibrinopeptide A levels in maternal and newborn plasma. Eur J
389 Obstet Gynecol Reprod Biol 1989;30:239-44.
- 390 7. [de Boer K](#), [ten Cate JW](#), [Sturk A](#), [Borm JJ](#), [Treffers PE](#). Enhanced thrombin generation in
391 normal and hypertensive pregnancy. Am J Obstet Gynecol 1989;160:95-100.
- 392 8. Chaiworapongsa T, Espinoza J, Yoshimatsu J, Kim YM, Bujold E, Edwin S, Yoon BH,
393 Romero R. Activation of coagulation system in preterm labor and preterm premature
394 rupture of membranes. J Matern Fetal Neonatal Med 2002;11:368-73.
- 395 9. Yoneyama Y, Suzuki S, Sawa R, Otsubo Y, Power GG, Araki T. Plasma adenosine levels
396 increase in women with normal pregnancies. Am J Obstet Gynecol 2000;182:1200-3.
- 397 10. Sheu JR, Hsiao G, Luk HN, Chen YW, Chen TL, Lee LW, Lin CH, Chou DS.
398 Mechanisms involved in the antiplatelet activity of midazolam in human platelets.
399 Anesthesiology 2002;96:651-8.
- 400 11. Bremme KA. Haemostatic changes in pregnancy. Best Pract Res Clin Haematol
401 2003;16:153-68.
- 402 12. Eichinger S, Weltermann A, Philipp K, Hafner E, Kaider A, Kittl EM, Brenner B,
403 Mannhalter C, Lechner K, Kyrle PA. Prospective evaluation of hemostatic system
404 activation and thrombin potential in healthy pregnant women with and without factor V
405 Leiden. Thromb Haemost. 1999;82:1232-6.
- 406 13. Ku DH, Arkel YS, Paidas MP, Lockwood CJ. Circulating levels of inflammatory
407 cytokines (IL-1 beta and TNF-alpha), resistance to activated protein C, thrombin and
408 fibrin generation in uncomplicated pregnancies. Thromb Haemost 2003;90:1074-9.
- 409 14. Lockwood CJ. Pregnancy-associated changes in the hemostatic system. Clin Obstet
410 Gynecol 2006;49:836-43.
- 411 15. Erlich J, Parry GC, Fearn C, Muller M, Carmeliet P, Luther T, Mackman N. Tissue factor
412 is required for uterine hemostasis and maintenance of the placental labyrinth during
413 gestation. Proc Natl Acad Sci U S A 1999;96:8138-43.
- 414 16. Kuczyński J, Uszyński W, Zekanowska E, Soszka T, Uszyński M. Tissue factor (TF) and
415 tissue factor pathway inhibitor (TFPI) in the placenta and myometrium. Eur J Obstet
416 Gynecol Reprod Biol 2002;105:15-9.
- 417 17. Lockwood CJ, Krikun G, Schatz F. Decidual cell-expressed tissue factor maintains
418 hemostasis in human endometrium. Ann N Y Acad Sci 2001;943:77-88.
- 419 18. Lockwood CJ, Krikun G, Schatz F. The decidua regulates hemostasis in human
420 endometrium. Semin Reprod Endocrinol 1999;17:45-51.
- 421 19. Edstrom CS, Calhoun DA, Christensen RD. Expression of tissue factor pathway inhibitor
422 in human fetal and placental tissues. Early Hum Dev 2000;59:77-84.

- 423 20. Uszyński M, Zekanowska E, Uszyński W, Kuczyński J. Tissue factor (TF) and tissue
424 factor pathway inhibitor (TFPI) in amniotic fluid and blood plasma: implications for the
425 mechanism of amniotic fluid embolism. *Eur J Obstet Gynecol Reprod Biol* 2001;95:163-
426 6.
- 427 21. Lockwood CJ, Bach R, Guha A, Zhou XD, Miller WA, Nemerson Y. Amniotic fluid
428 contains tissue factor, a potent initiator of coagulation. *Am J Obstet Gynecol*
429 1991;165:1335-41.
- 430 22. Omsj  IH, Oian P, Malt  JM, Osterud B. Thromboplastin activity in amniotic fluid.
431 *Gynecol Obstet Invest* 1985;19:1-5.
- 432 23. Creter D. Amnioplastin: new reagent for coagulation tests. *Lancet* 1977;2:251.
- 433 24. Carmeliet P, Mackman N, Moons L, Luther T, Gressens P, Van Vlaenderen I, Demunck H,
434 Kasper M, Breier G, Evrard P, M ller M, Risau W, Edgington T, Collen D. Role of tissue
435 factor in embryonic blood vessel development. *Nature* 1996;383:73-5.
- 436 25. Holmes VA, Wallace JM. Haemostasis in normal pregnancy: a balancing act? *Biochem*
437 *Soc Trans* 2005;33:428-32.
- 438 26. Reber G, Amiral J, de Moerloose P. Modified antithrombin III levels during normal
439 pregnancy and relationship with prothrombin fragment F1 + 2 and thrombin-antithrombin
440 complexes. *Thromb Res* 1998;91:45-7.
- 441 27. Uszyński M. Generation of thrombin in blood plasma of non-pregnant and pregnant
442 women studied through concentration of thrombin-antithrombin III complexes. *Eur J*
443 *Obstet Gynecol Reprod Biol* 1997;75:127-31.
- 444 28. Reinthaller A, Mursch-Edlmayr G, Tatra G. Thrombin-antithrombin III complex levels in
445 normal pregnancy with hypertensive disorders and after delivery. *Br J Obstet Gynaecol*
446 1990;97:506-10.
- 447 29. Andersson T, Lorentzen B, Hogdahl H, Clausen T, Mowinkel MC, Abildgaard U.
448 Thrombin-inhibitor complexes in the blood during and after delivery. *Thromb Res*
449 1996;82:109-17.
- 450 30. Schjetlein R, Abdelnoor M, Haugen G, Husby H, Sandset PM, Wisl ff F. Hemostatic
451 variables as independent predictors for fetal growth retardation in preeclampsia. *Acta*
452 *Obstet Gynecol Scand* 1999;78:191-7.
- 453 31. Chaiworapongsa T, Yoshimatsu J, Espinoza J, Kim YM, Berman S, Edwin S, Yoon BH,
454 Romero R. Evidence of in vivo generation of thrombin in patients with small-for-
455 gestational-age fetuses and pre-eclampsia. *J Matern Fetal Neonatal Med* 2002;11:362-7.
- 456 32. Hayashi M, Numaguchi M, Ohkubo N, Yaoi Y. Blood macrophage colony-stimulating
457 factor and thrombin-antithrombin III complex concentrations in pregnancy and
458 preeclampsia. *Am J Med Sci* 1998;315:251-7.
- 459 33. Kobayashi T, Terao T. Preeclampsia as chronic disseminated intravascular coagulation.
460 Study of two parameters: thrombin-antithrombin III complex and D-dimers. *Gynecol*
461 *Obstet Invest* 1987;24:170-8.
- 462 34. Hayashi M1, Inoue T, Hoshimoto K, Negishi H, Ohkura T, Inaba N. Characterization of
463 five marker levels of the hemostatic system and endothelial status in normotensive
464 pregnancy and pre-eclampsia. *Eur J Haematol* 2002;69:297-302.
- 465 35. Hayashi M, Ohkura T. Elevated levels of serum macrophage colony-stimulating factor in
466 normotensive pregnancies complicated by intrauterine fetal growth restriction. *Exp*
467 *Hematol* 2002;30:388-93.
- 468 36. Ballard HS, Marcus AJ. Primary and secondary platelet aggregation in uraemia. *Scand J*
469 *Haematol* 1972;9:198-203.
- 470 37. Erez O, Romer R, Vaisbuch E, Chaiworapongsa T, Kusanovic JP, Mazaki-Tovi S, Gotsch
471 F, Gomez R, Maymon E, Pacora P, Edwin SS, Kim CJ, Than NG, Mittal P, Yeo L, Dong

- 472 Z, Yoon BH, Hassan SS, Mazor M. Changes in amniotic fluid concentration of thrombin-
473 antithrombin III complexes in patients with preterm labor: evidence of an increased
474 thrombin generation. *J Matern Fetal Neonatal Med* 2009;22:971-82.
- 475 38. Elovitz MA, Baron J, Phillippe M. The role of thrombin in preterm parturition. *Am J*
476 *Obstet Gynecol* 2001;185:1059-63.
- 477 39. Rosen T, Kuczynski E, O'Neill LM, Funai EF, Lockwood CJ. Plasma levels of thrombin-
478 antithrombin complexes predict preterm premature rupture of the fetal membranes. *J*
479 *Matern Fetal Med* 2001;10:297-300.
- 480 40. Kobayashi T, Tokunaga N, Sugimura M, Suzuki K, Kanayama N, Nishiguchi T, Terao T.
481 Coagulation/fibrinolysis disorder in patients with severe preeclampsia. *Semin Thromb*
482 *Hemost* 1999;25:451-4.
- 483 41. Kobayashi T, Sumimoto K, Tokunaga N, Sugimura M, Nishiguchi T, Kanayama N, Terao
484 T. Coagulation index to distinguish severe preeclampsia from normal pregnancy. *Semin*
485 *Thromb Hemost* 2002;28:495-500.
- 486 42. Lockwood CJ, Toti P, Arcuri F, Paidas M, Buchwalder L, Krikun G, Schatz F.
487 Mechanisms of abruption-induced premature rupture of the fetal membranes: thrombin-
488 enhanced interleukin-8 expression in term decidua. *Am J Pathol* 2005;167:1443-9.
- 489 43. Gómez R, Romero R, Nien JK, Medina L, Carstens M, Kim YM, Chaiworapongsa T,
490 Espinoza J, González R. Idiopathic vaginal bleeding during pregnancy as the only clinical
491 manifestation of intrauterine infection. *J Matern Fetal Neonatal Med* 2005;18:31-7.
- 492 44. Gervasi MT, Chaiworapongsa T, Naccasha N, Pacora P, Berman S, Maymon E, Kim JC,
493 Kim YM, Yoshimatsu J, Espinoza J, Romero R. Maternal intravascular inflammation in
494 preterm premature rupture of membranes. *J Matern Fetal Neonatal Med* 2002;11:171-5.
- 495 45. Østerud B, Bjørklid E. Sources of tissue factor. *Semin Thromb Hemost* 2006;32:11-23.
- 496 46. Rosen T, Schatz F, Kuczynski E, Lam H, Koo AB, Lockwood CJ. Thrombin-enhanced
497 matrix metalloproteinase-1 expression: a mechanism linking placental abruption with
498 premature rupture of the membranes. *J Matern Fetal Neonatal Med* 2002;11:11-7.
- 499 47. Mackenzie AP, Schatz F, Krikun G, Funai EF, Kadner S, Lockwood CJ. Mechanisms of
500 abruption-induced premature rupture of the fetal membranes: Thrombin enhanced
501 decidual matrix metalloproteinase-3 (stromelysin-1) expression. *Am J Obstet Gynecol*
502 2004;191:1996-2001.
- 503 48. Elovitz MA, Ascher-Landsberg J, Saunders T, Phillippe M. The mechanisms underlying
504 the stimulatory effects of thrombin on myometrial smooth muscle. *Am J Obstet Gynecol*
505 2000;183:674-81.
- 506 49. Elovitz MA, Saunders T, Ascher-Landsberg J, Phillippe M. Effects of thrombin on
507 myometrial contractions in vitro and in vivo. *Am J Obstet Gynecol* 2000;183:799-804.
- 508 50. Bilsel AS, Onaran N, Moini H, Emerk K. Long-term effect of 17beta-estradiol and
509 thrombin on tissue factor pathway inhibitor release from HUVEC. *Thromb Res*
510 2000;99:173-8.
- 511 51. VanWijk MJ, Boer K, Berckmans RJ, Meijers JC, van der Post JA, Sturk A, VanBavel E,
512 Nieuwland R. Enhanced coagulation activation in preeclampsia: the role of APC
513 resistance, microparticles and other plasma constituents. *Thromb Haemost* 2002;88:415-
514 20.
- 515 52. Erez O, Romero R, Vaisbuch E, Kusanovic JP, Mazaki-Tovi S, Chaiworapongsa T, Gotsch
516 F, Fareed J, Hoppensteadt D, Than NG, Yoon BH, Edwin S, Dong Z, Espinoza J, Mazor
517 M, Hassan SS. High tissue factor activity and low tissue factor pathway inhibitor
518 concentrations in patients with preterm labor. *J Matern Fetal Neonatal Med* 2010;23:23-
519 33.

- 520 53. Erez O, Romero R, Hoppensteadt D, Than NG, Fareed J, Mazaki-Tovi S, Espinoza J,
521 Chaiworapongsa T, Kim SS, Yoon BH, Hassan SS, Gotsch F, Friel L, Vaisbuch E,
522 Kusanovic JP. Tissue factor and its natural inhibitor in pre-eclampsia and SGA. *J Matern*
523 *Fetal Neonatal Med* 2008;21:855-69.
- 524 54. Erez O, Gotsch F, Mazaki-Tovi S, Vaisbuch E, Kusanovic JP, Kim CJ, Chaiworapongsa T,
525 Hoppensteadt D, Fareed J, Than NG, Nhan-Chang CL, Yeo L, Pacora P, Mazor M, Hassan
526 SS, Mittal P, Romero R. Evidence of maternal platelet activation, excessive thrombin
527 generation, and high amniotic fluid tissue factor immunoreactivity and functional activity
528 in patients with fetal death. *J Matern Fetal Neonatal Med* 2009;22:672-87.
- 529 55. Beller FK, Ebert C. The coagulation and fibrinolytic enzyme system in pregnancy and in
530 the puerperium. *Eur J Obstet Gynecol Reprod Biol* 1982;13:177-97.
- 531 56. Stirling Y, Woolf L, North WR, Seghatchian MJ, Meade TW. Haemostasis in normal
532 pregnancy. *Thromb Haemost* 1984;52:176-82.
- 533 57. Brenner B. Haemostatic changes in pregnancy. *Thromb Res* 2004;114:409-14.
- 534 58. Romero R, Gonçalves LF, Kusanovic JP, Devesa R, Espinoza J. Mechanisms of preterm
535 labor and preterm premature rupture of the membranes. In: Kurjak A, Chervenak F, eds.
536 *Textbook of Perinatal Medicine 2nd Edition*, 2006.
- 537 59. Lockwood CJ, Krikun G, Rahman M, Caze R, Buchwalder L, Schatz F. The role of
538 decidualization in regulating endometrial hemostasis during the menstrual cycle,
539 gestation, and in pathological states. *Semin Thromb Hemost* 2007;33:111-7.
- 540 60. Broze GJ Jr, Warren LA, Novotny WF, Higuchi DA, Girard JJ, Miletich JP. The
541 lipoprotein-associated coagulation inhibitor that inhibits the factor VII-tissue factor
542 complex also inhibits factor Xa: insight into its possible mechanism of action. *Blood*
543 1988;71:335-43.
- 544 61. Laskowski M, Kato I. Protein inhibitors of proteinases. *Annu Rev Biochem* 1980;49:593-
545 626.
- 546 62. Broze GJ, Girard TJ, Novotny WF. Regulation of coagulation by a multivalent Kunitz-
547 type inhibitor. *Biochemistry* 1990;29:7539-46.
- 548 63. Sarig G, Blumenfeld Z, Leiba R, Lanir N, Brenner B. Modulation of systemic hemostatic
549 parameters by enoxaparin during gestation in women with thrombophilia and pregnancy
550 loss. *Thromb Haemost* 2005;94:980-5.
- 551 64. Tay SP, Cheong SK, Boo NY. Circulating tissue factor, tissue factor pathway inhibitor and
552 D-dimer in umbilical cord blood of normal term neonates and adult plasma. *Blood Coagul*
553 *Fibrinolysis* 2003;14:125-9.
- 554 65. Hubé F, Reverdiau P, Iochmann S, Trassard S, Thibault G, Gruel Y. Demonstration of a
555 tissue factor pathway inhibitor 2 messenger RNA synthesis by pure villous
556 cytotrophoblast cells isolated from term human placentas. *Biol Reprod* 2003;68:1888-94.
- 557 66. Iino M, Foster DC, Kisiel W. Quantification and characterization of human endothelial
558 cell-derived tissue factor pathway inhibitor-2. *Arterioscler Thromb Vasc Biol* 1998;18:40-
559 6.
- 560 67. Sprecher CA, Kisiel W, Mathewes S, Foster DC. Molecular cloning, expression, and
561 partial characterization of a second human tissue-factor-pathway inhibitor. *Proc Natl Acad*
562 *Sci U S A* 1994;91:3353-7.
- 563 68. Udagawa K, Miyagi Y, Hirahara F, Miyagi E, Nagashima Y, Minaguchi H, Misugi K,
564 Yasumitsu H, Miyazaki K. Specific expression of PP5/TFPI2 mRNA by
565 syncytiotrophoblasts in human placenta as revealed by in situ hybridization. *Placenta*
566 1998;19:217-23.

- 567 69. Kamei S, Kazama Y, Kuijper JL, Foster DC, Kisiel W. Genomic structure and promoter
568 activity of the human tissue factor pathway inhibitor-2 gene. *Biochim Biophys Acta*
569 2001;1517:430-5.
- 570 70. Kisiel W, Sprecher CA, Foster DC. Evidence that a second human tissue factor pathway
571 inhibitor (TFPI-2) and human placental protein 5 are equivalent. *Blood* 1994;84:4384-5.
- 572 71. Bützow R, Virtanen I, Seppälä M, Närvänen O, Stenman UH, Ristimäki A, Bohn H.
573 Monoclonal antibodies reacting with placental protein 5: use in radioimmunoassay,
574 Western blot analysis, and immunohistochemistry. *J Lab Clin Med* 1988;111:249-56.
- 575 72. Chand HS, Foster DC, Kisiel W. Structure, function and biology of tissue factor pathway
576 inhibitor-2. *Thromb Haemost* 2005;94:1122-30.
- 577 73. Seppälä M, Wahlström T, Bohn H. Circulating levels and tissue localization of placental
578 protein five (PP5) in pregnancy and trophoblastic disease: absence of PP5 expression in
579 the malignant trophoblast. *Int J Cancer* 1979;24:6-10.
- 580 74. Obiewke BC, Chard T. Placental protein 5: circulating levels in twin pregnancy and some
581 observations on the analysis of biochemical data from multiple pregnancy. *Eur J Obstet*
582 *Gynecol Reprod Biol* 1981;12:135-41.
- 583 75. Shimura M, Wada H, Wakita Y, Nakase T, Hiyoyama K, Nagaya S, Mori Y, Shiku H.
584 Plasma tissue factor and tissue factor pathway inhibitor levels in patients with
585 disseminated intravascular coagulation. *Am J Hematol* 1997;55:169-74.
- 586 76. Abdel Gader AM, Al-Mishari AA, Awadalla SA, Buyuomi NM, Khashoggi T, Al-Hakeem
587 M. Total and free tissue factor pathway inhibitor in pregnancy hypertension. *Int J*
588 *Gynaecol Obstet* 2006;95:248-53.
- 589 77. Erez O, Espinoza J, Chaiworapongsa T, Gotsch F, Kusanovic JP, Than NG, Mazaki-Tovi
590 S, Vaisbuch E, Papp Z, Yoon BH, Han YM, Hoppensteadt D, Fareed J, Hassan SS,
591 Romero R. A link between a hemostatic disorder and preterm PROM: a role for tissue
592 factor and tissue factor pathway inhibitor. *J Matern Fetal Neonatal Med* 2008;21:732-44.
- 593 78. Xiong Y, Zhou Q, Jiang F, Zhou S, Lou Y, Guo Q, Liang W, Kong D, Ma D, Li X.
594 Changes of plasma and placental tissue factor pathway inhibitor-2 in women with
595 preeclampsia and normal pregnancy. *Thromb Res* 2010;125:e317-22.
- 596 79. Aharon A, Lanir N, Drugan A, Brenner B. Placental TFPI is decreased in gestational
597 vascular complications and can be restored by maternal enoxaparin treatment. *J Thromb*
598 *Haemost* 2005;3:2355-7.
- 599 80. Castoldi E, Hackeng TM. Regulation of coagulation by protein S. *Curr Opin Hematol*
600 2008;15:529-36.
- 601 81. Hackeng TM, Seré KM, Tans G, Rosing J. Protein S stimulates inhibition of the tissue
602 factor pathway by tissue factor pathway inhibitor. *Proc Natl Acad Sci U S A*
603 2006;103:3106-11.
- 604 82. Ndonwi M, Broze G. Protein S enhances the tissue factor pathway inhibitor inhibition of
605 factor Xa but not its inhibition of factor VIIa-tissue factor. *J Thromb Haemost*
606 2008;6:1044-6.
- 607 83. Faught W, Garner P, Jones G, Ivey B. Changes in protein C and protein S levels in normal
608 pregnancy. *Am J Obstet Gynecol* 1995;172:147-50.
- 609 84. Dahlbäck B, Carlsson M, Svensson PJ. Familial thrombophilia due to a previously
610 unrecognized mechanism characterized by poor anticoagulant response to activated
611 protein C: prediction of a cofactor to activated protein C. *Proc Natl Acad Sci U S A*
612 1993;90:1004-8.
- 613 85. Rodger MA, Paidas M, McLintock C, Middeldorp S, Kahn S, Martinelli I, Hague W,
614 Rosene Montella K, Greer I. Inherited thrombophilia and pregnancy complications
615 revisited. *Obstet Gynecol* 2008;112:320-4.

- 616 86. Yalinkaya A, Erdemoglu M, Akdeniz N, Kale A, Kale E. The relationship between
617 thrombophilic mutations and preeclampsia: a prospective case-control study. *Ann Saudi*
618 *Med* 2006;26:105-9.
- 619 87. Preston FE, Rosendaal FR, Walker ID, Briët E, Berntorp E, Conard J, Fontcuberta J,
620 Makris M, Mariani G, Noteboom W, Pabinger I, Legnani C, Scharrer I, Schulman S, van
621 der Meer FJ. Increased fetal loss in women with heritable thrombophilia. *Lancet*
622 1996;348:913-6.
- 623 88. Kupfermanc MJ, Many A, Bar-Am A, Lessing JB, Ascher-Landsberg J. Mid-trimester
624 severe intrauterine growth restriction is associated with a high prevalence of
625 thrombophilia. *BJOG* 2002;109:1373-6.
- 626 89. Han X, Fiehler R, Broze GJ. Isolation of a protein Z-dependent plasma protease inhibitor.
627 *Proc Natl Acad Sci U S A* 1998;95:9250-5.
- 628 90. Han X, Huang ZF, Fiehler R, Broze GJ Jr. The protein Z-dependent protease inhibitor is a
629 serpin. *Biochemistry* 1999;38:11073-8.
- 630 91. Han X, Fiehler R, Broze GJ. Characterization of the protein Z-dependent protease
631 inhibitor. *Blood* 2000;96:3049-55.
- 632 92. Yin ZF, Huang ZF, Cui J, Fiehler R, Lasky N, Ginsburg D, Broze GJ Jr. Prothrombotic
633 phenotype of protein Z deficiency. *Proc Natl Acad Sci U S A* 2000;97:6734-8.
- 634 93. Taylor FB, Chang AC, Peer G, Li A, Ezban M, Hedner U. Active site inhibited factor VIIa
635 (DEGR VIIa) attenuates the coagulant and interleukin-6 and -8, but not tumor necrosis
636 factor, responses of the baboon to LD100 *Escherichia coli*. *Blood* 1998;91:1609-15.
- 637 94. Kusanovic JP, Espinoza J, Romero R, Hoppensteadt D, Nien JK, Kim CJ, Erez O, Soto E,
638 Fareed J, Edwin S, Chaiwerapongsa T, Than NG, Yoon BH, Gomez R, Papp Z, Hassan
639 SS. Plasma protein Z concentrations in pregnant women with idiopathic intrauterine
640 bleeding and in women with spontaneous preterm labor. *J Matern Fetal Neonatal Med*
641 2007;20:453-63.
- 642 95. Bretelle F, Arnoux D, Shojai R, D'Ercole C, Sampol J, Dignat F, Camoin-Jau L. Protein Z
643 in patients with pregnancy complications. *Am J Obstet Gynecol* 2005;193:1698-702.
- 644 96. Nien JK, Romero R, Hoppensteadt D, Erez O, Espinoza J, Soto E, Kusanovic JP, Gotsch
645 F, Kim CJ, Mittal P, Fareed J, Santolaya J, Chaiworapongsa T, Edwin S, Pineles B,
646 Hassan S. Pyelonephritis during pregnancy: a cause for an acquired deficiency of protein
647 Z. *J Matern Fetal Neonatal Med* 2008;21:629-37.
- 648 97. Sood R, Kalloway S, Mast AE, Hillard CJ, Weiler H. Fetomaternal cross talk in the
649 placental vascular bed: control of coagulation by trophoblast cells. *Blood* 2006;107:3173-
650 80.
- 651 98. Sood R, Sholl L, Isermann B, Zogg M, Coughlin SR, Weiler H. Maternal Par4 and
652 platelets contribute to defective placenta formation in mouse embryos lacking
653 thrombomodulin. *Blood* 2008;112:585-91.
- 654 99. Lanir N, Aharon A, Brenner B. Procoagulant and anticoagulant mechanisms in human
655 placenta. *Semin Thromb Hemost* 2003;29:175-84.
- 656 100. Isermann B, Sood R, Pawlinski R, Zogg M, Kalloway S, Degen JL, Mackman N, Weiler
657 H. The thrombomodulin-protein C system is essential for the maintenance of pregnancy.
658 *Nat Med* 2003;9:331-7.
- 659 101. Aharon A, Brenner B, Katz T, Miyagi Y, Lanir N. Tissue factor and tissue factor pathway
660 inhibitor levels in trophoblast cells: implications for placental hemostasis. *Thromb*
661 *Haemost* 2004;92:776-86.
- 662 102. Butenas S, Orfeo T, Brummel-Ziedins KE, Mann KG. Tissue factor in thrombosis and
663 hemorrhage. *Surgery* 2007;142:S2-14.

103. Midderdorp S. Thrombophilia and pregnancy complications: cause or association? *J Thromb Haemost* 2007;5 Suppl 1:276-82.
104. Martinelli I, Legnani C, Bucciarelli P, Grandone E, De Stefano V, Mannucci PM. Risk of pregnancy-related venous thrombosis in carriers of severe inherited thrombophilia. *Thromb Haemost* 2001;86:800-3.
105. Kovo M, Schreiber L, Bar J. Placental vascular pathology as a mechanism of disease in pregnancy complications. *Thromb Res* 2013;131 Suppl 1:S18-21.
106. Redline RW, Heller D, Keating S, Kingdom J. Placental diagnostic criteria and clinical correlation--a workshop report. *Placenta* 2005;26 Suppl A:S114-7.
107. Bar J, Schreiber L, Ben-Haroush A, Ahmed H, Golan A, Kovo M. The placental vascular component in early and late intrauterine fetal death. *Thromb Res* 2012;130:901-5.
108. Cromi A, Ghezzi F, Dürig P, Di Naro E, Raio L. Sonographic umbilical cord morphometry and coiling patterns in twin-twin transfusion syndrome. *Prenat Diagn* 2005;25:851-5.
109. Salafia CM, Pezzullo JC, Ghidini A, Lopèz-Zeno JA, Whittington SS. Clinical correlations of patterns of placental pathology in preterm pre-eclampsia. *Placenta* 1998;19:67-72.
110. Roberts DJ, Post MD. The placenta in pre-eclampsia and intrauterine growth restriction. *J Clin Pathol* 2008;61:1254-60.
111. Mayhew TM, Ohadike C, Baker PN, Crocker IP, Mitchell C, Ong SS. Stereological investigation of placental morphology in pregnancies complicated by pre-eclampsia with and without intrauterine growth restriction. *Placenta* 2003;24:219-26.
112. Ogge G, Chaiworapongsa T, Romero R, Hussein Y, Kusanovic JP, Yeo L, Kim CJ, Hassan SS. Placental lesions associated with maternal underperfusion are more frequent in early-onset than in late-onset preeclampsia. *J Perinat Med* 2011;39:641-52.
113. Kovo M, Schreiber L, Ben-Haroush A, Wand S, Golan A, Bar J. Placental vascular lesion differences in pregnancy-induced hypertension and normotensive fetal growth restriction. *Am J Obstet Gynecol* 2010;202:561.e1-5.
114. Teng Y, Jiang R, Lin Q, Ding C, Ye Z. The relationship between plasma and placental tissue factor, and tissue factor pathway inhibitors in severe pre-eclampsia patients. *Thromb Res* 2010;126:e41-5.
115. Redline RW. Placental pathology: a systematic approach with clinical correlations. *Placenta* 2008;29 Suppl A:S86-91.
116. Salafia CM, Minior VK, Pezzullo JC, Popek EJ, Rosenkrantz TS, Vintzileos AM. Intrauterine growth restriction in infants of less than thirty-two weeks' gestation: associated placental pathologic features. *Am J Obstet Gynecol* 1995;173:1049-57.
117. Kovo M, Schreiber L, Ben-Haroush A, Gold E, Golan A, Bar J. The placental component in early-onset and late-onset preeclampsia in relation to fetal growth restriction. *Prenat Diagn* 2012;32:632-7.
118. Romero R, Espinoza J, Kusanovic JP, Gotsch F, Hassan S, Erez O, Chaiworapongsa T, Mazar M. The preterm parturition syndrome. *BJOG* 2006;113 Suppl 3:17-42.
119. Arias F, Rodriguez L, Rayne SC, Kraus FT. Maternal placental vasculopathy and infection: two distinct subgroups among patients with preterm labor and preterm ruptured membranes. *Am J Obstet Gynecol* 1993;168:585-91.
120. Nath CA, Ananth CV, Smulian JC, Shen-Schwarz S, Kaminsky L; New Jersey-Placental Abruptio Study Investigators. Histologic evidence of inflammation and risk of placental abruptio. *Am J Obstet Gynecol* 2007;197:319.e1-6.

- 711 121. Kim YM, Chaiworapongsa T, Gomez R, Bujold E, Yoon BH, Rotmensch S, Thaler HT,
712 Romero R. Failure of physiologic transformation of the spiral arteries in the placental bed
713 in preterm premature rupture of membranes. *Am J Obstet Gynecol* 2002;187:1137-42.
- 714 122. Salafia CM, Vogel CA, Vintzileos AM, Bantham KF, Pezzullo J, Silberman L. Placental
715 pathologic findings in preterm birth. *Am J Obstet Gynecol* 1991;165:934-8.
- 716 123. Stephenson CD, Lockwood CJ, Ma Y, Guller S. Thrombin-dependent regulation of matrix
717 metalloproteinase (MMP)-9 levels in human fetal membranes. *J Matern Fetal Neonatal*
718 *Med* 2005;18:17-22.

719 Figure 1. Thrombin–antithrombin III (TAT) levels in control patients, patients with preterm labor
 720 who delivered between 21 and 7 days, and patients with preterm labor who delivered within 7
 721 days. Open diamonds, Mean levels; black error bars, SD. *P <.05, Student-Newman-Keuls
 722 method (from Elovitz MA, Baron J, Phillippe M. The role of thrombin in preterm parturition. Am
 723 J Obstet Gynecol 2001 Nov;185(5):1059-1063. With permission).

724 Figure 2. (a) Comparison of median maternal plasma TF concentration between patients with
 725 normal pregnancy (n=79), pre-eclampsia (n=133), and women who delivered an SGA neonate
 726 (n=61). (b) Comparison of median maternal plasma TFPI concentration between patients with
 727 normal pregnancy (n=86), pre-eclampsia (n=133), and women who delivered an SGA neonate
 728 (n=61). (c) Comparison of maternal plasma TFPI/TF ratio between women with normal
 729 pregnancy (n=79), pre-eclampsia (n=133), and women who delivered an SGA neonate (n=61).
 730 (From [Erez O](#), [Romero R](#), [Hoppensteadt D](#), [Than NG](#), [Fareed J](#), [Mazaki-Tovi S](#), [Espinoza](#)
 731 [J](#), [Chaiworapongsa T](#), [Kim SS](#), [Yoon BH](#), [Hassan SS](#), [Gotsch F](#), [Friel L](#), [Vaisbuch E](#), [Kusanovic](#)
 732 [JP](#). Tissue factor and its natural inhibitor in pre-eclampsia and SGA. [J Matern Fetal Neonatal](#)
 733 [Med](#). 2008 Dec;21(12):855-69. With permission).

734 Figure 3. Factor X activation and protein Z/protein Z-dependent protease inhibitor (ZPI)
 735 inhibition of activated factor X. (a) Then formation of the complex of tissue factor (TF) and
 736 factor VIIa (FVIIa) at the site of injury and activation of extrinsic coagulation cascade. (b)
 737 Activation of circulating factor X by the TFpFVIIa complex in the presence of exposed
 738 phospholipids and Ca²⁺. (c) Inhibition of factor Xa (FXa) by the protein Z/ZPI complex by
 739 binding to its active site. Modified from Broze JG, Lancet 2001;357:900–901.

740 Figure 4. Histologic features of maternal vessel and implantation site reaction patterns: a. Acute
 741 atherosclerosis of maternal arterioles in the placental membranes: a cluster of decidual arterioles shows
 742 varying stages of fibrinoid necrosis. The vessel at the upper right shows full histologic expression
 743 with dark homogenous fibrinoid replacement of the vessel wall accompanied by occasional
 744 foamy macrophages ([original magnification is indicated for all panels] X 20). b. Mural

745 hypertrophy of decidual arterioles in the placental membranes: a cluster of arterioles shows
 746 medial hypertrophy with the vessel wall occupying greater than one third of total vessel diameter
 747 (X 10). c. Muscularized basal plate arteries with accompanying implantation site abnormalities:
 748 maternal spiral arteries in the basal plate lack normal trophoblast remodeling and retain their pre-
 749 pregnancy muscular media. Clusters of immature intermediate trophoblast and increased
 750 placental giant cells are seen above and below the muscular arteries, respectively (X 10). d. Acute
 751 atherosclerosis of muscularized basal plate arteries with accompanying implantation site abnormalities:
 752 three cross sections of a basal plate artery are seen. The two on the left show persistence of the
 753 muscular media while the one on the right has undergone fibrinoid necrosis of the media with
 754 foamy macrophages (acute atherosclerosis). Clusters of immature intermediate trophoblast are also
 755 seen overlying the arteries (X 4). e. Immature intermediate trophoblast: clusters of abnormally
 756 small intermediate trophoblast with focal vacuolation are surrounded by an excessive amount of
 757 basal plate fibrin. Increased placental site giant cells are also seen at the lower margin (X 10). f.
 758 Increased placental site giant cells: numerous multinucleate placental site giant cells, not usually
 759 seen in the delivered placenta, are scattered in loose decidual tissue which is devoid of normal
 760 intermediate trophoblast and fibrinoid (X 10).
 761 From: Redline RW, et al. Maternal vascular underperfusion: nosology and reproducibility of
 762 placental reaction patterns. *Pediatr Dev Pathol.* 2004;7(3):237-49. With permission.

763 Figure 5. Histologic features of villous and intervillous lesions; a. Increased syncytial knots:
 764 aggregates of syncytiotrophoblast nuclei cluster at one or more poles of distal villi in the vicinity
 765 of larger stem villi (arrowhead) at the periphery of the lobule ([original magnification is indicated
 766 for all panels] X 10). b. Villous agglutination: clusters of degenerating distal villi are adherent to
 767 one another and focally enmeshed in fibrin (X 4). c. Distal villous hypoplasia: a long, thin, non-
 768 branching stem villus is surrounded by a markedly reduced number of small hypoplastic distal
 769 villi (X 10). d. Increased intervillous fibrin: stem villi are surrounded by a mantle of fibrin-type
 770 fibrinoid that does not extend to distal villi at the center of the lobule (X 2). e. Nodular
 771 intervillous (and intravillous) fibrin: small aggregates of intervillous fibrin adhere to, and are
 772 focally reepithelialized by, distal villous trophoblast (X 20). f. Increased intervillous fibrin with

773 intermediate trophoblast (X-cells): stem and distal villi are enmeshed in a matrix of fibrin and
 774 fibrinoid containing prominent intermediate trophoblast (arrowhead) (X 10).
 775 From: Redline RW, et al. Maternal vascular underperfusion: nosology and reproducibility of
 776 placental reaction patterns. *Pediatr Dev Pathol.* 2004;7(3):237-49. With permission.

777 Figure 6. Amniotic fluid tissue factor concentration among women with normal pregnancies
 778 (median 3710.4 pg/ml, range 2198.8–6268) and patients with a fetal demise (median 8535.4
 779 pg/ml, range 2208.2–125,990.0); (b) Amniotic fluid tissue factor activity among women with
 780 normal pregnancies (median 28.4 pM, range 10.2–84.9) and patients with a fetal demise (median
 781 81.6 pM, range 7.2–1603.4). From EREZ O, GOTSCH F, MAZAKI-TOVI S, et al. Evidence of
 782 maternal platelet activation, excessive thrombin generation, and high amniotic fluid tissue factor
 783 immunoreactivity and functional activity in patients with fetal death. *J Matern Fetal Neonatal*
 784 *Med* 2009;22:672-87, with permission.

785 Figure 7. Maternal plasma TAT III concentration in women with preterm labor (PTL) and those
 786 with a Normal pregnancy (From Chaiworapongsa T, Espinoza J, Yoshimatsu J, Kim YM, Bujold
 787 E, Edwin S, et al. Activation of coagulation system in preterm labor and preterm premature
 788 rupture of membranes. *J Matern Fetal Neonatal Med* 2002 11(6):368-373, with permission).

Table 1 (on next page)

Table 1

Table 1. Concentration and activity in maternal plasma of coagulating and anticoagulating factors and their relation with thrombin generation in the great obstetrical syndromes.

Table 1. Concentration and activity in maternal plasma of coagulating and anticoagulating factors and their relation with thrombin generation in the great obstetrical syndromes.

	TF concentration and/or activity	TFPI concentration and/or activity	TAT III complexes concentration	Protein Z concentration	Thrombin generation	References
Premature rupture of membranes	Activity ↑ Concentration ↑	Concentration ↓	Concentration ↑	Concentration ↓	↑	1-5
Preterm labor	Activity ↑ Concentration =	Activity = Concentration ↓	Concentration ↑	Concentration ↓	↑	1-5
Fetal demise	Activity = Concentration =	Activity = Concentration ↓	Concentration ↑	Concentration ↓	↑	1-5
Preeclampsia	Activity ↑ Concentration ↑	Concentration ↑	Concentration ↑	Concentration ↓	↑	1-5
Intrauterine growth retardation	Concentration ↓	Concentration =	Concentration ↑	Concentration ↓	↑	1-5

1. Erez O, Romero R, Vaisbuch E, Kusanovic JP, Mazaki-Tovi S, Chaiworapongsa T, Gotsch F, Fareed J, Hoppensteadt D, Than NG, Yoon BH, Edwin S, Dong Z, Espinoza J, Mazor M, Hassan SS. High tissue factor activity and low tissue factor pathway inhibitor concentrations in patients with preterm labor. *J Matern Fetal Neonatal Med.* 2010 Jan;23(1):23-33
2. Erez O, Gotsch F, Mazaki-Tovi S, Vaisbuch E, Kusanovic JP, Kim CJ, Chaiworapongsa T, Hoppensteadt D, Fareed J, Than NG, Nhan-Chang CL, Yeo L, Pacora P, Mazor M, Hassan SS, Mittal P, Romero R. Evidence of maternal platelet activation, excessive thrombin generation, and high amniotic fluid tissue factor immunoreactivity and functional activity in patients with fetal death. *J Matern Fetal Neonatal Med.* 2009 Aug;22(8):672-87
3. Kusanovic JP, Espinoza J, Romero R, Hoppensteadt D, Nien JK, Kim CJ, et al. Plasma protein Z concentrations in pregnant women with idiopathic intrauterine bleeding and in women with spontaneous preterm labor. *J Matern Fetal Neonatal Med* 2007 Jun;20(6):453-463.
4. Gris JC, Quere I, Dechaud H, Mercier E, Pincon C, Hoffet M, Vasse M, Mares P. High frequency of protein Z deficiency in patients with unexplained early fetal loss. *Blood* 2002;99:2606–2608
5. Paidas MJ, Ku DH, Lee MJ, Manish S, Thurston A, Lockwood CJ, Arkel YS. Protein Z, protein S levels are lower in patients with thrombophilia and subsequent pregnancy complications. *J Thromb Haemost* 2005;3:497–501.

Figure 1(on next page)

Figure 1

Thrombin-antithrombin III (TAT) levels in control patients, patients with preterm labor who delivered between 21 and 7 days, and patients with preterm labor who delivered within 7 days. Open diamonds, Mean levels; black error bars, SD. *P <.05, Student-Newman-Keuls method (from Elovitz MA, Baron J, Phillippe M. The role of thrombin in preterm parturition. Am J Obstet Gynecol 2001 Nov;185(5):1059-1063. With permission).






Figure 2 (on next page)

Figure 2


Figure 2. (a) Comparison of median maternal plasma TF concentration between patients with normal pregnancy (n=79), pre-eclampsia (n=133), and women who delivered an SGA neonate (n=61). (b) Comparison of median maternal plasma TFPI concentration between patients with normal pregnancy (n=86), pre-eclampsia (n=133), and women who delivered an SGA neonate (n=61). (c) Comparison of maternal plasma TFPI/TF ratio between women with normal pregnancy (n=79), pre-eclampsia (n=133), and women who delivered an SGA neonate (n=61). (From Erez O , Romero R , Hoppensteadt D , Than NG , Fareed J , Mazaki-Tovi S , Espinoza J , Chaiworapongsa T , Kim SS , Yoon BH , Hassan SS , Gotsch F , Friel L , Vaisbuch E , Kusanovic JP . Tissue factor and its natural inhibitor in pre-eclampsia and SGA. J Matern Fetal Neonatal Med. 2008 Dec;21(12):855-69. With permission).



(a)



(b)



(c)

Figure 3 (on next page)

Figure 3

Factor X activation and protein Z/protein Z-dependent protease inhibitor (ZPI) inhibition of activated factor X. (a) Then formation of the complex of tissue factor (TF) and factor VIIa (FVIIa) at the site of injury and activation of extrinsic coagulation cascade. (b) Activation of circulating factor X by the TFpFVIIa complex in the presence of exposed phospholipids and Ca^{2+} . (c) Inhibition of factor Xa (FXa) by the protein Z/ZPI complex by binding to its active site. Modified from Broze JG, Lancet 2001;357:900-901.

Figure 4 (on next page)

Figure 4

Histologic features of maternal vessel and implantation site reaction patterns: a. Acute atherosclerosis of maternal arterioles in the placental membranes: a cluster of decidual arterioles shows varying stages of fibrinoid necrosis. The vessel at the upper right shows full histologic expression with dark homogenous fibrinoid replacement of the vessel wall accompanied by occasional foamy macrophages ([original magnification is indicated for all panels] X .(20 b. Mural hypertrophy of decidual arterioles in the placental membranes: a cluster of arterioles shows medial hypertrophy with the vessel wall occupying greater than one third of total vessel diameter (X 10). c. Muscularized basal plate arteries with accompanying implantation site abnormalities: maternal spiral arteries in the basal plate lack normal trophoblast remodeling and retain their pre-pregnancy muscular media. Clusters of immature intermediate trophoblast and increased placental giant cells are seen above and below the muscular arteries, respectively (X 10). d. Acute atherosclerosis of muscularized basal plate arteries with accompanying implantation site abnormalities: three cross sections of a basal plate artery are seen. The two on the left show persistence of the muscular media while the one on the right has undergone fibrinoid necrosis of the media with foamy macrophages (acute atherosclerosis). Clusters of immature intermediate trophoblast are also seen overlying the arteries (X 4). e. Immature intermediate trophoblast: clusters of abnormally small intermediate trophoblast with focal vacuolation are surrounded by an excessive amount of basal plate fibrin. Increased placental site giant cells are also seen at the lower margin (X .(10 f. Increased placental site giant cells: numerous multinucleate placental site giant cells, not usually seen in the delivered placenta, are scattered in loose decidual tissue which is devoid of normal intermediate trophoblast and fibrinoid (X .(10 From: Redline RW, et al. Maternal vascular underperfusion: nosology and reproducibility of placental reaction patterns. *Pediatr Dev Pathol.* 2004;7(3):237-49. With permission.




Figure 5(on next page)

Figure 5

Histologic features of villous and intervillous lesions; a. Increased syncytial knots: aggregates of syncytiotrophoblast nuclei cluster at one or more poles of distal villi in the vicinity of larger stem villi (arrowhead) at the periphery of the lobule ([original magnification is indicated for all panels] X 10). b. Villous agglutination: clusters of degenerating distal villi are adherent to one another and focally enmeshed in fibrin (X 4). c. Distal villous hypoplasia: a long, thin, non-branching stem villus is surrounded by a markedly reduced number of small hypoplastic distal villi (X 10). d. Increased intervillous fibrin: stem villi are surrounded by a mantle of fibrin-type fibrinoid that does not extend to distal villi at the center of the lobule (X 2). e. Nodular intervillous (and intravillous) fibrin: small aggregates of intervillous fibrin adhere to, and are focally reepithelialized by, distal villous trophoblast (X 20). f. Increased intervillous fibrin with intermediate trophoblast (X-cells): stem and distal villi are enmeshed in a matrix of fibrin and fibrinoid containing prominent intermediate trophoblast (arrowhead) (X 10). From: Redline RW, et al. Maternal vascular underperfusion: nosology and reproducibility of placental reaction patterns. *Pediatr Dev Pathol*. 2004;7(3):237-49. With permission.




Figure 6 (on next page)

Figure 6

Amniotic fluid tissue factor concentration among women with normal pregnancies (median 3710.4 pg/ml, range 2198.8–6268) and patients with a fetal demise (median 8535.4 pg/ml, range 2208.2–125,990.0); (b) Amniotic fluid tissue factor activity among women with normal pregnancies (median 28.4 pM, range 10.2–84.9) and patients with a fetal demise (median 81.6 pM, range 7.2–1603.4). From Erez O, Gotsch F, Mazaki-Tovi S, et al. Evidence of maternal platelet activation, excessive thrombin generation, and high amniotic fluid tissue factor immunoreactivity and functional activity in patients with fetal death. *J Matern Fetal Neonatal Med* 2009;22:672-87, with permission.





Figure 7 (on next page)

Figure 7

Maternal plasma TAT III concentration in women with preterm labor (PTL) and those with a Normal pregnancy (From Chaiworapongsa T, Espinoza J, Yoshimatsu J, Kim YM, Bujold E, Edwin S, et al. Activation of coagulation system in preterm labor and preterm premature rupture of membranes. J Matern Fetal Neonatal Med 2002 11(6):368-373, with permission).

