***Loancorhynchus catrillancai* gen. et sp. nov., a new swordfish (Xiphiidae, Blochiidae) from the Middle Eocene of central Chile**

Rodrigo A Otero Corresp. 1, 2

¹ Laboratorio de Ontogenia y Filogenia, Departamento de Biología, Facultad de Ciencias, Universidad de Chile, Santiago, Chile

² Consultora Paleosuchus Ltda., Santiago, Chile

Corresponding Author: Rodrigo A Otero

Email address: rotero@paleosuchus.cl

This contribution describes the skull remains of a swordfish (Perciformes, Xiphiidae), recovered from Middle Eocene beds of central Chile. Comparison with known fossil and extant xiphioids reveals derived traits only present in the Neogene swordfish *Xiphias gladius* (Xiphiidae, Xiphiinae), these being a elongated rostrum composed of premaxillaries and possible prenasals, a dorsoventrally high and slender hyomandibular-metapterygoid complex, and a rounded, convex operculum. Also, strong ridges and sulci are present in the dorsal part of the rostrum, a feature only present in the billfish genera *Aglyptorhynchus* and *Xiphiorhynchus*, and in the swordfish genus *Blochius*. In addition, the specimen also has villiform teeth in the rostrum and lower jaw, a feature previously documented only in the Paleocene genus *Hemingwaya*. Such a unique combination of characters in the specimen allows classifying it as a new genus and species, *Loancorhynchus catrillancai*. Phylogenetic analyses obtained *Xiphiorhynchus* outside the Xiphiidae, suggesting instead narrow relationships to the Paleocene-Eocene genera *Hemingwaya* + *Palaeorhynchus* + *Homorhynchus*. *Loancorhynchus* is obtained as an intermediate form between *Xiphias* and *Blochius*. The specimen represents the first Paleogene swordfish described in the southeastern Pacific.

***Loancorhynchus catrillancai* gen. et sp. nov., a new swordfish (Xiphiidae, Blochiidae) from the Middle Eocene of central Chile**

Rodrigo A. Otero^{1,2}

¹Laboratorio de Ontogenia y Filogenia, Departamento de Biología, Facultad de Ciencias, Universidad de Chile., Santiago, Chile

²Consultora Paleosuchus Ltda., Santiago, Chile

Corresponding author:

Rodrigo Otero

otero2112@gmail.com

-

ABSTRACT

This contribution describes the skull remains of a swordfish (Perciformes, Xiphioidae), recovered from Middle Eocene beds of central Chile. Comparison with known fossil and extant xiphioids reveals derived traits only present in the Neogene swordfish *Xiphias gladius* (Xiphiidae, Xiphiinae), these being a elongated rostrum composed of premaxillaries and possible prenasals, a dorsoventrally high and slender hyomandibular-metapterygoid complex, and a rounded, convex operculum. Also, strong ridges and sulci are present in the dorsal part of the rostrum, a feature only present in the billfish genera *Aglyptorhynchus* and *Xiphiorhynchus*, and in the swordfish genus *Blochius*. In addition, the specimen also has villiform teeth in the rostrum and lower jaw, a feature previously documented only in the Paleocene genus *Hemingwaya*. Such a unique combination of characters in the specimen allows classifying it as a new genus and species, *Loancorhynchus catrillancai*. Phylogenetic analyses obtained *Xiphiorhynchus* outside the Xiphiidae, suggesting instead narrow relationships to the Paleocene-Eocene genera *Hemingwaya* + *Palaeorhynchus* + *Homorhynchus*. *Loancorhynchus* is obtained as an intermediate form between *Xiphias* and *Blochius*. The specimen represents the first Paleogene swordfish described in the southeastern Pacific.

INTRODUCTION

The Xiphioidae (Osteichthyes, Perciformes) is a group of fishes considered to be monophyletic, being characterized by elongated premaxillaries forming a rigid rostrum, and by having villiform teeth (Fierstine, 2006), although the latter can be secondarily lost during the ontogeny of some taxa (e.g., *Xiphias gladius*). To date, five family-level groups are known: Hemingwayidae, Palaeorhynchidae, Blochiidae, Xiphiidae and Istiophoridae (Fierstine, 2006; Gottfried et al.,

2013). The two latter clades, Xiphiidae and Istiophoridae, are the only lineages with extant representatives.

Late Cretaceous–Paleogene records of xiphioids are scarce and restricted to the Weddellian Province. *Zinsmeister* (1979) defined it as a large province including southern South America, Antarctica, New Zealand and part of Australia, isolated between the Late Cretaceous–Paleogene by the geography of Gondwana landmass and prevailing oceanic circulation, resulting in the development of highly endemic marine faunas (*Zinsmeister*, 1982). The first mention of a Paleogene Xiphioidei from the Weddellian Province concerns a large, isolated vertebra from lower Eocene levels of La Meseta Formation, in Marambio (=Seymour) Island, Antarctica, referred to cf. *Xiphiorhynchus* (*Cione et al.*, 2001). *Friedman and Otero* (2009) added to this record the preliminary description of a fragmentary rostrum and dentaries referred to as Xiphiorhynchine indet., which was recovered from Paleogene levels of central Chile. This material is part of the same specimen described here. After that, *Gottfried et al.* (2013) described a new species of palaeorhynchid, *Aglyptorynchus hakataramea*, from late Oligocene levels exposed in South Canterbury, South Island, New Zealand. A fourth specimen (GS 13924; GNS Science, Lower Hutt, New Zealand), still awaiting description, belongs to an articulated skull embedded in a concretion, recovered from Eocene levels of Hampden Beach, Otago, South Island, New Zealand (*Campbell et al.*, 2013). Locally, Neogene records of xiphiids and istiophorids are well-known in the Miocene-Pliocene of Chile (*Long, 1993; Walsh, 2001; Pyenson et al.*, 2014). A mention of cf. *Xiphiorhynchus* from the Miocene-Pliocene of Peru was provided by *Muizon and DeVries* (1985), but the material was not figured neither a repository was provided.

This contribution describes the only xiphioid recovered to date from Paleogene beds of the southeastern Pacific. After further preparation, new skull elements are added to its first preliminary description by *Friedman and Otero (2009)*. In addition, the stratigraphic provenance of the specimen is clarified here. As a result, a new genus and species are here proposed. The fossil is found to be Middle Eocene in age, representing the first significantly diagnostic specimen from the Paleogene of South America, and adding to the knowledge of the sparse Paleogene xiphioids known along the Southern Hemisphere.

LOCALITY AND GEOLOGIC SETTING

Loanco is a small cove placed 350 km south from Santiago (Fig. 1A). In this locality, rich fossiliferous marine rocks crop out along the coastal cliffs and the intertidal surface. Beds exposed north of Loanco have been recognized as equivalent to the Late Cretaceous Quiriquina Formation (*Tavera, 1987*), assigned to the upper Maastrichtian based on biostratigraphy and stratigraphic correlations (*Otero, 2015; Castro et al., 2016*). A second sedimentary and also marine unit was recently found cropping out south of Loanco. From the latter, *Suárez and Otero (2008)* recovered associated teeth, vertebrae and mandibular cartilage belonging to a single individual referred to the genus *Isurolamna* (Chondrichthyes, Elasmobranchii). Previous geologic studies have referred the outcrops in south of Loanco as Late Cretaceous in age (*Sernageomin, 2003*). During 2012, new fieldwork in this locality has provided material referable to the nautiloid genus *Aturia* (*Zambrano et al., 2014*) and *Imaizula araucana* (recovered during fieldwork of this research), which are unequivocally Cenozoic taxa. More recently, *Otero (2015)* identified the presence of chondrichthyan teeth referable to *Macrorhizodus praecursor* (SGO.PV.6633; SGO, Paleontología de Vertebrados, Museo

Nacional de Historia Natural, Santiago, Chile). The combined biochron of this assemblage indicates a Middle Eocene age for the fossil-bearing levels.

The Quiriquina Formation and the Paleogene marine units of central Chile were deposited in a continental shelf which is part of a single realm known as the Arauco Basin (*Mordojovic, 1981*), extended at least between 33° to 37°S. In this context, Paleogene units have been recorded along the whole basin, being deposited in angular unconformity over Late Cretaceous beds of the Quiriquina Formation or equivalent units. In Cocholgüe and the Quiriquina Island, 120 km south from Loanco, the Quiriquina Formation is unconformably overlain by Paleogene sediments of the Concepción Group (*sensu Charrier et al., 2007*; =Lebu Group in *Cecioni, 1968*; *Le Roux et al., 2008*). Several Paleogene units have been characterized in the Arauco Basin, representing different stages of a transgressive-regressive cycle. After a Paleocene hiatus, from base to roof, a first shallow transgression is represented by the marine-to-continental, upper Paleogene-lower Eocene Pilpilco Formation (*Muñoz-Cristi, 1968*), conformed by ca. 150 m of fine-grained, greenish sandstones with clay intercalations. The unit unconformably overlies the metamorphic basement and the Quiriquina Formation. The Pilpilco Formation is covered by a conformably deposited, regressive unit represented by the continental, coal-rich sandstones of the Curanilahue Formation (*Tavera, 1942*; *Martínez-Pardo, 1968a*) assigned to the lower Eocene. Over the latter and in conformable contact lays the Boca Lebu Formation (*Muñoz-Cristi, 1946*; *1968*), which comprises a variable section between 250 to 600 m of marine, fossiliferous green sandstones with glauconite. The upper regression is represented by the continental Trihueco Formation (*Muñoz-Cristi, 1946*; *1968*) comprised by 3 to 10 m thick banks with blue-to-grey sandstones with intercalations of brown, coal-rich shales and coal seams. The final transgression is represented by the Millongue Formation (*Muñoz-Cristi, 1946*). This unit was

characterized by *García (1968)* as 270 m of thin beds of shales and limestones with leaf prints and frequent fragments of carbonized wood, representing a marine-continental environment. Following this geologic scheme (Fig. 1B), the fossil-bearing section studied here could be referred to one of these marine units (i.e., basal Pilpilco Formation; Boca Lebu Formation or Millongue Formation).

The studied section (Fig. 1C) is comprised from base to roof, by: 3 m of greenish sandstones with banks of serpulids; 0.5 m with no outcrop; 2 m of reddish sandstones; 2 m with no outcrop; 1.8 m of mid-grained sandstone, brown, with micaceous fragments, and having a basal fine conglomerate. Fossils include teeth of *Macrorhizodus praecursor*, a phragmocone of the nautiloid *Aturia* sp. (SGO.PI.6776), and an articulated individual of *Imaizula araucana*; 14.3 m of mid-grained, yellow-to-green sandstone with lenses including banks of terebratulid brachiopods. Isolated terebratulids, pectinids and carbonized wood fragments occur near the roof of the level; 6 m of mid-grained, reddish to brown sandstones, including few individuals of the bivalvian *Pinna* sp. in life position, as well as an isolated lamniform vertebra, serpulids, carbonized wood fragments, and *Teredolites* isp. Lithologically, these strata could be referred to the Boca Lebu or Millongue formations; however, the presence of *Pinna* sp. has been previously considered as a proxy for distinguishing both units, because the occurrence of this genus seems to be restricted only to the Millongue Formation (*Martínez-Pardo, 1968b*: p. 100). Adding to these facts, a Middle Eocene age is proposed based on the combined biochron of the faunal assemblage. Then, the studied unit is here proposed to be equivalent to the Millongue Formation. Several outcrops of the same unit in south Loanco were documented by the author during the extraction of SGO.PV.6635 (associated chondrichthyan remains referred to *Isurolamna* sp.); however, these were later covered by sand banks after the large earthquake and subsequent

tsunami of central Chile in February 27, 2010. Currently, the unit is only visible during low tides along the coastal line south from Loanco, but its extension under the recent sands reaches hundreds of meters. The sediment attached to SGO.PV.6634 is remarkably similar to the fossil-bearing level including *Aturia* sp. and *Imaizula araucana*; however, the lack of historical information regarding its collection (see text below) makes impossible to assure a precise stratigraphic provenance.

MATERIAL AND METHODS

Historical background of the studied specimen—SGO.PV.6634 (Fig. 2) was recovered by H. Fuenzalida, former director of the Museo Nacional de Historia Natural (MNHN, National Museum of Natural History, Santiago, Chile), during 1935 from rocks exposed south of Loanco, as is accounted by his personal field notes archived in the aforementioned institution. The collected material also included ammonoids, bivalvians and plesiosaur remains; however, these were recovered without stratigraphic data. Later, the collection of Fuenzalida was studied by *San Martín (1946)*, who assigned a 'Senonian' age to the assemblage based on ammonoids, and recognizing affinities with the fauna of the Quiriquina Formation. Among the determinations by *San Martín (1946)*, the plesiosaur remains were considered as Cenozoic remains of indeterminate cetaceans. The swordfish remains were not prepared at that moment, being interpreted as *Teredo* remains. Since then, the whole collection was housed in the MNHN. Probably because of its antiquity, the collection was not included in any catalog, being re-discovered during 2008 by S. Soto-Acuña (Universidad de Chile). After this, a first inspection of the 'wood remains' revealed that they are actually bone remains. Subsequently, part of the material was mechanically prepared by the author during 2008 and 2009. The rostrum and

dentaries were preliminary presented and identified as an indeterminate xiphiorhynchine by Friedman and Otero (2009). These authors also gave to the specimen, for the first time, a formal numeration and repository (SGO.PV.6634). Further preparation was carried out by the author during 2012 and 2013, uncovering new elements of the skull. Anatomical identifications were made during April, 2013 by direct comparison with the holotype of *Aglyptorhynchus hatakaramaea* Gottfried et al. 2013, housed in the Department of Geology of the Otago University, Dunedin, New Zealand. Further anatomical identifications follow the classical work of Gregory and Conrad (1937). Considering the osteological nomenclature of billfishes in general, there is a controversial issue regarding the presence of prenasal bone (Shultz, 1987; Fierstine, 1990; Fierstine and Voigt 1996). Resolving this issue is out of the scope of this contribution. However, this research follows the uncertain denomination of the dorsomedial rostrum element in question as prenasals, as was done by Fierstine (1990). The same element was considered by Gregory and Conrad (1937) as the nasal.

Phylogenetic analysis—Phylogenetic analysis used a modified version of the datamatrix with twelve taxa (fossil and extant) and 25 characters provided by Fierstine and Monsch (2002). Modifications here introduced include three new characters (26 to 28) and addition of new states for characters 4, 11 and 15 (see Supplementary Data). Original dataset of Istiophoridae was replaced with the scorings of *Istiophorus albicans* and *Makaira nigricans* (taken from Nakamura, 1985). Original dataset of *Aglyptorhynchus* was also replaced with the scores of *Aglyptorhynchus hatakaramaea*, which is a fairly complete specimen from the Eocene of New Zealand (Gottfried et al., 2013). Also, the scores of *Hemingwaya sarissa* Sytchevskaya and Prokofiev, 2002; and SGO.PV.6634 were added to the datamatrix. Few taxa in the original datamatrix of Fierstine and Monsch (2002) were pruned based in different criteria:

Cylindracanthus—This genus lacks a subdermal tooth base attachment (*Grandstaff et al., 2017*), leading to the conclusion that *Cylindracanthus* does not belong to any billfish (=Xiphiidae), which indeed have subdermal tooth base attachments. Previously, *Parris and Grandstaff (2001)* suggested that *Cylindracanthus* could be related to Acipenseriformes.

Hemirhabdorrhynchus—*Fierstine and Monsch (2002)* originally excluded this genus from phylogenetic analysis based in its ambiguity. This observation is here followed. Even more, no further clarification regarding the anatomy and taxonomy of *Hemirhabdorrhynchus* was provided since *Shultz (1987)*. *Fierstine and Monsch (2002)* mentioned *Shultz (1987)* as the source of information used by them for scoring of this genus.

'Blochius' moorheadi—This was detected as an unstable taxon with the IterPcr Script (*Pol and Escapa, 2009*). *Fierstine and Monsch (2002)* also found this taxon as unstable, pruning it from the returned reduced consensus tree (*Fierstine and Monsch, 2002: fig. 7*).

Pseudotetrapturus luteus—This taxon was removed for reducing ambiguity. Besides the rostrum, the skull of the holotype is naturally crushed, and the postcranial skeleton is currently lost (*Monsch and Bannikov, 2011*).

The modified datamatrix includes 11 taxa and 28 characters. This was analyzed with TNT software (*Goloboff et al., 2008*).

Permissions—The studied fossil was collected in 1935. At that time, field permissions for paleontologic excavation in Chile were not required. Further fieldwork in 2012 was granted by the Chilean authority in charge, Consejo de Monumentos Nacionales, in document Ord. CMN N° 2962/2009 and currently, in Ord. CMN N° 3793/2014.

Nomenclatural acts—New names contained in the electronic version of this article are effectively published under the International Commission on Zoological Nomenclature (ICZN)

from the electronic edition alone. This published work and the nomenclatural acts it contains have been registered in ZooBank, the online registration system for the ICZN. The ZooBank LSIDs (Life Science Identifiers) can be resolved and the associated information viewed through any standard web browser by appending the LSID to the prefix "http://zoobank.org/". The LSID for this publication are:

Genus name: LSID:urn:lsid:zoobank.org:act:FA566753-31D4-45B5-9E7A-03F8C7526E7D

Species name: LSID: urn:lsid:zoobank.org:act:8FFA9EE9-A81B-4468-9FBA-9D6C306D86C8

Publication: LSID:urn:lsid:zoobank.org:pub:ED22325A-7C78-469A-9985-365C31064652

The electronic version of this article in Portable Document Format (PDF) will represent a published work according to the International Commission on Zoological Nomenclature (ICZN), and hence the new names contained in the electronic version are effectively published under that Code from the electronic edition alone. This published work and the nomenclatural acts it contains have been registered in ZooBank, the online registration system for the ICZN. The ZooBank LSIDs (Life Science Identifiers) can be resolved and the associated information viewed through any standard web browser by appending the LSID to the prefix http://zoobank.org/. The LSID for this publication is: LSID:urn:lsid:zoobank.org:pub:ED22325A-7C78-469A-9985-365C31064652

The online version of this work is archived and available from the following digital repositories: PeerJ, PubMed Central and CLOCKSS.

SYSTEMATIC PALEONTOLOGY

221 Order PERCIFORMES *sensu Johnson and Patterson, 1993*

222 Suborder XIPHIODEI *sensu Gosline, 1968*

223 Family BLOCHIIDAE (*Bleeker, 1859*)

224

225 Genus *LOANCORHYNCHUS* nov.

226

227 **Type Species**—*Loancorhynchus catrillancai* gen. et sp. nov., by monotypy.

228 **Derivation of Name**—Genus name after its type locality, Loanco, Región del Maule,
229 central Chile.

230 **Diagnosis**—As for the single known species, below.

231

232 *LOANCORHYNCHUS CATRILLANCAI* gen. et sp. nov.

233 (Fig. 2-6)

234

235 *Teredo* sp.: *San Martín, 1946: p. 46.*

236 Xiphiorhynchinae indet.: In *Friedman and Otero, 2009.*

237 **Holotype**—SGO.PV.6634, a dissarticulated three-dimensional skull preserving most of
238 the rostrum (premaxillaries and prenasals?), fragments of both dentaries, a fragment of the left
239 maxilla, left frontal, left metapterygoid+hyomandibular, both operculi, and several branchiostegal
240 bones.

241 **Derivation of Name**—Honoring the memory of the Mapuche leader Camilo Catrillanca.

242 **Locality, Horizon and Age**—Loanco, Región del Maule. Indeterminate levels of the
243 Millongue Formation cropping out south Loanco. Middle Eocene.

Diagnosis—Rostrum much longer than the dentaries, with villiform teeth reaching the posterior part of the rostrum; rostrum and dentaries with villiform teeth over their occlusal and latero-occlusal surface; rostrum with clearly oval, dorsally compressed cross-section, intermediate between *Xiphias* and *Xiphiorhynchus*; operculum rounder and broader than in *Blochius longirostris* but dorsoventrally more compressed than in *Xiphias gladius*; hyomandibular/metaethmoid complex more gracile than in *Blochius* but broader than in *Xiphias*.

Other distinguishing characters—*Loancorhynchus catrillancai* skull has premaxillaries joined into a pointed rostrum and a rostrum usually longer than the lower jaw. Both features have been considered as diagnostic of the family Blochiidae by *Fierstine and Monsch (2002)*. Unique combination of characters include features found in Hemingwayidae (*Sytchevskaya and Prokofiev, 2002*), Xiphiorhynchinae (*Regan, 1909*), Xiphiinae (sensu *Fierstine, 2006*) and Blochiidae (*Bleeker, 1859*) *Loancorhynchus catrillancai* possesses villiform teeth along the premaxillaries and also in the occlusal surface of the dentaries. Such feature is only documented in the *Hemingwaya sarissa* *Sytchevskaya and Prokofiev, 2002*. *Loancorhynchus catrillancai* also possesses a thick, dorsoventrally compressed rostrum formed mostly by premaxillaries extended posteriorly on each side, while its dorsal midline is conformed by an anterior extension of the prenasals(?), and posteriorly, by a short extension of the dermethmoid, as it occurs in *Xiphias*; It also shares with *Xiphias* the presence of a metapterygoid fused to the hyomandibular, forming a dorsoventrally high complex with a prominent dorsal articulation, and a subrounded, laterally convex operculum (*Gregory and Conrad, 1937*). Traits shared with Xiphiorhynchinae include a lower jaw shorter than the rostrum but comparatively larger than that of *Xiphias*; posterior part of the rostrum with strong folds and sulci, as those present in *Aglyptorhynchus*, *Xiphiorhynchus* and

Blochius; additional traits include a lower jaw with dentaries contacted in a large craniocaudal symphysis as that present in *Pseudotetrapturus*, *Blochius* and all xiphiids.

Taxonomic placement of *Loancorhynchus catrillancai* —The phylogenetic analysis (see Discussion below) confirms the placement of Blochiidae as the sister taxon of *Xiphias gladius* (i.e., Xiphiidae), as previously obtained by *Fierstine and Monsch (2002)* and by *Fierstine (2006)*. *Loancorhynchus catrillancai* has a set of derived traits shared with *Xiphias*, including a rostrum formed by premaxillaries+prenasals(?), a dorsoventrally high metapterygoid-hyomandibular complex, and a large, rounded and convex operculum. On the other hand, the presence of dorsal rostral ridges and sulci seems to be a symplesiomorphic trait shared with *Blochius* spp. These characters support *Loancorhynchus catrillancai* within the node including *Xiphias* + *Blochius* spp.; it is also segregated from other lineages with dorsal ridges (but having more basal features), such as *Aglyptorhynchus* and *Xiphiorhynchus*. Finally, the presence of villiform teeth over the occlusal surface of the dentaries is remarkable. Such a feature was previously described only in hemingwayids (i.e., *Hemingwaya sarissa* *Sytchevskaya and Prokofiev, 2002*). The presence of villiform teeth in *Loancorhynchus catrillancai* dentaries could be atavic, or else, it could suggest a non-adult ontogenetic stage, as it occurs in the larval rostrum of *Xiphias* that bears villiform teeth, but these are absent in its adult stage (*Nakamura, 1985; Johnson, 1986; Fierstine and Monsch, 2002*). However, the latter hypothesis is unprobably for a large specimen such as *Loancorhynchus catrillancai*, with a length estimated in 2.7 m (see further text).

DESCRIPTION OF *LOANCORHYNCHUS CATRILLANCAI* HOLOTYPE

General remarks— *Loancorhynchus catrillancai* is three-dimensionally preserved.

Besides the rostrum, all the available postorbital elements are well-preserved and none of them is severely deformed. Both operculi even preserve their convexity. The rostrum is partially crushed dorsoventrally. Its anteriormost fragment is slightly recurved to the right side. On the contrary, the left dentary (the most complete) is recurved to the left side, which indicates that both deformations were caused likely by taphonomic condition instead of occurring in the living animal. During its preparation, it was evident that the skull was naturally dissarticulated and embedded in a soft, reddish, micaceous sandstone (consistent with those observed in the base of the studied section; see Geologic Setting). The length of the cranial elements in anatomic position, represents approx. 60 cm. Based on the affinities of *Loancorhynchus catrillancai* with Blochiidae, the skull could reach a conservative length of 90 cm, assuming a rostrum with a blunt tip instead a gradually reduced, sharp tip. The body length of *Loancorhynchus catrillancai* can be estimated based on the proportions of complete skeletons of *Blochius longirostris*, where the skull is ca. one third of the whole skeleton (Fierstine and Monsch, 2002: fig. 1). Then, *Loancorhynchus catrillancai* body length can be estimated in 2.7 meters.

Premaxillaries—The premaxillaries form most of the anteriormost part of the

rostrum. The anterior tip of the rostrum is lost; however, this was likely formed by the union of both premaxillaries, suggested by their large participation in the anteriormost available rostrum fragment. In the latter, premaxillaries are strongly fused in the mid and ventral part, being interrupted in the dorsal midline by a thin anterior extension of the prenasals(?). Both fused premaxillaries show a dorsoventrally compressed oval cross-section. Nutrient canals are not observed, probably due to taphonomic conditions. Profuse villiform teeth cover all the ventral and lateral surface of both premaxillaries, over the whole length of the preserved rostrum (Fig. 3A,

B). The posterior end of each premaxillary bears profuse craniocaudal striations. Two main interior canals are visible in the anterior fragment of the rostrum (Fig. 3E,F, G). Smaller canals are also present, being paired with respect to the rostrum midline.

Maxillaries—A small fragment of the left maxillary is preserved (Fig. 4). This lies crushed under the left premaxillary. This bone is elongated with a sub-squared cross-section. Based on the available fragment, the lateral exposure of each maxillary seems to be precluded by the posterior extension of the premaxillaries.

Prenasals(?)—In the anteriormost available rostrum fragment, two thin bones in its dorsal midline are here interpreted as both prenasals(?) sensu *Fierstine (1990)* (Fig. 4). In cross-section, their contact with the premaxillaries is obscured due to preservation. The posteriormost available rostrum fragment preserves the posterior part of both prenasals(?). These laterally diverge and they have a posterolateral notch (Fig. 4D), which marks the anterior margin of the orbit. Then, the anteroventral orbit is laterally conformed by the posterior extension of the premaxillary, and ventrally, by the posterior extension of the maxillary. The posterior end of both prenasals(?) have strong folds and sulci over their dorsal surface.

Dermethmoid—In the posterior part of the rostrum, a partially open sutural contact is visible besides the dorsal midline. This marks the contact of the prenasals(?) with the dermethmoid, which is placed in the midline and tapers anteriorly. Dorsally, the dermethmoid bears strong folds and sulci (Fig. 4). Its posterior part is missing.

Hyomandibular—Only the left hyomandibular is available. This element is complete and well preserved (Fig. 5A-D). In lateral (external) view, this shows a prominent pterotic facet which is medially recurved. In the same view, the sphenotic facet is high and diverged from the pterotic facet under ca. 90°. A large opercular process is extended over the posterior margin of

the hyomandibular. In posterior view, this process appears as diagonally oriented with respect to the vertical. Over the lateral (external) surface, the hyomandibular bears a dorsoventral scar, consistent with the preopercular groove. In both anterior and internal views, there is a large foramen under the pterotic facet, here interpreted as the foramen for the *truncus hyoideomandibularis* of the facial nerve (VII).

Metapterygoid—This bone remains strongly articulated to the hyomandibular (Fig. 5A-D), although, its suture is still visible from the internal and lateral views. This element diagonally overlaps the hyomndibular. Laterally, it contacts with the latter through a suture adjacent to the preopercular groove. In internal view, the dorsal part of the metapterygoid is comparatively broader than its external exposure. Its ventral margin has a squared outline.

Frontal—The left frontal is preserved (Fig. 5E, F), having the medial and posterior margins missing. This element has a posterolateral process. This indicates that the skull becomes laterally broader immediately behind the orbit. In ventral (internal) view, the frontal shows a thick ridge that broadens posteriorly.

Operculum—Only the left operculum is preserved (Fig. 5G) This shows the articular process for the hyomandibular. It has a well-marked convexity immediately ventral to the articulation.

Subopercular—Only the right subopercular is preserved (Fig. 5H). This seems to be complete. It has a dorsal rounded contour. This element remains articulated to the right interopercular.

interopercular—It is crushed together with the subopercular due to taphonomic conditions, but it can be distinguished as a bony element separated from the subopercular (Fig. 5H). It has a triangular outline with a convex posterior margin.

Dentaries—Fragments of both dentaries are preserved (Fig. 6). The left dentary is the most complete. However, its anterior part as well as its posterior end are both missing. The right dentary is represented by two separated fragments. The complete occlusal and laterodorsal surface of the left dentary are covered by profuse, small villiform teeth. Villiform teeth are also present in the two fragments of the right dentary. The left dentary is very straight, having a slight lateral divergence in its posteriormost preserved end. Two separated fragments of the right dentary can be attached to the left dentary in anatomic position. This shows that the jaw had a large symphysis. The anterior cross-section of the preserved jaw has a sub-squared outline. The anterior part of both dentaries narrows anteriorly. This suggests that the length of the missing anterior part could be approximately 5 cm. Both mandibular rami diverge posteriorly. Based on this divergence, a tentative position of the jaw elements with respect to the rostrum can be estimated. Even if this is not accurate, it is clear that the lower jaw is much shorter than the rostrum.

Fin rays—Remains of least nine fin rays were recovered. All of them were included in a single block. A large incomplete fragment has an articular head, recurved with respect to the shaft. It also has a small bulk in its anterior margin. These features are present in the first pectoral ray of *Xiphias gladius*, reason by why it is identified as the latter element.

DISCUSSION

Historical background in billfish phylogeny—The fossil billfish record was extensively reviewed by *Fierstine (1972)*, recognizing five families (Blochiidae, Istiophoridae, Palaeorhynchidae, Xiphiidae and Xiphiorhynchidae). To these, *Schultz (1987)* added a sixth family, Tetrapturidae, including the genera *Hemirhabdorrhynchus*, *Aglyptorrhynchus*,

Pseudoistriophorus, and *Tetrapturus*. The latter author also considered the genus *Acestrus* (an istiophorid in *Fierstine, 1972*) as *incertae sedis*. Later, *Fierstine and Monsch (2002)* recovered the five families of *Fierstine (1972)*, but considered the genera *Aglyptorhynchus*, *Cylindracanthus* and *Hemirhabdorrhynchus* as unsolved. The taxonomic status of *Cylindracanthus* was discussed by *Parris and Grandstaff (2001)* who suggested affinities with Acipenseriformes instead to Xiphioids. *Fierstine (2006)* reviewed again the fossil record of billfishes, considering a large set of different specimens. This author listed the families Blochiidae, Palaeorhynchidae, Xiphiidae and Istiophoridae, and the family Hemingwayidae (*Sytchevskaya and Prokofiev, 2002*). In addition, the former family 'Xiphiorhynchidae' (*sensu Fierstine, 1972*) was re-ranked to subfamily level (Xiphiorhynchinae), and included within the Xiphiidae. After, *Fierstine and Weems (2009)* distinguished *Aglyptorhynchus* from all other Palaeorhynchidae, erecting a new subfamily, Aglyptorhynchinae.

Phylogenetic analysis—The datamatrix of *Fierstine and Monsch (2002)* with the modifications introduced here, was first tested with Traditional Search (Wagner; 1,000 replicates; 1,000 trees to save per replication), recovering nine most parsimonious cladograms. *Acanthocybium solandri* was selected as the outgroup. Strict consensus cladogram returned a polytomy between *Loancorhynchus catrillancai*, *Xiphias gladius* and *Blochius* spp. After, IterPcr script (*Pol and Escapa, 2009*) was used for identifying 'Blochius' *moorheadi* and *Xiphiorhynchus* as unstable taxa. Thus, 'Blochius' *moorheadi* was subsequently pruned, while *Xiphiorhynchus* was kept because its phylogenetic placement is relevant for assessing any eventual relationship to *Loancorhynchus catrillancai*. As a result, *Xiphias gladius* was obtained as the sister taxon of a polytomic group conformed by *Blochius* spp. + *Loancorhynchus catrillancai*. Further pruning of non-xiphioid taxa *Cylindracanthus* and *Hemirhabdorrhynchus*,

returned the same relationships between *Xiphias* and *Blochius* spp. + *Loancorhynchus* *catrillancai*, the latter in polytomy. *Pseudotetrapturus* was later pruned from the previous datamatrix version, and tested with Implied Weighting, obtaining *Loancorhynchus catrillancai* as the sister taxon of *Blochius*. As obtained synapomorphies, the presence of dorsal longitudinal ridges on the rostrum (Char. 7) groups *Loancorhynchus catrillancai* with the Blochiidae, while the presence of villiform teeth in the rostrum and dentary (Char. 27) segregates *Loancorhynchus catrillancai* from *Blochius longirostris* and *Blochius macropterus*. The topology of this cladogram is shown in Fig. 7.

In addition, the phylogenetic analysis returned *Xiphiorhynchus* as a basal lineage with respect to *Xiphias*. Previous to this research, *Xiphiorhynchus* was obtained as the sister taxon of *Xiphias* (Fierstine and Monsch, 2002). Another novel result is the obtention of *Hemingwaya* as the sister taxon of *Palaeorhynchus*+*Homorhynchus*.

Paleobiogeography—Previous to this research, the known records of Blochiidae were restricted to the Middle Eocene of Monte Bolca, Italy. *Loancorhynchus catrillancai* represents the first occurrence of a blochiid in the southern hemisphere, and the fourth record of a Paleogene xiphioid in the Weddellian Province, adding to cf. *Xiphiorhynchus* from the Eocene of Marambio (=Seymour) Island, Antarctica (Cione et al., 2001), *Aglyptorynchus hakataramea*, from the late Oligocene of New Zealand (Gottfried et al., 2013), and the still undescribed GS 13924 skull from the Eocene of Otago, New Zealand (Campbell et al., 2013). This sparse record reveals a diversity of Paleogene xiphioids in the southern hemisphere, restricted to the families Xiphiorhynchidae, Palaeorhynchidae, and now Blochiidae.

CONCLUSIONS

Loancorhynchus catrillancai represents the unique Middle Eocene xiphioid known to date in the southeastern Pacific. Its first available elements (rostrum and dentaries) were initially considered as an indeterminate xiphiorhynchine (Friedman and Otero, 2009), while its stratigraphic provenance remained dubious since 1935, when this specimen was recovered. New preparation revealed basal features found among blochiids, xiphiorhynchines and hemingwayids. Also, derived traits exclusively known in Neogene xiphiines were found in SGO.PV.6634. Based on these facts, this research reassesses the taxonomical determination of SGO.PV.6634, being now identified as a new genus and species, *Loancorhynchus catrillancai*. This new genus and species represents the first record of the clade Blochiidae in the southern hemisphere. The new occurrence of a Middle Eocene swordfish now in the southeastern Pacific, helps to fill the Eocene-Middle Miocene gap in the xiphioid fossil record. It also represents the fourth record of a Paleogene billfish in the Southern Hemisphere, showing that this group already reached a wide distribution along the Southern Hemisphere previous to the Neogene, with Eocene records known in New Zealand (Gottfried et al., 2013; Campbell et al., 2013), Antarctica (Cione et al., 2001) and the new record here presented, now in South America.

Acknowledgments

Special thanks are due to H. Fierstine (California Polytechnic State University) who provided valuable literature during early stages of this research. Thanks also to M. Friedman (University of Oxford) for his contribution in the preliminary study of SGO.PV.6634. A.M. Prokofiev (A.N. Severtsov's Institute of Ecology and Evolution, Russian Academy of Sciences) is also thanked by valuable literature provided. K. Monsch (Naturalis Biodiversity Center, Netherlands) is acknowledged for the revision of a early version of the manuscript. M. Gottfried (Geological

Sciences and Museum, Michigan State University, USA), P. Brito (Universidade do Estado do Rio de Janeiro, Brasil), D. Parris (New Jersey State Museum, USA) and G. Piñeiro (Universidad de la República, Uruguay) are especially thanked for their reviews and all the comments that improved this paper. R.E. Fordyce (Otago University, Dunedin, New Zealand) is acknowledged for allowing access to the holotype of *Aglyptorhynchus hakataramea* during 2013.

Institutional Abbreviations

GS, Geological Survey, GNS Science, Lower Hutt, Wellington, New Zealand.
SGO.PI, Paleontología de Invertebrados, Museo Nacional de Historia Natural, Santiago, Chile.
SGO.PV, Paleontología de Vertebrados, Museo Nacional de Historia Natural, Santiago, Chile.

New Species Registration

The following information was supplied regarding the registration of newly described taxa:

Genus name: LSID:urn:lsid:zoobank.org:act:FA566753-31D4-45B5-9E7A-03F8C7526E7D
 Species name: LSID: urn:lsid:zoobank.org:act:8FFA9EE9-A81B-4468-9FBA-9D6C306D86C8
 Publication: LSID:urn:lsid:zoobank.org:pub:ED22325A-7C78-469A-9985-365C31064652

REFERENCES

Bleeker PE. 1859. Enumeratio Specierum Piscium hueusque in Archipelago Indico observatarum. Verhandelingen der Naturkundige Vereeninging in Nederlandsch Indië **6**:1–276.

471 **Campbell H, Beu A, Crampton J, Kennedy L, Terezow M. 2013.** A Photographic Guide to
472 Fossils of New Zealand. New Holland Publishers, New Zealand. 144 p.

473 **Castro N, Salazar C, Otero RA, Stinnesbeck W, Escare A. 2016.** V Simposio de
474 Paleontología en Chile, Libro de Resúmenes, p. 26-28. November 7-11, 2016. Concepción.

475 **Cecioni G. 1968.** El Terciario de Chile. Zona central. Sociedad Geológica de Chile. Editorial
476 Andrés Bello, Santiago, Chile. 280 p.

477 **Charrier R, Pinto L, Rodríguez MP. 2007.** Tectonostratigraphic evolution of the Andean
478 Orogen in Chile. In: Moreno T, Gibbons W, eds., *The Geology of Chile*, The Geological
479 Society, London, 21–114.

480 **Cione AL, Reguero MA, Elliot DH. 2001.** A large osteichthyan vertebra from the Eocene of
481 Antarctica. *Neues Jahrbuch Fur Geologie Und Palaontologie Monatshefte* **9**:543-552.

482 **Fierstine H. 1972.** The paleontology of billfish —The state of the art. In: Shomura R, Williams
483 F, eds., *Contributed papers, Proceeding of the International Billfish Symposium*, Kailua-
484 Kona, Hawaii, 9-12 August, 1972, 34–44.

485 **Fierstine H. 1990.** A paleontological review of three billfish families (Istiophoridae, Xiphiidae,
486 and Xiphiorhynchidae). In: stroud RH, ed. *Planning the future of billfishes. Part 2.*
487 *contributed papers*. Proceedings of the Second International Billfish Symposium, p. 11–19.
488 Kailua-Kona, Hawaii, August 1–5, 1988.

489 **Fierstine H. 2006.** Fossil history of billfishes (Xiphiidae). *Bulletin of Marine Science*
490 **79**:433–453.

491 **Fierstine HL, Monsch KA. 2002.** Redescription and phylogenetic relationships of the Family
492 Blochiidae (Perciformes: Scombroidei), Middle Eocene, Monte Bolca, Italy. *Miscellanea*

493 Paleontologia, Studi e Ricerche sui Giacimenti Terziari de Bolca, Museo Civico di Storia
494 Naturale di Verona **9**:121–163.

495 **Fierstine, HL, Voigt NL. 1996.** Use of rostral characters for identifying adult billfishes
496 (Teleostei: Perciformes: Istiophoridae and Xiphiidae). *Copeia* **1996**:148–161.

497 **Fierstine HL, Weems RE, 2009.** Paleontology of the Oligocene Ashley and Chamber Bridge
498 formations of South Carolina, 4:analysis and new records of billfishes (Perciformes:
499 Xiphioidei). *Palaeoichthyologica* **11**:43–88.

500 **Friedman M, Otero RA. 2009.** First fossil billfish (Perciformes: Xiphiidae) from central Chile.
501 Congreso Geológico Chileno XII, Symposium 10, Abstract 29, p. 1-4. Santiago, November
502 22-26, 2009.

503 **García F. 1968.** Estratigrafía del Terciario de Chile central. In: Cecioni G, ed. *El Terciario de*
504 *Chile. Zona central.* Sociedad Geológica de Chile. Editorial Andrés Bello, Santiago, Chile, p.
505 25–57.

506 **Goloboff PA, Farris JS, Nixon KC. 2008.** TNT, a free program for phylogenetic analysis.
507 *Cladistics* **24**:774–786.

508 **Gosline WA. 1968.** The suborders of Perciform Fishes. *Proceedings of the United States*
509 *National Museum* **126**:1–78.

510 **Gottfried M, Fordyce RE, Rust S. 2013.** A new billfish (Perciformes, Xiphioidei) from the late
511 Oligocene of New Zealand. *Journal of Vertebrate Paleontology* **32**:27–34.

512 **Grandstaff BS, Pellegrini RA, Parris DC, Clements D. 2017.** Thin section microscopy of the
513 fossil fish *Cylindracanthus*. *Proceedings of the South Dakota Academy of Science*
514 **96**:130–143.

- 515 **Gregory WK, Conrad GM. 1937.** The comparative osteology of the swordfish (*Xiphias*) and
516 the sailfish (*Istiophorus*). American Museum Novitates **952**:1–28.
- 517 **Johnson GD, Patterson C. 1993.** Percomorph phylogeny: a survey of acanthomorphs and a new
518 proposal. Bulletin of Marine Science **52**:554–626.
- 519 **Johnson GD. 1986.** Scombroid phylogeny: An alternative hypothesis. Bulletin of Marine
520 Science **39**:1–41.
- 521 **Le Roux JP, Nielsen SN, Henríquez A. 2008.** Depositional environment of *Stelloglyphus*
522 *llicoensis* isp. nov.: a new radial trace fossil from the Neogene Ranquil Formation, south-
523 central Chile. Revista Geológica de Chile **35**:307-319.
- 524 **Long DJ. 1993.** Late Miocene and Early Pliocene fish assemblages from the north central coast
525 of Chile. Tertiary Research **14**:117–126.
- 526 **Martínez-Pardo R. 1968a.** Edad y paleoecología del "Piso de Millongue". In: Cecioni G, ed. *El*
527 *Terciario de Chile. Zona central*. Sociedad Geológica de Chile. Editorial Andrés Bello,
528 Santiago, Chile, p. 125–141.
- 529 **Martínez-Pardo R. 1968b.** Necesidad de una nueva comprensión de los esquemas estratigráfico
530 clásicos sobre el Terciario de Chile Central. In: Cecioni G, ed. *El Terciario de Chile. Zona*
531 *central*. Sociedad Geológica de Chile. Editorial Andrés Bello, Santiago, Chile, p. 95–103.
- 532 **Monsch K, Bannikov AF. 2011.** New taxonomic synopses and revision of the scombroid fishes
533 (Scombroidei, Perciformes), including billfishes, from the Cenozoic of territories of the
534 former USSR. Earth and Environmental Science Transactions of the Royal Society of
535 Edinburgh **102**:253–300.
- 536 **Mordojovic C. 1981.** Sedimentary basins of the Chilean Pacific Basin. American Association of
537 Petroleum Geologists, Bulletin **12**:63–82.

- 538 **Muizon C. de, DeVries TJ. 1985.** Geology and Paleontology of late Cenozoic marine deposits
539 in the Sacaco area (Peru). *Geologische Rundschau* **74**:547–563.
- 540 **Muñoz-Cristi J. 1946.** Estado actual del conocimiento sobre la geología de la provincia de
541 Arauco. *Anales Facultad de Ciencias Físicas y Matemáticas, Universidad de Chile* **3**:30–63.
- 542 **Muñoz-Cristi J. 1968.** Contribución al conocimiento geológico de la región situada al sur de
543 Arauco y participación del material volcánico en los sedimentos eocenos. In: Cecioni G, ed.
544 *El Terciario de Chile. Zona central*. Sociedad Geológica de Chile. Editorial Andrés Bello,
545 Santiago, Chile, p. 63–93.
- 546 **Nakamura I. 1985.** an annotated and illustrated catalogue of marlins, sailfishes, spearfishes, and
547 swordfishes known to date. *Fao Fisheries synopsis No. 125* **5**:1–65.
- 548 **Otero RA. 2015.** Síntesis preliminar del registro fósil de vertebrados en Faro Carranza (Región
549 del Maule) y su aporte al conocimiento geológico del área. XIV Congreso Geológico Chileno,
550 Libro de Actas, p. 708-711. La Serena, October 2015.
- 551 **Parris DC, Grandstaff BS. 2001.** Reassessment of the affinities of the extinct genus
552 *Cylindracanthus* (Osteichthyes). *Proceedings of the South Dakota Academy of Science*
553 **80**:161–172.
- 554 **Pol D, Escapa IH. 2009.** Unstable taxa in cladistics analysis: identification and the assessment
555 of relevant characters. *Cladistics* **25**:515–527.
- 556 **Pyenson ND, Gutstein CS, Parham JF, Le Roux JP, Carreño-Chavarría C, Little H,**
557 **Metallo A, Rossi V, Valenzuela-Toro AM, Velez-Juarbe J, Santelli CM, Rubilar Rogers**
558 **D, Cozzuol MA, Suárez ME. 2014.** Repeated mass strandings of Miocene marine mammals
559 from Atacama Region of Chile point to sudden death at sea. *Proceedings of the Royal Society*
560 *B* **281**: 9 p. DOI: 10.1098/rspb.2013.3316

- 561 **Regan CT. 1909.** On the anatomy and classification of the scombroid fishes. *Annals and*
562 *Magazine of Natural History* **8**:66–75.
- 563 **San Martín R. 1946.** Los sedimentos del Cretáceo y Terciario de Faro Carranza y Constitución.
564 Memoria para optar al título de Profesor de Estado en Ciencias Biológicas y Químicas,
565 Universidad de Chile. 76 p.
- 566 **Schultz O. 1987.** Taxonomische Neugruppierung der Oberfamilie Xiphoidea (Pisces,
567 Osteichthyes). *Annalen des Naturhistorischen Museums in Wien Serie A. für Mineralogie and*
568 *Petrographie, Geologie und Palaontologie, Anthropologie und Prahistorie.* **89**:95–202.
- 569 **Sernageomin, 2003.** Mapa Geológico de Chile. Versión Digital. Servicio Nacional de Geología
570 y Minería (Chile), Publicación geológica digital N° 4, Escala 1:1.000.000.
- 571 **Suárez ME, Otero RA. 2008.** Nuevos hallazgos de vertebrados marinos en el Campaniano-
572 Maastrichtiano de Loanco, VII Región. I Simposio Paleontología en Chile, November 2008.
573 Santiago, Actas 78-82.
- 574 **Sytchevskaya EK, Prokofiev AM. 2002.** First Findings of Xiphoidea (Perciformes) in the Late
575 Paleocene of Turkmenistan. *Journal of Ichthyology* **42**: 227–237.
- 576 **Tavera J. 1942.** Contribucion al estudio de la estratigrafía y paleontología del Terciario de
577 Arauco. *Anales del Primer Congreso Panamericano de Ingeniería en Minas y Geología*
578 (Santiago), Tomo 2, pp. 580–632.
- 579 **Tavera J. 1987.** Noticia sobre hallazgo de una extremidad de *Pliosaurus chilensis* Gay en la
580 localidad para la Formación Quiriquina de Faro Carranza (latitud 35°36'). Monografía del
581 Departamento de Geología, Universidad de Chile. 18 pp.

- Walsh AA. 2001. The Bahía Inglesa Formation Bonebed: Genesis and Palaeontology of a Neogene Konzentrat Lagerstätte from north-central Chile. University of Portsmouth, Doctoral Thesis, 440 p.
- Zambrano P, Nielsen S, Stinensbeck W. 2014. Taxonomía y relaciones tafonómicas en los nautiloideos paleógenos de Sudamérica. IV Simposio Paleontología en Chile. Universidad Austral de Chile, Valdivia, Chile. Libro de Resúmenes, p. 69.
- Zinsmeister W. 1979. Biogeographic significance of the Upper Mesozoic and early Tertiary molluscan faunas of Seymour Island (Antarctic Peninsula) to the final break-up of Gondwanaland. In: Gray J, Boucot AJ, eds., *Historical Biogeography, Plate Tectonics and the Changing Environment*. Oregon State University Press, Eugene, Oregon, 349–355.
- Zinsmeister W. 1982. Late Cretaceous-Early Tertiary Molluscan Biogeography of the Southern Circum-Pacific Journal of Paleontology 56:84–102.

FIGURE CAPTIONS

Figure 1: Locality and stratigraphy of SGO.PV.6634. (A) Map indicating the locality of Loanco, where SGO.PV.6634 was recovered. (B) Scheme of the transgressive-regressive deposits in the Arauco Basin, indicating the respective units. Modified from Charrier et al., 2007. (C) Stratigraphic column of the best Paleogene outcrop currently available (2016) in the coast south of Loanco. The unit is larger, however, this is covered by recent sands and by tsunami deposits after February 27, 2010. Sediment attached to SGO.PV.6634 is coincident with the fossil-bearing level containing *Aturia* sp. and *Imaizula araucana*.

Figure 2: Skull elements of SGO.PV.6634, holotype of *Loancorhynchus catrillancai* gen. et sp. nov. (A) Estimated body outline (based in *Fierstine and Monsch, 2002*). (B) Detail of the available skull elements of SGO.PV.6634. (C) Dorsal view of the rostral elements (D) Fin rays, with detail of the proximal part of the first right ray. **Anatomical abbreviations:** **ah**, articular head; **at**, anterior tuberosity; **dr**, dorsal ridges; **fr**, frontal; **ld**, left dentary; **lmx**, left maxillary; **lop**, left operculum; **pmx**, premaxillary; **pn?**, prenasals?; **rhy**, right hyomandibular; **rop**, right operculum.

Figure 3: *Loancorhynchus catrillancai* gen. et sp. nov. SGO.PV.6634, holotype. Distribution of villiform teeth and canali. (A) posterior part of the rostrum in lateral view. (B) detail of the villiform teeth present in the posterior part of the right premaxillary. (C) Left dentary in left lateral view. (D) detail of the villiform teeth in the anterior part of the left dentary. (E) Ventral view of the anterior rostrum fragment. (F, G) Detail of the internal paired canali in the rostrum.

619

620 **Figure 4: *Loancorhynchus catrillancai* gen. et sp. nov. SGO.PV.6634, holotype. Anatomy of**
 621 **the rostrum.** (A) Right view. (B) Line drawing interpretation of the anatomical elements. (C)
 622 same elements in dorsal view. (D) Line drawing interpretation of the anatomical elements. (E)
 623 anterior view of the anteriormost rostral fragment. (F) Scheme of the previous. **Anatomical**
 624 **abbreviations:** **deth**, dermethmoid; **lmx**, left maxillary; **lpm**, left premaxillary; **pln**,
 625 posterolateral notch; **pn?**, prenasals(?); **rpm**, right premaxillary; **vt**, villiform teeth.

626

627 **Figure 5: *Loancorhynchus catrillancai* gen. et sp. nov. SGO.PV.6634, holotype. Skull**
 628 **elements.** (A) Right hyomandibular/metapterygoid.in lateral (external) view. (B) Posterior view.
 629 (C) Anterior view. (D) Internal view. (E) Left frontal in dorsal view. (F) Same in ventral view.
 630 (G) Right opercle and supraopercular in right lateral view. (H) Left opercle in left lateral view.
 631 **Anatomical abbreviations:** **fhy**, facet for the hyomandibular; **hyo**, hyomandibular; **mpt**,
 632 metapterygoid; **opp**, opercular process; **or**, orbit; **pog**, preopercular groove; **ptf**, pterotic facet;
 633 **riop**, right interoperculum; **rsop**, right supraopercular; **spf**, sphenotic facet; **vii**, foramen for the
 634 *truncus hyoideomandibularis* of the facial nerve (VII).

635

636 **Figure 6: *Loancorhynchus catrillancai* gen. et sp. nov. SGO.PV.6634, holotype. Dentaries.**
 637 (A) Dentaries in ventral view. (B) Scheme of the previous. (C) Dentaries in occlusal view. (D)
 638 Left dentary (the most complete) in left lateral view. (E) Scheme of the previous. **Anatomical**
 639 **abbreviations:** **ld**, left dentary; **os**, occlusal surface; **rd**, right dentary; **vt**, villiform teeth.

640

Figure 7: Cladogram of the Xiphiioidei, including *Loancorhynchus catrillancai* gen. et sp. nov. Cladogram is based on the phylogenetic analysis performed here (Single MPC; CI=0.750; RI=0.600; Implied Weighting (K=3); New Technology Search; Ratchet. Pruned taxa: ‘*Blochius*’ *moorheadi*, *Cylindracanthus*, *Hemirhabdorrhynchus* and *Pseudotetrapturus luteus*). Biochrons are based on *Fierstine (2006)*. Body outlines are based in *Nakamura (1985)*. Skull schemes based in *Gregory and Conrad (1937)* and *Fierstine and Monsch (2002)*. Dorsal longitudinal ridges on the rostrum (Char. 7) supports *Loancorhynchus catrillancai* within the Blochiidae. Villiform teeth in the rostrum and dentary (Char. 27). separates *L. catrillancai* from *Blochius longirostris* and *Blochius macropterus*.

Figure 1

Locality and stratigraphy of SGO.PV.6634.

(A) Map indicating the locality of Loanco, where SGO.PV.6634 was recovered. (B) Scheme of the transgressive-regressive deposits in the Arauco Basin, indicating the respective units. Modified from Charrier et al., 2007. (C) Stratigraphic column of the best Paleogene outcrop currently available (2016) in the coast south of Loanco. The unit is larger, however, this is covered by recent sands and by tsunami deposits after February 27, 2010. Sediment attached to SGO.PV.6634 is coincident with the fossil-bearing level containing *Aturia* sp. and *Imaizula araucana*.

Figure 2

Skull elements of SGO.PV.6634, holotype of *Loancorhynchus catrillancai* gen. et sp. nov.

(A) Estimated body outline (based in *Fierstine and Monsch, 2002*). (B) Detail of the available skull elements of SGO.PV.6634. (C) Dorsal view of the rostral elements (D) Fin rays, with detail of the proximal part of the first right ray. **Anatomical abbreviations:** **ah**, articular head; **at**, anterior tuberosity; **dr**, dorsal ridges; **fr**, frontal; **ld**, left dentary; **lmx**, left maxillary; **lop**, left operculum; **pmx**, premaxillary; **pn?**, prenasals?; **rhy**, right hyomandibular; **rop**, right operculum.

Figure 3

Loancorhynchus catrillancai gen. et sp. nov. SGO.PV.6634, holotype. Distribution of villiform teeth and canali.

(A) posterior part of the rostrum in lateral view. (B) detail of the villiform teeth present in the posterior part of the right maxillary. (C) Left dentary in left lateral view. (D) detail of the villiform teeth in the anterior part of the left dentary. (E) Ventral view of the anterior rostrum fragment. (F, G) Detail of the internal paired canali in the rostrum.

Figure 4

Loancorhynchus catrillancai gen. et sp. nov. SGO.PV.6634, holotype. Anatomy of the rostrum.

(A) Right view. (B) Line drawing interpretation of the anatomical elements. (C) same elements in dorsal view. (D) Line drawing interpretation of the anatomical elements. (E) anterior view of the anteriormost rostral fragment. (F) Scheme of the previous. **Anatomical abbreviations:** **deth**, dermethmoid; **lmx**, left maxillary; **lpm**, left premaxillary; **pln**, posterolateral notch; **pn?**, prenasals(?); **rpm**, right premaxillary; **vt**, villiform teeth.

Figure 5

Loancorhynchus catrillancai gen. et sp. nov. SGO.PV.6634, holotype. Skull elements.

(A) Right hyomandibular/metapterygoid.in lateral (external) view. (B) Posterior view. (C) Anterior view. (D) Internal view. (E) Left frontal in dorsal view. (F) Same in ventral view. (G) Right opercle and supraopercular in right lateral view. (H) Left opercle in left lateral view.

Anatomical abbreviations: **fhy**, facet for the hyomandibular; **hyo**, hyomandibular; **mpt**, metapterygoid; **opp**, opercular process; **or**, orbit; **pog**, preopercular groove; **ptf**, pterotic facet; **riop**, right interoperculum; **rsop**, right supraopercular; **spf**, sphenotic facet; **vii**, foramen for the *truncus hyoideomandibularis* of the facial nerve (VII).

Figure 6

Loancorhynchus catrillancai gen. et sp. nov. SGO.PV.6634, holotype. Dentaries.

(A) Dentaries in ventral view. (B) Scheme of the previous. (C) Dentaries in occlusal view. (D) Left dentary (the most complete) in left lateral view. (E) Scheme of the previous.

Anatomical abbreviations: **ld**, left dentary; **os**, occlusal surface; **rd**, right dentary; **vt**, villiform teeth.

Figure 7

Cladogram of the Xiphioidae, including *Loancorhynchus catrillancai* gen. et sp. nov.

Cladogram is based on the phylogenetic analysis performed here (Single MPC; CI=0.750;

RI=0.600; Implied Weighting (K=3); New Technology Search; Ratchet. Pruned taxa:

'*Blochius*' *moorheadi*, *Cylindracanthus*, *Hemirhabdorrhynchus* and *Pseudotetrapturus luteus*).

Biochrons are based on *Fierstine (2006)*. Body outlines are based in *Nakamura (1985)*. Skull

schemes based in *Gregory and Conrad (1937)* and *Fierstine and Monsch (2002)*. Dorsal

longitudinal ridges on the rostrum (Char. 7) supports *Loancorhynchus catrillancai* within the

Blochidae. Villiform teeth in the rostrum and dentary (Char. 27). separates *L. catrillancai* from

Blochius longirostris and *Blochius macropterus*.