

Genome-wide characterization of the auxin response factor (ARF) gene family of litchi (*Litchi chinensis* Sonn.): phylogenetic analysis, miRNA regulation and expression changes during fruit abscission

Yan Q Zhang^{1,2,3}, Zao H Zeng^{1,2,3}, Cheng J Chen^{1,2,3}, Cai Q Li^{1,2,3}, Rui Xia^{Corresp., 1,2,3}, Jian G Li^{Corresp. 1,2,3}

¹ State Key Laboratory for Conservation and Utilization of Subtropical Agro-Bioresources, Guangzhou, China

² China Litchi Research Center, Guangzhou, China

³ Guangdong Litchi Engineering Research Center, College of Horticulture, South China Agricultural University, Guangzhou, China

Corresponding Authors: Rui Xia, Jian G Li
Email address: rxia@scau.edu.cn, jianli@scau.edu.cn

Auxin response factors (ARFs) play fundamental roles in modulating various biological processes including fruit development and abscission via regulating the expression of auxin response genes. Currently, little is known about roles of ARFs in litchi (*Litchi chinensis* Sonn.), an economical important subtropical fruit tree whose production is suffering from fruit abscission. In this study, a genome-wide analysis of ARFs was conducted for litchi, thirty-nine ARF genes (*LcARFs*) were identified. Conserved domain analysis showed that all the *LcARFs* identified have the signature B3 DNA-binding (B3) and ARF (Aux_rep) domains, with only 23 members having the dimerization domain (Aux_IAA). Number of exons in *LcARF* genes ranges from 2 to 16, suggesting a large variation for the gene structure of *LcARFs*. Phylogenetic analysis showed that the 39 *LcARFs* could be divided into three main groups: class I, II, III. Totally, twenty-three *LcARFs* were found to be potential targets of small RNAs, with three conserved and one novel miRNA-ARF (miRN43-ARF9) regulatory pathways discovered in litchi. Expression patterns were used to evaluate candidate *LcARFs* involved in various developmental processes, especially in flower formation and organ abscission. The results revealed that most ARF genes likely acted as repressors in litchi fruit abscission, i.e. *ARF2D/2E*, *7A/7B*, *9A/9B*, *16A/16B*, while a few *LcARFs*, such as *LcARF5A/B*, might positively involve in this process. These findings provide useful information and resources for further studies on the roles of ARF genes in litchi growth and development, especially in the process of fruit abscission.

Genome-wide characterization of the auxin response factor (ARF) gene family of litchi (*Litchi chinensis* Sonn.): phylogenetic analysis, miRNA regulation and expression changes during fruit abscission

Yanqing Zhang^{1,2,3}, Zaohai Zeng^{1,2,3}, Chengjie Chen^{1,2,3}, Caiqin Li^{1,2,3}, Rui Xia^{1,2,3*}, Jianguo Li^{1,2,3*}

¹ State Key Laboratory for Conservation and Utilization of Subtropical Agro-Bioresources

² China Litchi Research Center, Guangzhou, China

³ Guangdong Litchi Engineering Research Center, College of Horticulture, South China Agricultural University, Guangzhou, China

Corresponding Author:

Rui Xia^{1,2,3}

Yuehan Road 483, Wushan, Tianhe, Guangzhou, Guangdong Province, 510642, China

Email address: rxia@scau.edu.cn

Jianguo Li^{1,2,3}

Yuehan Road 483, Wushan, Tianhe Guangzhou, Guangdong Province, 510642, China

Email address: jianli@scau.edu.cn

Abstract

Auxin response factors (ARFs) play fundamental roles in modulating various biological processes including fruit development and abscission via regulating the expression of auxin response genes. Currently, little is known about roles of ARFs in litchi (*Litchi chinensis* Sonn.), an economical important subtropical fruit tree whose production is suffering from fruit abscission. In this study, a genome-wide analysis of ARFs was conducted for litchi, thirty-nine ARF genes (*LcARFs*) were identified. Conserved domain analysis showed that all the *LcARFs* identified have the signature B3 DNA-binding (B3) and ARF (Aux_rep) domains, with only 23 members having the dimerization domain (Aux_IAA). Number of exons in *LcARF* genes ranges from 2 to 16, suggesting a large variation for the gene structure of *LcARFs*. Phylogenetic analysis showed that the 39 *LcARFs* could be divided into three main groups: class I, II, III. Totally, twenty-three *LcARFs* were found to be potential targets of small RNAs, with three conserved and one novel miRNA-*ARF* (miRN43-*ARF9*) regulatory pathways discovered in litchi. Expression patterns were used to evaluate candidate *LcARFs* involved in various developmental processes, especially in flower formation and organ abscission. The results revealed that most ARF genes likely acted as repressors in litchi fruit abscission, i.e. *ARF2D/2E*, *7A/7B*, *9A/9B*, *16A/16B*, while a few *LcARFs*, such as *LcARF5A/B*, might positively involve in this process. These findings provide useful information and resources for further studies on the roles of ARF genes in litchi growth and development, especially in the process of fruit abscission.

Introduction

Auxin plays a central role in numerous aspects of plant developmental and physiological processes, including embryogenesis, apical dominance, vascular elongation, flowering, fruit development, and lateral root initiation (Woodward & Bartel, 2005; Fleming, 2006). Auxin response factors (ARFs) are a group of important transcription factors in the auxin signaling pathway, which can activate or repress the expression of early/primary auxin response genes by binding to the auxin response element (AuxRE) site in their promoter regions (Liscum & Reed, 2002; Guilfoyle & Hagen, 2007). A typical ARF is characterized by a highly conserved N-terminal B3-type DNA binding domain (DBD) that recognized the AuxRE motif, an activation domain (AD) or repression domain (RD), and a carboxy-terminal dimerization domain (CTD: domain III/IV), which is involved in protein-protein interactions by dimerizing with auxin/indole-3-acetic acid (Aux/IAA) family genes (Kim, Harter & Theologis, 1997; Guilfoyle & Hagen, 2007; Piya et al., 2014).

ARFs exert pivotal function in the regulation of plant growth and development through the auxin signaling pathway (Kepinski and Leyser, 2005; Li et al., 2016). Due to the significance of ARFs, genome-wide characterization of ARFs have been completed in many species such as model plants *Arabidopsis* (Hagen & Guilfoyle, 2002) and *Solanum lycopersicum* (Zouine et al., 2014), important crops such as soybean (Ha et al., 2013; Le et al., 2016) and maize (Xing et al., 2011; Wang et al., 2012), fruit trees like citrus (*Citrus sinensis*) (Xie et al., 2015) and apple (*Malus domestica*) (LUO et al., 2014), and so on. From embryogenesis to flowering, mutants in members of ARFs exhibit diverse phenotypes, which show their unique and redundant functions for plant development. For instance, in *Arabidopsis thaliana*, *arf1* and *arf2* mutations affect leaf senescence and floral organ abscission (Ellis et al., 2005) and the loss of *AtARF3* causes defectives in gynoecium patterning (Nemhauser, Feldman & Zambryski, 2000; Liu et al., 2014a). *AtARF5* influences embryo, root and shoot development (Krogan et al., 2012; Crawford et al., 2015). *AtARF9* acts in suspensor cells to mediate hypophysis specification (Rademacher et al., 2012), and *AtARF10/16/17* play vital roles in negatively regulating seed germination and post-germination activities (Liu et al., 2007).

Recently, small RNAs, especially miRNAs, have been emerging as critical regulators in almost all aspects of plant growth and development. Many members of the ARF family have been reported to be targets of miRNAs. In *Arabidopsis*, *AtARF6* and *AtARF8* are targets of miR167 (Wu, Tian & Reed, 2006), *AtARF10/16/17* are targets of miR160 (Liu et al., 2007, 2010), and

ARF2/3/4 are targets of trans-acting siRNAs (tasiRNAs) generated from miR390-targeted TAS3 gene (trans-acting siRNA gene 3) (Allen et al., 2005; Axtell et al., 2006). These targeting relationships of miRNA on ARF genes are widely conserved in land plants (Xia, Xu & Meyers, 2017) and also in horticultural plants (Chen et al., 2018b). It has been reported that down-regulation of *ARF6* and *ARF8* by miR167 leads to floral development defects and female sterility in tomatoes (Liu et al., 2014b). Down-regulation of sly-miR160, increasing the expression of its targets *SLARF10/16/17*, regulates auxin-mediated ovary patterning as well as floral organ abscission and lateral organ lamina outgrowth (Damodharan, Zhao & Arazi, 2016). In *Arabidopsis* and tomato, the tasiRNA-mediated regulation of *ARF2* involved in controlling the onset of leaf senescence and floral organ abscission (Ellis et al., 2005; Lim et al., 2010; Guan et al., 2014; Ren et al., 2017).

Litchi (*Litchi chinensis* sonn.), an important economic fruit trees in southern China, usually undergo serious fruit abscission before harvest, leading to a low yield. Many studies demonstrate *ARFs* play critical roles in regulating plant organ abscission (Ellis et al., 2005; Kuang et al., 2012; Guan et al., 2014; Xie et al., 2015; Xu et al., 2015), while which *ARFs* are involved or more important than the other in the fruit abscission in litchi remains elusive. Here, we identified 39 ARF genes in litchi. Gene structure, phylogeny and targeting relationship with miRNAs were characterized. The expression of *LcARFs* was examined in diverse tissues and in the process of fruit abscission which was induced by three different treatments. Among them, *ARF2D/2E*, *7A/7B*, *9A/9B*, *16A/16B* and *LcARF5A/B*, were found to be associated with litchi fruit abscission. Our results offer new knowledge and resources to study the function of plant ARF genes and their roles in the fruit abscission in litchi.

Materials & Methods

Plant materials and treatments

The young fruitlet used for RNA-seq in our study were collected from 9-year-old litchi trees (*Litchi. chinensis* Sonn. cv. ‘Feizixiao’) in an orchard located at South China Agricultural University (Guangzhou, China). Three treatments have been performed 25 day after anthesis. The three treatments included ethephon (ETH), girdling plus defoliation (GPD) and dipping in 20mg/L 2, 4-D for one minute after girdling plus defoliation (GPDD). Details can refer to Li et al (Li et al., 2015) and Peng et al (Peng et al., 2013). Samples of ‘Feizixiao’ (FZX) for qRT-PCR were obtained from the orchard of Guangzhou Fruit Research Institute (Guangdong, China). Tissues including fruit-bearing shoots (FBS), young leaves (YL), mature leaves (ML), female flower (FF), male flower (MF), sex undetermined flower (USF) and young fruit (YF) of 25 days after fertilization were collected from different directions of each tree. After separation, all tissues were quickly frozen in liquid nitrogen and stored at -80°C before usage.

Identification and phylogenetic analyses of LcARFs

Amino acid sequences of 25 and 23 ARFs from rice (Wang et al., 2007) and *Arabidopsis* (Okushima et al., 2005; Wang et al., 2007) were downloaded from UniProt (<http://www.uniprot.org/>) and 22 ARFs from tomato (Zouine et al., 2014) were downloaded from Sol Genomics Network (<https://solgenomics.net/>). These ARFs were used as bait to identify potential ARFs in litchi genome and obtained by Blast analysis in TBtools (Chen et al., 2018a), with the E-value $1e^{-10}$. All sequences were then further validated by conserved domain search against the CDD (<http://www.ncbi.nlm.nih.gov/cdd/>) and PFAM (<http://pfam.xfam.org/>) databases. Based on the optimized alignment of amino acid sequences of LcARFs proteins, achieved by TrimAL 1.3 (<http://phylemon2.bioinfo.cipf.es/index.html>) after the initial sequence alignment in ClustalX 2.1, a maximum likelihood (ML) tree was constructed in MEGA 7.0 software with a bootstrap of 1000 replicates. Intron and exon distribution patterns and genome structure were analyzed and visualized in TBtools (Chen et al., 2018a). The detail of predicted protein sequences of LcARFs were shown in Data S1 and S2.

LcARF targets of miRNAs and RNA-seq analysis

Most of the *ARF*-targeting miRNA/tasiRNAs were identified in our previous study (Ma et al., 2018). To further validate the targeting relationship, target sites were verified by psRNATarget (<http://plantgrn.noble.org/psRNATarget>) (Dai & Zhao, 2011) and miRBase (<http://www.mirbase.org/>).

RNA-seq analysis was carried on as described previously (Li et al., 2013). These RNA-seq data was normalized using TPM method (Wagner, Kin & Lynch, 2012). Based on the sequences of the identified *LcARFs*, a BLASTN search was conducted to find their gene counterparts in previous data (Li et al., 2013). Finally, a global gene expression profiles were visualized by heatmap via TBtools software (Chen et al., 2018a).

Quantitative RT-PCR

Total RNA of above seven tissue samples was extracted by column plant RNAout kit (TIANDZ, China) according to the manufacturer's instructions. About 2 µg total RNA was applied to synthesized first strand cDNA using reverse transcriptase AT311-03 (TRANSGEN BIOTECH, China). PCR primers (Table S1) of *LcARFs* and reference genes *GAPDH* and *EF* (Zhong et al., 2011) were designed by Primer Premier 5.0. qRT-PCR was performed according to the manufacturer's specifications of THUNDERBIRD qPCR MIX QPS-201 (TOYOBO, China) on a LightCycler 480 (Roche, Switzerland). Each expression profile was independently verified in three biological replicates. Relative expression level of each gene was calculated by the $2^{-\Delta\Delta Ct}$ method (Livak & Schmittgen, 2001). Significance analysis was conducted in SPSS vision 22.0 and visulized by SigmaPlot 12.5.

Results

Identification and phylogenetic analysis of LcARF genes

To identify all ARF members in litchi, protein sequences of ARFs in rice, *Arabidopsis* and tomato were used as queries in BLASTP to search against litchi annotated gene database. Sixty-eight potential ARF genes were identified. After redundant result elimination and further conserved domain validation, 39 LcARFs were obtained ultimately. Among them, 23 protein sequences contained B3, ARF and the Aux/IAA domains. Length of these litchi ARF proteins ranged from 260 (LcARF1B) to 1200 (LcARF4A) and the relative molecular mass of them varied from 29522.5 (LcARF1B) to 144936.9 Da (LcARF16C), with PIs in the range of 5.28 (LcARF16C) to 8.64 (LcARF19A) (Table S2). According to the subcellular localization predictor ([CELLO v.2.5; http://cello.life.nctu.edu.tw](http://cello.life.nctu.edu.tw)) prediction, most LcARFs were predicted to be located in the nucleus.

A phylogenetic tree was generated using the maximum likelihood method based on alignment of litchi ARFs with their orthologs from rice, *Arabidopsis* and tomato (Fig. 1). All 109 ARFs fell into three broad groups: class I, II and III, which contained 53, 34 and 22 members, respectively. Obviously, most ARFs of the four species were clustered together in the first two classes. Litchi ARFs were named according to their positions with orthologs from the other three species in the tree. Thus, the 39 LcARFs could also be assigned to three separate clusters as well. Class I contained LcARF1/2/3/4/9/18 (1A/B/C/D, 2A/B/C/D/E/F, 3A/B, 4A/B/C, 9A/B, 18A/B); Class II included LcARF5/6/7/8/19 (5A/B, 6A/B/C/D, 7A/B, 8A/B, 19A/B); Class III included LcARF10/16/17 (10A/B, 16A/B/C/D, 17A/B).

Gene structure and conserved domains of litchi ARFs

To better understand the structure evolution of LcARF genes, their gene structure (intron/exon number and positions) and functional domains were analyzed. As shown in Fig. 2, 23 *LcARFs* harbored the typical ARF protein structure which composed of a highly conserved DNA-binding domain (DBD) in the N-terminal region with a plant specific B3-type subdomain, an Auxin-resp subdomain, and an AUX_IAA dimerization subdomains. The remaining 16 *LcARFs* contained only B3 and Auxin_resp subdomains. Gene structure analysis revealed that the exon number of class III (2-4 exons) was significantly less than the other two groups (8-17 exons for class II and 10-16 exons for class I). These results provided additional evidence confirming the phylogenetic relationships among LcARFs.

Analyses of miRNA targeting *LcARFs*

Twenty-two out of the 39 *LcARF* genes were found to be the targets of miRNA (Fig. 3). All group members of class III (*LcARF10A/B*, *16A/B/C/D* and *17A/B*) were found to be targeted by Lc-miR160. *LcARF6A/B/C/D* and *8A/B* were members of class II and all of them were collectively targeted by Lc-miR167. Two kinds of miRNA targeting patterns were observed in the class I. One group comprising *LcARF2E/2F/4B/4C/3A/3B* were targeted by tasiARFs. In litchi, *LcTAS3* are divided into two subgroups, long TAS3 genes(*LcTAS3_1* and *LcTAS3_2*) and short TAS3 genes (*LcTAS3_3* and *LcTAS3_4*), which trigger to produce two or one tasiARFs when cleaved by miR390 (Ma et al., 2018). Additionally, *LcARF2E/2F* incorporated one target site of tasiARF, while *LcARF3A/3B/4B/4C* contained two, which is in accordance with our previous study (Xia, Xu & Meyers, 2017). Interestingly, in the other group, a novel miRNA-*ARF* pathway was discovered, in which *LcARF9A/B* was targeted by the Lc-miRN43.

Expression of *LcARF* Genes in different organs and tissues

A large body of evidence support the importance of ARF genes in plant growth and development (Ellis et al., 2005; Guilfoyle & Hagen, 2007; Lim et al., 2010; Li et al., 2016). To explore how *LcARF* genes function in the development of litchi, we examined their expression levels in various litchi organs/tissues by qRT-PCR. Seven organs/tissues samples, composing fruit-bearing shoots (FBS), male flower (MF), female flower (FF), sex undetermined flower (USF), mature leaves (ML), young leaves (YL) and young fruits (YF) in ‘FZX’, were collected and tested. As shown in Fig. 4A, expression of all *ARFs* was detected at least in several tissues. Generally, most *LcARFs* show low expression in YL and YF but high in USF. *LcARFs* from different clades showed various expression patterns. *LcARFs* from class I and III were detected to be higher expressed in ML while those from class II were lowly accumulated, suggesting that these *LcARFs* in different classes may perform different functions in mature leaves development (Fig. 4A). Additionally, as is shown in Fig. 4B, ten *LcARFs* (*LcARF2B*, *3A*, *4A/B*, *5A/B*, *6D*, *9A/B*, *18B*) were significantly higher expressed in FF than MF, implying their potential function in ovule and ovary derived from female flowers. Remarkably, three *LcARFs* (*LcARF8A*, *LcARF10A/B*) were of significantly higher expression in MF than FF, indicated that they might play roles in male flower formation. Additionally, a table for *LcARFs* and their potential function was concluded in Data S3 according to early study which may be useful for further functional validation.

Expression profiles of *LcARF* genes in response to ETH, GPD, and GPDD treatment

Auxin is a critical signal in the abscission of fruits in plants (Blanusa et al., 2005; Meir et al., 2010; Xie et al., 2013). To explore the role of *LcARFs* in fruitlet abscission in litchi, RNA-seq was carried on and transcription levels of *LcARF* genes in the abscission zone were investigated under three treatments (ETH, GPD, GPDD). Both the ETH and GPD treatments promote fruitlet abscission while GPDD delays the process (Peng et al., 2013; Li et al., 2015). As the results in Fig. 5, the transcript expression of most *LcARFs* were decreased after treated by ETH and GPD, but upsurge after GPDD treatment, which was corresponding to the process of promoting or inhibiting of fruitlet shedding, respectively. It seemed that there was a negative correlation between *LcARFs* and the fruit abscission and among these *LcARFs*, those from two groups (group 2 and group 3) were particularly representative. Moreover, *LcARFs* from group 3 were more sensitive to GPDD treatment along with stronger expression than those from group 2. Thus, we could deduce that *LcARFs* from group 3, including *ARF2D/2E*, *7A/7B*, *9A/9B*, *16A/16B*, played major roles on the litchi fruitlet abscission. In contrast, a few *LcARFs*, such as group 1 including *LcARF5A/5B*, showed opposite expression pattern, suggesting that they might function to accelerate the process of abscission. There were as well some *LcARFs* with no significant expression change and seemed to be unrelated to fruitlet abscission.

Discussions

Litchi is an important tropic fruit trees and massive fruit abscission before harvest usually leads to low and even no production. Auxin is proposed to be one of the endogenous hormones playing significant roles in the regulation of fruit abscission in litchi (Yuan, 1988; Stern & Gazit, 2000). In this study, 39 LcARF genes were identified, which was more than the other model plants, such as *Arabidopsis* (23) (Okushima et al., 2005; Wang et al., 2007), rice (25) (Wang et al., 2007) and tomato (22) (Zouine et al., 2014), implying extensive duplication and diversification of the ARF gene family in litchi. Analysis of conserved motifs revealed that all LcARFs had a typical DBD domain required for efficient binding to AuxRE and a Auxin_resp (Fig. 2) (Hagen & Guilfoyle, 2002; Ha et al., 2013). However, only 23 of 39 LcARFs contain AUX_IAA domain, which can mediate the dimerization of ARFs or ARF and Aux/IAA protein (Guilfoyle & Hagen, 2007). Lack of the AUX_IAA domain for dimerization makes it interesting to address questions like how these ARFs function and whether they need dimerization with other proteins. In plants, ARFs can function as transcription activators (ADs) or repressors (RDs), according to the amino acid composition of Auxin_resp domain (Guilfoyle & Hagen, 2007). In *Arabidopsis thaliana*, ARF ADs and RDs were proposed to contain biased amino acid sequences, which ARF ADs were enriched in glutamine (Q), while RDs were enriched in serine (S), serine and proline (SP), serine glycine (SG) (Guilfoyle & Hagen, 2001; Tiwari, Hagen & Guilfoyle, 2003). Intriguingly, no ARFs in litchi were enriched in Q, but with SPL and SP/SG enrichment; then none of LcARF proteins seems to be an activator (reviewed in (Guilfoyle & Hagen, 2007)). Further experiments are needed to verify this observation.

Much evidence demonstrates that miRNAs play essential roles in post-transcriptional gene regulation in plants (Jones-Rhoades, Bartel & Bartel, 2006; Li & Zhang, 2016). It has been found that several ARF genes are regulated by a few miRNAs. In our work, 22 out of 39 LcARF genes were found to be targets of miRNAs (Fig. 3) and *LcARFs* from different classes were displayed different miRNA targeting pattern. Members of class III (*LcARF10A/B*, *16A/B/C/D* and *17A/B*) were found to be targeted by Lc-miR160, which might affect flower development of litchi, as down-regulation of *ARF10/16/17* by miRNA160 is reported to regulate floral organ abscission in tomato (Damodharan, Zhao & Arazi, 2016). Consistent with *Arabidopsis*, *LcARF6/8* were collectively targeted by Lc-miR167. Interestingly, even though there has been reported *LcARF8B* to be targeted by Lc-miR167 (Ma et al., 2018), *LcARF8A* was a novel target by Lc-miR167. Notably, in another group, a novel miRNA-ARF pathway was discovered in litchi, in which *LcARF9A/B* was targeted by Lc-miRN43. In fact, miRN43 is an innovative miRNA as well, for it

is unable to find any ortholog in the database of miRase, which may provide a new idea for us to study the specific functions of litchi.

The expression profiles can help us screen for candidate *LcARF* genes with potentially distinct functions. Most *LcARFs* were higher accumulated in mature leaves than young leaves including *LcARF2*, and its Arabidopsis homolog AtARF2 has been reported to regulate leaf senescence with high expression in mature leaves (Ellis et al., 2005). Thus we can deduce that *LcARF2* is likely to involve in leaf senescence in litchi. It has been reported that Arabidopsis ARF genes regulate flower formation, especially gametophyte development, which are critical for sexual reproduction. AtARF2–AtARF4 and AtARF5 play significant roles in regulating both female and male gametophyte development in Arabidopsis (Liu et al., 2017) and ARF6 and ARF8 regulate both stamen and gynoecium maturation (Nagpal, 2005; Ru et al., 2006; Wu, Tian & Reed, 2006; Liu et al., 2014b). Thus their litchi homologs *LcARF2B*, 3A, 4A/B, 5A/B, 6D and 8A may be important for female and male gametophyte development in litchi, along with significantly higher accumulation in female or male flower. Moreover, *LcARF4*, 5 and 9 were prominent higher expressed in female flower than male flower, implying their potential function in ovule and ovary development.

ARF genes have been reported to be involved in plant organ abscission. Overexpression of *SlARF2* in tomato results in flower organ senescence (Ren et al., 2017) and *SlARF1*, 2, 7, 11 and 19 showed overlapping functions in tomato abscission (Guan et al., 2014). Similar roles of ARFs in abscission were observed in *Arabidopsis* (Ellis et al., 2005). Our previous studies show that the treatment of girdling plus defoliation (GPD) in litchi could reduce the transcript level of auxin response factor (*LcARF1*) mRNA, along with the increase of fruitlet abscission (Kuang et al., 2012). Here in our RNA-seq survey of ARF genes expression, we found that most of the ARF genes show opposite correlation with the fruit abscission, i.e., ARF genes were down-regulated by abscission induced treatments (ETH and GPD), but up-regulated by GPDD, in which the abscission of GPD was inhibited by the addition of 2, 4-D, suggesting that the majority of *LcARF* genes are negatively involved in litchi fruit abscission, especially *LcARFs* from group 3 (*ARF2D/2E*, 7A/7B, 9A/9B, 16A/16B), with more prominent expression after GPDD treatment. By contrast, a few *LcARFs*, such *LcARF5A/5B*, show positive correlation with fruit abscission, indicating that they might serve as contributing factors to fruit shedding.

Conclusions

In this study, a total of 39 ARF genes were identified from the litchi genome. Comprehensive analyses, including phylogenetic relationship, exon-intron structure, conserved domain, and potential targets for small RNAs, revealed that the ARF gene family was expanded in litchi with species-specific features. A novel miRNA-*ARF* (miRN43-*ARF9*) regulatory pathway was discovered, which is likely specific in litchi. Expression profiles in various organs and under different abscission-related treatments (ETH, GPD and GPDD) uncovered the expression diversity of these ARF genes in litchi. Some ARF genes, including *ARF2D/2E*, *7A/7B*, *9A/9B*, *16A/16B* and *5A/5B*, likely play predominant roles in the process of litchi fruit abscission. These findings provide new knowledge and resources for further functional characterization of ARF genes in litchi.

References:

- Addicott FT, Lynch RS. 1955. Physiology of abscission. *Annual Review of Plant Physiology and Plant Molecular Biology* 6:211–238. DOI: 10.1146/annurev.pp.06.060155.001235.
- Allen E, Xie Z, Gustafson AM, Carrington JC. 2005. microRNA-directed phasing during trans-acting siRNA biogenesis in plants. *Cell* 121:207–221. DOI: 10.1016/j.cell.2005.04.004.
- Axtell MJ, Jan C, Rajagopalan R, Bartel DP. 2006. A Two-Hit Trigger for siRNA Biogenesis in Plants. *Cell* 127:565–577. DOI: 10.1016/j.cell.2006.09.032.
- Blanusa T, Else MA, Atkinson CJ, Davies WJ. 2005. The regulation of sweet cherry fruit abscission by polar auxin transport. *Plant Growth Regulation* 45:189–198. DOI: 10.1007/s10725-005-3568-9.
- Chen C, Xia R, Chen H, He Y. 2018a. TBtools, a Toolkit for Biologists integrating various HTS-data handling tools with a user-friendly interface. *bioRxiv*:289660. DOI: 10.1101/289660.
- Chen C, Zeng Z, Liu Z, Xia R. 2018b. Small RNAs, emerging regulators critical for the development of horticultural traits. *Horticulture Research* 5:6–8. DOI: 10.1038/s41438-018-0072-8.
- Crawford BCW, Sewell J, Golembeski G, Roshan C, Long JA, Yanofsky MF. 2015. Genetic control of distal stem cell fate within root and embryonic meristems. *Science* 347:655–660.
- Dai X, Zhao PX. 2011. PsRNATarget: A plant small RNA target analysis server. *Nucleic Acids Research* 39:155–159. DOI: 10.1093/nar/gkr319.
- Damodharan S, Zhao D, Arazi T. 2016. A common miRNA160-based mechanism regulates ovary patterning, floral organ abscission and lamina outgrowth in tomato. *Plant Journal* 86:458–471. DOI: 10.1111/tpj.13127.
- Ellis CM, Nagpal P, Young JC, Hagen G, Guilfoyle TJ, Reed JW. 2005. AUXIN RESPONSE FACTOR1 and AUXIN RESPONSE FACTOR2 regulate senescence and floral organ abscission in *Arabidopsis thaliana*. *Development* 132:4563–4574. DOI: 10.1242/dev.02012.
- Fleming AJ. 2006. Plant signalling: the inexorable rise of auxin. *Trends in Cell Biology* 16:397–402. DOI: 10.1016/j.tcb.2006.06.005.
- Guan X, Xu T, Gao S, Qi M, Wang Y, Liu X, Li T. 2014. Temporal and Spatial Distribution of Auxin Response Factor Genes During Tomato Flower Abscission. *Journal of Plant Growth Regulation* 33:317–327. DOI: 10.1007/s00344-013-9377-x.
- Guilfoyle TJ, Hagen G. 2001. Auxin response factors. *Journal of Plant Growth Regulation* 20:281–291. DOI: 10.1007/s003440010026.
- Guilfoyle TJ, Hagen G. 2007. Auxin response factors. *Current Opinion in Plant Biology* 10:453–460. DOI: 10.1016/j.pbi.2007.08.014.

351 Guilfoyle TJ, Key JL. 1986. Auxin-regulated gene expression in higher plants. *Critical Reviews*
352 *in Plant Sciences* 4:247–276. DOI: 10.1080/07352688609382226.

353 Ha C Van, Le DT, Nishiyama R, Watanabe Y, Sulieman S, Tran UT, Mochida K, Dong N Van,
354 Yamaguchi-Shinozaki K, Shinozaki K, Tran L-SP. 2013. The auxin response factor transcription
355 factor family in soybean: genome-wide identification and expression analyses during
356 development and water stress. *DNA research : an international journal for rapid publication of*
357 *reports on genes and genomes* 20:511–24. DOI: 10.1093/dnares/dst027.

358 Hagen G, Guilfoyle T. 2002. Auxin-responsive gene expression: Genes, promoters and
359 regulatory factors. *Plant Molecular Biology* 49:373–385. DOI: 10.1023/A:1015207114117.

360 Jones-Rhoades MW, Bartel DP, Bartel B. 2006. MicroRNAs AND THEIR REGULATORY
361 ROLES IN PLANTS. *Annual Review of Plant Biology* 57:19–53. DOI:
362 10.1146/annurev.arplant.57.032905.105218.

363 Kepinski S, Leyser O. 2005. The Arabidopsis F-box protein TIR1 is an auxin receptor. *Nature*
364 435:446–451. DOI: 10.1038/nature03542.

365 Kim J, Harter K, Theologis A. 1997. Protein-protein interactions among the Aux/IAA proteins.
366 *Proceedings of the National Academy of Sciences of the United States of America* 94:11786–91.
367 DOI: 10.1073/pnas.94.22.11786.

368 Krogan NT, Ckurshumova W, Marcos D, Caragea AE, Berleth T. 2012. Deletion of MP/ARF5
369 domains III and IV reveals a requirement for Aux/IAA regulation in Arabidopsis leaf vascular
370 patterning. *New Phytologist* 194:391–401. DOI: 10.1111/j.1469-8137.2012.04064.x.

371 Kuang JF, Wu JY, Zhong HY, Li CQ, Chen JY, Lu WJ, Li JG. 2012. Carbohydrate stress
372 affecting fruitlet abscission and expression of genes related to auxin signal transduction pathway
373 in litchi. *International Journal of Molecular Sciences* 13:16084–16103. DOI:
374 10.3390/ijms131216084.

375 Le B, Nawaz MA, Rehman HM, Le T, Yang SH, Golokhvast KS, Son E, Chung G. 2016.
376 Genome-wide characterization and expression pattern of auxin response factor (ARF) gene
377 family in soybean and common bean. *Genes and Genomics* 38:1165–1178. DOI:
378 10.1007/s13258-016-0462-y.

379 Li C, Wang Y, Huang X, Li J, Wang H, Li J. 2013. De novo assembly and characterization of
380 fruit transcriptome in Litchi chinensis Sonn and analysis of differentially regulated genes in fruit
381 in response to shading. *BMC Genomics* 14:1. DOI: 10.1186/1471-2164-14-552.

382 Li C, Wang Y, Ying P, Ma W, Li J. 2015. Genome-wide digital transcript analysis of putative
383 fruitlet abscission related genes regulated by ethephon in litchi. *Frontiers in Plant Science* 6:1–
384 16. DOI: 10.3389/fpls.2015.00502.

385 Li S-B, Xie Z-Z, Hu C-G, Zhang J-Z. 2016. A Review of Auxin Response Factors (ARFs) in
386 Plants. *Frontiers in plant science* 7:47. DOI: 10.3389/fpls.2016.00047.

387 Li C, Zhang B. 2016. MicroRNAs in Control of Plant Development. *Journal of Cellular*
388 *Physiology* 231:303–313. DOI: 10.1002/jcp.25125.

389 Lim PO, Lee IC, Kim J, Kim HJ, Ryu JS, Woo HR, Nam HG. 2010. Auxin response factor 2
390 (ARF2) plays a major role in regulating auxin-mediated leaf longevity. *Journal of Experimental*
391 *Botany* 61:1419–1430. DOI: 10.1093/jxb/erq010.

392 Liscum E, Reed JW. 2002. Genetics of Aux/IAA and ARF action in plant growth and
393 development. *Plant Molecular Biology* 49:387–400. DOI: 10.1023/A:1015255030047.

394 Liu X, Dinh TT, Li D, Shi B, Li Y, Cao X, Guo L, Pan Y, Jiao Y, Chen X. 2014a. AUXIN
395 RESPONSE FACTOR 3 integrates the functions of AGAMOUS and APETALA2 in floral
396 meristem determinacy. *Plant Journal* 80:629–641. DOI: 10.1111/tpj.12658.

397 Liu X, Huang J, Wang Y, Khanna K, Xie Z, Owen HA, Zhao D. 2010. The role of floral organs
398 in carpels, an Arabidopsis loss-of-function mutation in MicroRNA160a, in organogenesis and
399 the mechanism regulating its expression. *Plant Journal* 62:416–428. DOI: 10.1111/j.1365-
400 313X.2010.04164.x.

401 Liu Z, Miao L, Huo R, Song X, Johnson C, Kong L, Sundaresan V, Yu X. 2017. ARF2–ARF4
402 and ARF5 are Essential for Female and Male Gametophyte Development in Arabidopsis. *Plant*
403 *and Cell Physiology* 59:179–189. DOI: 10.1093/pcp/pcx174.

404 Liu PP, Montgomery TA, Fahlgren N, Kasschau KD, Nonogaki H, Carrington JC. 2007.
405 Repression of AUXIN RESPONSE FACTOR10 by microRNA160 is critical for seed
406 germination and post-germination stages. *Plant Journal* 52:133–146. DOI: 10.1111/j.1365-
407 313X.2007.03218.x.

408 Liu N, Wu S, Houten J Van, Wang Y, Ding B, Fei Z, Clarke TH, Reed JW, Van Der Knaap E.
409 2014b. Down-regulation of AUXIN RESPONSE FACTORS 6 and 8 by microRNA 167 leads to
410 floral development defects and female sterility in tomato. *Journal of Experimental Botany*
411 65:2507–2520. DOI: 10.1093/jxb/eru141.

412 Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using real-time
413 quantitative PCR and the $2^{-\Delta\Delta CT}$ method. *Methods* 25:402–408. DOI: 10.1006/meth.2001.1262.

414 Louie DS, Addicott FT. 1970. Applied auxin gradients and abscission in explants. *Plant*
415 *physiology* 45:654–657.

416 LUO X-C, SUN M-H, XU R-R, SHU H-R, WANG J-W, ZHANG S-Z. 2014. Genomewide
417 identification and expression analysis of the ARF gene family in apple. *Journal of Genetics*
418 93:785–797. DOI: 10.1007/s12041-014-0462-0.

419 Ma W, Chen C, Liu Y, Zeng M, Meyers BC, Li J, Xia R. 2018. Coupling of microRNA-directed
420 phased small interfering RNA generation from long noncoding genes with alternative splicing
421 and alternative polyadenylation in small RNA-mediated gene silencing. *New Phytologist*
422 217:1535–1550. DOI: 10.1111/nph.14934.

423 Meir S, Philosoph-Hadas S, Sundaresan S, Selvaraj KS V., Burd S, Ophir R, Kochanek B, Reid
424 MS, Jiang C-Z, Lers A. 2010. Microarray Analysis of the Abscission-Related Transcriptome in
425 the Tomato Flower Abscission Zone in Response to Auxin Depletion. *Plant Physiology*
426 154:1929–1956. DOI: 10.1104/pp.110.160697.

427 Nagpal P. 2005. Auxin response factors ARF6 and ARF8 promote jasmonic acid production and
428 flower maturation. *Development* 132:4107–4118. DOI: 10.1242/dev.01955.

429 Nemhauser JL, Feldman LJ, Zambryski PC. 2000. Auxin and ETTIN in Arabidopsis gynoecium
430 morphogenesis. *Development* 127:3877–3888.

431 Okushima Y, Overvoorde PJ, Arima K, Alonso JM, Chan A, Chang C, Ecker JR, Hughes B, Lui
432 A, Nguyen D, Onodera C, Quach H, Smith A. 2005. Functional Genomic Analysis of the
433 AUXIN RESPONSE FACTOR Gene Family Members in Arabidopsis thaliana. *The Plant Cell*
434 17:444–463. DOI: 10.1105/tpc.104.028316.2.

435 Pattison RJ, Csukasi F, Catalá C. 2014. Mechanisms regulating auxin action during fruit
436 development. *Physiologia Plantarum* 151:62–72. DOI: 10.1111/ppl.12142.

437 Peng G, Wu J, Lu W, Li J. 2013. A polygalacturonase gene clustered into clade E involved in
438 lychee fruitlet abscission. *Scientia Horticulturae* 150:244–250. DOI:
439 10.1016/j.scienta.2012.10.029.

440 Peng M, Ying P, Liu X, Li C, Xia R, Li J, Zhao M. 2017. Genome-Wide Identification of
441 Histone Modifiers and Their Expression Patterns during Fruit Abscission in Litchi. *Frontiers in*
442 *Plant Science* 8:1–16. DOI: 10.3389/fpls.2017.00639.

443 Piya S, Shrestha SK, Binder B, Stewart CN, Hewezi T. 2014. Protein-protein interaction and
444 gene co-expression maps of ARFs and Aux/IAs in Arabidopsis. *Frontiers in Plant Science* 5:1–
445 9. DOI: 10.3389/fpls.2014.00744.

446 Rademacher EH, Lokerse AS, Schlereth A, Llavata-Peris CI, Bayer M, Kientz M, FreireRios A,
447 Borst JW, Lukowitz W, Jürgens G, Weijers D. 2012. Different Auxin Response Machineries
448 Control Distinct Cell Fates in the Early Plant Embryo. *Developmental Cell* 22:211–222. DOI:
449 10.1016/j.devcel.2011.10.026.

450 Ren Z, Liu R, Gu W, Dong X. 2017. The Solanum lycopersicum auxin response factor SlARF2
451 participates in regulating lateral root formation and flower organ senescence. *Plant Science*
452 256:103–111. DOI: 10.1016/j.plantsci.2016.12.008.

453 Ru P, Xu L, Ma H, Huang H. 2006. Plant fertility defects induced by the enhanced expression of
454 microRNA167. *Cell Research* 16:457–465. DOI: 10.1038/sj.cr.7310057.

455 Stern RA, Gazit S. 2000. Application of the polyamine putrescine increased yield, of “Mauritius”
456 litchi (*Litchi chinensis* Sonn.). *Journal of Horticultural Science and Biotechnology* 75:612–614.
457 DOI: 10.1080/14620316.2000.11511295.

458 Sundberg E, Østergaard L. 2009. Distinct and dynamic auxin activities during reproductive
459 development. *Cold Spring Harbor perspectives in biology* 1:1–15. DOI:
460 10.1101/cshperspect.a001628.

461 Tiwari SB, Hagen G, Guilfoyle T. 2003. The Roles of Auxin Response Factor Domains in
462 Auxin-Responsive Transcription. *Plant Cell* 15:533–543. DOI: 10.1105/tpc.008417.All.

463 Wagner GP, Kin K, Lynch VJ. 2012. Measurement of mRNA abundance using RNA-seq data:
464 RPKM measure is inconsistent among samples. *Theory in Biosciences* 131:281–285. DOI:
465 10.1007/s12064-012-0162-3.

466 Wang Y, Deng D, Shi Y, Miao N, Bian Y, Yin Z. 2012. Diversification, phylogeny and
467 evolution of auxin response factor (ARF) family: Insights gained from analyzing maize ARF
468 genes. *Molecular Biology Reports* 39:2401–2415. DOI: 10.1007/s11033-011-0991-z.

469 Wang D, Pei K, Fu Y, Sun Z, Li S, Liu H, Tang K, Han B, Tao Y. 2007. Genome-wide analysis
470 of the auxin response factors (ARF) gene family in rice (*Oryza sativa*). *Gene* 394:13–24. DOI:
471 S0378-1119(07)00032-7 [pii]r10.1016/j.gene.2007.01.006.

472 Woodward AW, Bartel B. 2005. Auxin: Regulation, action, and interaction. *Annals of Botany*
473 95:707–735. DOI: 10.1093/aob/mci083.

474 Wu M-F, Tian Q, Reed JW. 2006. Arabidopsis microRNA167 controls patterns of ARF6 and
475 ARF8 expression, and regulates both female and male reproduction. *Development* 133:4211–
476 4218. DOI: 10.1242/dev.02602.

477 Xia R, Xu J, Meyers BC. 2017. The Emergence, Evolution, and Diversification of the miR390-
478 TAS3-ARF Pathway in Land Plants. *The Plant Cell* 29:tpc.00185.2017. DOI:
479 10.1105/tpc.17.00185.

480 Xie RJ, Deng L, Jing L, He SL, Ma YT, Yi SL, Zheng YQ, Zheng L. 2013. Recent advances in
481 molecular events of fruit abscission. 57:201–209. DOI: 10.1007/s10535-012-0282-0.

482 Xie R, Pang S, Ma Y, Deng L, He S, Yi S, Lv Q, Zheng Y. 2015. The ARF, AUX/IAA and GH3
483 gene families in citrus: genome-wide identification and expression analysis during fruitlet drop
484 from abscission zone A. *Molecular Genetics and Genomics* 290:2089–2105. DOI:
485 10.1007/s00438-015-1063-1.

486 Xing H, Pudake RN, Guo G, Xing G, Hu Z, Zhang Y, Sun Q, Ni Z. 2011. Genome-wide

487 identification and expression profiling of auxin response factor (ARF) gene family in maize.
 488 *BMC Genomics* 12:178. DOI: 10.1186/1471-2164-12-178.
 489 Xu T, Wang Y, Liu X, Lv S, Feng C, Qi M, Li T. 2015. Small RNA and degradome sequencing
 490 reveals microRNAs and their targets involved in tomato pedicel abscission. *Planta* 242:963–984.
 491 DOI: 10.1007/s00425-015-2318-0.
 492 Yuan R. 1988. Litchi Fruit Abscission: its Patterns, Effect of Shading and Relation to
 493 Endogenous Absciscic Acid 1. 36:281–292.
 494 Zhong H-Y, Chen J-W, Li C-Q, Chen L, Wu J-Y, Chen J-Y, Lu W-J, Li J-G. 2011. Selection of
 495 reliable reference genes for expression studies by reverse transcription quantitative real-time
 496 PCR in litchi under different experimental conditions. *Plant cell reports* 30:641–53. DOI:
 497 10.1007/s00299-010-0992-8.
 498 Zouine M, Fu Y, Chateigner-Boutin AL, Mila I, Frasse P, Wang H, Audran C, Roustan JP,
 499 Bouzayen M. 2014. Characterization of the tomato ARF gene family uncovers a multi-levels
 500 post-transcriptional regulation including alternative splicing. *PLoS ONE* 9:1–12. DOI:
 501 10.1371/journal.pone.0084203.
 502

Figure 2

Phylogenetic relationship, exon-intron structure, conserved domains analyses of LcARFs.

(A) Phylogenetic relationship among the litchi ARF proteins. The unrooted tree was generated using the maximum likelihood method by JTT matrix-based model. The reliability was assessed using 1000 bootstrap replicates. Three clusters are labeled as Class I, Class II and Class III. **(B) Exon- intron structure and conserved domains of LcARFs.**

Information of exon, intron, and functional domain was obtained from model gene annotation and results of NCBI CDD search and visualized by TBtools. B3: B3 DNA-binding domain; Auxin-resp: ARF domain; AUX_IAA: C-terminal dimerization domain. Lengths of exons and introns and domains of each LcARF protein were exhibited proportionally.




Figure 3

LcARFs targeted by some *LcmiRNAs*.

miR160 targets *LcARF10/16/17*, miR167 targets *LcARF6/8* and miRN43 targets *LcARF9*. Additionally, the *LcTAS3* was targeted by miR390 and then triggered the production of tasiARF which target *LcARF2E/2F/4B/4C/3A/3B*.




Figure 4

Expression Profiles of LcARF Genes in various tissues of 'FZX' by qRT-PCR.

A) Expression of all *LcARFs* in different tissues. The heatmap was generated based on the relative expression values of 39 *LcARFs* obtained by qRT-PCR in seven different tissues and organs. Red and blue were represented relatively high and lower expression (\log_2 ratio), respectively. Every sample has three biological replicates. YF: young fruit (25 days after fertilization), FF: female flower, MF: male flower, USF: undetermined sex flowers, ML: mature leaves, YL: young leaves, FBS: fruit-bearing shoots. **B) Relative abundance of *LcARFs* significantly expressed in FF.** Data from three independent biological replicates are shown with standard deviation (SD). Asterisks on the top of bars indicate significant differences as determined by Student's t-test (*P, 0.05).




Figure 5

Expression Profiles of LcARF Genes in Response to ETH, GPD, and GPDD treatments.

Fruit-bearing shoots of 'FZX' litchi were obtained at 25 days after anthesis and then carried on ETH, GPD, and GPDD treatment from 0, 1, 2 and 3 days, respectively. CK: control; ETH: treated by ethylene; GPD: girdling plus defoliation; GPDD: dipping in 2, 4-D after GPD treatment. The heatmap was created based on the TPM values of *LcARFs* from the transcriptome data. In the heatmap, red and blue were represented higher and lower expression (\log_2 ratio), respectively. Heatmap and hierarchical clustering were performed by average linkage (default) method.

