

Using *Ipomoea aquatic* as an environmental-friendly alternative to *Elodea nuttallii* for the aquaculture of Chinese mitten crab

Linlin Shi^{1, 2}, Meijuan Jin², Shixue Yin^{Corresp., 3}, Changying Lu², Haihou Wang², Xingwei Zhou², Lijuan Mei³, Mingxing Shen^{Corresp. 2}

¹ College of Agriculture, Yangzhou University, Yangzhou, Jiangsu, China

² Research Center of Agricultural Resource and Environment, Institute of Agricultural Science in Taihu Lake District, Suzhou, Jiangsu, China

³ College of Environmental Science and Engineering, Yangzhou University, Yangzhou, Jiangsu, China

Corresponding Authors: Shixue Yin, Mingxing Shen
Email address: sxyin@yzu.edu.cn, smx@jaas.ac.cn

Elodea nuttallii (EN) is widely used in Chinese mitten crab (CMC) rearing practice, but it is not a native aquatic plant and cannot endure high temperature. Thus, large EN mortality and water deterioration events could occur during high temperature seasons. The aim of this study was to identify the use of local macrophytes in CMC rearing practice, including *Ipomoea aquatic* (IA) and *Oryza sativa* (OS). A completely randomized field experiment was conducted to investigate the crab yield, water quality, bacterioplankton community and functions in the three different systems (EN, IA, and OS). Average crab yields in the different macrophyte systems did not differ significantly. The IA and OS systems significantly decreased the TN and NO₃⁻-N quantities in the outflow waters during the rearing period compared to the EN system, and the IA and OS plants assimilated more nitrogen than the EN plant. Moreover, the significant changes of bacterioplankton abundances and biodiversity in the three systems implied that cleanliness of rearing waters was concomitantly attributed to the differential microbial community and functions. In addition, principle component analysis successfully differentiated the bacterioplankton communities of the three macrophytes systems. Environmental factor fitting and the co-occurrence network analyses indicated that pH was the driver of bacterioplankton community structure. Functional predictions using PICRUST (v.1.1.3) software based on evolutionary modeling indicated a higher potential for microbial denitrification in the IA and OS systems. Notably, the OS plants stopped growing in the middle of the rearing period. Thus, the IA system rather than the OS system could be a feasible and environmental-friendly alternative to the EN system in CMC rearing practice.

Using *Ipomoea aquatic* as an environmental-friendly alternative to *Elodea nuttallii* for the aquaculture of Chinese mitten crab

Linlin Shi^{1,2}, Meijuan Jin², Shixue Yin^{3,#}, Changying Lu², Haihou Wang², Xingwei Zhou², Lijuan Mei³, Mingxing Shen^{2,#}

¹College of Agriculture, Yangzhou University, Jiangsu, China

²Research Center of Agricultural Resource and Environment, Institute of Agricultural Science in Taihu Lake District, Jiangsu, China

³College of Environmental Science and Engineering, Yangzhou University, Jiangsu, China

#Corresponding author: Shixue Yin

Email address: sxyin@yzu.edu.cn

#Corresponding author: Mingxing Shen

Email address: smx@saas.ac.cn

Abstract

Elodea nuttallii (EN) is widely used in Chinese mitten crab (CMC) rearing practice, but it is not a native aquatic plant and cannot endure high temperature. Thus, large EN mortality and water deterioration events could occur during high temperature seasons. The aim of this study was to identify the use of local macrophytes in CMC rearing practice, including *Ipomoea aquatic* (IA)

and *Oryza sativa* (OS). A completely randomized field experiment was conducted to investigate the crab yield, water quality, bacterioplankton community and functions in the three different systems (EN, IA, and OS). Average crab yields in the different macrophyte systems did not differ significantly. The IA and OS systems significantly decreased the TN and NO₃⁻-N quantities in the outflow waters during the rearing period compared to the EN system, and the IA and OS plants assimilated more nitrogen than the EN plant. Moreover, the significant changes of bacterioplankton abundances and biodiversity in the three systems implied that cleanliness of rearing waters was concomitantly attributed to the differential microbial community and functions. In addition, principle component analysis successfully differentiated the bacterioplankton communities of the three macrophytes systems. Environmental factor fitting and the co-occurrence network analyses indicated that pH was the driver of bacterioplankton community structure. Functional predictions using PICRUSt (v.1.1.3) software based on evolutionary modeling indicated a higher potential for microbial denitrification in the IA and OS systems. Notably, the OS plants stopped growing in the middle of the rearing period. Thus, the IA system rather than the OS system could be a feasible and environmental-friendly alternative to the EN system in CMC rearing practice.

Introduction

The Chinese mitten crab (CMC), *Eriocheir sinensis*, is considered an invasive species in Europe and North America (Brodin & Drotz 2014; Hanson & Sytsma 2008), but it is an expensive delicacy in Asia (Chen & Zhang 2007). In 2014, 796,621 tons of farmed Chinese

mitten crabs were produced (Zeng et al. 2013); crabs were primarily bred in ponds and lakes (Zeng et al. 2013). The mitten crabs produced in Yangcheng Lake, Suzhou, China, are of high quality and have high economic value (Gu et al. 2013). Most of the crabs produced in Yangcheng Lake are exported to Shanghai, Hong Kong, and high-profit foreign markets.

Aquatic plants are required for mitten crab farming. The plants provide shelter for the crabs during exuviation, which is an important part of crab growth (Meng et al. 2013). In addition, aquatic plants assimilate excess nutrients, improve water cleanliness, and absorb solar radiation to maintain cool water temperatures. These properties increase crab growth, yield, and quality (Zhan & Yang 2015).

Elodea nuttallii (EN), a perennial aquatic plant native to North America, provides these benefits, and is thus widely used in CMC aquaculture (Wang et al. 2016). However, EN cannot withstand high temperatures (Zhan & Yang 2015). In Yangcheng Lake, summer air and water temperatures typically reach 35–40°C and 26–34°C, respectively, which frequently leads to massive EN die off, resulting in serious water quality deterioration (Wu et al. 2016). Under such conditions, crab growth is negatively affected due to loss of shelter for exuviation hiding places and poor water quality (Zhan & Yang 2015). Consequently, alternative aquatic plants are required to facilitate mitten crab aquaculture in areas such as Yangcheng Lake.

Local plants that have adapted to local conditions are the best candidates for EN replacement. For example, *Ipomoea aquatica* (IA) is a semiaquatic, tropical/subtropical plant that might be applicable to mitten crab aquaculture (Zhang et al. 2014). IA grows well in shallow waters,

withstands high temperatures, and efficiently removes nutrients (e.g., nitrogen and phosphorous) from water bodies (Liu et al. 2007; Wei et al. 2017). Furthermore, the tender shoots and leaves of IA are consumable, and providing additional economic value. Alternatively, *Oryza sativa* (OS) is a submerged grain crop that is common in Asia, and its assimilation ability of nitrogen and phosphorus is similar to IA (Nawaz & Farooq 2017). OS is thus another candidate for EN replacement.

Therefore, in this study, we aimed to answer three questions. Which of the two locally adapted plants would most adequately replace EN, similarly improving crab yield and quality? How do the candidate plant systems affect water quality and the associated environmental characteristics? Considering microorganisms play important roles in nutrient conversion in wastewater (Daims et al. 2006), are the two plant systems associated with different bacterioplankton communities that differentially affect water quality? Answers to these questions are of critical importance to crab farmers, and those who are concerned with water quality. Despite the importance of these questions, they remain unanswered. We thus aimed to address the above questions by evaluating candidate aquatic plant systems, and exploring the associated bacterioplankton communities.

Material and methods

Pond construction, seedling preparation, and floating system construction

All of the experiments were performed in Lianzigang, Suzhou, China (31°27'40.18"N, 120°43.5'5.32"E). This region has a subtropical monsoon climate, with an annual average

rainfall of 1076.2 mm (<http://www.pmsc.cn/>). Annual average temperature in Lianzigang is 15–17°C, with high temperatures of 35–40°C in July and August (<http://www.pmsc.cn/>). All of the experiments were conducted between May 1 and November 30, 2017.

The crab-rearing system used here consisted of a water inlet, a water outlet, a micro-porous aeration system, a pond, floating macrophytes, and vertical posts to anchor the macrophytes (Fig. 1). The pond, which was mechanically excavated, had a total area of 1,000 m² and was 1.5-m deep. The pond was separated into nine sub-ponds with cement walls that were 0.4 m wide and 1.5 m high. Each sub-pond had an area of ~100 m² and an independent staff gauge. Sub-pond bottoms were left in a natural state to allow crabs to burrow, and to facilitate EN root establishment and growth. IA, OS, and EN were grown in separate sub-ponds with three replicates per plant using a completely randomized design.

Sub-ponds were disinfected using sodium hypochlorite. After disinfection, water was pumped into each sub-pond to a depth of ~10 cm. Basal fertilizer (45 kg ha⁻¹ of compound-fertilizer; N:P₂O₅:K₂O = 15:15:15) was applied 2–3 days before seedling transplantation.

OS and IA seedlings were prepared by sowing seeds on a patented nutritional matrix (0.16 m²) containing sufficient nutrients for seedling growth. The matrix had sufficient buoyancy to allow seedlings to float on the surface of the water. Seeds were germinated on the matrix, grown for 30 days in a greenhouse at 25°C, and then transferred to the bottom of the appropriate sub-ponds. Seedling matrices were arranged side by side until seedling coverage reached 60% of the total water surface area. Additional water was then pumped into each sub-pond to increase the water

depth to 20 cm, and seedlings were allowed to re-establish growth for 5–7 days. Upon growth re-establishment, water was again gradually and gently pumped into the sub-ponds to increase water depth to 1.0 m. At this depth, all of the seedlings were floating. Seedlings were fixed in place using ropes attached to buoys, which were fastened to posts in the sides of the cement dividers. EN seedlings were directly transplanted from other ponds. EN cluster spacing was 0.5 m × 0.5 m, with forty seedlings per cluster. The transplanted EN plants covered ~60% of the total water surface area. Upon growth re-establishment, water was pumped into the EN sub-ponds to a depth of 1.0 m.

Crab pond management, nutrient measurements, and crab yield determination

Crab ponds were managed using standard Chinese mitten crab rearing methods (Zhan & Yang 2015). Juvenile crabs, with an average weight of 15 g, were purchased from a local company (Su'an Fishery Co., Ltd., Nantong, China) and added to the experimental sub-ponds at a density of about 12,000 individuals per hectare. Crab feed (bait) was purchased from the Tongwei Group (<http://www.tongwei.com/>). The nutrient composition of the feed was varied to meet the different needs at each of the growth stages. Crabs were fed twice per day dependent on growth stage, as recommended by the bait manufacturer. Crabs were not dosed with antibiotics or chemical fishery drugs. A micro-porous aeration system was used nightly to ensure sufficient aeration. Approximately 5 cm of water was pumped out of all ponds each week, and replaced with an equal volume of fresh water from Yangcheng Lake (depending on local precipitation).

Outflow water samples were taken every 7–10 days during the rearing period, and

immediately frozen at -20°C . At the end of the experiment, nutrients in the outflow samples were measured using an auto analyzer (SKALAR SAN⁺⁺, Breda, the Netherlands); the nutrients measured included total nitrogen (TN), total phosphorous (TP), ammonium-N ($\text{NH}_4^{+}\text{-N}$), nitrate-N ($\text{NO}_3^{-}\text{-N}$), and nitrite-N ($\text{NO}_2^{-}\text{-N}$). Absolute cumulative nutrient quantity in discharged water was measured by nutrient concentration in discharged water and outflow water quantity estimated by staff gauge. Outflow pH was measured using a WTW portable pH meter (ProfLine 3310, Weilheim, Germany). Inlet water samples were taken and measured every month during the rearing period, and average nutrient concentrations were finally calculated. Dissolved oxygen (DO) concentrations were not measured, because preliminary studies identified large spatial and temporal variations in DO concentrations (Fig. S1) due to uncontrollable factors (i.e., air temperature, air pressure, water disturbances, aeration, activities of aquatic organisms, and photosynthesis of plants and algae) (Dai et al. 2013).

The biomass yields (TBM) of EN, IA, and OS were estimated by dry matter productivity in unit area (DM) and aquatic plant areas (APA) in each pond. The fresh plant tissues in 1 m^2 area were collected and oven-dried with 3 replicates in each pond at the end of the experiment, and average dry weight in 1 m^2 area was considered as DM. APA was equal to the 60% of total area of each pond. Thus, the TBM could be calculated by the equation: $\text{TBM} = \text{DM} \times \text{APA}$. Trimmed plant tissues were included in the biomass production estimates. As IA and EN has a sprawling growth pattern, it was periodically cut back outside of the rope-restricted area to maintain ~60% coverage to keep constant aquatic plant coverage (Trimming details see Fig. S1-S2). Nutrients

assimilated by the plants were calculated based on plant biomass and nutrient concentrations. The tissue-mixed plant samples that were collected in different growth stages were used for nutrient concentration measurement. Table S1 showed the average nutrient concentrations of plants.

To assess crab production, mature crabs were harvested using crab traps, and remaining crabs were captured by hand at night, after the pond water was completely drained. Males and females were manually separated and weighed.

Characterization of bacterioplankton communities

As crab rearing is sensitive to high air temperatures (Yuan et al. 2017), the bacterioplankton communities during periods of high temperatures were more interesting. Thus, the bacterioplankton community was assessed on July 2, 2017, when the maximum air temperature reached 37°C. To obtain sufficient bacterioplankton biomass for community profiling via DNA sequencing, we collected 10 L of water at 20 cm depth from each replicate pond. Water samples were filtered through a 0.22-µm polycarbonate membrane (Millipore, Billerica, MA, USA). DNA was extracted from the filtered biomass using a FastDNA Spin Kit for soil (MP bio, Solon, OH, USA), following the manufacturer's protocols. Extracted DNA concentration was determined with a NanoDrop 2000 UV-vis spectrophotometer (Thermo Scientific, Wilmington, NC, USA); DNA quality was assessed with gel electrophoresis on a 1% agarose gel. The V4 hypervariable region of the bacterial 16S rRNA gene was amplified using PCR, with the primers 563F (5'-AYTGGGYDTAAAGVG-3') and 802R (5'-TACNVGGGTATCTAATCC-3')

(Cardenas et al. 2010), following previously described protocols (Wang et al. 2017). The amplified PCR products were purified using an AxyPrep DNA Gel Extraction Kit (Axygen Biosciences, Union City, CA, USA), and quantified using a QuantiFluor-ST kit (Promega, Madison, WI, USA), following the manufacturer's instructions. Purified amplicons were pooled in equimolar concentrations and sequenced on an Illumina MiSeq platform (Illumina, San Diego, CA, USA) at Majorbio Bio-Pharm Technology Co. Ltd. (Shanghai, China), following standard protocols.

Raw sequence reads were demultiplexed using QIIME (v.1.9.1) (Caporaso et al. 2010a). Barcoding adapters and PCR primers were cleaved using cutadapt (v.1.16) (Martin 2011). Low-quality reads were removed from the dataset with USEARCH10 (Edgar & Flyvbjerg 2015), using the "fastq_filter" command with the parameters maxee=1 and truncqual=15. The remaining paired-end reads were merged using the "fastq_mergepairs" command in USEARCH10 (Edgar & Flyvbjerg 2015). Merged read abundances were normalized across samples by randomly subsampling 28,000 sequences from each sample. A zero-radius OTU (zOTU) table was produced from the sequence reads using the Unoise3 algorithm (Edgar 2018). The taxonomic classification of each zOTU was assigned using the UCLUST algorithm against the Silva (SSU123) 16S *r*RNA database with default parameters (Caporaso et al. 2010b; Edgar 2010).

An additional OTU table was generated to predict bacterioplankton functions. The abundance-normalized sequences were clustered at the 97% nucleotide similarity cutoff level into OTUs

using the “pick_closed_reference_otus.py” function in QIIME (v.1.9.1) (Caporaso et al. 2010b). OTUs were taxonomically classified using the UCLUST algorithm (Edgar 2010) against the Greengenes (gg_13_5) reference database (DeSantis et al. 2006) with default parameters. The resulting OTU table was analyzed using PICRUSt (v.1.1.3) (scripts “normalize_by_funtion.py”, “predict_metagenomes.py”, “categorize_by_function.py”, and “metagenome_contributions.py”) (Langille et al. 2013). The PICRUSt algorithm produced a table of functional Kyoto Encyclopedia of Genes and Genomes (KEGG) orthologs (KOs). To obtain OTU-specific counts of genes (da Fonseca et al. 2019; Fan et al. 2018) associated with nitrification and denitrification, the PICRUSt script “metagenome_contributions.py” with -l option was applied to selected KOs (K00370, K00371, K00374, K02567, K02568, K00368, K15864, K04561, K02305, K00376, K10535, K10944, K10945, and K10946) (Script S1).

Statistical analysis

Statistical analyses were primarily performed in R (v.3.3.2) (Team 2013). To identify significant differences among treatments, the Levene and Kolmogorov-Smirnov tests were used to check the homogeneity of variances and data normality, respectively. One-way ANOVA was used to determine the significance among the treatments, and Tukey’s HSD test was then applied for multiple comparisons. If the measurement variable did not meet the normality assumption, a Kruskal-Wallis test was performed instead of one-way ANOVA. We considered $P < 0.05$ to be statistically significant, unless otherwise noted.

The bacterioplankton communities were analyzed using the vegan package (Dixon 2009) in R.

The matrix of zOTU abundances was transformed prior to distance-based analyses using the “decostand” function with the Hellinger method (Legendre & Gallagher 2001). Principal component analyses (PCAs) were performed using the "rda" function in vegan to visualize differences among the bacterioplankton communities from the different macrophyte systems. Environmental variables were then fitted and projected onto an ordination using “envfit” function in vegan based on 1,000 permutations.

To investigate the correlation of microbial taxa and environmental variables, we constructed a co-occurrence network using CoNet (Faust & Raes 2016), based on the zOTU table and the environmental variables. In the co-occurrence analysis, the read count matrix was first filtered, and only those zOTUs with at least seven minimum occurrence values across the nine samples were retained. Pair-wise associations among zOTUs and environmental factors were calculated using the Pearson, Spearman, Kendall, Bray-Curtis, and Kullback-Leibler correlation methods. The initial top and bottom edge numbers were set at 1,000. For each edge and each measure of association, 1,000 permutation scores and 1,000 bootstrap scores were computed. The resultant networks were visualized using Cytoscape (v.3.2.1) (Shannon et al. 2003).

Results

System performance, crab yield, and water quality

The macrophyte systems floated steadily throughout the whole experiment despite two moderate windstorms. Thus, the floating systems described here were suitable for plant growth

on water surfaces. All of the plants grew well initially (between May and August). However, OS began to go to seed (indicating the start of the reproductive stage) in early August, about 25 days earlier than the normal (September, if OS is grown in a field). After early August, OS plants yellowed and the roots darkened. In contrast, IA and EN grew well until the end of the experiment.

Crab yields among the three macrophyte systems did not differ significantly. The average yields of male and female crabs were $\sim 740 \text{ kg ha}^{-1}$ and $\sim 267 \text{ kg ha}^{-1}$, respectively (Fig. 2). Individual weight is one of the most important factors determining the value of commercial crabs. Here, the individual weight distributions did not differ significantly among plant systems ($P > 0.05$). The median weights of male and female crabs were 168 g and 109 g, respectively (Fig. 2).

The nitrogen assimilated by IA was estimated to be 118 kg ha^{-1} , based on macrophyte biomass weight and nutrient concentrations. This level of assimilation was 3.5 times that of EN, and 1.5 times that of OS (Fig. 3). In contrast, the phosphorous assimilated by IA, EN, and OS did not differ significantly.

The average TN, TP, $\text{NH}_4^+\text{-N}$ and $\text{NO}_3^-\text{-N}$ concentrations of inlet waters in the rearing period were 1.25 ± 0.44 (Mean \pm SD) mg L^{-1} , $0.03 \pm 0.02 \text{ mg L}^{-1}$, $0.25 \pm 0.43 \text{ mg L}^{-1}$ and $0.57 \pm 0.47 \text{ mg L}^{-1}$, respectively (Fig. 4). The TN content in 31.3% of the samples from the EN system exceeded the Environmental Quality Standards for Surface Water (GB3838-2002) limit for type III water. In contrast, the TN content in 9.4% and 15.6% samples from the OS and IA systems, respectively, exceeded the type III water quality limit (GB3838-2002) (Fig. 4). The number of

samples that exceeded the TP and $\text{NH}_4^+\text{-N}$ type III water quality limits did not differ substantially among treatments. No samples exceeded the water quality limit for $\text{NO}_3^-\text{-N}$ (10 mg L^{-1}). Cumulative curve analysis indicated that amount of TP and $\text{NH}_4^+\text{-N}$ accumulated over the course of the experiment (up to November 27) did not differ significantly among different plant systems. However, the levels of accumulated TN and $\text{NO}_3^-\text{-N}$ were significantly higher in the EN system than in the IA or OS systems (Fig. 5). Average pH was 7.48 in the IA system, 8.16 in the EN system, and 7.9 in the OS system (Fig. 6).

Bacterioplankton communities and predicted functions associated with nitrification/denitrification

The abundances of nearly all of the bacterioplankton phyla differed significantly among macrophyte systems (Fig. 7 A). The abundances of *Acidobacteria*, *Chloroflexi*, *Firmicutes*, β -*Proteobacteria*, and δ -*Proteobacteria* were significantly higher in the OS and IA systems as compared to the EN system, while the abundances of *Actinobacteria*, *Armatimonadetes*, and γ -*Proteobacteria* were significantly lower. OTU-based diversity (α -diversity) was higher in the OS and IA systems than in the EN system (Fig. 7 B), suggesting that bacterioplankton communities were more complex in the OS and IA systems than in the EN system.

PCA indicated that the bacterioplankton communities were discrete among the different plants, with the first two axes explaining 83.67% of the total variation (Fig. 8 A). After fitting environmental variables (Table S2) and bacterioplankton community ordination, we found that pH was significantly associated with community differences (Bonferroni-adjusted $P < 0.05$) (Fig.

8 A). The co-occurrence network analysis also identified pH as the only environmental factor that co-varied with certain taxa. These pH-correlated taxa included zOTUs belonging to the phyla *Acidobacteria*, *Chloroflexi*, *Firmicutes*, *Proteobacteria*, and *Planctomycetes* (Fig. 8 B). The total average abundance of co-occurring zOTUs was 7.7%, suggesting that pH influenced the relative abundance of the most abundant members of each community.

Potential functions of the bacterioplankton communities were predicted using PICRUSt. We focused specifically on functional genes associated with nitrification and denitrification, as these processes are highly related to nitrogen cycling and are likely to affect nitrogen concentration in pond water. Genes associated with all of the steps of denitrification (*napA*, *napB*, *nirK*, *norB*, *norC*, and *nosZ*; (Kanehisa et al. 2017)) were generally more abundant in the IA and/or OS systems than in the EN system. However, the abundances of *narI*, *narG*, and *narH*, which are only involved in the reduction of nitrate to nitrite (Kanehisa et al. 2017), were lower in the IA and/or OS systems (Fig. 9). The abundances of the nitrifying genes *pmoA-amoA*, *pmoB-amoB*, and *pmoC-amoC*, which are involved in the oxidation of ammonia to hydroxylamine (Kanehisa et al. 2017), were significantly higher in the OS system than in the IA and EN systems. The abundance of *hao* was not significantly different among the three macrophyte systems, although the abundances of *narG* and *narH* were lower in the IA and OS systems than in the EN system (Fig. 9).

Discussion

The macrophyte system described here floated steadily throughout the whole experiment. In

addition, mitten crab yield and quality did not differ significantly among plant systems (Fig. 2). Thus, our results demonstrated that local plants were a feasible alternative to EN. Indeed, our data indicated that IA was be the best local replacement for EN, as this plant grew well in ponds and provided sufficient shade. In contrast, OS is not a suitable alternative to EN, as it stopped growing in the middle of the experimental period. The mechanisms underlying the different performances of OS and IA were related to the different growth behaviors and nutrient requirements of the two species. OS has several distinctive growth stages, including tillering, heading, ripening and et al. (Zhang et al. 2018), which typically require different levels of nutrients (Fairhurst et al. 2007). For example, during the tillering stage, OS biomass increases much faster than during other stages of OS growth, and consequently requires a more intensive nutrient supply (Nawaz & Farooq 2017). If the nutrient supply needs are not met (i.e., due to low nutrient concentrations in the water), many tillers become non-productive, and the few remaining productive tillers enter the reproductive stage earlier than normal (Nawaz & Farooq 2017). In submerged paddy fields, the intensity of nutrient supply is manipulated by top-dressing fertilizers (Fairhurst et al. 2007). However, this technique cannot be used in crab-rearing ponds because it would pollute the water.

Unlike OS, IA does not exhibit obvious physiological differences among growth stages. IA has sprawling growth, with biomass increasing over rearing time (from May to November) (Shaltout et al. 2010). Thus, low concentrations of nutrients in the water are sufficient. We consequently concluded that IA is an attractive macrophytic alternative to EN in crab-rearing

296 ponds.

297 Based on the Chinese national water quality standards (GB3838-2002) (SEPA), 31.3% of the
 298 outflow samples from the EN system contained TN levels in excess of limits for type III waters
 299 (Fig. 4). Thus, the outflow from ponds using the EN system is likely to pollute the downstream
 300 environment. Fewer outflow samples from the IA and OS systems, as compared to the EN
 301 system, had TN levels above the type III water quality limits. These data suggest IA or OS
 302 systems would generate less environmental pollution than EN systems. This suggestion was
 303 reinforced by the absolute cumulative nutrient quantities in outflow samples (Fig. 5), which
 304 showed that the TN and $\text{NO}_3\text{-N}$ concentrations were significantly higher in the EN system, as
 305 compared to the IA and OS systems. Indeed, IA and OS assimilated more nitrogen from the
 306 water than did EN (Fig. 3). Thus, it was possible that IA and OS more effectively improved
 307 water quality than EN, which is a desirable property for macrophytes grown in crab-rearing
 308 ponds. An alternative explanation was that the differences in water quality among plant systems
 309 were a result of the activities of macrophyte-specific bacterioplankton communities.

310 Bacterioplankton communities differed significantly among the three macrophyte systems (Fig.
 311 6), possibly because of the different root deposits produced by the three plants. Terrestrial and
 312 macrophytic plants release unique root exudates that drastically alter bacterioplankton
 313 community structure (Baudoin et al. 2003; Casamatta & Wickstrom 2000; Nelson et al. 2013;
 314 Tanaka et al. 2012; Zhao et al. 2013). Bacterioplankton community composition might also be
 315 regulated by pH, as pH was the only environmental factor that was significantly associated with

the relative abundances of specific bacterioplankton taxa (Fig. 8). These results were consistent with previous reports, which indicated that pH affects the community structures of both terrestrial bacteria (Fierer & Jackson 2006; Lauber et al. 2009; Rousk et al. 2010) and bacterioplankton (Ren et al. 2015). Moreover, the significant increases in bacterioplankton diversity with lower water nutrients in the IA and OS treatments (Figs. 5, 7 B) indicated that biodiversity helped improve water quality, as observed in other aquatic systems (Cardinale 2011; Gregoracci et al. 2012).

The predicted functions of the bacterioplankton community also differed among the macrophyte systems, particularly those related to nitrification and denitrification. Most denitrifying genes, especially those associated with the reduction of nitrite, nitric oxide, and nitrous oxide, were more abundant in the IA and OS bacterioplankton communities, as compared to the EN bacterioplankton community (Fig. 9). Moreover, abiotic environmental factors usually control the denitrification process, i.e., pH, temperature and organic carbons (OC). The IA and OS systems that could provide labile OC derived from root exudates (Wang et al. 2018; Xu et al. 2008; Zhu & Cheng 2011) favored denitrification. Overall, the biotic and abiotic influences on denitrification in the IA and OS systems were thought to be co-occurring, consistent with the conclusion in riparian wetlands (Xiong et al. 2017). In addition, concentrations of NO_3^- -N (and TN) were lower in the IA and OS systems than in the EN system. Thus, the IA and OS systems might have a higher denitrification potential than the EN system. Meanwhile, there were no obvious differences in the abundance patterns of nitrification-associated genes among

macrophyte systems. The genes responsible for the oxidation of ammonium to hydroxylamine (*pmoA-amoA*, *pmoB-amoB*, and *pmoC-amoC*) were significantly more abundant in the OS system (but not the IA system) as compared to the EN system, but the abundances of these genes were relatively low across all of the plant systems, with only ~2–20 copies per sample. In addition, the gene abundances in nitrification process usually could not accurately predict nitrification potential, which can be affected by other abiotic factors (Francis et al. 2005; Rocca et al. 2015; Yao et al. 2018). Thus, nitrification potential may not vary significantly among these three macrophytes, and further studies are required to validate these coupled nitrification-denitrification processes in future.

Conclusions

EN is routinely cultivated in ponds used for mitten crab aquaculture. However, EN temperature sensitivity often leads to plant deterioration and decreased water quality at high ambient temperatures, negatively affecting crab production. Here, we successfully designed a floating system to support the growth of IA and OS on the surfaces of crab-rearing ponds. We then compared the crab yield, outflow water quality, and bacterioplankton communities among ponds with EN, IA, and OS macrophyte systems. Our results indicated that IA growth behavior was preferable to that of OS. Crab yields did not differ significantly among systems. Moreover, outflow water quality, as indicated by TN and NO₃⁻-N concentrations, was better in the IA and OS systems than in the EN system, due to the greater nitrogen assimilation of IA and OS as compared to EN. In addition, the microbial communities associated with IA and OS had a greater

denitrification potential than the microbial community associated with EN. Thus, our results indicated mitten crabs could be successfully reared using native aquatic plant. Specifically, IA was a suitable and environmental-friendly replacement for EN, but OS was not.

Data Availability

Demultiplexed sequences and metadata are available from the NCBI Sequence Read Archives (SRA) under accession number SRP136316.

Acknowledgments

Authors would like to thank the anonymous reviewers and Kiran Liversage for their good suggestions, which have helped us to improve this manuscript. We also thank LetPub (www.letpub.com) for its linguistic assistance during the preparation of this manuscript.

References

Fisheries and aquaculture software. FishStatJ - software for fishery statistical time series. In: FAO Fisheries and Aquaculture Department. Available at <http://www.fao.org/fishery/>.

Baudoin E, Benizri E, and Guckert A. 2003. Impact of artificial root exudates on the bacterial community structure in bulk soil and maize rhizosphere. *Soil Biology and Biochemistry* 35:1183-1192. [https://doi.org/10.1016/S0038-0717\(03\)00179-2](https://doi.org/10.1016/S0038-0717(03)00179-2)

Brodin T, and Drotz MK. 2014. Individual variation in dispersal associated behavioral traits of the invasive Chinese mitten crab (*Eriocheir sinensis*, H. Milne Edwards, 1854) during initial invasion of Lake Vanern, Sweden. *Current*

374 *Zoology* 60:410-416. 10.1093/czoolo/60.3.410

375 Caporaso JG, Bittinger K, Bushman FD, DeSantis TZ, Andersen GL, and Knight R. 2010a. PyNAST: a flexible tool
376 for aligning sequences to a template alignment. *Bioinformatics* 26:266-267. 10.1093/bioinformatics/btp636

377 Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, Fierer N, Pena AG, Goodrich JK,
378 Gordon JI, Huttley GA, Kelley ST, Knights D, Koenig JE, Ley RE, Lozupone CA, McDonald D, Muegge BD,
379 Pirrung M, Reeder J, Sevinsky JR, Tumbaugh PJ, Walters WA, Widmann J, Yatsunenko T, Zaneveld J, and
380 Knight R. 2010b. QIIME allows analysis of high-throughput community sequencing data. *Nature Methods* 7:335-
381 336. 10.1038/nmeth.f.303

382 Cardenas E, Wu WM, Leigh MB, Carley J, Carroll S, Gentry T, Luo J, Watson D, Gu B, Ginder-Vogel M, Kitanidis
383 PK, Jardine PM, Zhou J, Criddle CS, Marsh TL, and Tiedje JM. 2010. Significant association between sulfate-
384 reducing bacteria and uranium-reducing microbial communities as revealed by a combined massively parallel
385 sequencing-indicator species approach. *Appl Environ Microbiol* 76:6778-6786. 10.1128/AEM.01097-10

386 Cardinale BJ. 2011. Biodiversity improves water quality through niche partitioning. *Nature* 472:86-89.
387 10.1038/nature09904

388 Casamatta DA, and Wickstrom CE. 2000. Sensitivity of two disjunct bacterioplankton communities to exudates
389 from the cyanobacterium *Microcystis aeruginosa* Kützing. *Microbial Ecology* 40:64-73. 10.1007/s002480000035

390 Chen DW, and Zhang M. 2007. Non-volatile taste active compounds in the meat of Chinese mitten crab (*Eriocheir*
391 *sinensis*). *Food Chemistry* 104:1200-1205. <https://doi.org/10.1016/j.foodchem.2007.01.042>

da Fonseca MMB, Minnicelli CF, Silva-Portela RdCB, de Farias MF, dos Santos PRS, Fernandes GJT, and Agnez-
Lima LF. 2019. Unlocking and functional profiling of the bacterial communities in diesel tanks upon additive
treatment. *Fuel* 236:1311-1320. <https://doi.org/10.1016/j.fuel.2018.09.107>

Dai H, Li Y, Ma X, Wang W, Zhu X, and Li Z. 2013. The research of dissolved oxygen distribution in the crab
ecological culture ponds (in Chinese). *Journal of Shanghai Ocean University* 22:66-73.

Daims H, Taylor MW, and Wagner M. 2006. Wastewater treatment: a model system for microbial ecology. *Trends
in Biotechnology* 24:483-489. <https://doi.org/10.1016/j.tibtech.2006.09.002>

DeSantis TZ, Hugenholtz P, Larsen N, Rojas M, Brodie EL, Keller K, Huber T, Dalevi D, Hu P, and Andersen GL.
2006. Greengenes, a Chimera-Checked 16S rRNA Gene Database and Workbench Compatible with ARB. *Applied
and Environmental Microbiology* 72:5069.

Dixon P. 2009. VEGAN, a package of R functions for community ecology. *Journal of Vegetation Science* 14:927-
930. [10.1111/j.1654-1103.2003.tb02228.x](https://doi.org/10.1111/j.1654-1103.2003.tb02228.x)

Edgar RC. 2010. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* 26:2460-2461.
[10.1093/bioinformatics/btq461](https://doi.org/10.1093/bioinformatics/btq461)

Edgar RC. 2018. Updating the 97% identity threshold for 16S ribosomal RNA OTUs. *Bioinformatics* 34:2371-2375.
[10.1093/bioinformatics/bty113](https://doi.org/10.1093/bioinformatics/bty113)

Edgar RC, and Flyvbjerg H. 2015. Error filtering, pair assembly and error correction for next-generation sequencing
reads. *Bioinformatics* 31:3476-3482. [10.1093/bioinformatics/btv401](https://doi.org/10.1093/bioinformatics/btv401)

410 Fairhurst TH, Witt C, Buresh RJ, and Dobermann A. 2007. Rice: A Practical Guide to Nutrient Management (2nd
 411 edition). Metro Manila, Philippines: International Rice Research Institute (IRRI). p 136.

412 Fan XY, Gao JF, Pan KL, Li DC, Dai HH, and Li X. 2018. Functional genera, potential pathogens and predicted
 413 antibiotic resistance genes in 16 full-scale wastewater treatment plants treating different types of wastewater.
 414 *Bioresource Technology* 268:97-106. <https://doi.org/10.1016/j.biortech.2018.07.118>

415 Faust K, and Raes J. 2016. CoNet app: inference of biological association networks using Cytoscape.
 416 *F1000Research* 5:1519-1519. 10.12688/f1000research.9050.2

417 Fierer N, and Jackson RB. 2006. The diversity and biogeography of soil bacterial communities. *Proc Natl Acad Sci*
 418 *U S A* 103:626-631. 10.1073/pnas.0507535103

419 Francis CA, Roberts KJ, Beman JM, Santoro AE, and Oakley BB. 2005. Ubiquity and diversity of ammonia-
 420 oxidizing archaea in water columns and sediments of the ocean. *Proc Natl Acad Sci U S A* 102:14683.
 421 10.1073/pnas.0506625102

422 Gregoracci GB, Nascimento JR, Cabral AS, Paranhos R, Valentin JL, Thompson CC, and Thompson FL. 2012.
 423 Structuring of bacterioplankton diversity in a large tropical bay. *PLoS One* 7:e31408.
 424 10.1371/journal.pone.0031408

425 Gu SQ, Wang XC, Tao NP, and Wu N. 2013. Characterization of volatile compounds in different edible parts of
 426 steamed Chinese mitten crab (*Eriocheir sinensis*). *Food Research International* 54:81-92.
 427 <https://doi.org/10.1016/j.foodres.2013.05.018>

428 Hanson E, and Sytsma M. 2008. The potential for mitten crab *Eriocheir sinensis* H. Milne Edwards, 1853
429 (Crustacea: Brachyura) invasion of Pacific Northwest and Alaskan estuaries. *Biological Invasions* 10:603-614.
430 10.1007/s10530-007-9156-3

431 Kanehisa M, Furumichi M, Tanabe M, Sato Y, and Morishima K. 2017. KEGG: new perspectives on genomes,
432 pathways, diseases and drugs. *Nucleic Acids Research* 45:353-361. 10.1093/nar/gkw1092

433 Langille MGI, Zaneveld J, Caporaso JG, McDonald D, Knights D, Reyes JA, Clemente JC, Burkepile DE, Thurber
434 RLV, Knight R, Beiko RG, and Huttenhower C. 2013. Predictive functional profiling of microbial communities
435 using 16S rRNA marker gene sequences. *Nature Biotechnology* 31:814-821. 10.1038/nbt.2676

436 Lauber CL, Hamady M, Knight R, and Fierer N. 2009. Pyrosequencing-based assessment of soil pH as a predictor of
437 soil bacterial community structure at the continental scale. *Appl Environ Microbiol* 75:5111-5120.
438 10.1128/aem.00335-09

439 Legendre P, and Gallagher ED. 2001. Ecologically meaningful transformations for ordination of species data.
440 *Oecologia* 129:271-280. 10.1007/s004420100716

441 Liu J, Dong Y, Xu H, Wang D, and Xu J. 2007. Accumulation of Cd, Pb and Zn by 19 wetland plant species in
442 constructed wetland. *Journal of Hazardous Materials* 147:947-953. <https://doi.org/10.1016/j.jhazmat.2007.01.125>

443 Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnetjournal: Next
444 Generation Sequencing Data Analysis* 17:10-12.

445 Meng X, Song X, Chen G, Feng Y, Wu L, and Shen Z. 2013. Utilization of natural feed resources in a macrophyte-

dominated lake for aquaculture of Chinese mitten crab (*Eriocheir sinensis*) and its purification effects on water environment (in Chinese). *Journal of Lake Sciences* 25:723-728.

Nawaz A, and Farooq M. 2017. Rice Physiology. In: Chauhan BS, Jabran K, and Mahajan G, eds. *Rice Production Worldwide*. Cham: Springer International Publishing, 455-485.

Nelson CE, Goldberg SJ, Wegley Kelly L, Haas AF, Smith JE, Rohwer F, and Carlson CA. 2013. Coral and macroalgal exudates vary in neutral sugar composition and differentially enrich reef bacterioplankton lineages. *ISME J* 7:962-979. 10.1038/ismej.2012.161

Ren L, Jeppesen E, He D, Wang J, Liboriussen L, Xing P, and Wu QL. 2015. pH influences the importance of niche-related and neutral processes in lacustrine bacterioplankton assembly. *Appl Environ Microbiol* 81:3104-3114. 10.1128/aem.04042-14

Rocca JD, Hall EK, Lennon JT, Evans SE, Waldrop MP, Cotner JB, Nemergut DR, Graham EB, and Wallenstein MD. 2015. Relationships between protein-encoding gene abundance and corresponding process are commonly assumed yet rarely observed. *ISME J* 9:1693-1699. 10.1038/ismej.2014.252

Rousk J, Baath E, Brookes PC, Lauber CL, Lozupone C, Caporaso JG, Knight R, and Fierer N. 2010. Soil bacterial and fungal communities across a pH gradient in an arable soil. *ISME J* 4:1340-1351. 10.1038/ismej.2010.58

SEPA. Environmental Quality Standard for Surface Water (GB3838-2002) (in Chinese). *State Environmental Protection Administration of China, Beijing*.

Shaltout KH, Al-Sodany YM, and Eid EM. 2010. Growth behaviour of the invasive species *Ipomoea carnea* in the

464 Nile Delta, Egypt. *Hydrobiologia* 656:187-197. 10.1007/s10750-010-0443-0

465 Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N, Schwikowski B, and Ideker T. 2003.

466 Cytoscape: A software environment for integrated models of biomolecular interaction networks. *Genome Research*

467 13:2498-2504. 10.1101/gr.1239303

468 Tanaka Y, Tamaki H, Matsuzawa H, Nigaya M, Mori K, and Kamagata Y. 2012. Microbial community analysis in

469 the roots of aquatic plants and isolation of novel microbes including an organism of the candidate phylum OP10.

470 *Microbes and Environments* 27:149-157. 10.1264/jsme2.ME11288

471 Team RC. 2013. R: A Language and Environment for Statistical Computing. R Foundation for Statistical

472 Computing, Vienna, Austria. Available at <https://cran.r-project.org/>.

473 Wang H, Wei Z, Mei L, Gu J, Yin S, Faust K, Raes J, Deng Y, Wang Y, Shen Q, and Yin S. 2017. Combined use of

474 network inference tools identifies ecologically meaningful bacterial associations in a paddy soil. *Soil Biology and*

475 *Biochemistry* 105:227-235. <https://doi.org/10.1016/j.soilbio.2016.11.029>

476 Wang Q, Liu J, Zhang S, Lian Y, Ding H, Du X, Li Z, and De Silva SS. 2016. Sustainable farming practices of the

477 Chinese mitten crab (*Eriocheir sinensis*) around Hongze Lake, lower Yangtze River Basin, China. *Ambio* 45:361-

478 373. 10.1007/s13280-015-0722-0

479 Wang X, Chen R, Jing Z, Yao T, Feng Y, and Lin X. 2018. Root derived carbon transport extends the rhizosphere of

480 rice compared to wheat. *Soil Biology and Biochemistry* 122:211-219. <https://doi.org/10.1016/j.soilbio.2018.03.024>

481 Wei H, Meng T, Li C, Xu K, Huo Z, Wei H, Guo B, Zhang H, and Dai Q. 2017. Comparisons of grain yield and

482 nutrient accumulation and translocation in high-yielding japonica/indica hybrids, indica hybrids, and japonica
483 conventional varieties. *Field Crops Research* 204:101-109. 10.1016/j.fcr.2017.01.001

484 Wu K, Ma X, Wang Y, Wang W, and Lang Y. 2016. Effect of three water plants decomposition on water quality (in
485 Chinese). *Journal of Shanghai Ocean University* 25:726-734.

486 Xiong Z, Guo L, Zhang Q, Liu G, and Liu W. 2017. Edaphic Conditions Regulate Denitrification Directly and
487 Indirectly by Altering Denitrifier Abundance in Wetlands along the Han River, China. *Environmental Science &*
488 *Technology* 51:5483-5491. 10.1021/acs.est.6b06521

489 Xu X, Kuzyakov Y, Wanek W, and Richter A. 2008. Root-derived respiration and non-structural carbon of rice
490 seedlings. *European Journal of Soil Biology* 44:22-29. <https://doi.org/10.1016/j.ejsobi.2007.09.008>

491 Yao L, Chen C, Liu G, Li F, and Liu W. 2018. Environmental factors, but not abundance and diversity of nitrifying
492 microorganisms, explain sediment nitrification rates in Yangtze lakes. *RSC Advances* 8:1875-1883.
493 10.1039/C7RA11956A

494 Yuan Q, Wang Q, Zhang T, Li Z, and Liu J. 2017. Effects of water temperature on growth, feeding and molting of
495 juvenile Chinese mitten crab *Eriocheir sinensis*. *Aquaculture* 468:169-174.
496 <https://doi.org/10.1016/j.aquaculture.2016.10.007>

497 Zeng QF, Gu XH, Chen X, and Mao ZG. 2013. The impact of Chinese mitten crab culture on water quality,
498 sediment and the pelagic and macrobenthic community in the reclamation area of Guchenghu Lake. *Fisheries*
499 *Science* 79:689-697. 10.1007/s12562-013-0638-1

500 Zhan J, and Yang X. 2015. *Standard aquaculture techniques for Chinese mitten crab* (in Chinese). Beijing:
501 Chemical Industry Press.

502 Zhang J, Zhang N, Liu Y-X, Zhang X, Hu B, Qin Y, Xu H, Wang H, Guo X, Qian J, Wang W, Zhang P, Jin T, Chu
503 C, and Bai Y. 2018. Root microbiota shift in rice correlates with resident time in the field and developmental stage.
504 *Science China Life Sciences* 61:613-621. 10.1007/s11427-018-9284-4

505 Zhang Q, Achal V, Xu Y, and Xiang W-N. 2014. Aquaculture wastewater quality improvement by water spinach
506 (*Ipomoea aquatica* Forsskal) floating bed and ecological benefit assessment in ecological agriculture district.
507 *Aquacultural Engineering* 60:48-55. <https://doi.org/10.1016/j.aquaeng.2014.04.002>

508 Zhao D-Y, Liu P, Fang C, Sun Y-M, Zeng J, Wang J-Q, Ma T, Xiao Y-H, and Wu QL. 2013. Submerged
509 macrophytes modify bacterial community composition in sediments in a large, shallow, freshwater lake. *Canadian*
510 *Journal of Microbiology* 59:237-244. 10.1139/cjm-2012-0554

511 Zhu J, and Cheng K. 2011. Effect of organic acids exuded from hydroponic plants roots on nitrogen cycling bacteria
512 (in Chinese). *Chinese Journal of Environmental Engineering* 5:2139-2143.

Figure 1

Experimental design and schematics for three macrophyte systems.

Elodea nuttallii (EN), *Oryza sativa* (OS), and *Ipomoea aquatica* (IA). Each macrophyte was planted in a separate sub-pond (three replicate ponds per species). The sub-ponds were connected by a series of PVC pipes. Pond water was replenished from the Yangcheng Lake water, and discharged water flowed into a constructed wetland. The growth of all three macrophytes was restricted to 60% of the pond areas. A micro-porous aeration system was used nightly to ensure sufficient aeration. Photo credit: Linlin Shi.




Figure 2 (on next page)

The yield and individual weight distributions of the Chinese mitten crabs did not differ significantly among macrophyte systems.

Error bars denote the standard deviations of the mean (n=3). EN, *Elodea nuttallii*; OS, *Oryza sativa*; IA, *Ipomoea aquatica*.




Figure 3 (on next page)

Assimilation of nitrogen and phosphorous by *Elodea nuttallii* (EN), *Ipomoea aquatica* (IA), and *Oryza sativa* (OS).

Different letters above bars represent significant differences among plant systems ($P < 0.05$; Tukey's HSD test). Error bars denote the standard deviations of the means ($n=3$).




Figure 4 (on next page)

Figure 4 (A-D) separately shows the TN, TP, $\text{NH}_4^+\text{-N}$ and $\text{NO}_3^-\text{-N}$ concentrations in the rearing pond.

Error bars denote the standard deviations of the means ($n=3$). The brown dashed lines represent the average nutrient concentrations in the inlet waters during the rearing period. The grey dashed lines represent the type III water quality limits from the Chinese Environmental Quality Standard for Surface Water (GB3838-2002). $\text{NO}_3^-\text{-N}$ concentrations were all below the type III water quality limit (10 mg L^{-1}). EN, *Elodea nuttallii*; OS, *Oryza sativa*; IA, *Ipomoea aquatica*.




Figure 5 (on next page)

Figure 5 (A-D) separately shows the absolute cumulative TN, TP, NH_4^+ -N and NO_3^- -N quantities in rearing period.

The statistical significance of differences in cumulative nutrient quantity among macrophyte systems was determined with a Kruskal-Wallis test ($n=3$) at the end of the rearing periods (November 27). Different lowercase letters represent significant differences ($P < 0.05$; Tukey's HSD test). EN, *Elodea nuttallii*; OS, *Oryza sativa*; IA, *Ipomoea aquatica*.




Figure 6(on next page)

pH of the outflow water samples during the rearing period.

EN, *Elodea nuttallii*; OS, *Oryza sativa*; IA, *Ipomoea aquatica*.




Figure 7 (on next page)

Bacterioplankton abundances and diversity indices of the three macrophyte systems.

(A) The relative abundances of the dominant bacterial phyla in the bacterioplankton communities, and (B) the associated alpha diversity. Asterisks indicate significant differences in abundances among the macrophyte systems ($P < 0.05$ level; one-way ANOVA; $n=3$). Error bars denote the standard errors of the means. Different letters above bars represent significant differences among treatments ($P < 0.05$; Tukey's HSD test; one-way ANOVA; $n=3$). EN, *Elodea nuttallii*; OS, *Oryza sativa*; IA, *Ipomoea aquatica*.




Figure 8(on next page)

Bacterioplankton community structure and environmental factors.

(A) Principal component analysis (PCA) of bacterioplankton community composition and environmental variables. The significant environmental variable ($P < 0.05$) is shown with a black arrow; other factors are shown in gray. EN, *Elodea nuttallii*; OS, *Oryza sativa*; IA, *Ipomoea aquatica*. (B) Subnetwork showing the correlation between bacterial taxa and pH. Squared nodes correspond to environmental parameters and circle nodes correspond to zOTUs. Circle nodes not assignable to genus are labeled with the names of higher taxonomic ranks, and node colors represent phyla. Red and blue colors of edge represent negative and positive correlations, respectively.





Figure 9(on next page)

The abundances of the bacterioplankton genes associated with nitrification and denitrification among macrophyte systems, as predicted by PICRUSt.

(A) and (B) shows the denitrification and nitrification predicted gene abundances, respectively. Error bars denote the standard deviations of the means (n=3). Different lowercase letters above bars represent statistically significant differences ($P < 0.05$; one-way ANOVA followed by Tukey's HSD tests). EN, *Elodea nuttallii*; OS, *Oryza sativa*; IA, *Ipomoea aquatica*.

(A) Denitrification pathway



(B) Nitrification pathway

