

Automated pupillometry to detect command following in neurological patients: A proof-of-concept study

Alexandra Vassilieva¹, Markus Harboe Olsen², Costanza Peinkhofer^{1,3}, Gitte Moos Knudsen^{1,4,5}, Daniel Kondziella

Corresp. 1, 4, 6

¹ Department of Neurology, Rigshospitalet, University of Copenhagen, Copenhagen, Denmark

² Department of Neuroanesthesiology, Rigshospitalet, University of Copenhagen, Copenhagen, Denmark

³ Medical Faculty, University of Trieste, Trieste, Italy

⁴ Faculty of Health Sciences and Medicine, University of Copenhagen, Copenhagen, Denmark

⁵ Neurobiology Research Unit and Center for Integrated Molecular Brain Imaging, University of Copenhagen, Copenhagen, Denmark

⁶ Department of Neuroscience, Norwegian University of Science and Technology, Trondheim, Norway

Corresponding Author: Daniel Kondziella

Email address: daniel_kondziella@yahoo.com

Background: Levels of consciousness in patients with acute and chronic brain injury are notoriously underestimated. Paradigms based on electroencephalography (EEG) and functional magnetic resonance imaging (fMRI) may detect covert consciousness in clinically unresponsive patients but are subject to logistical challenges and the need for advanced statistical analysis.

Methods: To assess the feasibility of automated pupillometry for the detection of command following, we enrolled 20 healthy volunteers and 48 patients with a wide range of neurological disorders, including 7 patients in the intensive care unit (ICU), who were asked to engage in mental arithmetic.

Results: Fourteen of 20 (70%) healthy volunteers and 17 of 43 (39.5%) neurological patients, including 1 in the ICU, fulfilled prespecified criteria for command following by showing pupillary dilations during ≥ 4 of 5 arithmetic tasks. None of the 5 sedated and unconscious ICU patients passed this threshold.

Conclusions: Automated pupillometry combined with mental arithmetic appears to be a promising paradigm for the detection of covert consciousness in people with brain injury. We plan to build on this study by focusing on non-communicating ICU patients in whom the level of consciousness is unknown. If some of these patients show reproducible pupillary dilation during mental arithmetic, this would suggest that the present paradigm can reveal covert consciousness in unresponsive patients in whom standard investigations have failed to detect signs of consciousness.

Automated pupillometry to detect command following in neurological patients: A proof-of-concept study

Alexandra Vassilieva ¹, Markus Harboe Olsen ², Costanza Peinkhofer ^{1,3}, Gitte Moos Knudsen ^{1,4,5}, Daniel Kondziella ^{1,5,6}

Department of Neurology ¹, Rigshospitalet, University of Copenhagen, Copenhagen, Denmark

Department of Neuroanesthesiology ², Rigshospitalet, University of Copenhagen, Copenhagen, Denmark

Medical Faculty ³, University of Trieste, Trieste, Italy

Neurobiology Research Unit ⁴, Rigshospitalet, Copenhagen University Hospital and Center for Integrated Molecular Brain Imaging, Copenhagen, Denmark, Copenhagen, Denmark

Faculty of Health and Medical Sciences, University of Copenhagen ⁵, Copenhagen, Denmark

Department of Neuroscience ⁶, Norwegian University of Science and Technology, Trondheim, Norway

Corresponding author: Daniel Kondziella, MD, MSc, dr.philos., FEBN
Department of Neurology,
Rigshospitalet, Copenhagen University Hospital
Blegdamsvej 9
DK-2100 Copenhagen
daniel_kondziella@yahoo.com
+45 60 13 11 00

Manuscript type: Original Research

Running title: Pupillometry to detect command following

Word count: 2893 (abstract 222); **tables:** 2; **figures:** 3; **references:** 41

Abstract

Background: Levels of consciousness in patients with acute and chronic brain injury are notoriously underestimated. Paradigms based on electroencephalography (EEG) and functional magnetic resonance imaging (fMRI) may detect covert consciousness in clinically unresponsive patients but are subject to logistical challenges and the need for advanced statistical analysis.

Methods: To assess the feasibility of automated pupillometry for the detection of command following, we enrolled 20 healthy volunteers and 48 patients with a wide range of neurological disorders, including 7 patients in the intensive care unit (ICU), who were asked to engage in mental arithmetic.

Results: Fourteen of 20 (70%) healthy volunteers and 17 of 43 (39.5%) neurological patients, including 1 in the ICU, fulfilled prespecified criteria for command following by showing pupillary dilations during ≥ 4 of 5 arithmetic tasks. None of the 5 sedated and unconscious ICU patients passed this threshold.

Conclusions: Automated pupillometry combined with mental arithmetic appears to be a promising paradigm for the detection of covert consciousness in people with brain injury. We plan to build on this study by focusing on non-communicating ICU patients in whom the level of consciousness is unknown. If some of these patients show reproducible pupillary dilation during mental arithmetic, this would suggest that the present paradigm can reveal covert consciousness in unresponsive patients in whom standard investigations have failed to detect signs of consciousness.

Introduction

It can be difficult to assess if patients with acute brain injury are conscious by means of standard clinical examinations alone because these patients must be sufficiently aroused and able to mobilize motor function (Schnakers et al., 2009b,a; Laureys et al., 2010; Di et al., 2014; Kondziella et al., 2016). Thus, standard neurological assessment often misclassifies unresponsive patients as being in a vegetative state (VS, aka. unresponsive wakefulness syndrome, UWS) (Schnakers et al., 2009b). This has important implications for prognosis and puts these patients at risk of unjustified withdrawal of life-sustaining therapy (Demertzi et al., 2011; Turgeon et al., 2011; Ong, Dhand & Diringer, 2016; Harvey et al., 2018).

Our limited knowledge of disorders of consciousness contributes to this dilemma. It is still not widely recognized that up to 15% patients are entirely unable to interact with their environment because of complete motor paralysis, despite being minimally conscious (minimal conscious state, MCS) or even fully conscious (Kondziella et al., 2016). This state of covert consciousness in completely paralyzed patients has been termed cognitive motor dissociation (Schiff, 2015). Owen and co-workers were the first to document cognitive motor dissociation in a landmark paper from 2006 (Owen et al., 2006). Herein, the authors showed that a young traffic accident victim without any signs of consciousness at the bedside, thereby fulfilling clinical criteria of VS/UWS, was able to follow commands simply by modulating her brain's metabolic activity as measured by functional magnetic resonance imaging (fMRI) (Owen et al., 2006). Thus, in the past 15 years consciousness paradigms based on fMRI and electroencephalography (EEG) that circumvent the need for motor function have been developed (Monti et al., 2010; Cruse et al., 2011; King et al., 2013; Sitt et al., 2014; Stender et al., 2014; Rohaut et al., 2017; Vanhaudenhuyse et al., 2018). However, although these technologies may detect covert consciousness, fMRI- and EEG-based paradigms are labor-intensive, expensive, logistically challenging and not readily available in the intensive care unit (ICU) (Weijer et al., 2015; Kondziella et al., 2016). A cheap and fast, easy-to-interpret point-of-care test for consciousness assessment at the bedside is clearly needed.

Portable infrared pupillometry is a new technology that may prove useful for the determination of covert consciousness in the clinically unresponsive patient. The pupillary reflex is a polysynaptic brainstem reflex under cortical modulation, i.e. cognitive processes such as decision making and mental arithmetic produce pupillary dilation (Kahneman & Beatty, 1966; Kahneman, 1973; Loewenfeldt, 1999; Marquart & de Winter, 2015; Steinhauer, Condray & Pless, 2015; Quirins et al., 2018). Hence, pupillary responses following mental arithmetic have been used to establish communication with patients with the locked-in syndrome and to detect command-following in 1 patient in MCS (Stoll et al., 2013). However, these were patients with chronic brain injury many months or years after the injury, and the technical equipment used was complex, involving a fixed bedside camera and a computer screen for the display of visual instructions (Stoll et al., 2013). Here, we wished to assess whether a convenient hand-held pupillometer and a simpler paradigm with vocal instructions allow reliable measurements of pupillary dilation during mental arithmetic as a sign of command following, and hence consciousness, in a wide range of neurological patients admitted for in-hospital care.

Methods

Objectives

We aimed to evaluate a paradigm for the assessment of consciousness and command following in patients with neurological disorders admitted to in-patient hospital care. To this end, we assessed pupillary dilation following mental arithmetic in neurological patients in the ICU and neurological ward, as well as in healthy volunteers. Sedated unconscious patients served as negative controls. We hypothesized that most (but not necessarily all) healthy volunteers and conscious neurological patients would be able to cooperate and show pupillary dilation during mental arithmetic, whereas unconscious and sedated patients would not.

Study population

We collected a convenience sample of 48 neurological patients admitted to the neuro-ICU and neurological wards at Rigshospitalet, Copenhagen University Hospital, including unsedated ICU patients with spontaneous eye opening in MCS minus (i.e. evidence of visual pursuit; $n=1$, pupillary dilation measured twice at 7 days interval) or better (conscious state; $n=1$). Five unconscious and deeply sedated/comatose patients in the ICU were recruited for negative control (Richmond Agitation-Sedation Scale score -4 , i.e. deep sedation without response to voice, but possibly movement to physical stimulation). Levels of consciousness were estimated following standard neurological bedside examination by a board-certified neurologist experienced in neurocritical care and according to established criteria (Giacino et al., 2018). Twenty healthy volunteers served as positive controls.

Automated pupillometry and mental arithmetic paradigm

The integrity of the pupillary light reflex of both eyes was checked using the NPi®-200 Pupillometer (NeuroOptics, Laguna Hills, CA 92653 USA). We documented the neurological pupil index (NPI), which is a proprietary pupillometry sum score from 0-5, with ≥ 3 indicating physiological limits (including a maximal difference between the 2 eyes of <0.7) (Chen et al., 2011; Larson & Behrends, 2015; Peinkhofer et al., 2018), pupillary diameters before and after light exposure, percentage change of pupillary diameters, and pupillary constriction and dilation velocities. Patients with non-physiological values were excluded. Then, we used the PLR®-3000 pupillometer (NeuroOptics, Laguna Hills, CA 92653 U.S.A) to track pupillary size of the right eye over time (approximately 3-5 minutes in total), while asking the participants to engage in mental arithmetic. During the examination, the examiner held the pupillometer in one hand and covered the opposite eye with the hand to avoid that changes in ambient light intensities would influence pupil size in different subjects. The set-up was identical for healthy volunteers and patients, except that patients sometimes were examined in the supine position (**Figure 1**). Each participant was asked to calculate a series of 5 arithmetic problems of moderate difficulty (21×22 , 33×32 , 55×54 , 43×44 , 81×82 ; approx. 30 seconds each) with rest periods (30 seconds) in-between. A subgroup of patients was given arithmetic problems of lesser difficulty (4×46 , 8×32 , 3×67 , 6×37 , 7×43 ; approx. 15 seconds each, with 15 seconds rest periods). We carefully explained all participants that pupillary dilation is induced solely by the efforts associated with mental arithmetic, and that it was irrelevant for our study if their calculations were correct

or not. Hence, participants were instructed not to reveal the results of their mental arithmetic but to pay attention to the task and make an honest effort.

Outcome measures

Outcome measures included pupillary diameters during periods of mental arithmetic (intervention) and relaxation (rest periods).

Statistical analysis

Data were analysed using R (R 3.4.1, R Development Core Team [2008], Vienna, Austria) by a blinded investigator. Pupillary measurements were visually assessed for quality control in a run chart. Pupillary diameter changes in each of the five mental arithmetic tasks (intervention) were assessed by comparing the period of intervention with the periods of relaxation (rest periods) before and after. Successful pupillary dilation during intervention was defined as a significantly larger median pupillary size during mental arithmetic compared to the immediate rest periods prior and after (p-value < 0.01; Wilcoxon signed-rank test, followed by Bonferroni correction). We deemed command following to be successful when a participant showed pupillary dilation in at least 4 of the 5 mental arithmetic tasks (80%).

Ethics

The Ethics Committee of the Capital Region of Denmark approved the study (journal-nr.:H-18045266). Written consent was obtained from all participants or their next-of-kin (unconscious or minimally conscious ICU patients). Data were anonymized and handled according to the European Union's Data Protection Law. The pupillometry device used in the present study (PLR®-3000 Pupillometer; NeuroOptics, Laguna Hills, CA 92653 USA) was on loan from the manufacturer; however, neither the manufacturer, nor the vendor were involved in the design or conduct of the study, data analysis or writing of the manuscript, and the authors did not receive any other monetary or non-monetary benefits.

Results

We examined 70 participants, 2 of which were excluded because of a NPi below 3, suggesting abnormal physiological pupillary function. Hence, we enrolled 68 participants: 20 healthy controls, 41 neurological patients on the ward and 7 patients in the neuro-ICU. (One patient in the neuro-ICU was measured twice, resulting in 69 assessments in total). Diagnoses reflected a wide spectrum of neurological disorders, including cerebrovascular, neuromuscular, epilepsy, trauma, neuroinfections, and multiple sclerosis. Baseline pupillary function was normal in all participants and did not differ between healthy controls (mean NPi score 4.3 ± 0.39) and neurological patients (4.3 ± 0.32 ; $p=0.812$). **Table 1** shows demographic data.

Pupillary dilation was seen in 65 of 100 (65%) measurements in healthy controls; in 58 of 100 (58%), respectively, 72 of 105 (68.6%, simpler tasks) measurements in neurological patients on the ward; and in 7 of 15 (46.67%) measurements in unsedated ICU patients. By contrast, larger pupillary diameters were

noted during 7 out of 25 (28%) measurements in comatose/sedated ICU patients (negative control group), consistent with chance occurrence.

Fourteen of 20 (70%) healthy volunteers fulfilled the prespecified criteria for successful command following, whereas this was the case for only 16 of 41 (39%) neurological patients on the ward, 1 (conscious) of 2 unsedated patients in the ICU, and 0 of 5 comatose/sedated patients in the ICU. Healthy controls had higher rates of command following than neurological patients (risk ratio 1.81, 95% CI 1.13-2.99; z-statistic 2.48; p=0.013; excluding sedated ICU patients). Reducing the degree of difficulty of the mental arithmetic task did not change the proportion of neurological patients passing criteria for command following (8/20 patients, 40% vs. 8/21 patients, 38%).

Table 2 provides details. Examples from healthy controls and neurological patients are given in **Figure 2** and **Figure 3**. Anonymized raw data are available in the *online supplemental files (DatasetS1, DatasetS2, DatasetS3)*.

Discussion

Cognitive and emotional processes evoke pupillary dilation in both humans and non-human primates, reflecting vigilance, arousal and attention (Laeng, Sirois & Gredebäck, 2012; Schneider M. et al., 2016; Becket Ebitz & Moore, 2017; McGarrigle et al., 2017; Foroughi, Sibley & Coyne, 2017). Hence, pupillary diameters may serve as an index of brain activity and mental efforts (or lack thereof) (Quirins et al., 2018). Here, we employed mental arithmetic as a paradigm for patients and healthy volunteers to control and maximize pupil dilation to signal command following. We found that a short session of mental arithmetic with simple verbal instructions, without prior training, revealed command following as detected by a handheld automated pupillometry device in 70% of healthy volunteers and 40% of conscious neurological patients.

Seventy % command following in healthy people may seem low, but this figure is very consistent with what has been reported with active EEG- and fMRI-based paradigms. For instance, in one study, 9 of 12 (75%) healthy controls produced EEG data that could be classified significantly above chance (Cruse et al., 2011); in another study, 12 of 16 healthy subjects had EEG responses to motor imagery [75.0% (95% CI: 47.6–92.7%)] (Edlow et al., 2017). Similarly, 11 of 16 healthy volunteers [68.8% (95% CI: 41.3–89.0%)] demonstrated responses within supplementary motor areas and premotor cortices when examined by a motor imagery fMRI paradigm (Edlow et al., 2017). Again, 7 of 10 (70%) healthy subjects were able to demonstrate covert command following in another motor imagery fMRI study (Bodien, Giacino & Edlow, 2017). Thus, many healthy people cannot cooperate in active paradigms. Of note, however, mental arithmetic seems to generate the most robust activation in the majority of healthy subjects for both EEG and fMRI (Harrison et al., 2017). Obviously, in the present study we used relatively difficult multiplication tasks, and easier ones such as serial 7's (Steinhauer, Condray & Pless, 2015) might have resulted in a greater fraction of participants being able to comply with our paradigm. Indeed, a few participants who were unable to comply stated that they gave up because they felt stressed by the calculations (although

reducing the level of abstraction from 2x2-differed to 1x2-differed calculations did not improve compliance). It is interesting in this regard that a recent report showed that it is possible to probe for consciousness using pupillometry also without interfering with participants' stream of consciousness by questioning them, albeit using a much more complex 'local global' auditory paradigm (Quirins et al., 2018).

To assess the feasibility of mental arithmetic and pupillometry in the clinical setting, we pragmatically enrolled a large variety of patients with neurological disorders. Not surprisingly, the rate of command following was substantially lower - around 40% - either because of mild cognitive dysfunction related to the underlying neurological condition, the mental stress associated with being admitted to hospital, the higher median age, a lower level of education or a combination of these factors, although we did not examine this specifically. It is likely that allowing training sessions until patients feel they can cooperate might have yielded a higher success rate. Although we enrolled only two unsedated ICU patients, one of them successfully participated in our paradigm, which corroborates the feasibility of our approach. Also, as expected, none of the sedated ICU patients met our prespecified criteria of command following despite spontaneous fluctuations in pupillary diameters.

Our paradigm has a few limitations that should be discussed. First, one should be aware that, in principle, pupillary dilations can occur simply as a sign of arousal rather than evidence for mathematical calculations and command following. However, to reduce the likelihood for false positive results we employed 5 serial arithmetic tasks requiring 4 correct dilations (80%) as evidence for command following. Second, physiological pupillary unrest is greater in darkness, in particular with sleepiness (Wilhelm et al., 2001), and covering the opposite eye of the participants while performing pupillometry may have induced greater fluctuations of pupillary diameters. Also, in darkness the pupils will usually be relatively large, which limits the range of pupillary movement (dilation) that might follow a command. Adding a dim light into the pupillometer would constrict the pupil into a range where dilations might be clearer. Third, participants received mental arithmetic tasks not by standardized recorded instructions but by oral instructions delivered by the experimenter; however, we believe that the convenience of oral instructions at the bedside by far outweighs the very slight bias that may possibly arise due to minor deviations in intonations, etc. Of note, statistical evaluations of pupillary data were assessed by a blinded investigator, which strengthens the robustness of our paradigm. Fourth, it should be remembered that medications such as opioids influence pupillary function (and, obviously, consciousness levels). Finally, we did not use an algorithm to remove noise from artifacts caused by head movements, blinking, and partial lid closures (Larson & Behrends, 2015; Neice et al., 2017; Behrends et al., 2018). However, the purpose of our paradigm was not to study pupillary function per se but to identify evidence for covert consciousness using a very simple bedside technique, and we therefore considered algorithms to remove noise unnecessary (and detrimental to the simplicity of the paradigm). Importantly, artifacts from eye or head movements may be more frequent in patients in the MCS than in patients with the VS/UWS and might deserve assessment as consciousness markers in their own right; this should be evaluated by future research.

Conclusions

Here we have shown that a fast and easy paradigm based on automated pupillometry and mental arithmetic is able to detect command following in healthy volunteers and conscious patients with neurological disorders admitted to in-hospital care. As a next step, we plan to focus on unsedated non-communicating ICU patients in whom the level of consciousness is unknown. Patients who show reliable pupillary dilation following mental arithmetic are likely to be conscious, whereas absence of pupillary dilation is a poor predictor of lack of consciousness. We suggest that our paradigm can be helpful to identify consciousness in non-communicating patients with acute brain injury for whom traditional bedside examination and laboratory investigations have failed to detect signs of covert consciousness. Compared to EEG and fMRI, pupillometry for the detection of command following, and ultimately covert consciousness, would offer several advantages in the clinical setting, including quick and convenient assessment at the bedside, simple analysis and low costs.

References

- Becket Ebitz R., Moore T. 2017. Selective modulation of the pupil light reflex by microstimulation of prefrontal cortex. *Journal of Neuroscience* 37:5008–5018. DOI: 10.1523/JNEUROSCI.2433-16.2017.
- Behrends M., Larson MD., Neice AE., Bokoch MP. 2018. Suppression of pupillary unrest by general anesthesia and propofol sedation. *Journal of Clinical Monitoring and Computing*. DOI: 10.1007/s10877-018-0147-y.
- Bodien YG., Giacino JT., Edlow BL. 2017. Functional MRI Motor Imagery Tasks to Detect Command Following in Traumatic Disorders of Consciousness. *Frontiers in Neurology* 8:688. DOI: 10.3389/fneur.2017.00688.
- Chen J., Gombart Z., Rogers S., Gardiner S., Cecil S., Bullock R. 2011. Pupillary reactivity as an early indicator of increased intracranial pressure: The introduction of the neurological pupil index. *Surgical Neurology International* 2:82. DOI: 10.4103/2152-7806.82248.
- Cruse D., Chennu S., Chatelle C., Bekinschtein TA., Fernández-Espejo D., Pickard JD., Laureys S., Owen AM. 2011. Bedside detection of awareness in the vegetative state: a cohort study. *Lancet* 378:2088–94. DOI: 10.1016/S0140-6736(11)61224-5.
- Demertzi A., Ledoux D., Bruno M-A., Vanhaudenhuyse A., Gosseries O., Soddu A., Schnakers C., Moonen G., Laureys S. 2011. Attitudes towards end-of-life issues in disorders of consciousness: a European survey. *Journal of Neurology* 258:1058–1065. DOI: 10.1007/s00415-010-5882-z.
- Di H., Nie Y., Hu X., Tong Y., Heine L., Wannez S., Huang W., Yu D., He M., Thibaut A., Schnakers C., Laureys S. 2014. Assessment of visual fixation in vegetative and minimally conscious states. *BMC neurology* 14:147. DOI: 10.1186/1471-2377-14-147.
- Edlow BL., Chatelle C., Spencer CA., Chu CJ., Bodien YG., O'Connor KL., Hirschberg RE., Hochberg LR., Giacino JT., Rosenthal ES., Wu O. 2017. Early detection of consciousness in patients with acute severe traumatic brain injury. *Brain* 140:23–31. DOI: 10.1093/brain/awx176.
- Foroughi CK., Sibley C., Coyne JT. 2017. Pupil size as a measure of within-task learning. *Psychophysiology* 54:1436–1443. DOI: 10.1111/psyp.12896.
- Giacino JT., Katz DI., Schiff ND., Whyte J., Ashman EJ., Ashwal S., Barbano R., Hammond FM., Laureys S., Ling GSF., Nakase-Richardson R., Seel RT., Yablon S., Getchius TSD., Gronseth GS., Armstrong MJ. 2018. Practice guideline update recommendations summary: Disorders of consciousness. *Neurology*:10.1212/WNL.0000000000005926. DOI: 10.1212/WNL.0000000000005926.
- Harrison AH., Noseworthy MD., Reilly JP., Guan W., Connolly JF. 2017. EEG and fMRI agree: Mental arithmetic is the easiest form of imagery to detect. *Consciousness and Cognition* 48:104–116. DOI: 10.1016/j.concog.2016.10.006.
- Harvey D., Butler J., Groves J., Manara A., Menon D., Thomas E., Wilson M. 2018. Management of perceived devastating brain injury after hospital admission: a consensus statement from stakeholder professional organizations. *British Journal of Anaesthesia* 120:138–145. DOI: 10.1016/j.bja.2017.10.002.
- Kahneman D. 1973. *Attention and Effort*. Prentice-Hall.

283 Kahneman D., Beatty J. 1966. Pupil diameter and load on memory. *Science (New York, N.Y.)* 154:1583–5.

284 King JR., Faugeras F., Gramfort A., Schurger A., El Karoui I., Sitt JD., Rohaut B., Wacongne C., Labyt E.,
285 Bekinschtein T., Cohen L., Naccache L., Dehaene S. 2013. Single-trial decoding of auditory novelty
286 responses facilitates the detection of residual consciousness. *NeuroImage* 83:726–38. DOI:
287 10.1016/j.neuroimage.2013.07.013.

288 Kondziella D., Friberg CK., Frokjaer VG., Fabricius M., Møller K. 2016. Preserved consciousness in
289 vegetative and minimal conscious states: systematic review and meta-analysis. *Journal of*
290 *neurology, neurosurgery, and psychiatry* 87:485–92. DOI: 10.1136/jnnp-2015-310958.

291 Laeng B., Sirois S., Gredebäck G. 2012. Pupillometry: A Window to the Preconscious? *Perspectives on*
292 *psychological science : a journal of the Association for Psychological Science* 7:18–27. DOI:
293 10.1177/1745691611427305.

294 Larson MD., Behrends M. 2015. Portable infrared pupillometry: A review. *Anesthesia and Analgesia*
295 120:1242–1253. DOI: 10.1213/ANE.0000000000000314.

296 Laureys S., Celesia GG., Cohadon F., Lavrijsen J., León-Carrión J., Sannita WG., Szabon L., Schmutzhard E.,
297 von Wild KR., Zeman A., Dolce G., European Task Force on Disorders of Consciousness. 2010.
298 Unresponsive wakefulness syndrome: a new name for the vegetative state or apallic syndrome.
299 *BMC medicine* 8:68. DOI: 10.1186/1741-7015-8-68.

300 Loewenfeldt I. 1999. *The Pupil: Anatomy, Physiology, and Clinical Applications*. Butterworth-Heinemann.

301 Marquart G., de Winter J. 2015. Workload assessment for mental arithmetic tasks using the task-evoked
302 pupillary response. *PeerJ* 1. DOI: 10.7717/peerj-cs.16.

303 McGarrigle R., Dawes P., Stewart AJ., Kuchinsky SE., Munro KJ. 2017. Pupillometry reveals changes in
304 physiological arousal during a sustained listening task. *Psychophysiology* 54:193–203. DOI:
305 10.1111/psyp.12772.

306 Monti MM., Vanhaudenhuyse A., Coleman MR., Boly M., Pickard JD., Tshibanda L., Owen AM., Laureys S.
307 2010. Willful Modulation of Brain Activity in Disorders of Consciousness. *The New England journal*
308 *of medicine* 362:579–589. DOI: 10.1056/NEJMoa0905370.

309 Neice AE., Behrends M., Bokoch MP., Seligman KM., Conrad NM., Larson MD. 2017. Prediction of Opioid
310 Analgesic Efficacy by Measurement of Pupillary Unrest. *Anesthesia & Analgesia* 124:915–921. DOI:
311 10.1213/ANE.0000000000001728.

312 Ong CJ., Dhand A., Diringer MN. 2016. Early Withdrawal Decision-Making in Patients with Coma After
313 Cardiac Arrest: A Qualitative Study of Intensive Care Clinicians. *Neurocritical Care* 25:258–265. DOI:
314 10.1007/s12028-016-0275-5.

315 Owen AM., Coleman MR., Boly M., Davis MH., Laureys S., Pickard JD. 2006. Detecting awareness in the
316 vegetative state. *Science (New York, N.Y.)* 313:1402. DOI: 10.1126/science.1130197.

317 Peinkhofer C., Martens P., Grand J., Truelsen T., Knudsen GM., Kjaergaard J., Kondziella D. 2018.
318 Influence of Strategic Cortical Infarctions on Pupillary Function. *Frontiers in Neurology* 9:916. DOI:
319 10.3389/fneur.2018.00916.

- Quirins M., Marois C., Valente M., Seassau M., Weiss N., El Karoui I., Hochmann J-R., Naccache L. 2018. Conscious processing of auditory regularities induces a pupil dilation. *Scientific reports* 8:14819. DOI: 10.1038/s41598-018-33202-7.
- Rohaut B., Raimondo F., Galanaud D., Valente M., Sitt JD., Naccache L. 2017. Probing consciousness in a sensory-disconnected paralyzed patient. *Brain injury*:1–6. DOI: 10.1080/02699052.2017.1327673.
- Schiff ND. 2015. Cognitive Motor Dissociation Following Severe Brain Injuries. *JAMA Neurology* 72:1413. DOI: 10.1001/jamaneurol.2015.2899.
- Schnakers C., Perrin F., Schabus M., Hustinx R., Majerus S., Moonen G., Boly M., Vanhaudenhuyse A., Bruno M-A., Laureys S. 2009a. Detecting consciousness in a total locked-in syndrome: an active event-related paradigm. *Neurocase* 15:271–7. DOI: 10.1080/13554790902724904.
- Schnakers C., Vanhaudenhuyse A., Giacino J., Ventura M., Boly M., Majerus S., Moonen G., Laureys S. 2009b. Diagnostic accuracy of the vegetative and minimally conscious state: clinical consensus versus standardized neurobehavioral assessment. *BMC neurology* 9:35. DOI: 10.1186/1471-2377-9-35.
- Schneider M., Hathway P., Leuchs L., Samann P.G., Czisch M., Spoormaker V.I. 2016. Spontaneous pupil dilations during the resting state are associated with activation of the salience network. *NeuroImage* 139:189–201. DOI: 10.1016/j.neuroimage.2016.06.011.
- Sitt JD., King J-RR., El Karoui I., Rohaut B., Faugeras F., Gramfort A., Cohen L., Sigman M., Dehaene S., Naccache L., Karoui I El., Rohaut B., Faugeras F., Gramfort A., Cohen L., Sigman M., Dehaene S., Naccache L. 2014. Large scale screening of neural signatures of consciousness in patients in a vegetative or minimally conscious state. *Brain* 137:2258–2270. DOI: 10.1093/brain/awu141.
- Steinhauer SR., Condray R., Pless ML. 2015. Pharmacological Isolation of Cognitive Components Influencing the Pupillary Light Reflex. *Journal of Ophthalmology* 2015:1–7. DOI: 10.1155/2015/179542.
- Stender J., Gosseries O., Bruno M-A., Charland-Verville V., Vanhaudenhuyse A., Demertzi A., Chatelle C., Thonnard M., Thibaut A., Heine L., Soddu A., Boly M., Schnakers C., Gjedde A., Laureys S. 2014. Diagnostic precision of PET imaging and functional MRI in disorders of consciousness: a clinical validation study. *Lancet* 384:514–22. DOI: 10.1016/S0140-6736(14)60042-8.
- Stoll J., Chatelle C., Carter O., Koch C., Laureys S., Einhäuser W. 2013. Pupil responses allow communication in locked-in syndrome patients. *Current Biology* 23:R647–R648. DOI: 10.1016/j.cub.2013.06.011.
- Turgeon AF., Lauzier F., Simard J-F., Scales DC., Burns KEA., Moore L., Zygun DA., Bernard F., Meade MO., Dung TC., Ratnapalan M., Todd S., Harlock J., Fergusson DA. 2011. Mortality associated with withdrawal of life-sustaining therapy for patients with severe traumatic brain injury: a Canadian multicentre cohort study. *CMAJ : Canadian Medical Association journal = journal de l'Association medicale canadienne* 183:1581–8. DOI: 10.1503/cmaj.101786.
- Vanhaudenhuyse A., Charland-Verville V., Thibaut A., Chatelle C., Tshibanda JFL., Maudoux A., Faymonville ME., Laureys S., Gosseries O. 2018. Conscious while being considered in an unresponsive wakefulness syndrome for 20 years. *Frontiers in Neurology* 9:1–9. DOI: 10.3389/fneur.2018.00671.

360 Weijer C., Bruni T., Gofton T., Young GB., Norton L., Peterson A., Owen AM. 2015. Ethical considerations
 361 in functional magnetic resonance imaging research in acutely comatose patients. *Brain : a journal*
 362 *of neurology*. DOI: 10.1093/brain/awv272.

363 Wilhelm B., Giedke H., Lüdtke H., Bittner E., Hofmann A., Wilhelm H. 2001. Daytime variations in central
 364 nervous system activation measured by a pupillographic sleepiness test. *Journal of sleep research*
 365 10:1–7.

366

Table 1

This table shows demographic data, including neurological diagnoses. “Stroke” includes hemorrhagic and ischemic stroke, “neuromuscular” includes Guillain-Barré syndrome, chronic inflammatory demyelinating polyradiculopathy, Isaacs’ syndrome, Pompe’s disease and multifocal motor neuropathy, “epilepsy” denotes epilepsy with or without structural cause on magnetic resonance imaging. CS = conscious, ICU = intensive care unit, IQR = interquartile range, MCS = minimally conscious state, N = number of subjects, SAH = subarachnoid hemorrhage, TBI = traumatic brain injury. * All mental arithmetic tasks involved 2x2-differed calculations (e.g. 33 x 32), except for 1x2-differed calculations (e.g. 8 x 32) in neurological patients indicated with (*). ** This unsedated ICU patient in MCS with a pontine hemorrhagic stroke was examined twice at 7 days interval but failed to show command following during mental arithmetic in both sessions. *** Other diagnoses, not listed above, include relapsing remitting multiple sclerosis, unspecified sensory disturbances, brain abscess, anoxic-ischemic encephalopathy and hemangioblastoma

Table 2

This table depicts the rate of successful command following by mental arithmetic in healthy volunteers, conscious neurological patients on the ward, minimally or fully conscious patients in the ICU, and comatose/sedated ICU patients. Successful command following was defined by ≥4 significant pupillary dilations during 5 mental arithmetic tasks. CS = conscious, ICU = intensive care unit, MCS = minimally conscious state, N = number of subjects. * All mental arithmetic tasks involved 2x2-differed calculations (e.g. 33 x 32), except for 1x2-differed calculations (e.g. 8 x 32) in neurological patients indicated with (*). ** This unsedated ICU patient in MCS with a pontine hemorrhagic stroke was examined twice at 7 days interval but failed to show command following during mental arithmetic in both sessions

Figure 1

We used the PLR®-3000 pupillometer (NeuroOptics, Laguna Hills, CA 92653 U.S.A), an automated handheld device (**A** and **B**), to track pupillary size, while asking patients and healthy volunteers (one shown here; permission obtained) to perform mental arithmetic. The examiner holds the pupillometer in one hand and covers the other eye with the hand (**A**). The set-up is identical for healthy volunteers and patients, except that patients may be better examined in the supine position.

Figure 2

This figure shows results from participants with successful command following, detected by automated pupillometry, during a mental arithmetic paradigm: two patients admitted to the neurological ward with diagnoses of multifocal motor neuropathy (**A**) and Guillain-Barré syndrome (**B**), respectively, a healthy participant (**C**), and a conscious 34-year old male with the pharyngeal-cervical-brachial variant of Guillain-Barré syndrome admitted to the ICU (**D**). Data from each subject are presented twice and in 2 different formats (raw measurements above, annotated data below). Minor artifacts due to blinking or eye movements are seen in A-C, but not D (probably because of facial and oculomotor nerve palsies). Color code: Periods with mental arithmetic are shown in green, rest periods in yellow. Numbers on the x-axis (“0-100-200-300”) denote time in seconds. Pupillary sizes during mental arithmetic were significantly larger ($p\text{-value} < 0.0001$) than during rest periods, consistent with pupillary dilation, in all five tasks for each of the four participants. # = $p\text{-value} < 0.0001$; ✓ = pupillary dilation; n = number of measurements; m = median pupillary size, mm = millimeter

Figure 3

This figure depicts results from a healthy volunteer with unsuccessful command following (**A**) and a 62-year old male in the ICU with subarachnoid hemorrhage and deep sedation (Richmond Agitation-Sedation Scale score of -4) who served as a negative control (**B**). Data from each subject are presented twice and in 2 different formats (raw measurements above, annotated data below). Minor artifacts due to blinking or eye movements are seen in A, but not B (because of sedation-induced impairment of the blink reflex). In the healthy volunteer, pupillary dilation was noted in only three out of five mental arithmetic tasks, which did not meet our prespecified criteria for successful command following (≥ 4 pupillary dilations, 80%). Minor random fluctuations in the pupillary diameter are seen in the unconscious sedated ICU patient. # = $p\text{-value} < 0.0001$; ✓ = pupillary dilation; ÷ = absence of pupillary dilation; n = number of measurements; m = median pupillary size, mm = millimeter

Figure 1

Pupillary dilation during mental arithmetic assessed by automated pupillometry

We used the PLR®-3000 pupillometer (NeuroOptics, Laguna Hills, CA 92653 U.S.A), an automated handheld device (**A** and **B**), to track pupillary size, while asking patients and healthy volunteers (one shown here; permission obtained) to perform mental arithmetic. The examiner holds the pupillometer in one hand and covers the other eye with the hand (**A**). The set-up is identical for healthy volunteers and patients, except that patients may be better examined in the supine position.




Figure 2

Pupillometry data from four patients with successful command following

This figure shows results from participants with successful command following, detected by automated pupillometry, during a mental arithmetic paradigm: two patients admitted to the neurological ward with diagnoses of multifocal motor neuropathy (**A**) and Guillain-Barré syndrome (**B**), respectively, a healthy participant (**C**), and a conscious 34-year old male with the pharyngeal-cervical-brachial variant of Guillain-Barré syndrome admitted to the ICU (**D**). Data from each subject are presented twice and in 2 different formats (raw measurements above, annotated data below). Minor artifacts due to blinking or eye movements are seen in A-C, but not D (probably because of facial and oculomotor nerve palsies). Color code: Periods with mental arithmetic are shown in green, rest periods in yellow. Numbers on the x-axis ("0-100-200-300") denote time in seconds. Pupillary sizes during mental arithmetic were significantly larger ($p\text{-value} < 0.0001$) than during rest periods, consistent with pupillary dilation, in all five tasks for each of the four participants. # = $p\text{-value} < 0.0001$; ✓ = pupillary dilation; n = number of measurements; m = median pupillary size, mm = millimeter




Figure 3

Pupillometry data from two patients without command following

This figure depicts results from a healthy volunteer with unsuccessful command following (**A**) and a 62-year old male in the ICU with subarachnoid hemorrhage and deep sedation (Richmond Agitation-Sedation Scale score of -4) who served as a negative control (**B**). Data from each subject are presented twice and in 2 different formats (raw measurements above, annotated data below). Minor artifacts due to blinking or eye movements are seen in A, but not B (because of sedation-induced impairment of the blink reflex). In the healthy volunteer, pupillary dilation was noted in only three out of five mental arithmetic tasks, which did not meet our prespecified criteria for successful command following (≥ 4 pupillary dilations, 80%). Minor random fluctuations in the pupillary diameter are seen in the unconscious sedated ICU patient. # = p-value < 0.0001 ; ✓ = pupillary dilation; ÷ = absence of pupillary dilation; n = number of measurements; m = median pupillary size, mm = millimeter




Table 1 (on next page)

Demographic data

This table shows demographic data, including neurological diagnoses. “Stroke” includes hemorrhagic and ischemic stroke, “neuromuscular” includes Guillain-Barré syndrome, chronic inflammatory demyelinating polyradiculopathy, Isaacs’ syndrome, Pompe’s disease and multifocal motor neuropathy, “epilepsy” denotes epilepsy with or without structural cause on magnetic resonance imaging. CS = conscious, ICU = intensive care unit, IQR = interquartile range, MCS = minimally conscious state, N = number of subjects, SAH = subarachnoid hemorrhage, TBI = traumatic brain injury. * All mental arithmetic tasks involved 2x2-difficult calculations (e.g. 33 x 32), except for 1x2-difficult calculations (e.g. 8 x 32) in neurological patients indicated with (*). ** This unsedated ICU patient in MCS with a pontine hemorrhagic stroke was examined twice at 7 days interval but failed to show command following during mental arithmetic in both sessions. *** Other diagnoses, not listed above, include relapsing remitting multiple sclerosis, unspecified sensory disturbances, brain abscess, anoxic-ischemic encephalopathy and hemangioblastoma

	Healthy volunteers	Neurological patients	Neurological patients *	ICU patients, MCS or CS	ICU patients, coma
N	20	20	21	2	5
Female	10 (50%)	9 (45%)	8 (38%)	2 (66%)	2 (40%)
Age in years, median (IQR)	34.5 (29-47)	60.5 (51-68)	50 (41-70)	34 (34-34)	62 (55-64)
Stroke	-	2	2	1 **	0
SAH	-	0	1	0	2
TBI	-	0	0	0	2
Epilepsy	-	5	2	0	0
Neuromuscular	-	9	10	1	0
Other ***	-	4	6	0	1

1

Table 2 (on next page)

Command following by mental arithmetic

This table depicts the rate of successful command following by mental arithmetic in healthy volunteers, conscious neurological patients on the ward, minimally or fully conscious patients in the ICU, and comatose/sedated ICU patients. Successful command following was defined by ≥ 4 significant pupillary dilations during 5 mental arithmetic tasks. CS = conscious, ICU = intensive care unit, MCS = minimally conscious state, N = number of subjects. * All mental arithmetic tasks involved 2x2-differed calculations (e.g. 33 x 32), except for 1x2-differed calculations (e.g. 8 x 32) in neurological patients indicated with (*). ** This unsedated ICU patient in MCS with a pontine hemorrhagic stroke was examined twice at 7 days interval but failed to show command following during mental arithmetic in both sessions

	Healthy volunteers	Neurological patients	Neurological patients *	ICU patients, MCS or CS	ICU patients, coma/sedation
N	20	20	21	2	5
0 significant	2	3	1	1**	1
1 significant	2	1	1	0	2
2 significant	0	2	1	1**	1
3 significant	2	6	10	0	1
4 significant	13	5	4	0	0
5 significant	1	3	4	1	0
Successful	70% (n=14/20)	40% (n=8/20)	38% (n=8/21)	33% (n=1/3)	0% (n=0/5)

1