

# Gene expression in *Tribolium castaneum* life stages: Identifying a species-specific target for pest control applications

Lindsey C Perkin <sup>Corresp., 1</sup>, Brenda Oppert <sup>2</sup>

<sup>1</sup> Southern Plains Agricultural Research Center, USDA, Agricultural Research Service, College Station, TX, United States

<sup>2</sup> Center for Grain and Animal Health Research, USDA, Agricultural Research Service, Manhattan, KS, United States

Corresponding Author: Lindsey C Perkin  
Email address: lindsey.perkin@ars.usda.gov

The red flour beetle, *Tribolium castaneum*, is a major agricultural pest of post-harvest products and stored grain. Control of *T. castaneum* in stored products and grain is primarily by fumigants and sprays, but insecticide resistance is a major problem, and new control strategies are needed. *T. castaneum* is a genetic model for coleopterans, and the reference genome can be used for discovery of candidate gene targets for molecular-based control, such as RNA interference. Gene targets need to be pest specific, and ideally, they are expressed at low levels for successful control. Therefore, we sequenced the transcriptome of four major life stages of *T. castaneum*, sorted data into groups based on high or low expression levels, and compared relative gene expression among all life stages. We narrowed our candidate gene list to a cuticle protein gene (CPG) for further analysis. We found that the CPG sequence was unique to *T. castaneum* and expressed only in the larval stage. RNA interference targeting CPG in newly-emerged larvae caused a significant ( $p < 0.05$ ) decrease in CPG expression (1,491-fold) compared to control larvae and 64% mortality over 18 d. RNA-Seq of survivors after 18 d identified changes in the expression of other genes as well, including 52 long noncoding RNAs. Expression of three additional cuticle protein genes were increased and two chitinase genes were decreased in response to injection of CPG dsRNA. The data demonstrate that RNA-Seq can identify genes important for insect survival and thus may be used to develop novel biologically-based insect control products.

**Gene expression in *Tribolium castaneum* life stages: Identifying a species-specific target for pest control applications**

Lindsey C. Perkin<sup>§</sup>, and Brenda Oppert

USDA, Agricultural Research Service, Southern Plains Agricultural Research Center, 2881 F&B Road, College Station, TX 77845  
USDA, Agricultural Research Service, Center for Grain and Animal Health Research, 1515 College Avenue, Manhattan, KS 66502

Running title: *Tribolium castaneum* life stage gene expression

<sup>§</sup>Corresponding author

Email addresses:

LCP: [lindsey.perkin@ars.usda.gov](mailto:lindsey.perkin@ars.usda.gov)

BO: [brenda.oppert@ars.usda.gov](mailto:brenda.oppert@ars.usda.gov)

# Abstract

The red flour beetle, *Tribolium castaneum*, is a major agricultural pest of post-harvest products and stored grain. Control of *T. castaneum* in stored products and grain is primarily by fumigants and sprays, but insecticide resistance is a major problem, and new control strategies are needed. *T. castaneum* is a genetic model for coleopterans, and the reference genome can be used for discovery of candidate gene targets for molecular-based control, such as RNA interference. Gene targets need to be pest specific, and ideally, they are expressed at low levels for successful control. Therefore, we sequenced the transcriptome of four major life stages of *T. castaneum*, sorted data into groups based on high or low expression levels, and compared relative gene expression among all life stages. We narrowed our candidate gene list to a cuticle protein gene (CPG) for further analysis. We found that the CPG sequence was unique to *T. castaneum* and expressed only in the larval stage. RNA interference targeting CPG in newly-emerged larvae caused a significant ( $p<0.05$ ) decrease in CPG expression (1,491-fold) compared to control larvae and 64% mortality over 18 d. RNA-Seq of survivors after 18 d identified changes in the expression of other genes as well, including 52 long noncoding RNAs. Injection of CPG dsRNA also resulted in increased expression of three additional cuticle protein genes and reduced expression of two chitinase genes in *T. castaneum* larvae. The data demonstrate that RNA-Seq can identify genes important for insect survival and thus may be used to develop novel biologically-based insect control products.

# Introduction

The red flour beetle, *Tribolium castaneum*, is a pest of stored grain commodities. Traditional control methods for *T. castaneum* and other stored product beetles are rapidly becoming less effective, mostly because insect populations are developing resistance to pesticide treatments (Boyer *et al.* 2012). For example, storage pests around the globe are developing high resistance levels to one of the most common grain fumigants, phosphine (Opit *et al.*, 2012; Pimentel *et al.*, 2010). Thus, there is a need for new pest control strategies, and we are evaluating genetic-based treatments with target specificity and less damage to the environment, including the application of RNA interference (RNAi; Baum *et al.*, 2007; Noh *et al.*, 2012).

*T. castaneum* is a useful model to identify candidate genes because it has a sequenced genome (Tribolium Genome Sequencing Consortium, 2008) and responds well to RNAi when dsRNA is injected (RNAi; Brown *et al.*, 1999; Tomoyasu & Denell, 2004; Aronstein *et al.*, 2011; Miller *et al.*, 2012). Oral RNAi was successful in *T. castaneum* larvae against genes encoding vATPase (Whyard *et al.*, 2009), inhibitor of apoptosis (Cao *et al.*, 2018), and a voltage-gated sodium ion channel (Haim *et al.*, 2016). However, we and others have not had success with oral RNAi in *T. castaneum* (unpublished data, Palli, 2014). Many factors may influence RNAi efficacy in insects, such as target sequence specificity, concentration and length of dsRNA, persistence of silencing effect in the target pest, and nucleases counteracting the effect of dsRNA (Huvenne & Smagghe 2010; Allen & Walker, 2012; Lomate & Bonning, 2016; Guan *et al.*, 2018, Cao *et al.*, 2018). In the meantime, we have focused on the identification of gene targets with low expression requiring lower doses of dsRNA, and those that are expressed in critical feeding stages (larvae and adults) to improve the efficacy of oral RNAi.

63

64 The iBeetle project (<http://ibeetle-base.uni-goettingen.de>) conducted a large-scale RNAi screen  
 65 in *T. castaneum* larvae and pupae with injected dsRNA, and various phenotypes were observed,  
 66 including mortality and developmental abnormalities (Schmitt-Engel *et al.*, 2015). From this  
 67 screen, eleven genes were identified as potential pest control targets (Donitz *et al.*, 2015; Ulrich  
 68 *et al.*, 2015). These genes encode mostly products with GO terms related to the proteasome, and  
 69 mortality was observed after injection of larvae, pupae and adults. Some of these genes have  
 70 orthologs in other species, such as *Drosophila melanogaster*, *Aedes aegypti*, *Apis mellifera*, and  
 71 *Acyrtosiphon pisum*, and therefore RNAi or other molecular strategies based on these genes  
 72 may not be species-specific and may not be practical for insect control. Knorr et al. (2018)  
 73 selected 50 genes from the iBeetle study that resulted in  $\geq 90\%$  mortality to evaluate for RNAi in  
 74 *Diabrotica vigifera vigifera* and found that feeding adults dsRNA targeting 20 genes resulted in  
 75 mortality, and 36 retarded growth. While this work is important to highlight genes that have  
 76 application across species, the usefulness in the field may be limited due to off-target effects.

77

78 Previously, we applied RNA-Seq to the major developmental stages of *T. castaneum* to identify  
 79 highly expressed cysteine peptidase genes in feeding stages (larvae and adults) with likely  
 80 digestive functions (Perkin *et al.*, 2016). A similar approach was used across the life stages of *D.*  
 81 *melanogaster* to reveal new transcribed regions and small non-coding RNAs, demonstrating that  
 82 transcriptome sequencing can discover new elements and help to annotate genomes of even the  
 83 most well studied insects (Graveley *et al.*, 2011). Transcriptome sequencing of developmental  
 84 stages of the non-model oriental fruit fly, *Bactocera dorsalis*, and the harlequin bug, *Murgantia*

*histrionica*, led to a list of candidate developmental and insecticide resistance genes for downstream studies (Shen *et al.*, 2011; Sparks *et al.*, 2017).

In the present study, we examined gene expression patterns in *T. castaneum* life stages (adults, eggs, larvae, and pupae), and used statistical analysis to sort expression patterns into groups of low and high expression. Our hypothesis was that genes with low expression can be more successfully targeted with RNAi or other applied control methods because the dose of dsRNA required to knockdown transcript expression will be lower, and highly expressed genes and gene expression patterns may be useful in understanding the physiology of the insect, or to identify promoters for transgenic applications. Previously, we found that some genes with high expression (i.e. those encoding cysteine peptidases) were important to insect physiology and were duplicated in the genome, and complete knockdown of these genes induced a compensation response and upregulation of other closely related genes (Perkin *et al.*, 2016). Through screening of potential targets in the present study, we identified a unique cuticle protein gene (CPG) expressed in larvae, and we provide functional evidence for its potential as a molecular target for pest control. We further evaluated the effect of CPG RNAi on the expression of other genes, defining new potential metabolic connections between CPG and other genes.

## Materials & Methods

### *Insects*

Insect strains and collection protocols were as previously described (Perkin *et al.*, 2016). Briefly, the *T. castaneum* lab colony used in this study was originally collected from a grain bin in Kansas and has been maintained on 95% wheat flour and 5% brewer's yeast at 28°C, 75%

R.H., 0L:24D. Insects were subcultured from the laboratory colony, and specific life stages were removed. Adults were collected at 3-7 days post-eclosion and were mixed sex. Eggs were sifted from diet 24-48 hours after oviposition and were separated from the fine particulate with a brush. Larvae were collected at a late instar stage that was actively feeding (approximately 14 days post hatch). Pupae included those with pigmentation in the eye, but not the elytra.

#### *Library preparation and RNA sequencing of life stages*

RNA was collected as three independent biological replicates from mixed sex groups of each life stage (10 adults, approximately 500 eggs, 10 larvae, and 10 pupae per replicate). Tissue was pulverized in TRIZOL (BulletBlender, Next Advance Inc., Averill Park, NY, USA) at speed 8 for 2 min with RNase-free zirconium oxide beads. RNA extraction and purification were with a Zymo mini prep kit (Irvine, CA, USA). From the total RNA, DIRECTbeads (Agilent, Santa Clara, CA, USA) were used to isolate polyA mRNA, and libraries were made with a 200 bp RNA-Seq v2 kit (Life Technologies, Grand Island, NY, USA). Samples were sequenced on 318v2 chips on the Ion Torrent Personal Genome Machine (PGM, Life Technologies). Each run provided approximately 1-5 million reads, with a total of 5-12 million reads per life stage (Perkin *et al.*, 2016). Life stage sequences were deposited at NCBI SRA as part of BioProject PRJNA299695.

#### *Data analysis*

Relative gene expression was analyzed using ArrayStar (Lasergene Genomics Suite v14, DNASTAR, Madison, WI, USA) by mapping reads to the NCBI Tcas5.2 genome build. Read counts were normalized by Reads Per Kilobase of template per Million mapped reads (RPKM,

Mortazavi *et al.*, 2008). All data were filtered to groups with low expression (RPKM greater than 2 less than 8  $\log_2$ ; i.e., the  $\log_2$  of the RPKM value was between 2 and 8, not inclusive) and high expression (RPKM greater than 8  $\log_2$ ) for comparisons and biological screening. We use a cutoff of RPKM greater than 2 to avoid underrepresented genes that can skew data.

Gene ontology (GO) terms were obtained and enrichment of GO terms was analyzed in BLAST2GO PRO (version 4.1.5, Valencia, Spain, Götz *et al.*, 2008). Over-representation tests were performed with GO terms associated with genes in low and high categories in each life stage, using the program g:Profiler and the g:GOST function, which provides statistical enrichment analysis (<http://biit.cs.ut.ee/gprofiler/index.cgi>; (Reimand *et al.*, 2011). In g:Profiler, results were limited to *T. castaneum* and the main GO categories: biological process (BP), molecular function (MF) and cellular component (CC). Significance was determined using the False Discovery Rate (FDR) multiple-testing correction threshold of 0.05 (Benjamini and Hochberg 1995). Enriched GO terms also were obtained from the pairwise comparison of gene expression in larvae injected with cuticle protein gene (CPG, LOC103313766) dsRNA compared to mock-injected, where the test set was the significant ( $p < 0.05$ ) differentially expressed genes, and the reference set was expression from all genes. GO terms were limited to  $p < 0.05$  (after FDR correction) and collapsed to the most specific description of each term.

### *Candidate gene selection*

Genes were first filtered to only include those with expression levels between RPKM 2 and 8 ( $\log_2$ ). The resulting gene list included 181 genes. We then filtered to only include genes expressed at low levels in larvae because a field application would rely on genes expressed



during the most active feeding stage, reducing the list to 139 genes. Our next filter was a manually curated gene set and was based on GO analysis and gene annotations, selecting genes with a function potentially necessary for survival.

The final filter submitted potential genes to the program OfftargetFinder (Good *et al.*, 2016). This program checks gene specificity by searching 101 arthropod transcriptomes, 21mers at a time, to the 1000 Insect Transcriptome Evolution (1KITE) dataset (<http://www.1kite.org>) and is specifically designed to identify potential problems in designing dsRNA for insect control. The algorithm provides visualization of specific regions of the target gene that may cause off-target effects and allows for dsRNA to be designed to regions with increased specificity to the target pest. This filter limited our gene list to only three genes. We selected CPG as our primary target because the 3' region of CPG was unique to *T. castaneum*, and this region was used to design primers for dsRNA.

# *RNAi*

Primers for dsRNA targeting CPG were designed similar to Perkin *et al.* (2017a) via Primer-BLAST (<http://www.ncbi.nlm.nih.gov/tools/primer-blast/>) using default parameters, and primer specificity was evaluated against the *T. castaneum* genome. A primary set of primers was designed to amplify CPG, and secondary primers were designed to amplify the 3' region to minimize off-target effects..

The primary set of primers (Forward: ATAATCAAGCCCGTTTCCAACA, Reverse: AATCACGACTACAAACATTCTTAGG) amplified CPG in 25 µl PCR reactions using

genomic DNA template from the lab strain (2.5  $\mu$ l 10X buffer, 2.0  $\mu$ l 2.5 mM dNTP, 0.5  $\mu$ l 10  $\mu$ M forward primer, 0.5  $\mu$ l 10  $\mu$ M reverse primer, 0.125  $\mu$ l JumpStart AccuTaq DNA polymerase (Sigma-Aldrich, Saint Louis, MO, USA), 2  $\mu$ l genomic DNA, 17.375  $\mu$ l nuclease-free water) and thermal cycle conditions as specified by the JumpStart AccuTaq DNA polymerase product information guide (denaturation for 30 sec at 95°C, 30 cycles of 95°C for 30 sec, primer annealing at  $T_m$  55.5°C for 30 sec, and extension at 68°C for 30 sec, final extension at 68°C for 5 min, and hold at 4°C). The product was assessed on a 1% agarose E-gel (ThermoFisher, Waltham, MA, USA) to ensure the amplified region was the correct size.

A second round of PCR was done with the secondary set of primers (Forward: TATTCGTCTGTCGTCGCTCC, Reverse: AATCACGACTACAAACATTCTTAGG) for specific targeting of CPG. The secondary PCR reaction was 100  $\mu$ l and the product from the primary PCR reaction was used as the template (10  $\mu$ l 10X buffer, 8  $\mu$ l 2.5 mM dNTP, 2  $\mu$ l T7 forward primer, 2  $\mu$ l T7 reverse primer, 0.5  $\mu$ l JumpStart AccuTaq DNA polymerase, 10  $\mu$ l primary PCR product, 67.5  $\mu$ l nuclease-free water). The thermal reaction profile was as follows: denaturation for 30 sec at 95°C, 5 cycles of 95°C for 30 sec, primer  $T_m$  for 30 sec, 72°C for 30 sec, 29 cycles of 95°C for 30 sec, primer  $T_m$  plus 5°C for 30 sec, and 72°C for 30 sec, extension for 5 min at 68°C and hold at 4°C. The secondary PCR primers had a T7 construct attached to the 5' end (TAATACGACTCACTATAGGG), and thus 5°C was added to the second round of amplification as suggested in the kit manual. The secondary PCR products were checked again via 1% agarose E-gel for appropriate length (150 bp) and sufficient amplification.

PCR products from the secondary amplification were used to make dsRNA via MEGAscript T7 kit (ThermoFisher) according to the kit protocol (8 µl dNTP mix, 2 µl 10X reaction buffer, 8 µl secondary PCR products with T7 construct, 2 µl enzyme mix), and after a 6 h incubation period at 37°C with constant mixing, products were purified via MEGAclean kit (ThermoFisher). Size and quantity were verified on a digital nanophotometer (Implen, Westlake Village, CA, USA) and TapeStation (Agilent).

# *Micro-injected dsRNA*

Immediately before injection, dsRNA solution was mixed with blue dye (1:20) to aid in visualization of the injected liquid as in Perkin *et al.* (2017a). Larvae used in injections were briefly put on ice, transferred to a microscope slide with double-sided tape for stabilization, and the entire slide was placed on top of a small ice block. Thirty individuals were injected with 200 ng of dsRNA in triplicate. CPG-dsRNA was injected into larvae weighing approximately 1 mg. Each treatment injection had a corresponding control injection of dye only (Mock) and a non-injection control (Control). Injections were with a Drummond Nanoject (Drummond Scientific Co., Broomall, PA, USA) set at 69 nl and fast injecting speed with a “bee-stinger” needle. Needles were made with 3.5 Drummond glass capillary tubes (3-000-203-G) and a micropipette puller (Sutter Instrument Co. Model P-97, Novato, CA, USA) using the program: heat 750, pull 100, velocity 8, time 250, and pressure 500. After injection, each group was allowed to recover for 2 h at room temperature, and then were covered with diet (95% wheat flour, 5% brewer’s yeast) and kept at 28°C, 75% R.H., 0L:24D. All treatments and controls were followed for 18 days to monitor mortality and/or developmental abnormalities.

222 *Library preparation and RNA sequencing and data analysis after RNAi*

223 We validated knockdown of CPG with RNA-Seq as previously (Perkin et al., 2017b). Eight  
 224 individuals from each treatment and control were flash frozen in liquid nitrogen on day 18 post  
 225 injection. Total RNA was extracted by Trizol (Thermo Fisher Scientific, Waltham, MA USA)  
 226 and Quick-RNA Mini Prep kit (Zymo Research, Irvine, CA USA). Libraries were made on the  
 227 NeoPrep (200 bp insert) and sequenced on a Mi-Seq (2x300, Illumina, San Diego, CA USA).  
 228 Three independent biological replicates of all groups and 3-4 technical replicates of CPG  
 229 dsRNA-injected larvae were sequenced (metrics are found in Supplemental File 5). Sequence  
 230 reads were deposited at NCBI SRA as PRJNA520884.  
 231 Arraystar (DNASTar, Madison, WI USA) was used to calculate RPKM and fold change between  
 232 treatment and control groups by mapping reads to the Tcas5.2 genome. We also used Arraystar  
 233 for statistical analysis, student t-tests between treatments and ANOVA among treatments. Genes  
 234 were filtered to only include those that had >8-fold difference between treatments, and  
 235 significant at  $p < 0.01$  after FDR correction. To classify ncRNA, sequences were submitted to  
 236 RNAcon (website: <http://crdd.osdd.net/raghava/rnacon/submit.html>) for prediction using SVM  
 237 scores (default of 0.0; Panwar *et al.*, 2014).

238

239 *Cuticle protein tree*

240 NCBI RefseqRNA was searched with term “cuticle protein” and limited to *T. castaneum*, which  
 241 returned a list of 160 genes which we then manually curated to only include relevant genes with  
 242 a chitin domain (pfam00379). This resulted in a total of 125 sequences in the final data set.  
 243 Protein sequences from the selected genes were used as input in MEGA X (version 10.0.5;  
 244 Kumar et al., 2018). Protein sequences were aligned using ClustalW. The evolutionary history

of cuticle proteins was inferred using the Maximum Likelihood method, JTT matrix-based model, and 500 bootstrap iterations (Jones et al., 1992), using the tree with the highest log likelihood.

## Results

In the following sections, transcriptome datasets of gene expression in four developmental stages of *T. castaneum* were analyzed by sorting genes into categories of high and low expression levels. Gene Ontology (GO) terms were associated with each group, and overrepresented terms were identified by enrichment analysis. A cuticle protein gene was selected as a gene expressed at low levels only in larvae and was evaluated via RNAi as a gene with critical function for insect survival.

### *Life stage expression analysis – genes expressed at high levels*

Analysis of *T. castaneum* genes with high expression levels (RPKM >8, log<sub>2</sub>) in all life stages indicated there were 101, 290, 238, and 54 genes uniquely expressed in adults, eggs, larvae, and pupae, respectively (Fig. 1, Supplemental File 1). Eggs and larvae stages shared the most commonly expressed genes (162), followed by adults and larvae (48), and adults and eggs (29). Genes common to eggs and larvae included 44 ribosomal proteins (large and small subunits), indicative of the high degree of protein production occurring in these stages. Adults and larvae feeding stages shared two digestive cathepsin L genes (LOC659441 and LOC659502; Martynov et al., 2015; Perkin et al., 2016) and two chitinase precursor genes (*Chit8* and *Chit11*). Only LOC663906 encoding a single-stranded DNA-binding protein was commonly expressed in eggs and pupae. Twelve genes were highly expressed in all stages, including those encoding two

elongation factors, three ribosomal proteins, a cathepsin L (26-29-p), glyceraldehyde-3-phosphate dehydrogenase 2, NADH dehydrogenase subunit 2, ATP synthase, cytochrome c and cytochrome b, and I-rRNA (Table 1).

In the highly-expressed gene group, the top Molecular Function (MF) GO terms for each life stage varied, but adults and larvae had shared term “nucleotide binding”, whereas eggs and larvae shared “hydrolase activity” (Fig. 2). “Oxidoreductase activity” was common to larvae, pupae and adults. An enrichment analysis found 16 MF terms belonging to “structural constituent of the ribosome” ( $p=1.07 \times 10^{-11}$ ) and 62 with “catalytic activity” ( $p=0.028$ ) enriched only in adults.

We also examined the most highly expressed genes in each developmental stage. The top 10 most highly expressed genes in adults included those encoding an odorant binding protein, protamine, and cytochrome c oxidase subunit II (RPKM 7,694-20,300), and the highest expressed genes in eggs had similar function, including chemosensory protein 2, I-rRNA and cytochrome c oxidase subunit I and II (RPKM 7,960-26,611; Supplemental File 1). These genes also were highly expressed in larvae, along with cathepsin L genes involved in digestion of cereal proteins (RPKM 13,248-43,202; Martynov *et al.*, 2015; Perkin *et al.*, 2016). Highly expressed genes in pupae encoded two cuticular proteins and a conserved cathepsin L, 26-29-p (RPKM 3,576-13,818).

*Life stage expression analysis – genes expressed at low levels*

290 A comparison of genes with low expression levels (RPKM between 2-8 log<sub>2</sub>) revealed  
 291 expression of 273 genes unique to adults, 167 to eggs, 139 to larvae, and 14 to pupae (Fig. 3;  
 292 Supplemental File 2). Adults and larvae shared 138 genes including chitin deacetylase 7 (*Cda7*)  
 293 and two chitinase precursors (*Cht4* and *Cht9*). Larvae and pupae shared a single gene,  
 294 LOC664054 encoding pathogenesis-related protein 5, which also was expressed in adults but at  
 295 higher levels and thus was not part of the low expression group. Nine genes expressed at low  
 296 levels were common across all life stages, four of which had uncharacterized functions (Table 1).  
 297 The other five included genes encoding maternal protein tudor (*Tctud*), poly(A) polymerase type  
 298 3, THO complex subunit 2, transmembrane protein 35, and sentrin-specific protease.

299

300 A GO analysis of each group of unique genes indicated differences in top level functions among  
 301 stages, with more similar expression in the adults and larval stages, sharing MF terms “serine-  
 302 type endopeptidase activity”, “oxidoreductase activity”, and “transition metal ion binding” (Fig.  
 303 4). However, an enrichment analysis indicated that serine-type endopeptidases were  
 304 overrepresented only in the adults stage (p=0.005), and substrate-specific transporter activity was  
 305 overrepresented in the larval stage (p=0.04). Still, these two life stages shared 138 genes  
 306 expressed at low levels, not unexpected since both stages are active and feeding. Pupae had the  
 307 least number of genes and GO terms and included those with MF term “serine-type  
 308 endopeptidase activity”, and two unique terms: “NADH dehydrogenase activity”, and “structural  
 309 constituent of cuticle”. Enrichment analysis for pupae indicated overrepresentation of negative  
 310 regulation of programmed cell death (BP; p=0.002), which was due to reads mapping to the gene  
 311 LOC663274 that encodes fas apoptotic inhibitory molecule 1, consistent with cell division and  
 312 tissue rearrangement occurring during this stage. Genes expressed in eggs were associated with

terms “sequence-specific DNA binding” and “transcription factor activity sequence-specific DNA binding”. The most enriched categories in eggs were BP anatomical structure development (p=0.002), sex determination (p=0.011), and cell differentiation (p=0.006), consistent with highly complex developmental processes during embryogenesis.

### *Selection of candidate gene*

To find candidate genes for RNAi, we manually curated genes unique to each life stage and expressed at low levels (Supplemental File 2). Six cuticle genes were expressed only in larvae, but unfortunately many cuticle genes have orthologs in other species. However, a BLAST of CPG (LOC103313766) indicated no hits to other species (data not shown). In examining previous data (Morris *et al.*, 2009), CPG was found to be moderately expressed in the larval gut (relative to other gut-expressed genes). Unfortunately, this gene was not analyzed by the iBeetle project (Schmitt-Engel *et al.*, 2015). We then submitted the sequence to OffTargetFinder and found hits to only four other insect species which were focused to the 5’ and middle regions, leaving the 3’ as a unique portion of the sequence (Supplemental Fig. 1). So, we chose LOC103313766 CPG to evaluate as a potential candidate, since it met our criteria of uniquely expressed in *T. castaneum* larvae, contained regions where dsRNA would minimize off target effects, and was expressed in the larval gut.

### *RNAi validation of CPG*

To demonstrate that this was a biological target, we used RNAi to knockdown expression of CPG in *T. castaneum* larvae. We injected dsRNA into early stage larvae (2<sup>nd</sup>-3<sup>rd</sup> instar), which we hypothesized would have the most impact on transcript abundance. Knockdown and off target effects were evaluated by another round of RNA-Seq on RNA isolated from the injected



larvae and controls (mock-injected “Mock” and noninjected “Control”) 18 d post injection. The RNAi treatment significantly reduced the expression of CPG target transcript 1,491-fold compared to Control ( $p=2.17 \times 10^{-7}$ ) and 284-fold compared to Mock ( $p=7.14 \times 10^{-7}$ ). An ANOVA comparing the RPKM values of Mock, Control, and CPG treatments also was significantly different (Fig. 5A;  $p=2.64 \times 10^{-5}$ ). The large decrease in expression was biologically significant because 64% mortality was observed in the CPG dsRNA treatment group compared to the non-injected control at day 18 post-injection (Fig. 5B;  $p=0.057$ ). We also observed 29% mortality in Mock, but it was not significant when compared to Control mortality (19%;  $p=0.79$ ).

RNA-Seq analysis of larvae injected with CPG dsRNA compared to Mock resulted in 449 significantly differentially expressed genes ( $p < 0.01$  and at least 8-fold change; Supplemental Fig. 2 and Supplemental File 3). Of those genes, 100 were uncharacterized, and 20 were annotated as hypothetical proteins. Most (62%) had decreased expression, and the 10 most highly repressed genes were uncharacterized except the target gene and LOC100142553 adenylate kinase. Of those with increased expression, the most highly increased was *Jtb*, which encodes an orphan receptor with unknown function, LOC662244 F-box SPRY that contributes to ubiquitin-protein transferase activity, and *BBIP1* which is a chromosome associated protein. Other up regulated genes related to transcription/translation included LOC107398339 (splicing factor for mRNA), LOC103313738 (a repressor of developmentally regulated gene expression), and LOC664045, involved in chromatin silencing. In response to knockdown of CPG, three other cuticle protein genes were significantly increased: LOC100141875 cuticle 19-like, LOC103313752 cuticle protein 70, and LOC655183 cuticle protein.

There were 52 genes in this dataset that were annotated as noncoding RNA (ncRNA), 25 with increased and 27 with decreased expression (Supplemental File 4). All were preliminarily characterized as long ncRNAs since they were greater than 200 nt. All ncRNAs were screened through RNAcon (Panwar *et al.*, 2014) to predict the classification of each. There were 10 SSU-rRNA5, three Intron-GP-1, and one IRESe; the remaining sequences were below the default Support Vector Machines (SVM) threshold (0.0) and were classified as coding mRNA. However, these sequences were characterized by NCBI as ncRNA, probably due to unusual structure. Regardless, our transcriptome data suggests that ncRNA are actively transcribed in response to RNAi, as has been described previously (Ji *et al.*, 2015).

A GO term enrichment analysis of significantly decreased gene expression in response to CPG RNAi identified functions involved in chitin breakdown (BP GO:0006032 and MF GO:0004568;  $p=0.04$  and  $0.04$ , respectively). LOC107398196 and *Cht13*, both encoding chitinases, were responsible for these enriched GO terms. In contrast, the GO term aspartic-type endopeptidase activity ( $p=8.28 \times 10^{-5}$ ) was significantly enriched in up regulated genes.

### *Characterization of CPG*

To gain insight into the function of the CPG target gene, a maximum likelihood tree was constructed with *T. castaneum* protein sequences annotated as ‘cuticle protein’ (Fig. 6). CPG predicted protein (XP\_008196095, highlighted with a yellow box) was most similar to cuticle proteins 16.5 (XP\_015833062) and 19.8 (XP\_976285). Interestingly, the target CPG protein also was found in the same major clade as the up regulated cuticle protein 70 gene LOC103313752 (XP\_008196069, highlighted in red), and found in close proximity on LG3 (Table 2). Another

cuticle protein gene with increased expression, LOC655183 (XP\_008190752) also on LG3, was related to those encoding resilin proteins. The protein product of the cuticle gene with highest increased expression, LOC100141875 (XP\_001809825), was related to cuticle protein 7 (XP\_008193006) encoded by LOC656347, and both genes were found on LG5.

## Discussion

Current control methods for *T. castaneum* and other stored product pests are becoming less effective due to development of insecticide resistance (Collins, 1990; Jagadeesan *et al.*, 2012; Opit *et al.*, 2012). Alternative management strategies are limited, and new molecular-based controls are needed because they offer increased efficacy and selectivity, and targeted approaches to combatting resistance. There are major hurdles in developing molecular-based strategies in *T. castaneum*. The first step is to identify a target gene that can be effectively knocked-down. The target gene should be specific to the pest or a narrow range of pests. The most difficult challenge is delivery, whether it is topical or oral RNAi, or another mechanism. In this study, we evaluated molecular targets by injection of dsRNA prior to attempts of topical or oral delivery.

An RNA-Seq analysis of the four major life stages of *T. castaneum* identified twelve genes that were highly expressed in all developmental stages. One gene was a cathepsin L named 26-29-p. Cathepsin L genes in *T. castaneum* are part of a gene expansion group that are expressed at high levels in adults and larvae for the digestion of cereal proteins (Goptar *et al.*, 2012; Oppert *et al.*, 2010; Oppert *et al.*, 2003; Oppert *et al.*, 1993; Perkin *et al.*, 2016; Perkin *et al.*, 2017a; Vinokurov *et al.*, 2009). Based on phylogenetic analysis, 26-29-p is conserved across insects

and may be the ancestral cathepsin L gene, but it most likely functions in development and immunity (Perkin *et al.*, 2016). While this gene may not be useful for species-specific pest control because of the conservation of sequence among insects, its function in *T. castaneum* and other insects is of biological interest, as it appears to provide critical function(s) throughout development.

Nine genes were commonly expressed at low levels in all developmental stages, including *Tctud* (maternal protein tudor). *Tctud* has been found in many eukaryotes and has been implicated in protein-protein interactions, where methylated substrates bind to the tudor domain. In *D. melanogaster*, tudor proteins are found in chromatin, small nuclear RNA assembly, RNA-induced silencing complex, and germ granules (Ying & Chen, 2012). Similar to *Tctud*, temporal expression data shows low to moderately high expression of *Dmtud* in early fly embryos through 30-day old adults in both males and females (flybase.org, modENCODE temporal expression data, Gramates *et al.*, 2017). Additionally, mutants of various alleles caused lethality of early embryos through maternal effects, and both male and female sterility (Thomson & Lasko, 2004; flybase.org, Summary of Phenotypes, Gramates *et al.*, 2017). We submitted *Tctud* nucleotide sequence to OffTargetFinder and found multiple 21mer hits to 35 other species (data not shown), and therefore the gene was not selected for RNAi.

We have found that validation of RNAi knockdown through RNA-Seq can provide unique insights into function. Previously, we used RNA-Seq to validate knockdown of aspartate 1-decarboxylase (*ADC*) in *T. castaneum* larvae, resulting in the identification of additional gene interconnectivity (Perkin *et al.*, 2017b). Not only was the significant knockdown of *ADC*

confirmed through RNA-Seq, a previously unknown interaction between *ADC* and dopamine receptor 2 also was discovered. This information led to biological assays that determined that the reduction of *ADC* transcripts via RNAi resulted in adults with decreased mobility. Therefore, we used the same approach in the present study to evaluate knock down of a target gene and also understand the overall impact to the transcriptome of reduced gene expression.

To isolate a candidate gene that could be used as a molecular-based pest control target, we sought genes unique to a feeding stage (i.e. larvae or adults) and expressed at low levels. Through manual curation of life stage transcriptome data, we found CPG was uniquely expressed in larvae at low levels and lacked an ortholog in other species at the 3' end of the gene; in fact, BLAST did not return any hits at the mRNA level. We were encouraged that this target also has potential as an oral RNAi product, since the gene was moderately expressed in the larval gut. The protein product of this gene, XP\_008196095, is a peptide of 61 amino acids, with 41% identity to a neuropeptide-like precursor from the flesh fly, *Sacrophaga crassipalpis* (Li *et al.*, 2009). Additionally, Bhatia and Bhattacharya (2018) found that knockdown of a related cuticle protein gene in the green peach aphid (*Myzus persicae*) through oral delivery of dsRNA expressed in *Arabidopsis thaliana* resulted in reduced fecundity. We did not investigate whether the *T. castaneum* target cuticle protein also had neuropeptide properties or whether reduced gene expression caused a reduction in fecundity, but our functional test with RNAi demonstrated that decreased gene expression in early stage larvae resulted in significant mortality compared to Control and Mock. Thus, this gene is a candidate target for oral delivery of dsRNA.

RNA-Seq analysis of the transcriptome after RNAi knockdown of CPG identified three other cuticle genes with increased expression and two chitinase genes with decreased expression compared to Mock. The cuticle genes with increased expression included one encoding cuticle protein 70 (XP\_008196069), clustering in the same major clade as the target cuticle protein gene, suggesting similar function and perhaps an indication of redundancy in function as has previously been noted with gene expansion groups. In fact, this redundancy may explain survivors in the dsRNA-injected larvae, albeit a low percentage (32%), and it may complicate oral RNAi if there is compensation response similar to that previously identified in attempting to target cysteine protease genes via RNAi (Perkin et al., 2017a). Another cuticle protein gene LOC655183 also was up regulated after RNAi treatment, and this gene product (XP\_008190752) clustered with resilin proteins. Resilin is a specialized cuticle protein that is found in soft parts of the adult insect allowing for movement of wings (Andersen & Weis-Fogh, 1964). LOC655183 is an ortholog to a gene in *Drosophila melanogaster* encoding cuticular protein 56F, which is expressed in the imaginal discs of late larvae (flybase.org, Gramates *et al.*, 2017). The protein product of another cuticle gene with increased expression, LOC100141875 (XP\_001809825), clustered with cuticle protein 7 (XP\_008193006), an ortholog to one found in the carapace cuticle of juvenile horseshoe crab, *Limulus Polyphemus* with a high tyrosine content (Ditzel *et al.*, 2003).

Opposite in response to the increased expression of cuticle protein genes, *Chit13*, a chitinase, was decreased in response to the injection of CPG dsRNA. *Chit13* is predicted to be part of group IV chitinases expressed in the gut or fat body and expressed in response to feeding in larval and adult stages (Zhu *et al.*, 2008). All differentially expressed chitin-related genes are apparently

interconnected through a regulatory pathway that responds to the loss of function of CPG by RNAi.

We were surprised at the number of long ncRNAs (52) that were differentially expressed at a significant level ( $p < 0.01$ ) in surviving *T. castaneum* larvae that were injected with CPG dsRNA. Most ncRNA were not classified as typical ncRNAs and may represent new classes, or alternatively may be miss-annotated. However, differentially expressed long ncRNA have been implicated in human disease states, particularly cancer (Ma *et al.*, 2015). Of relevance to this study, differentially-expressed long ncRNAs can impose epigenetic changes that alter the transcription of other genes, including silencing (Tufarelli *et al.*, 2003). Therefore, we can speculate that the large number of differentially expressed genes in this study (449) may have been regulated in part by ncRNAs. While not to the same degree, ncRNAs were implicated in a previous RNAi study, in which LOC107398253 ncRNA (decreased 12.3-fold,  $p = 0.001$ , in the current study) was decreased 593-fold in response to RNAi of *ADC* (Perkin *et al.*, 2017b). This particular ncRNA was annotated as an U3 snoRNA, predicted to be involved in site-specific cleavage of ribosomal RNA (rRNA) during pre-rRNA processing (Clery *et al.*, 2007). More work is needed to understand the relationship of ncRNAs and RNAi in insects.

## Conclusions

In summary, these data add to the sparse stage-specific studies in *T. castaneum* and other beetles. We demonstrate that by using this approach, we were able to identify a candidate gene, CPG, that may be useful in developing an insect control product based on dsRNA. Because the gene

496 was expressed at low levels overall and moderate levels in the larval gut, oral delivery methods  
497 for CPG dsRNA have the potential to control damage by *T. castaneum* larvae.

498

# 499 **Acknowledgements**

500 The authors would like to thank technicians Ken Friesen and Tom Morgan for their contribution,  
501 injecting larvae and RNA extraction and library preparation, respectively. Mention of trade  
502 names or commercial products in this publication is solely for the purpose of providing specific  
503 information and does not imply recommendation or endorsement by the U.S. Department of  
504 Agriculture. USDA is an equal opportunity provider and employer.



# References

- Allen, M.L. and Walker, W.B., 3rd. 2012. Saliva of *Lygus lineolaris* digests double stranded ribonucleic acids. *Journal of Insect Physiology* 58: 391-396.
- Andersen, S. O. and Weis-Fogh, T. 1964. Reslin. A rubberlike protein in arthropod cuticle. *Advances in Insect Physiology* 2:1-65.
- Aronstein, K., Oppert, B. and Lorenzen, M. 2011. RNAi in agriculturally-important arthropods. In: *RNA Processing* (Grabowski, PP, ed.), pp. 157-180. InTech.
- Baum, J.A., Bogaert, T., Clinton, W., Heck, G.R., Feldmann, P., Ilagan, O., Johnson, S., Plaetinck, G., Munyikwa, T., Pleau, M., Vaughn, T. and Roberts, J. 2007. Control of coleopteran insect pests through RNA interference. *Nature Biotechnology* 25: 1322-1326.
- Benjamini, Y. and Hochberg, Y. 1995. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of Royal Statistics Society B Methodology* 57:289-300.
- Bhatia, V. and Bhattacharya, R. 2018. Host mediated RNAi of cuticular protein gene impaired fecundity in green peach aphid *Myzus persicae*. *Pest Management Science*. 10.1002/ps.4900.
- Boyer, S., Zhang, H. and Lemperiere, G. 2012. A review of control methods and resistance mechanisms in stored-product insects. *Bull Entomology Resources* 102:213-29.
- Brown, S.J., Mahaffey, J.P., Lorenzen, M.D., Denell, R.E., and Mahaffey, J.W. 1999. Using RNAi to investigate orthologous homeotic gene function during development of distantly related insects. *Evolution and Development* 1:11-15.
- Cao, M., Gatehouse, J.A., and Fitches, E.C. 2018. A systematic study of RNAi effects and dsRNA stability in *Tribolium castaneum* and *Acyrtosiphon pisum*, following injection

and ingestion of analogous dsRNAs. International Journal of Molecular Sciences. 19: 1079, doi:10.3390/ijms19041079.

Clery, A., Senty-Segault, V., Leclerc, F., Raue, H.A. and Branlant, C. 2007. Analysis of sequence and structural features that identify the B/C motif of U3 small nucleolar RNA as the recognition site for the Snr13p-Rps9p protein pair. Molecular and Cell Biology 27:1191-1206.

Collins, P.J. 1990. A new resistance to pyrethroids in *Tribolium castaneum* (herbst). Journal of Pesticide Science 28:101-115.

Ditzel, N., Andersen, S. O. and Højrup, P. 2003. Cuticular proteins from the horseshoe crab, *Limulus Polyphemus*. Comparative Biochemistry and Physiology Part B 134(3):489-497.

Donitz, J., Schmitt-Engel, C., Grossmann, D., Gerischer, L., Tech, M., Schoppmeier, M., Klingler, M. and Bucher, G. 2015. iBeetle-Base: a database for RNAi phenotypes in the red flour beetle *Tribolium castaneum*. Nucleic Acids Research 43: D720-725.

El Halim, H.M.A., Alshukri, B.M.H., Ahmad, M.S., Nakasu, E.Y.T., Awwad, M.H., Salama, E.M., Gatehouse, A.M.R., and Edwards, M.G. 2016. RNAi-mediated knockdown of the voltage gated sodium ion channel TcNa<sub>v</sub> causes mortality in *Tribolium castaneum*. Scientific Reports 6:29301, doi:10.1038/srep29301.

Good, R.T., Varghese, T., Golz, J.F., Russell, D.A., Papanicolaou, A., Edwards, O. and Robin, C. 2016. OfftargetFinder: a web tool for species-specific RNAi design. Bioinformatics 32:1232-1234.

Goptar, I.A., Semashko, T.A., Danilenko, S.A., Lysogorskaya, E.N., Oksenoit, E.S., Zhuzhikov, D.P., Belozersky, M.A., Dunaevsky, Y.E., Oppert, B., Filippova, I.Y. and Elpidina, E.N. 2012. Cysteine digestive peptidases function as post-glutamine cleaving enzymes in

tenebrionid stored-product pests. *Comp Biochem Physiol B Biochem Mol Biol* **161**: 148-154.

Götz, S., Garcia-Gomez, J.M., Terol, J., Williams, T.D., Nagaraj, S.H., Nueda, M.J., Robles, M., Talon, M., Dopazo, J. and Conesa, A. 2008. High-throughput functional annotation and data mining with the Blast2GO suite. *Nucleic Acids Research* 36:3420-3435.

Gramates, L.S., Marygold, S.J., Santos, G.D., Urbano, J.M., Antonazzo, G., Matthews, B.B., Rey, A.J., Tabone, C.J., Crosby, M.A., Emmert, D.B., Falls, K., Goodman, J.L., Hu, Y., Ponting, L., Schroeder, A.J., Strelets, V.B., Thurmond, J., Zhou, P. and the FlyBase, C. 2017. FlyBase at 25: looking to the future. *Nucleic Acids Research* 45:D663-D671.

Graveley, B.R., Brooks, A.N., Carlson, J.W., Duff, M.O., Landolin, J.M., Yang, L., Artieri, C.G., van Baren, M.J., Boley, N., Booth, B.W., Brown, J.B., Cherbas, L., Davis, C.A., Dobin, A., Li, R., Malone, J.H., Mattiuzzo, N.R., Miller, D., Sturgill, D., Tuch, B.B., Zaleski, C., Zhang, D., Blanchette, M., Dudoit, S., Eads, B., Green, R.E., Hammonds, A., Jiang, L., Kapranov, P., Langton, L., Perrimon, N., Sandler, J.E., Wan, K.H., Willingham, A., Zhang, Y., Zou, Y., Andrews, J., Bickel, P.J., Brenner, S.E., Brent, M.R., Cherbas, P., Gingeras, T.R., Hoskins, R.A., Kaufman, T.C., Oliver, B. and Celniker, S.E. 2011. The development transcriptome of *Drosophila melanogaster*. *Nature* 471:473-479.

Guan, R.B., Li, H.C., Fan, Y.J., Hu, S.R., Christiaens, O., Smagghe, G., and Miao, X.X 2018. A nuclease specific to lepidopteran insects suppresses RNAi. *Journal of Biological Chemistry* jbc.RA117.001553.

Huvenne, H. and Smagghe, G. 2010. Mechanisms of dsRNA uptake in insects and potential of RNAi for pest control: a review. *Journal of Insect Physiology* 56:227-235.

573 Jagadeesan, R., Collins, P.J., Daglish, G.J., Ebert, P.R., Schlipalius, D.I. 2012. Phosphine  
574 resistance in the rust red flour beetle, *Tribolium castaneum* (Coleoptera: Tenebrionidae):  
575 inheritance, gene interactions and fitness costs. PloS one 7:e31582.

576 Ji, Z., Song, R., Regev, A. and Struhl, K. 2015. Many lncRNAs, 5'UTRs, and pseudogenes are  
577 translated and some are likely to express functional proteins. eLife 4:e08890.

578 Jones D.T., Taylor W.R., and Thornton J.M. 1992. The rapid generation of mutation data  
579 matrices from protein sequences. Computer Applications in the Biosciences 8: 275-282.

580 Knorr, E., Fishilevich, E., Tenbusch, L., Frey, M.L.F., Rangasamy, M., Billion, A., Worden,  
581 S.E., Gandra, P., Arora, K., Lo, W., Schulenberg, G., Valverde-Garcia, P., Vilcinskas, A.  
582 and Narva, K.E. 2018. Gene silencing in *Tribolium castaneum* as a tool for the targeted  
583 identification of candidate RNAi targets in crop pests. Scientific Reports 8:2061,  
584 doi:10.1038/s41589-018-20416-y.

585 Kumar S., Stecher G., Li M., Knyaz C., and Tamura K. 2018. MEGA X: Molecular Evolutionary  
586 Genetics Analysis across computing platforms. Molecular Biology and Evolution  
587 35:1547-1549.

588 Li, A., Rinehart, J.P. and Denlinger, D.L. 2009. Neuropeptide-like precursor 4 is uniquely  
589 expressed during pupal diapause in the flesh fly. Peptides 30:518-521.

590 Lomate, P.R. and Bonning, B.C. 2016. Distinct properties of proteases and nucleases in the gut,  
591 salivary gland and saliva of southern green stink bug, *Nezara viridula*. Scientific Report  
592 6:27587.

593 Loktev, A.V., Zhang, Q., Beck, J.S., Searby, C.C., Scheetz, T.E., Bazan, J.F., Slusarski, D.C.,  
594 Sheffield, V.C., Jackson, P.K. and Nachury, M.V. 2008. A BBSome subunit links  
595 ciliogenesis, microtubule stability, and acetylation. Developmental Cell 15:854-865.

596 Ma, L., Li, A., Zou, D., Xu, X., Xia, L., Yu, J., Bajic V.B. and Zhang, Z. 2015. LncRNAWiki:  
597 harnessing community knowledge in collaborative curation of human long non-coding  
598 RNAs *Nucleic Acids Research* 43:187-192.

599 Martynov, A.G., Elpidina, E.N., Perkin, L. and Oppert, B. 2015. Functional analysis of C1  
600 family cysteine peptidases in the larval gut of *Tenebrio molitor* and *Tribolium castaneum*.  
601 *BMC Genomics* 16:75.

602 Morris, K., Lorenzen, M.D., Hiromasa, Y., Tomich, J.M., Oppert, C., Elpidina, E.N., Vinokurov,  
603 K., Jurat-Fuentes, J.L., Fabrick, J. and Oppert, B. 2009. *Tribolium castaneum* Larval Gut  
604 Transcriptome and Proteome: A Resource for the Study of the Coleopteran Gut. *Journal*  
605 *Proteome Research* 8:3889-3898.

606 Miller, S.C., Miyata, K., Brown, S.J. and Tomoyasu, Y. 2012. Dissecting systemic RNA  
607 interference in the red flour beetle *Tribolium castaneum*: parameters affecting the  
608 efficiency of RNAi. *PLoS One* 7:e47431.

609 Mortazavi, A., Williams, B.A., McCue, K., Schaeffer, L. and Wold, B. 2008. Mapping and  
610 quantifying mammalian transcriptomes by RNA-Seq. *Nature Methods* 5:621-628.

611 Noh, M.Y., Beeman, R.W. and Arakane, Y. 2012. RNAi-based functional genomics in *Tribolium*  
612 *castaneum* and possible application for controlling insect pests. *Entomology Research*  
613 42:1-10.

614 Noh, M.Y., Kramer, K.J., Muthukrishnan, S., Beeman, R.W., Kanost, M.R. and Arakane, Y.  
615 2015. Loss of function of the yellow-e gene causes dehydration-induced mortality of  
616 adult *Tribolium castaneum*. *Developmental Biology* 399:315-324.

- Opit, G.P., Phillips, T.W., Aikins, M.J. and Hasan, M.M. 2012. Phosphine resistance in *Tribolium castaneum* and *Rhyzopertha dominica* from stored wheat in Oklahoma. Journal of Economic Entomology 105:1107-1114.
- Oppert, B., Elpidina, E.N., Toutges, M. and Mazumdar-Leighton, S. 2010. Microarray analysis reveals strategies of *Tribolium castaneum* larvae to compensate for cysteine and serine protease inhibitors. Comparative Biochemistry and Physiology Part D Genomics Proteomics 5:280-287.
- Oppert, B., Morgan, T., Hartzer, K., Lenarcic, B., Galesa, K., Brzin, J., Turk, V., Yoza, K., Ohtsubo, K. and Kramer, K. 2003. Effects of proteinase inhibitors on digestive proteinases and growth of the red flour beetle, *Tribolium castaneum* (Herbst)(Coleoptera: Tenebrionidae). Comparative Biochemistry and Physiology Part C Toxicology and Pharmacology 134:481-490.
- Oppert, B., Morgan, T.D., Culbertson, C. and Kramer, K.J. 1993. Dietary mixtures of cysteine and serine proteinase inhibitors exhibit synergistic toxicity toward the red flour beetle, *Tribolium castaneum*. Comparative Biochemistry and Physiology Part C: Comparative Pharmacology 105:379-385.
- Palli, S. R. 2014. RNA interference in Colorado potato beetle: steps toward development of dsRNA as a commercial insecticide. Current Opinion Insect Science 6:1-8.
- Panwar, B., Arora, A. and Raghava, G.P.S. 2014. Prediction and classification of ncRNAs using structural information. BMC Genomics 15:127.
- Perkin, L., Elpidina, E.N. and Oppert, B. 2016. Expression patterns of cysteine peptidase genes across the *Tribolium castaneum* life cycle provide clues to biological function. PeerJ 4:e1581.

- Perkin, L.C., Elpidina, E.N. and Oppert, B. 2017a. RNA interference and dietary inhibitors induce a similar compensation response in *Tribolium castaneum* larvae. *Insect Molecular Biology* 26:35-45.
- Perkin, L.C., Gerken, A.R., Oppert, B. 2017b. RNA-Seq validation of RNAi identifies additional gene connectivity in *Tribolium castaneum* (Coleoptera: Tenebrionidae). *Journal of Insect Science* 17:1-7.
- Pimentel, M.A., D'A Faroni, L.R., Da Silva, F.H., Batista, M.D. and Guedes, R.N.C. 2010. Spread of phosphine resistance among Brazilian populations of three species of stored product insects. *Neotropical Entomology* 39:101-107.
- Reimand, J., Arak, T. and Vilo, J. 2011. g:Profiler--a web server for functional interpretation of gene lists (2011 update). *Nucleic Acids Research* 39:W307-315.
- Schmitt-Engel, C., Schultheis, D., Schwirz, J., Strohle, N., Troelenberg, N., Majumdar, U., Dao, V.A., Grossmann, D., Richter, T., Tech, M., Donitz, J., Gerischer, L., Theis, M., Schild, I., Trauner, J., Koniszewski, N.D., Kuster, E., Kittelmann, S., Hu, Y., Lehmann, S., Siemanowski, J., Ulrich, J., Panfilio, K.A., Schroder, R., Morgenstern, B., Stanke, M., Buchholz, F., Frasch, M., Roth, S., Wimmer, E.A., Schoppmeier, M., Klingler, M. and Bucher, G. 2015. The iBeetle large-scale RNAi screen reveals gene functions for insect development and physiology. *Nature Communications* 6:7822.
- Sparks, M.E., Rhoades, J.H., Nelson, D.R., Kuhar, D., Lancaster, J., Lehner, B., Tholl, D., Weber, D.C. and Gundersen-Rindal, D.E. 2017. A Transcriptome Survey Spanning Life Stages and Sexes of the Harlequin Bug, *Murgantia histrionica*. *Insects* 8.

661 Shen, G., Dou, W., Niu, J., Jiang, H., Yang, W., Jia, F., Hu, F., Cong, L. and Wang, J. 2011.  
662 Transcriptome analysis of the Oriental fruit fly (*Bactrocera dorsalis*). PLoS one  
663 6:e29127.

664 Thomson, T. and Lasko, P. 2004. Drosophila tudor is essential for polar granule assembly and  
665 pole cell specification, but not for posterior patterning. Genesis 40:164-170.

666 Tomoyasu, Y. and Denell, R.E. 2004. Larval RNAi in Tribolium (Coleoptera) for analyzing  
667 adult development. Developmental Genes and Evolution 214:575-578.

668 *Tribolium* Genome Sequencing Consortium. 2008. The genome of the model beetle and pest  
669 *Tribolium castaneum*. Nature 452:949-955.

670 Tufarelli, C., Stanley, J.A., Garrick D, Sharpe, J.A., Ayyub, H., Wood, W.G. and Higgs, D.R.  
671 2003. Transcription of antisense RNA leading to gene silencing and methylation as a  
672 novel cause of human genetic disease. Nature Genetics 34:157-165.

673 Ulrich, J., Dao, V.A., Majumdar, U., Schmitt-Engel, C., Schwirz, J., Schultheis, D., Strohle, N.,  
674 Troelenberg, N., Grossmann, D., Richter, T., Donitz, J., Gerischer, L., Leboulle, G.,  
675 Vilcinskas, A., Stanke, M. and Bucher, G. 2015. Large scale RNAi screen in Tribolium  
676 reveals novel target genes for pest control and the proteasome as prime target. BMC  
677 Genomics 16:674.

678 Vinokurov, K.S., Elpidina, E.N., Zhuzhikov, D.P., Oppert, B., Kodrik, D. and Sehnal, F. 2009.  
679 Digestive proteolysis organization in two closely related Tenebrionid beetles: red flour  
680 beetle (*Tribolium castaneum*) and confused flour beetle (*Tribolium confusum*). Archives  
681 in Insect Biochemistry 70:254-279.

682 Whyard, S., Singh, A.D., and Wong, S. 2009. Ingested double-stranded RNAs can act as species-  
683 specific insecticides. Insect Biochemistry and Molecular Biology 39:824-832.



- 684 Ying, M. and Chen, D. (2012) Tudor domain-containing proteins of *Drosophila melanogaster*.  
685 Developmental Growth & Differentiation 54:32-43.
- 686 Zhu, Q., Arakane, Y., Banerjee, D., Beeman, R.W., Kramer, K.J., and Muthukrishnan, S. 2008.  
687 Domain organization and phylogenetic analysis of the chitinase-like family of proteins in  
688 three species of insects. Insect Biochemistry and Molecular Biology 38: 452-466

**Table 1**(on next page)

Genes that were expressed at high and low levels in all life stages.

List of genes that were expressed at high (top) and low (bottom) levels in all life stages of *T. castaneum*.

# 1 Tables

2 Table 1. Genes that were expressed at high (top) and low (bottom) levels in all life stages of *T. castaneum*.

Name	Description	Average RPKM			
		Adult	Egg	Larvae	Pupae
LOC656235	Elongation factor 1-gamma	1,560	1,334	657	6,278
RpS3	Ribosomal protein S3	3,953	2,663	1,475	263
26-29-p	Cathepsin L	369	281	261	5,710
LOC659992	40S ribosomal protein S29	1,135	1,443	626	3,238
LOC663023	Glyceraldehyde-3-phosphate dehydrogenase 2	2,405	812	2,135	2,375
LOC660435	60S ribosomal protein L11	2,673	2,134	1,096	6,999
Efa1	Elongation factor 1-alpha	2,253	2,541	1,762	3,576
ND2	NADH dehydrogenase subunit 2	1,081	1,457	1,000	401
ATP6	ATP synthase F0 subunit 6	5,442	4,960	4,555	270
COX3	Cytochrome c oxidase subunit III	4,267	3,923	6,559	857
CYTB	Cytochrome b	3,181	4,829	7,123	558
I-rRNA	I-rRNA	16,635	11,824	34,216	11,020
LOC103315067	Uncharacterized	5.91	57.1	7.30	18.7
LOC103312214	Uncharacterized	10.3	89.0	8.11	24.6
LOC103315070	Uncharacterized	17.5	94.1	16.9	26.9
LOC107399196	Uncharacterized	25.5	181	9.69	22.7
LOC103312419	Maternal protein tudor	5.25	15.3	5.34	5.45
LOC103312455	poly(A) polymerase type 3	9.11	65.5	13.0	15.0
LOC663288	THO complex subunit 2	7.55	61.8	8.69	10.6
LOC659780	Transmembrane protein 35	240	7.72	241	179
LOC103313244	Sentrin-specific protease	10.5	74.6	35.0	9.45

3

4

5

## Table 2 (on next page)

Differentially expressed cuticle genes.

Differentially expressed cuticle genes in *T. castaneum* larvae injected with CPG dsRNA.

Shaded row indicates target gene.

Table 2. Differentially expressed cuticle genes in *T. castaneum* larvae injected with LOC103313766 cuticle protein dsRNA. Shaded row indicates target gene.

Gene	LG	Coordinate Start	Coordinate Stop	Protein	Annotation	Fold Change	Expression
LOC103313766	3	4099231	4115167	XP_008196095	Cuticle protein	1,491	↓
LOC658110	3	4097672	4112513	XP_015833062	Cuticle protein 16.5	2.51	↓
LOC658489	3	4077823	4109423	XP_976285	Cuticle protein 19.8	1.44	↓
LOC103313752	3	4114536	4126952	XP_008196069	Cuticle protein 70	10.8	↑
LOC655183	3	2446367	2620651	XP_008190752	Cuticle protein	4.01	↑
LOC100141875	5	3608978	3616499	XP_001809825	Cuticle protein	16.9	↑
LOC656347	5	3612511	3613210	XP_008193006	Cuticle protein 7	4.89	↑
LOC107399215	2	6796409	6962942	XP_015840495	Larval cuticle protein A2B-like	25.1	↓

# Figure 1

Venn diagram comparing genes expressed in *T. castaneum* life stages: adult, egg, larvae, and pupae with high expression values ( $>8 \log_2$  RPKM).


Shades of grey indicate the relative number of genes in each comparison. Dark grey indicates a higher number of genes and lighter shades represent comparisons with fewer genes.



# Figure 2

Top GO terms for genes with high expression in each life stage.


Each pie chart identifies the top GO terms for each life stage; (A) Egg, (B) Larvae, (C) Pupae, (D) Adult.



# Figure 3

Venn diagram comparing genes between all *T. castaneum* life stages: adult, egg, larvae, and pupae with low expression values ( $2-8 \log_2$  RPKM).

Shades of grey indicate the relative number of genes in each comparison. Dark grey indicates a higher number of genes and lighter shades represent comparisons with fewer genes.






# Figure 4

Top GO terms for genes with low expression in each *T. castaneum* life stage.


Each pie chart identifies the top GO terms for each life stage; (A) Egg, (B) Larvae, (C) Pupae, (D) Adult.



# Figure 5

Effects of injection of CPG dsRNA in *T. castaneum* larvae.

(A) Average expression of CPG in control, mock, and dsRNA-injected *T. castaneum* larvae 18 d following RNAi application. (B) Percent survival of *T. castaneum* larvae following RNAi treatment.



# Figure 6

*T. castaneum* cuticle protein maximum likelihood tree.

CPG (XP\_008196095) is highlighted with a yellow box, and proteins from the three cuticle genes with increased expression (XP\_008196069, XP\_001809825, XP\_008190752) are highlighted with a red box.

