

# Future CO<sub>2</sub>-induced seawater acidification mediates the physiological performance of a green alga *Ulva linza* in different photoperiods

Furong Yue<sup>1</sup>, Guang Gao<sup>1</sup>, Jing Ma<sup>1</sup>, Hailong Wu<sup>1</sup>, Xinshu Li<sup>1</sup>, Juntian Xu<sup>Corresp. 1, 2, 3</sup>

<sup>1</sup> Jiangsu Key Laboratory of Marine Bioresources and Environment, Huaihai Institute of Technology, Lianyungang, China

<sup>2</sup> Co-Innovation Center of Jiangsu Marine Bio-industry Technology, Huaihai Institute of Technology, Lianyungang, China

<sup>3</sup> Jiangsu Key Laboratory of Marine Biotechnology, Huaihai Institute of Technology, Lianyungang, China

Corresponding Author: Juntian Xu

Email address: jtxu@hhit.edu.cn

Photoperiods have an important impact on macroalgae living in the intertidal zone. Ocean acidification also influences the physiology of macroalgae. However, little is known about the interaction between ocean acidification and photoperiod on macroalgae. In this study, a green alga *Ulva linza* was cultured under three different photoperiods (L: D = 8:16, 12:12, 16:8) and two different CO<sub>2</sub> levels (LC, 400 ppm; HC, 1000 ppm) to investigate their responses. The results showed that relative growth rate of *U. linza* increased with extended light periods under LC but decreased at HC when exposed to the longest light period of 16h compared to 12h. Higher CO<sub>2</sub> levels enhanced the relative growth rate at a L: D of 8:16, had no effect at 12:12 but reduced RGR at 16:8. At LC, the L: D of 16:8 significantly stimulated maximum quantum yield (Yield). Higher CO<sub>2</sub> levels enhanced Yield at L: D of 12:12 and 8:16, had negative effect at 16:8. Non-photochemical quenching (NPQ) increased with increasing light period. High CO<sub>2</sub> levels did not affect respiration rate during shorter light periods but enhanced it at a light period of 16h. Longer light periods had negative effects on Chl *a* and Chl *b* content, and high CO<sub>2</sub> level also inhibited the synthesis of these pigments. Our data demonstrate the interactive effects of CO<sub>2</sub> and photoperiod on the physiological characteristics of the green tide macroalga *Ulva linza* and indicate that future ocean acidification may hinder the stimulatory effect of long light periods on growth of *Ulva* species.

# Future CO<sub>2</sub>-induced seawater acidification mediates the physiological performance of a green alga *Ulva linza* in different photoperiods

Furong Yue<sup>1</sup>, Guang Gao<sup>1</sup>, Jing Ma<sup>1</sup>, Hailong Wu<sup>1</sup>, Xinshu Li<sup>1</sup>, Juntian Xu<sup>1,2,3\*</sup>

<sup>1</sup>Jiangsu Key Laboratory of Marine Bioresources and Environment, Huaihai Institute of Technology, Lianyungang, 222005, China

<sup>2</sup>Co-Innovation Center of Jiangsu Marine Bio-industry Technology, Huaihai Institute of Technology, Lianyungang, 222005, China

<sup>3</sup>Jiangsu Key Laboratory of Marine Biotechnology, Huaihai Institute of Technology, Lianyungang, 222005, China

\*Corresponding author:

Juntian Xu<sup>1</sup>

Cangwu Road 59, Lianyungang, Jiangsu, 222005, China

[jtxu@hhit.edu.cn](mailto:jtxu@hhit.edu.cn), Phone/Fax: +86 (0)518 8558 5003

## Abstract

Photoperiods have an important impact on macroalgae living in the intertidal zone. Ocean acidification also influences the physiology of macroalgae. However, little is known about the interaction between ocean acidification and photoperiod on macroalgae. In this study, a green alga *Ulva linza* was cultured under three different photoperiods (L: D = 8:16, 12:12, 16:8) and two different CO<sub>2</sub> levels (LC, 400 ppm; HC, 1000 ppm) to investigate their responses. The results showed that relative growth rate of *U. linza* increased with extended light periods under LC but decreased at HC when exposed to the longest light period of 16h compared to 12h. Higher CO<sub>2</sub> levels enhanced the relative growth rate at a L: D of 8:16, had no effect at 12:12 but reduced RGR at 16:8. At LC, the L: D of 16:8 significantly stimulated maximum quantum yield (Yield). Higher CO<sub>2</sub> levels enhanced Yield at L: D of 12:12 and 8:16, had negative effect at 16:8. Non-photochemical quenching (NPQ) increased with increasing light period. High CO<sub>2</sub> levels did not affect respiration rate during shorter light periods but enhanced it at a light period of 16h. Longer light periods had negative effects on Chl *a* and Chl *b* content, and high CO<sub>2</sub> level also inhibited the synthesis of these pigments. Our data demonstrate the interactive effects of CO<sub>2</sub> and

photoperiod on the physiological characteristics of the green tide macroalga *Ulva linza* and indicate that future ocean acidification may hinder the stimulatory effect of long light periods on growth of *Ulva* species.

## Introduction

Due to the activities of humans, the concentration of atmospheric CO<sub>2</sub> has increased to 400 ppm from 278 ppm during the pre-industrial revolution (Gattuso et al., 2015). It is estimated that the oceans, as a CO<sub>2</sub> sink, have taken up approximately 48% of the fossil-fuel and cement-manufacturing emissions (Sabine et al., 2004). The concentration of atmospheric CO<sub>2</sub> has been predicted to reach almost 1000 ppm by the end of 21<sup>st</sup> century (Gattuso et al., 2015). This would result in the decline of surface ocean pH by 0.3-0.4 units, an increase of hydrogen ion concentration and bicarbonate concentration in seawater, which is termed ocean acidification (OA) (Orr et al., 2005; Koch et al., 2013 Albright et al., 2016). OA will be maintained for hundreds of years, and thus could cause a huge change in marine ecosystems (Kroeker et al., 2013; Wittmann & Pörtner, 2013).

Marine phytoplankton accounts for 50% of global primary production (Beardall et al., 2009) and macroalgae accounts for 10% of marine primary productivity (Smith, 1981; Charpy-Roubaud & Sournia, 1990; Heinrich et al., 2012). Macroalgae have evolved distinct strategies to capture inorganic carbon (Ci) for photosynthesis: (1) by the diffusive uptake of CO<sub>2</sub> (Giordano & Raven, 2005; Raven et al., 2005), (2) the active transport of HCO<sub>3</sub><sup>-</sup>/CO<sub>2</sub> (termed carbon concentrating mechanisms (CCMs) (Maberly et al., 1992; Raven, 2003; Giordano et al., 2005). The elevated CO<sub>2</sub> might enhance the growth and photosynthesis of algae that depend on CO<sub>2</sub> diffusion or have weak CCMs (Kübler et al., 1999; Gao et al., 2016a). Therefore, the physiological influences of OA on macroalgae might be species-specific. For example, OA has a positive influences on the growth of macroalgae, such as the red algae *Pyropia yezoensis*, *Pyropia haitanensis* and *Gracilaria lemaneiformis* (Gao et al., 1991; Xu & Gao, 2015; Chen et al., 2018), green algae *Ulva prolifera*, *Ulva lactuca*, *Caulerpa taxifolia* (Xu & Gao, 2012; Chen et al., 2015; Roth-Schulze et al., 2018) and brown algae *Hizikia fusiforme*, *Ecklonia radiata*, *Sargassum muticum* (Zou, 2005; Britton et al., 2016; Xu et al., 2017). However, research has shown that high CO<sub>2</sub> levels did not have a detectable effect on the relative growth rate of several species, for example, *Sargassum henslowianum*, *Ulva rigida*, *Ulva australis*, the giant kelp *Macrocystis pyrifera* and *Ulva australis* (Chen & Zou, 2014; Rautenberger et al., 2015; Fernández et al., 2015; Reidenbach et al., 2017). Furthermore, increasing seawater CO<sub>2</sub> concentration reduced the growth of *Gracilaria tenuistipitata*, *Porphyra leucosticte*, *Fucus vesiculosus*, *Halimeda opuntia* and *Ulva linza* (García-Sánchez et al., 1994; Mercado et al., 1999; Gutow et al., 2014; Johnson et al., 2014; Gao et al., 2018). In addition, the various effects of

ocean acidification may be also modulated by other environmental factors, such as temperature, nutrients and photoperiods (Gao et al., 2017a; Gao et al., 2019; Li et al., 2018).

Photoperiod is a key factor regulating seasonal responses of algae (Dring, 1998). Furthermore, daylength can influence algal affinity for dissolved inorganic carbon and cellular carbon demand, thus imposing an effect on CCMs regulation (Rost et al., 2006). Thus, the photoperiod may be critical for algae cultured under elevated CO<sub>2</sub> levels. Meanwhile, the interactive effects of OA and increased photoperiods could enhance the growth of *U. prolifera*, thus increase the opportunities for occurrence of green tides (Li et al., 2018). Elevated CO<sub>2</sub> levels enhanced the photosynthetic rate, but longer illumination periods reduced the photosynthetic efficiency in *E. huxleyi* (Nielsen, 1997; Rost et al., 2006). Therefore, at HC, the responses of algae to different photoperiods appear to be genera- and species-specific.

Green macroalgal bloom has occurred each year in the Yellow Sea (YS) since 2007, termed green tides (Xu et al., 2016). Green tides generate aesthetic problems and toxic gas when thalli decompose (Gao et al., 2016b). *Ulva* is the dominant genus contributing majority of green tides and *U. linza* is one of species causing green tides in the Yellow Sea (Fletcher, 1996; Kang et al., 2014; Gao et al., 2017a, b). However, little is known about the interactive effects of ocean acidification and photoperiod on *U. linza*. Previous study showed that the effect of ocean acidification on diatoms was related to light intensity (Gao et al., 2012). In this study, based on previous studies, we hypothesized the effect of ocean acidification on *U. linza* may dependent on photoperiod. To test our hypothesis, the physiological responses of the green macroalga *Ulva linza* under two different CO<sub>2</sub> levels and three different photoperiods were examined.

## Materials & Methods

### Thalli collection and culture conditions

*U. linza* was collected from the coastal water of Gaogong peninsula (119.3°E, 34.5°N), Lianyungang, Jiangsu Province, China. Gaogong peninsula is a public place and no approval is required for collecting naturally growing *Ulva* species in China because *Ulva* species cause green tides in coastal waters of the Yellow Sea. *U. linza* was identified by morphological characters (Ma et al., 2009). The samples were transferred to the laboratory in a portable cooler (4-6°C) box within one hour. Healthy thalli (first observed by color and then checked with maximum quantum yield of PSII) were selected and cleaned with filtered natural seawater to remove sediments, visible epiphytes and attached animals. Thalli were pre-cultured in a 500 ml flask in an illuminated incubator (GXZ-500B, Ningbo, China) at 20 °C, with the illumination intensity set at 150 μmol photons m<sup>-2</sup> s<sup>-1</sup> (12L:12D). Sterilized filtered seawater (salinity 30,

supplied with 8  $\mu\text{M}$   $\text{NaH}_2\text{PO}_4$  and 60  $\mu\text{M}$   $\text{NaNO}_3$ ) was used as culture medium and the medium was bubbled with air before being renewed every two days.

For these experiments, approximately three 1 cm long thalli were placed into a 550 ml flask filled with 500 ml sterile seawater aerated ambient outdoor air (400 ppm, LC) or with  $\text{CO}_2$ -enriched air (1000 ppm, HC) and randomly cultured in three separate incubators with different light levels (8L:16D, 12L:12D, 16L:8D) and continuous aeration at 20 °C with 150  $\mu\text{mol photons m}^{-2}\text{s}^{-1}$ . Different  $\text{CO}_2$  levels (400 ppm, LC; 1000 ppm, HC) were achieved by bubbling ambient air or  $\text{CO}_2$ -enriched air via a  $\text{CO}_2$  plant incubator (HP 1000 G-D, Ruihua Instruments, Wuhan, China).

To maintain the  $\text{pH}_{\text{NBS}}$  at about 8.12 (LC) and 7.78 (HC) under different photoperiods, the increased algal biomass were removed constantly and the medium was changed every two days and daily variations in pH were maintained at less than 0.05. The experiment lasted 9 days, the physiological factors were taken on the last 3 days.

## Estimate of carbonate system parameters

The seawater pH was monitored with a pH meter (pH 700, Eutech Instruments, Singapore) and total alkalinity (TA) was calculated by titrations (Gao et al., 2019). Other parameters of the carbonate system were obtained with CO2SYS software (Pierrot et al., 2006), the equilibrium constants  $K_1$  and  $K_2$  for carbonic acid dissociation (Roy et al., 1993).

## Measurement of growth

The length of *U. linza* was recorded every 2 days. The relative growth rate (RGR) of thalli were calculated across the 6-day period before other physiological parameters were measured during the following 3 days. RGR was calculated as follows:  $\text{RGR} = \ln(W_t / W_0) / t$ , where  $W_t$  is the length after  $t$  days culture,  $W_0$  is the initial length.

## Chlorophyll fluorescence measurements

Chlorophyll *a* fluorescence in *U. linza* was measured with a portable PAM (Pulse-amplitude-modulation; AquaPen-P AP-P 100, Chech). Algae were dark adapted for 15 min before the experiment. The parameters were calculated according to the following equations:

$$\text{rETR} = \text{PAR} \times Y(\text{II}) \times 0.84 \times 0.5 \text{ (Schreiber, 2004; Zhang et al., 2017),}$$

where rETR is the relative electron transport rate; PAR is the actinic light;  $Y(\text{II})$  is the maximum quantum yield of PSII.

$$\text{NPQ} = F_m - F_m' / F_m \text{ (Bilger \& Schreiber, 1986),}$$

where NPQ is the non-photochemical quenching;  $F_m$  is the maximum fluorescence value of *U.*

135 *linza* when they were adapted in the dark for 15 min;  $F_m'$  is the maximum fluorescence value of  
136 *U. linza* under actinic light conditions.

## 137 **Respiration rate measurements**

138 The respiration rate of *U. linza* was measured using a Clark-type oxygen electrode (YSI Model  
139 5300, Yellow Springs Instrument Co., USA). The thallus was cut into 1 cm long segments with  
140 scissors and the thalli were placed in culture conditions for at least 1h to decrease the effects of  
141 cutting damage. Approximately 0.01g fresh weight of thalli were placed in the reaction chamber  
142 with 8 ml medium. Temperature was controlled at 20 °C with a circulating water bath. The  
143 decrease of the oxygen content in the seawater in darkness with seven minutes was defined as the  
144 respiration rate.

## 145 **Measurement of photosynthetic pigments**

146 Chlorophyll *a* and *b* were extracted from thalli (about 10 mg FW) with 5 ml methanol at 4 °C for  
147 24h in the dark. The absorption values were obtained at 652 nm, 663 nm and 665 nm using an  
148 ultraviolet spectrophotometer (Ultrospect 3300 pro, Amersham Bioscience, Sweden). The  
149 contents of the Chl *a* and Chl *b* were estimated using the method of Porra (1989).

## 150 **Data analysis**

151 All the data are shown as mean  $\pm$  SD. Origin 9.0 and SPSS 18.0 were used to analyze data. Two-  
152 way ANOVA was used to assess the interactive effects of CO<sub>2</sub> levels and photoperiods on  
153 relative growth rate, chlorophyll fluorescence parameters, respiration rate and pigment content of  
154 *U. linza*. One-way ANOVA was used to analyze differences under the same conditions. Tukey  
155 HSD was conducted for *post hoc* investigation. Confidence intervals were set at 95%.

## 156 **Results**

157 Both the elevated CO<sub>2</sub> levels and photoperiod altered carbonate parameters in seawater, and they  
158 both had an interactive effect (Tables 1 and 2). The elevated CO<sub>2</sub> decreased pH and CO<sub>3</sub><sup>2-</sup>,  
159 increased *p*CO<sub>2</sub>, DIC, HCO<sub>3</sub><sup>-</sup> and CO<sub>2</sub> in the seawater. Increased photoperiod did not affect  
160 carbonate parameters at LC but the longest photoperiod increased DIC, HCO<sub>3</sub><sup>-</sup> and TA compared  
161 to shortest photoperiod.

162 The two-way ANOVA showed that elevated CO<sub>2</sub> and photoperiod had an interactive effect,  
163 and both elevated CO<sub>2</sub> levels and the photoperiods had a significant effect on the RGR of *U.*  
164 *linza* (Fig. 1 and Table 3). At LC, the RGR of adult *U. linza* increased with the extended light  
165 periods, and the highest RGR occurred at a L: D of 16:8. The effect of CO<sub>2</sub> also varied with  
166 photoperiod. Higher CO<sub>2</sub> levels enhanced RGR at L: D of 8:16, but had no effect at 12:12 and  
167 reduced it at 16:8.

The Yield and NPQ were measured under different CO<sub>2</sub> levels and photoperiod conditions (Fig. 2). Two-way ANOVA showed that elevated CO<sub>2</sub> and photoperiod had an interactive effect on Yield (Table 4). Higher CO<sub>2</sub> levels increased Yield when thalli were cultured under photoperiods of 8:16 and 12:12 but reduced it under 16:8. Photoperiod had the main effect on NPQ (Table 4). At LC, thalli cultured at L: D of 16:8 had higher NPQ compared to L: D of 8:16 while the difference between 8:16 and 12:12 was insignificant. At HC, NPQ increased with the increase in photoperiod although the increase was not statistically significant. The elevated CO<sub>2</sub> had neutral effect on NPQ of *U. linza*.

Maximum rETR (rETRmax), efficiency of electron transport ( $\alpha$ ), and saturating irradiance ( $I_k$ ) were calculated from the rapid light curves (Fig. 3, Table 5). Photoperiod and elevated CO<sub>2</sub> levels had an interactive effect, and elevated CO<sub>2</sub> levels had a main effect on light-saturated electron transport rate (rETRmax) (Table 6). Higher CO<sub>2</sub> levels increased rETRmax during the 8:16 and 12:12 photoperiods, but did not affect it at 16:8. A similar pattern was also found for  $\alpha$ . In contrast to rETRmax and  $\alpha$ , CO<sub>2</sub> did not affect  $I_k$  while photoperiod had the main effect on it. At LC,  $I_k$  increased when L: D rose from 8:16 to 12:12 but did not change with the further increase in photoperiod. At HC,  $I_k$  did not change when L: D rose from 8:16 to 12:12 but was enhanced when L: D increased to 16:8 (Table 5).

In addition to photosynthetic parameters, the effects of pCO<sub>2</sub> and photoperiod on the respiration rate of adult *U. linza* were also investigated (Fig. 4). Photoperiod and elevated CO<sub>2</sub> levels had an interactive effect, and both photoperiod and elevated CO<sub>2</sub> levels had the primary effect on the respiration rate of *U. linza* (Table 7). Higher CO<sub>2</sub> levels did not affect the respiration rate at photoperiods of 8:16 or 12:12 but increased it by 56.39% at a photoperiod of 16:8.

Changes in photosynthetic pigments of *U. linza* grown under various conditions are shown in Fig. 5. Two-way ANOVA showed that CO<sub>2</sub> and photoperiod had an interactive effect, and both CO<sub>2</sub> levels and the photoperiods had the main effect on the Chl *a* content of *U. linza* (Table 8). Prolonged light periods reduced the synthesis of Chl *a* in thalli although the difference between the photoperiods of 8:16 and 12:12 at HC was not statistically significant. Higher CO<sub>2</sub> levels reduced Chl *a* at the photoperiods of 8:16 and 12:12 but did not affect it at a L: D of 12:12. The same trend was found for Chl *b*. The Chl *a/b* ratios were all greater than 1.0, suggesting a higher synthesis of Chl *a* than Chl *b* under all culture conditions. Photoperiod and elevated CO<sub>2</sub> levels had an interactive effect on the Chl *a/b* ratio (Table 8); the higher CO<sub>2</sub> levels increased the ratio at a photoperiod of 12:12 but not at the other photoperiods.

## Discussion

In the present study, at LC, extended photoperiods had a positive effect on the relative growth rate of adult *U. linza*, similar to previous studies on *Laminaria sacharina*, *Porphyra umbilicalis*

and *Ulva prolifera* (Fortes & Lüning, 1980; Green & Neefus, 2016; Li et al., 2018). Carbon isotope fractionation experiments suggested that extended photoperiods could enhance growth by influencing inorganic carbon capture and fixation rate in algae (Rost et al., 2003). This hypothesis is supported by the present study where extended photoperiods increased the maximum quantum yield in PS II. On the other hand, the highest growth rates of *Compsopogon coeruleus* were obtained in shorter light periods (L: D = 8:16) (Zucchi & Necchi, 2001), the highest growth rates of *Porphyra umbilicalis* was found under neutral (L: D = 12:12) and longer (L: D = 16:8) light periods using 110  $\mu\text{mol photons m}^{-2}\text{s}^{-1}$  (Green & Neefus, 2016), and the relative growth rate of *Chlorella vulgaris* cultured at L: D = 20:4 was lower than the RGR of alga cultured at L: D = 16:8 (Kendirlioglu et al., 2015). Therefore, the effects of photoperiod on algae appear to be species-specific.

Although *Ulva* has efficient mechanisms for  $\text{CO}_2$  concentration, the growth of *Ulva* can be enhanced by elevated  $\text{CO}_2$  (Young et al., 2016; Gao et al., 2016a, 2017). However, in this study, we found that the effects of elevated  $\text{CO}_2$  levels on the relative growth rate of *U. linza* depended on photoperiod.  $\text{CO}_2$  enhanced the growth of adult *U. linza* under light/dark conditions of 8:16, had no effect on growth of *U. linza* under a L: D of 12:12, and reduced the relative growth rate under a L: D of 16:8. High  $\text{CO}_2$  levels can down-regulate algal  $\text{CO}_2$  concentration mechanisms (CCMs), meaning that energy would be saved, and thus enhancing the relative growth rate of algae (Gao et al., 2012, 2016a; Raven et al., 2017). This is supported by decreased pigment synthesis in thalli at higher  $\text{CO}_2$  levels. However, higher  $\text{CO}_2$  levels did not affect the growth rate of *U. linza* at medium photoperiods. The neutral effects of  $\text{CO}_2$  on the growth of *U. rigida* (Rautenberger et al., 2015) and *U. linza* (Gao et al., 2018) were also reported. We speculate that this is a compromise between the positive effects of elevated  $\text{CO}_2$  and negative effects of decreased pH. The negative effect of decreased pH on growth was documented for the brown alga *Sargassum muticum* (Xu et al., 2017). Algae might need to consume additional energy to act against acid-based perturbation caused by decreased pH, leading to reduced growth (Xu et al., 2017). This is supported by an enhanced respiration rate at the higher  $\text{CO}_2$  levels in this study. The phenomenon of an increased respiration rate of algae under elevated  $\text{CO}_2$  concentrations was found in *Hizikia fusiformis* (Zou et al., 2005), the microalgae *Phaeodactylum tricornutum* (Wu et al., 2010) and *Emiliania huxleyi* (Jin et al., 2015).

Furthermore, the higher  $\text{CO}_2$  level reduced the growth rate of *U. linza* under the longest photoperiod in the present study. This may be due to the combination of down-regulated CCMs and excess light energy. The operation of CCMs is an energy-consuming process and the down-regulation of CCMs can result in additional energy (Raven et al., 2017). Higher light can usually reduce algal photosynthetic activity (Singh and Singh, 2015; Gao et al., 2016a). The energy saved due to down-regulation of CCMs at HC combined with high light intensity could

synergistically damage the algal photosystem and photosynthetic rate (Gao et al., 2012, 2016a). Although light intensity did not change among different CO<sub>2</sub> treatments, the lengthened photoperiod may have similar effect to increased light intensity. This argument is supported by increased NPQ at longer photoperiods as NPQ is photo-protective process to dissipate excess light energy. Higher CO<sub>2</sub> levels also reduced maximum quantum yield and stimulated the respiration rate of *U. linza* during the longest photoperiod in this study, leading to the decrease in growth. Enhanced respiration rate is a signal that organisms are fighting against stress and damage (Xu et al., 2017).

In addition to growth, the interactive effect of CO<sub>2</sub> and photoperiod was also found in photosynthetic parameters. For instance, elevated CO<sub>2</sub> increased Yield, rETRmax and  $\alpha$  at shorter photoperiods (L: D of 8:16 and 12:12) but did not affect rETRmax and reduced Yield at longest photoperiod (L: D of 16:8). These findings indicate the close connection between growth and photosynthesis in terms of responding to the combination of CO<sub>2</sub> and photoperiod. It is worth noting that the interaction of CO<sub>2</sub> and photoperiod on growth of *U. linza* in this study is different from the findings in *U. prolifera* reported by Li et al (2018). Elevated CO<sub>2</sub> increased growth of *U. prolifera* at all three photoperiods (L: D of 12:12, 10:14 and 16:8). The different results may be due to differential physiological property between *U. linza* and *U. prolifera*. It has been documented that *U. prolifera* has a higher tolerance to high light intensity compared to *U. linza* (Cui et al., 2015). The strong capacity in dealing with high light intensity could contribute to the effect of elevated CO<sub>2</sub> and prolonged photoperiod on growth *U. linza* was still positive. Integrating our findings with Li et al's (2018) study, the interaction of CO<sub>2</sub> and photoperiod on *Ulva* species would be species dependent.

## Conclusions

This work is the first attempt to clarify the interaction between light/dark and elevated CO<sub>2</sub> levels on the physiological responses of *Ulva linza*. We found that the effect of OA on *U. linza* depended on photoperiod. Outbreaks of green tides during spring and summer in China occur when the photoperiod is reaching its peak. Our findings indicate future OA may hinder the occurrence of green tides dominated by *U. linza* in combination with extended photoperiods. More environmental factors, such as temperature and nutrient levels, need to be investigated to obtain a more comprehensive understanding on development of green tides in future oceans.

## Acknowledgements

# References

- Albright R. 2016. Reversal of ocean acidification enhances net coral reef calcification. *Nature* 531:362-365 DOI 10.1038/nature17155.
- Beardall J, Stojkovic S, Larsen S. 2009. Living in a high CO<sub>2</sub> world: impacts of global climate change on marine phytoplankton. *Plant Ecology & Diversity* 2:191-205 DOI 10.1080/17550870903271363.
- Bilger W, Schreiber U. 1986. Energy-dependent quenching of dark-level chlorophyll fluorescence in intact leaves. *Photosynthesis Research* 10:303-308 DOI 10.1007/bf00118295.
- Britton D, Cornwall CE, Revill AT, Hurd CL, Johnson CR. 2016. Ocean acidification reverses the positive effects of seawater pH fluctuations on growth and photosynthesis of the habitat-forming kelp, *Ecklonia radiata*. *Scientific Reports* 6:26036 DOI 10.1038/srep26036.
- Charpy-Roubaud C, Sournia A. 1990. The comparative estimation of phytoplanktonic, microphytobenthic and macrophytobenthic primary production in the oceans. *Marine Microbial Food Webs* 4:31-57
- Chen B, Zou D. 2014. Growth and photosynthetic activity of *Sargassum henslowianum* (Fucales, Phaeophyta) seedlings in responses to different light intensities, temperatures and CO<sub>2</sub> levels under laboratory conditions. *Marine Biology Research* 10:1019-1026 DOI 10.1080/17451000.2013.872798.
- Chen B, Zou D, Jiang H. 2015. Elevated CO<sub>2</sub> exacerbates competition for growth and photosynthesis between *Gracilaria lemaneiformis* and *Ulva lactuca*. *Aquaculture* 443:49-55 DOI 10.1016/j.aquaculture.2015.03.009.
- Chen B, Zou D, Du H, Ji Z. 2018. Carbon and nitrogen accumulation in the economic seaweed *Gracilaria lemaneiformis* affected by ocean acidification and increasing temperature. *Aquaculture* 482:176-182 DOI 10.1016/j.aquaculture.2017.09.042.
- Cui J, Zhang J, Huo Y, Zhou L, Wu Q, Chen L, Yu K, He P, 2015. Adaptability of free-floating green tide algae in the Yellow Sea to variable temperature and light intensity. *Marine Pollution Bulletin* 101:660-666 DOI 10.1016/j.marpolbul.2015.10.033.
- Dring MJ. 1988. Photocontrol of development in algae. *Annual Review of Plant Physiology and Plant Molecular Biology* 39:157-174 DOI 10.1146/annurev.pp.39.060188.001105.

- 308 Fernández PA, Roleda MY, Hurd CL. 2015. Effects of ocean acidification on the photosynthetic  
309 performance, carbonic anhydrase activity and growth of the giant kelp *Macrocystis pyrifera*.  
310 *Photosynthesis Research* 124:293-304  
311 DOI 10.1007/s11120-015-0138-5.
- 312 Fletcher RL. 1996. The occurrence of “green tides”—a review. In *Marine benthic vegetation* (pp.  
313 7-43) Springer, Berlin, Heidelberg.  
314 DOI 10.1007/978-3-642-61398-2\_2.
- 315 Fortes MD, Lüning K. 1980. Growth rates of North Sea macroalgae in relation to temperature,  
316 irradiance and photoperiod. *Helgoländer Meeresuntersuchungen* 34:15-29 DOI  
317 10.1007/BF01983538.
- 318 Gao G, Clare AS, Rose C, Caldwell GS. 2017a. Eutrophication and warming-driven green tides  
319 (*Ulva rigida*) are predicted to increase under future climate change scenarios. *Marine*  
320 *Pollution Bulletin* 114:439-447 DOI 10.1016/j.marpolbul.
- 321 Gao G, Beardall J, Bao M, Wang C, Ren W, Xu J. 2018. Ocean acidification and nutrient  
322 limitation synergistically reduce growth and photosynthetic performances of a green tide  
323 alga *Ulva linza*. *Biogeosciences* 15:3409-3420  
324 DOI 10.5194/bg-15-3409-2018.
- 325 Gao G, Gao Q, Bao M, Xu J, Li X. 2019. Nitrogen availability modulates the effects of ocean  
326 acidification on biomass yield and food quality of a marine crop *Pyropia yezoensis*. *Food*  
327 *Chemistry* 271:623-629  
328 DOI 10.1016/j.foodchem.2018.07.090.
- 329 Gao G, Liu Y, Li X, Feng Z, Xu J. 2016a. An ocean acidification acclimatized green tide alga is  
330 robust to changes of seawater carbon chemistry but vulnerable to light stress. *PloS one*  
331 11:e0169040 DOI 10.1371/journal.pone.0169040.
- 332 Gao G, Liu Y, Li X, Feng Z, Xu Z, Wu H, Xu J. 2017b. Expected CO<sub>2</sub>-induced ocean  
333 acidification modulates copper toxicity in the green tide alga *Ulva prolifera*. *Environmental*  
334 *and Experimental Botany* 135:63-72  
335 DOI 10.1016/j.envexpbot.2016.12.007.
- 336 Gao G, Zhong Z, Zhou X, Xu J. 2016b. Changes in morphological plasticity of *Ulva prolifera*  
337 under different environmental conditions: a laboratory experiment. *Harmful Algae* 59:51-58  
338 DOI 10.1016/j.hal.2016.09.004.
- 339 Gao K, Aruga Y, Asada K, Ishihara T, Akano T, Kiyohara M. 1991. Enhanced growth of the red  
340 alga *Porphyra yezoensis* Ueda in high CO<sub>2</sub> concentrations. *Journal of Applied Phycology*  
341 3:355-362 DOI 10.1007/bf02392889.
- 342 Gao K, Xu J, Gao G, Li Y, Hutchins DA, Huang B. 2012. Rising CO<sub>2</sub> and increased light  
343 exposure synergistically reduce marine primary productivity. *Nature Climate Change*

- 2:519-523 DOI 10.1038/nclimate1507.
- Gao Z, Xu D, Meng C, Zhang X, Wang Y, Li D. 2014. The green tide-forming macroalga *Ulva linza* outcompetes the red macroalga *Gracilaria lemaneiformis* via allelopathy and fast nutrients uptake. *Aquatic Ecology* 48:53-62 DOI 10.1007/s10452-013-9465-9.
- García-Sánchez MJ, Fernández JA, Niell X. 1994. Effect of inorganic carbon supply on the photosynthetic physiology of *Gracilaria tenuistipitata*. *Planta* 194:55-61 DOI 10.1007/bf00201034.
- Gattuso JP, Magnan A, Billé R, Cheung WW, Howes EL, Joos F. 2015. Contrasting futures for ocean and society from different anthropogenic CO<sub>2</sub> emissions scenarios. *Science* 349:aac4722 DOI 10.1126/science.aac4722.
- Giordano M, Beardall J, Raven JA. 2005. CO<sub>2</sub> concentrating mechanisms in algae: mechanisms, environmental modulation, and evolution. *Annual Review Plant Biology* 56:99-131 DOI 10.1146/annurev.arplant.56.032604.144052.
- Green LA, Neefus CD. 2016. Effects of temperature, light level, and photoperiod on the physiology of *Porphyra umbilicalis* Kützinger from the Northwest Atlantic, a candidate for aquaculture. *Journal of Applied Phycology* 28:1815-1826 DOI 10.1007/s10811-015-0702-6.
- Gutow L, Rahman MM, Bartl K, Saborowski R, Bartsch I, Wiencke C. 2014. Ocean acidification affects growth but not nutritional quality of the seaweed *Fucus vesiculosus* (Phaeophyceae, Fucales). *Journal of Experimental Marine Biology and Ecology* 453:84-90 DOI 10.1016/j.jembe.2014.01.005.
- Heinrich S, Valentin K, Frickenhaus S, John U, Wiencke C. 2012. Transcriptomic analysis of acclimation to temperature and light stress in *Saccharina latissima* (Phaeophyceae). *PLoS one* 7: e44342 DOI 10.1371/journal.pone.0044342.
- Jin P, Wang T, Liu N, Dupont S, Beardall J, Boyd PW. 2015. Ocean acidification increases the accumulation of toxic phenolic compounds across trophic levels. *Nature Communications* 6:8714 DOI 10.1038/ncomms9714.
- Johnson MD, Price NN, Smith JE. 2014. Contrasting effects of ocean acidification on tropical fleshy and calcareous algae. *PeerJ* 2:e411 DOI 10.7717/peerj.411.
- Kang EJ, Kim JH, Kim K, Choi HG, Kim KY. 2014. Re-evaluation of green tide-forming species in the Yellow Sea. *Algae* 29:49-59 DOI 10.4490/algae.2016.31.3.9.
- Kendirlioglu G, Agirman N, Cetin A. 2015. The effects of photoperiod on the growth, protein amount and pigment content of *Chlorella vulgaris*. *Turkish Journal of Science & Technology* 10:7-10.
- Kim JH, Kang EJ, Park MG, Lee BG, Kim KY. 2011. Effects of temperature and irradiance on

- 380 photosynthesis and growth of a green-tide-forming species (*Ulva linza*) in the Yellow Sea.
- 381 *Journal of Applied Phycology* 23:421-432
- 382 DOI 10.1007/s10811-010-9590-y.
- 383 Koch M, Bowes G, Ross C, Zhang XH. 2013. Climate change and ocean acidification effects on
- 384 seagrasses and marine macroalgae. *Global Change Biology* 19:103-132 DOI
- 385 10.1111/j.1365-2486.2012.02791.x.
- 386 Kroeker KJ, Kordas RL, Crim R, Hendriks IE, Ramajo L, Singh GS. 2013. Impacts of ocean
- 387 acidification on marine organisms: quantifying sensitivities and interaction with warming.
- 388 *Global Change Biology* 19:1884-1896 DOI 10.1111/gcb.12179.
- 389 Kübler JE, Johnston AM, Raven JA. 1999. The effects of reduced and elevated CO<sub>2</sub> and O<sub>2</sub> on
- 390 the seaweed *Lomentaria articulata*. *Plant, Cell & Environment* 22: 1303-1310 DOI
- 391 10.1046/j.1365-3040.1999.00492.x.
- 392 Li Y, Zhong J, Zheng M, Zhuo P, Xu N. 2018. Photoperiod mediates the effects of elevated CO<sub>2</sub>
- 393 on the growth and physiological performance in the green tide alga *Ulva prolifera*. *Marine*
- 394 *Environmental Research* 141:24-29
- 395 DOI 10.1016/j.marenvres.2018.07.015.
- 396 Ma J, J J, Xu R, He P, Zhang T, Wang X, Li Y, Ren S, Xu P, Lu Q. 2009. Preliminary study on
- 397 life history of *Ulva linza* Linnaeus [*Enteromorpha linza* (L.) J. Ag.]. *Journal of Fisheries of*
- 398 *China* 1 33:45-52Maberly SC, Raven JA, Johnston AM. 1992. Discrimination between <sup>12</sup>C
- 399 and <sup>13</sup>C by marine plants. *Oecologia* 91:481-492 DOI 10.1007/bf00650320.
- 400 Mercado JM, Javier F, Gordillo L, Niell FX, Figueroa FL. 1999. Effects of different levels of
- 401 CO<sub>2</sub> on photosynthesis and cell components of the red alga *Porphyra leucosticta*. *Journal of*
- 402 *Applied Phycology* 11:455-461
- 403 DOI 10.1023/A: 1008194223558.
- 404 Nielsen MV. 1997. Growth, dark respiration and photosynthetic parameters of the
- 405 coccolithophorid *Emiliania Huxleyi* (Prymnesiophyceae) acclimated to different day length-
- 406 irradiance combinations1. *Journal of Phycology* 33:818-822
- 407 DOI 10.1111/j.0022-3646.1997.00818.x.
- 408 Orr JC, Fabry VJ, Aumont O, Bopp L, Doney SC, Feely RA. 2005. Anthropogenic ocean
- 409 acidification over the twenty-first century and its impact on calcifying organisms. *Nature*
- 410 437:681-686 DOI 10.1038/nature04095.
- 411 Pierrot D, Lewis E, Wallace DWR. 2006. MS Excel program developed for CO<sub>2</sub> system
- 412 calculations. *ORNL/CDIAC-105a. Carbon Dioxide Information Analysis Center, Oak Ridge*
- 413 *National Laboratory, US Department of Energy, Oak Ridge, Tennessee.*
- 414 Porra RJ, Thompson WA, Kriedemann PE. 1989. Determination of accurate extinction
- 415 coefficients and simultaneous equations for assaying chlorophylls *a* and *b* extracted with

- 416 four different solvents: verification of the concentration of chlorophyll standards by atomic  
417 absorption spectroscopy. *Biochimica et Biophysica Acta (BBA)-Bioenergetics* 975:384-394  
418 DOI 10.1016/S0005-2728(89)80347-0.
- 419 Rautenberger R, Fernandez PA, Strittmatter M, Heesch S, Cornwall CE, Hurd CL, Roleda MY.  
420 2015. Saturating light and not increased carbon dioxide under ocean acidification drives  
421 photosynthesis and growth in *Ulva rigida* (Chlorophyta). *Ecology and Evolution* 5:874-888  
422 DOI 10.1002/ece3.1382.
- 423 Raven JA. 2003. Inorganic carbon concentrating mechanisms in relation to the biology of algae.  
424 *Photosynthesis Research* 77:155-171  
425 DOI 10.1023/A:1025877902752.
- 426 Raven JA, Ball LA, Beardall J, Giordano M, Maberly SC. 2005. Algae lacking carbon-  
427 concentrating mechanisms. *Canadian Journal of Botany* 83: 879-890 DOI 10.1139/b05-074.
- 428 Raven JA, Beardall J, Sánchez-Baracaldo P. 2017. The possible evolution and future of CO<sub>2</sub>-  
429 concentrating mechanisms. *Journal of Experimental Botany* 68:3701-3716 DOI  
430 10.1093/jxb/erx110.
- 431 Reidenbach LB, Fernandez PA, Leal PP, Noisette F, McGraw CM, Revill AT, 2017. Growth,  
432 ammonium metabolism, and photosynthetic properties of *Ulva australis* (Chlorophyta)  
433 under decreasing pH and ammonium enrichment. *PloS one* 12:e0188389 DOI  
434 10.1371/journal.pone.0188389.
- 435 Rost B, Riebesell U, Burkhardt S, Sültemeyer D. 2003. Carbon acquisition of bloom-forming  
436 marine phytoplankton. *Limnology and Oceanography* 48:55-67  
437 DOI 10.4319/lo.2003.48.1.0055.
- 438 Rost B, Riebesell U, Sültemeyer D. 2006. Carbon acquisition of marine phytoplankton: effect of  
439 photoperiod length. *Limnology and Oceanography* 51:12-20 DOI  
440 10.4319/lo.2006.51.1.0012.
- 441 Roth-Schulze AJ, Thomas T, Steinberg P, Deveney MR, Tanner JE, Wiltshire KH. 2018. The  
442 effects of warming and ocean acidification on growth, photosynthesis, and bacterial  
443 communities for the marine invasive macroalga *Caulerpa taxifolia*. *Limnology and*  
444 *Oceanography* 63:459-471 DOI 10.1002/lno.10739.
- 445 Roy RN, Roy LN, Vogel KM, Porter-Moore C, Pearson T, Good CE, Millero FJ, Campbell DM.  
446 1993. The dissociation constants of carbonic acid in seawater at salinities 5 to 45 and  
447 temperatures 0 to 45 C. *Marine Chemistry* 6: 249-267 DOI 10.1016/0304-4203(93)90207-5.
- 448 Sabine CL, Feely RA, Gruber N, Key RM, Lee K, Bullister JL. 2004. The oceanic sink for  
449 anthropogenic CO<sub>2</sub>. *Science* 305:367-371  
450 DOI 10.1126/science.1097403.

- Schreiber U. 2004. Pulse-amplitude-modulation (PAM) fluorometry and saturation pulse method: an overview. *Chlorophyll a Fluorescence* 19:279-319 DOI 10.1007/978-1-4020-3218-9\_11.
- Singh SP, Singh P. 2015. Effect of temperature and light on the growth of algae species: a review. *Renewable and Sustainable Energy Reviews* 50:431-444 DOI 10.1016/j.rser.2015.05.024.
- Smith, SV. 1981. Marine macrophytes as a global carbon sink. *Science* 211:838-840 DOI 10.1126/science.211.4484.838.
- Wittmann AC, Pörtner HO. 2013. Sensitivities of extant animal taxa to ocean acidification. *Nature Climate Change* 3:995-1001 DOI 10.1038/nclimate1982.
- Wu Y, Gao K, Riebesell U. 2010. CO<sub>2</sub>-induced seawater acidification affects physiological performance of the marine diatom *Phaeodactylum tricornutum*. *Biogeosciences* 7:2915-2923 DOI 10.5194/bg-7-2915-2010.
- Xu J, Gao K. 2012. Future CO<sub>2</sub>-induced ocean acidification mediates the physiological performance of a green tide alga. *Plant Physiology* 160:1762-1769 DOI 10.1104/pp.112.206961.
- Xu J, Gao K. 2015. Photosynthetic performance of the red alga *Pyropia haitanensis* during emersion, with special reference to effects of solar UV radiation, dehydration and elevated CO<sub>2</sub> concentration. *Photochemistry and Photobiology* 91:1376-1381 DOI 10.1111/php.12531.
- Xu Q, Zhang H, Cheng Y. 2016. Multi-sensor monitoring of *Ulva prolifera* blooms in the Yellow Sea using different methods. *Frontiers of Earth Science* 10:378-388 DOI 10.1007/s11707-015-0528-1.
- Xu Z, Gao G, Xu J, Wu H. 2017. Physiological response of a golden tide alga (*Sargassum muticum*) to the interaction of ocean acidification and phosphorus enrichment. *Biogeosciences* 4:671-681 DOI 10.5194/bg-14-671-2017.
- Young CS, Gobler CJ. 2016. Ocean acidification accelerates the growth of two bloom-forming macroalgae. *PloS one* 11:e0155152 DOI 10.1371/journal.pone.0155152.
- Zhang D, Zhang Q, Yang X. 2017. Seasonal dynamics of photosynthetic activity in the representative brown macroalgae *Sargassum thunbergii* (Sargassaceae Phaeophyta). *Plant Physiol Biochem* 120:88-94 DOI 10.1016/j.plaphy.2017.09.025.
- Zou D. 2005. Effects of elevated atmospheric CO<sub>2</sub> on growth, photosynthesis and nitrogen metabolism in the economic brown seaweed, *Hizikia fusiforme* (Sargassaceae, Phaeophyta). *Aquaculture* 250:726-735

487 DOI 10.1016/j.aquaculture.2005.05.014.  
 488 Zou D, Gao K, Ruan Z. 2007. Daily timing of emersion and elevated atmospheric CO<sub>2</sub>  
 489 concentration affect photosynthetic performance of the intertidal macroalga *Ulva lactuca*  
 490 (Chlorophyta) in sunlight. *Botanica Marina* 50:275-279 DOI 10.1515/BOT.2007.031.  
 491 Zucchi MR, Necchi O. 2001. Effects of temperature, irradiance and photoperiod on growth and  
 492 pigment content in some freshwater red algae in culture. *Phycological Research* 49:103-114  
 493 DOI 10.1046/j.1440-1835.2001.00230.x.

# Figure legend

Fig. 1. Relative growth rate (RGR) of *U. linza* thalli grown at different CO<sub>2</sub> levels (400 ppm, LC; 1000 ppm, HC) and different photoperiods (L: D = 8:16, 12:12, 16:8). Vertical bars are mean ± SD for triplicate samples. Different lowercase letters denote significant difference among the photoperiod levels at LC and different capital letters denote significant difference among the photoperiod levels at HC. Significant differences between two CO<sub>2</sub> levels under the same photoperiod are indicated by short lines.

Fig. 2. Yield (A) and NPQ (B) of *U. linza* thalli grown at different CO<sub>2</sub> levels (400 ppm, LC; 1000 ppm, HC) and different photoperiods (L: D = 8:16, 12:12, 16:8). Vertical bars are mean ± SD for triplicate samples. Different lowercase letters denote significant difference among the photoperiod levels at LC and different capital letters denote significant difference among the photoperiod levels at HC. Significant differences between two CO<sub>2</sub> levels under the same photoperiod are indicated by short lines.

Fig. 3. Rapid light curves (RLC) of *U. linza* thalli grown at ambient CO<sub>2</sub> levels (400 ppm, LC, A; 1000 ppm, HC, B) and different photoperiods (L: D = 8:16, 12:12, 16:8).


Fig. 4. Respiration rate of *U. linza* thalli grown at different CO<sub>2</sub> levels (1000 ppm, HC) and different photoperiods (L: D = 8:16, 12:12, 16:8). Vertical bars are mean ± SD for triplicate samples. Different lowercase letters denote significant difference among the photoperiod levels at LC and different capital letters denote significant difference among the photoperiod levels at HC. Significant differences between two CO<sub>2</sub> levels under the same photoperiod are indicated by short lines.

Fig. 5. Content of the pigments Chl *a* (A) and Chl *b* (B), and the Chl *a/b* ratio (C) of *U. linza* thalli grown at different CO<sub>2</sub> levels (400 ppm, LC; 1000 ppm, HC) and different photoperiods (L: D = 8:16, 12:12, 16:8). Vertical bars are mean ± SD for triplicate samples. Different lowercase letters denote significant difference among the photoperiod levels at LC and different capital letters denote significant difference among the photoperiod levels at HC. Significant differences between two CO<sub>2</sub> levels under the same photoperiod are indicated by short lines.

# Figure 1

Relative growth rate.


Relative growth rate (RGR) of *U. linza* thalli grown at different CO<sub>2</sub> levels (400 ppm, LC; 1000 ppm, HC) and different photoperiods (L: D = 8:16, 12:12, 16:8).



# Figure 2

Yield and NPQ.


Yield (A) and NPQ (B) of *U. linza* thalli grown at different CO<sub>2</sub> levels (400 ppm, LC; 1000 ppm, HC) and different photoperiods (L: D = 8:16, 12:12, 16:8).



# Figure 3

Rapid light curves.


Rapid light curves (RLC) of *U. linza* thalli grown at different CO<sub>2</sub> levels (400 ppm, LC; 1000 ppm, HC) and different photoperiods (L: D = 8:16, 12:12, 16:8).



# Figure 4

Respiration rate.


Respiration rate of *U. linza* thalli grown at different CO<sub>2</sub> levels (400 ppm, LC; 1000 ppm, HC) and different photoperiods (L: D = 8:16, 12:12, 16:8).



# Figure 5

Pigment contents and the Chl *a/b* ratio.

Pigment contents (Chl *a*, A; Chl *b*, B) and the Chl *a/b* ratio (C) of *U. linza* thalli grown at different CO<sub>2</sub> levels (400 ppm, LC; 1000 ppm, HC) and different photoperiods (L: D = 8:16, 12:12, 16:8).



**Table 1**(on next page)

Parameters of the seawater carbonate system in different cultures. DIC = dissolved inorganic carbon, TA =total alkalinity. Data are the mean  $\pm$  SD (n = 3).

TA = total alkalinity. Data are the mean  $\pm$  SD (n = 3).

- 1 Table 1. Parameters of the seawater carbonate system in different cultures. DIC = dissolved inorganic carbon, TA = total alkalinity.
- 2 Data are the mean  $\pm$  SD (n = 3).

|          | pH <sub>NBS</sub>             | pCO <sub>2</sub> (μatm)           | DIC (μmol kg <sup>-1</sup> )     | HCO <sub>3</sub> <sup>-</sup> (μmol kg <sup>-1</sup> ) | CO <sub>3</sub> <sup>2-</sup> (μmol kg <sup>-1</sup> ) | CO <sub>2</sub> (μmol kg <sup>-1</sup> ) | TA (μmol kg <sup>-1</sup> )      |
|----------|-------------------------------|-----------------------------------|----------------------------------|--------------------------------------------------------|--------------------------------------------------------|------------------------------------------|----------------------------------|
| 8:16 LC  | 8.14 $\pm$ 0.02 <sup>a</sup>  | 580.32 $\pm$ 35.74 <sup>a</sup>   | 2597.04 $\pm$ 57 <sup>a</sup>    | 2385.19 $\pm$ 56.15 <sup>a</sup>                       | 192.59 $\pm$ 7.23 <sup>a</sup>                         | 19.56 $\pm$ 1.19 <sup>a</sup>            | 2850.78 $\pm$ 56.26 <sup>a</sup> |
| 12:12 LC | 8.12 $\pm$ 0.01 <sup>a</sup>  | 613.13 $\pm$ 11.21 <sup>a</sup>   | 2594.32 $\pm$ 61.96 <sup>a</sup> | 2391.03 $\pm$ 55.85 <sup>a</sup>                       | 182.94 $\pm$ 6.07 <sup>a</sup>                         | 20.35 $\pm$ 0.37 <sup>a</sup>            | 2834.31 $\pm$ 67.99 <sup>a</sup> |
| 16:8 LC  | 8.13 $\pm$ 0.01 <sup>a</sup>  | 608.54 $\pm$ 14.83 <sup>a</sup>   | 2638.26 $\pm$ 40.24 <sup>a</sup> | 2427.94 $\pm$ 35.09 <sup>a</sup>                       | 190.12 $\pm$ 6.73 <sup>a</sup>                         | 20.19 $\pm$ 0.49 <sup>a</sup>            | 2886.86 $\pm$ 46.67 <sup>a</sup> |
| 8:16 HC  | 7.74 $\pm$ 0.03 <sup>b</sup>  | 1614.77 $\pm$ 100.05 <sup>b</sup> | 2780.28 $\pm$ 12.05 <sup>b</sup> | 2641.71 $\pm$ 12.23 <sup>b</sup>                       | 84.99 $\pm$ 5.08 <sup>b</sup>                          | 53.58 $\pm$ 3.32 <sup>b</sup>            | 2854.16 $\pm$ 13.32 <sup>a</sup> |
| 12:12 HC | 7.79 $\pm$ 0.02 <sup>cd</sup> | 1445.67 $\pm$ 58.75 <sup>c</sup>  | 2799.47 $\pm$ 14.61 <sup>b</sup> | 2655.70 $\pm$ 13.74 <sup>b</sup>                       | 95.80 $\pm$ 3.81 <sup>b</sup>                          | 47.97 $\pm$ 1.95 <sup>c</sup>            | 2893.15 $\pm$ 17.18 <sup>a</sup> |
| 16:8 HC  | 7.75 $\pm$ 0.01 <sup>bd</sup> | 1652.45 $\pm$ 43.38 <sup>b</sup>  | 2938.18 $\pm$ 47.63 <sup>c</sup> | 2790.86 $\pm$ 45.52 <sup>c</sup>                       | 92.48 $\pm$ 1.28 <sup>b</sup>                          | 54.83 $\pm$ 1.44 <sup>b</sup>            | 3019.14 $\pm$ 46.95 <sup>b</sup> |

3

## Table 2 (on next page)

Two-way ANOVA analysis of variance for the effects of CO<sub>2</sub> and photoperiod on pH, pCO<sub>2</sub>, dissolved inorganic carbon (DIC), HCO<sub>3</sub><sup>-</sup>, CO<sub>3</sub><sup>2-</sup>, CO<sub>2</sub> and total alkalinity (TA) in seawater.

CO<sub>2</sub>\*photoperiod means the interactive effects of CO<sub>2</sub> and photoperiod, df mean degree of freedom and F means the value of F statistic, and Sig. means *p*-value.

1 Table 2. Two-way ANOVA analysis of variance for the effects of CO<sub>2</sub> and photoperiod regimes on pH, *p*CO<sub>2</sub>, dissolved inorganic  
2 carbon (DIC), HCO<sub>3</sub><sup>-</sup>, CO<sub>3</sub><sup>2-</sup>, CO<sub>2</sub> and total alkalinity (TA) in seawater. CO<sub>2</sub>\*photoperiod means the interactive effects of CO<sub>2</sub> and  
3 photoperiod, df mean degree of freedom and F means the value of F statistic, and Sig. means *p*-value.

| Source                      | pH |          | <i>p</i> CO <sub>2</sub> |          | DIC    |         | HCO <sub>3</sub> <sup>-</sup> |         | CO <sub>3</sub> <sup>2-</sup> |          | CO <sub>2</sub> |          | TA     |       |       |
|-----------------------------|----|----------|--------------------------|----------|--------|---------|-------------------------------|---------|-------------------------------|----------|-----------------|----------|--------|-------|-------|
|                             | df | F        | Sig.                     | F        | Sig.   | F       | Sig.                          | F       | Sig.                          | F        | Sig.            | F        | Sig.   |       |       |
| Photoperiod                 | 2  | 1.333    | 0.3                      | 5.640    | 0.019  | 9.697   | 0.003                         | 10.112  | 0.003                         | .353     | 0.710           | 5.632    | 0.019  | 8.638 | 0.005 |
| CO <sub>2</sub>             | 1  | 2268.750 | <0.001                   | 1498.294 | <0.001 | 125.390 | <0.001                        | 236.530 | <0.001                        | 1453.669 | <0.001          | 1497.372 | <0.001 | 8.977 | 0.011 |
| Photoperiod*CO <sub>2</sub> | 2  | 7.750    | 0.007                    | 7.562    | 0.007  | 3.054   | 0.085                         | 3.184   | 0.078                         | 5.342    | 0.022           | 7.557    | 0.008  | 2.976 | 0.089 |
| Error                       | 12 |          |                          |          |        |         |                               |         |                               |          |                 |          |        |       |       |

4

# **Table 3**(on next page)

Two-way ANOVA analysis of variance for the effects of CO<sub>2</sub> and photoperiod regimes on RGR of *Ulva linza* cultured at different CO<sub>2</sub> levels and photoperiod regimes.

CO<sub>2</sub>\*photoperiod means the interactive effects of CO<sub>2</sub> and photoperiod, df mean degree of freedom and F means the value of F statistic, and Sig. means *p*-value.

1 Table 3. Two-way ANOVA analysis of variance for the effects of CO<sub>2</sub> and photoperiod regimes  
 2 on RGR of *Ulva linza* cultured at different CO<sub>2</sub> levels and photoperiod regimes.  
 3 CO<sub>2</sub>\*photoperiod means the interactive effects of CO<sub>2</sub> and photoperiod, df mean degree of  
 4 freedom and F means the value of F statistic, and Sig. means *p*-value.

| Source                       | df | F       | Sig.   |
|------------------------------|----|---------|--------|
| Photoperiod                  | 2  | 124.249 | <0.001 |
| CO <sub>2</sub>              | 1  | 42.292  | <0.001 |
| Photoperiod* CO <sub>2</sub> | 2  | 65.959  | <0.001 |
| Error                        | 12 |         |        |

5

# **Table 4**(on next page)

Two-way ANOVA analysis of variance for the effects of CO<sub>2</sub> and photoperiod regimes on Yield and NPQ of *Ulva linza* cultured at different CO<sub>2</sub> levels and photoperiod regimes.

CO<sub>2</sub>\*photoperiod means the interactive effects of CO<sub>2</sub> and photoperiod, df mean degree of freedom and F means the value of F statistic, and Sig. means *p*-value.

- 1 Table 4. Two-way ANOVA analysis of variance for the effects of CO<sub>2</sub> and photoperiod regimes
- 2 on Yield and NPQ of *Ulva linza* cultured at different CO<sub>2</sub> levels and photoperiod regimes.
- 3 CO<sub>2</sub>\*photoperiod means the interactive effects of CO<sub>2</sub> and photoperiod, df mean degree of
- 4 freedom and F means the value of F statistic, and Sig. means *p*-value.

| Source                       | Yield |        |        | NPQ |       |       |
|------------------------------|-------|--------|--------|-----|-------|-------|
|                              | df    | F      | Sig.   | df  | F     | Sig.  |
| Photoperiod                  | 2     | 0.919  | 0.425  | 2   | 8.453 | 0.005 |
| CO <sub>2</sub>              | 1     | 2.549  | 0.136  | 1   | 0.132 | 0.722 |
| Photoperiod* CO <sub>2</sub> | 2     | 19.786 | <0.001 | 2   | 0.339 | 0.719 |
| Error                        | 12    |        |        | 12  |       |       |

5

# **Table 5**(on next page)

Parameters of rapid light curves (RLC) of *U. linza* thalli cultured at different CO<sub>2</sub> levels and photoperiod regimes.

Data are the mean  $\pm$  SD ( $n \geq 3$ ). Different letters represent significant difference ( $p < 0.05$ ) among different treatments.

1 Table 5. Parameters of rapid light curves (RLC) of *U. linza* thalli cultured at different CO<sub>2</sub> levels  
 2 and photoperiod regimes. Data are the mean  $\pm$  SD ( $n \geq 3$ ). Different letters represent significant  
 3 difference ( $p < 0.05$ ) among different treatments.

|          | rETRmax                       | $\alpha$                      | I <sub>k</sub>                     |
|----------|-------------------------------|-------------------------------|------------------------------------|
| 8:16 LC  | 47.53 $\pm$ 3.63 <sup>a</sup> | 0.22 $\pm$ 0.04 <sup>a</sup>  | 215.67 $\pm$ 19.89 <sup>a</sup>    |
| 12:12 LC | 43.87 $\pm$ 3.60 <sup>c</sup> | 0.18 $\pm$ 0.02 <sup>c</sup>  | 248.87 $\pm$ 22.91 <sup>bd</sup>   |
| 16:8 LC  | 54.83 $\pm$ 2.65 <sup>b</sup> | 0.24 $\pm$ 0.03 <sup>ab</sup> | 231.58 $\pm$ 21.37 <sup>adef</sup> |
| 8:16 HC  | 56.51 $\pm$ 2.34 <sup>b</sup> | 0.25 $\pm$ 0.01 <sup>b</sup>  | 222.15 $\pm$ 13.41 <sup>ac</sup>   |
| 12:12 HC | 60.45 $\pm$ 1.38 <sup>d</sup> | 0.26 $\pm$ 0.01 <sup>b</sup>  | 236.58 $\pm$ 12.63 <sup>bce</sup>  |
| 16:8 HC  | 54.55 $\pm$ 3.77 <sup>b</sup> | 0.22 $\pm$ 0.01 <sup>a</sup>  | 251.72 $\pm$ 11.69 <sup>bf</sup>   |

4

# Table 6 (on next page)

Two-way ANOVA analysis of variance for the effects of CO<sub>2</sub> and photoperiod regimes on maximum rETR (rETR<sub>max</sub>), efficiency of electron transport ( $\alpha$ ), and saturating irradiance ( $I_k$ ) of *Ulva linza*.

CO<sub>2</sub>\*photoperiod means the interactive effects of CO<sub>2</sub> and photoperiod, df mean degree of freedom and F means the value of F statistic, and Sig. means *p*-value.

- 1 Table 6. Two-way ANOVA analysis of variance for the effects of CO<sub>2</sub> and photoperiod regimes on maximum rETR (rETRmax),
- 2 efficiency of electron transport ( $\alpha$ ), and saturating irradiance ( $I_k$ ) of *Ulva linza* cultured at different CO<sub>2</sub> levels and photoperiod
- 3 regimes. CO<sub>2</sub>\*photoperiod means the interactive effects of CO<sub>2</sub> and photoperiod, df mean degree of freedom and F means the value of
- 4 F statistic, and Sig. means *p*-value.

| Source                       | ETRmax |        |        | $\alpha$ |        |        | $I_k$ |       |       |
|------------------------------|--------|--------|--------|----------|--------|--------|-------|-------|-------|
|                              | Df     | F      | Sig.   | df       | F      | Sig.   | df    | F     | Sig.  |
| Photoperiod                  | 2      | 2.981  | 0.066  | 2        | 2.909  | 0.070  | 2     | 7.030 | 0.003 |
| CO <sub>2</sub>              | 1      | 69.993 | <0.001 | 1        | 15.225 | <0.001 | 1     | 0.664 | 0.421 |
| Photoperiod* CO <sub>2</sub> | 2      | 23.445 | <0.001 | 2        | 14.896 | <0.001 | 2     | 2.576 | 0.093 |
| Error                        | 30     |        |        | 30       |        |        | 30    |       |       |

# Table 7 (on next page)

Two-way ANOVA analysis of variance for the effects of CO<sub>2</sub> and photoperiod regimes on respiration rate of *Ulva linza* cultured at different CO<sub>2</sub> levels and photoperiod regimes.

CO<sub>2</sub>\*photoperiod means the interactive effects of CO<sub>2</sub> and photoperiod, df mean degree of freedom and F means the value of F statistic, and Sig. means *p*-value.

1 Table 7. Two-way ANOVA analysis of variance for the effects of CO<sub>2</sub> and photoperiod regimes  
 2 on respiration rate of *Ulva linza* cultured at different CO<sub>2</sub> levels and photoperiod regimes.  
 3 CO<sub>2</sub>\*photoperiod means the interactive effects of CO<sub>2</sub> and photoperiod, df mean degree of  
 4 freedom and F means the value of F statistic, and Sig. means *p*-value.

| Source                       | df | F      | Sig.   |
|------------------------------|----|--------|--------|
| Photoperiod                  | 2  | 16.006 | <0.001 |
| CO <sub>2</sub>              | 1  | 33.058 | <0.001 |
| Photoperiod* CO <sub>2</sub> | 2  | 9.566  | 0.003  |
| Error                        | 12 |        |        |

5

# Table 8 (on next page)

Two-way ANOVA analysis of variance for the effects of CO<sub>2</sub> and photoperiod regimes on Chl *a*, Chl *b*, and Chl *a/b* of *Ulva linza* cultured at different CO<sub>2</sub> levels and photoperiod regimes.

CO<sub>2</sub>\*photoperiod means the interactive effects of CO<sub>2</sub> and photoperiod, df mean degree of freedom and F means the value of F statistic, and Sig. means *p*-value.

1 Table 8. Two-way ANOVA analysis of variance for the effects of CO<sub>2</sub> and photoperiod regimes on Chl *a*, Chl *b*, and Chl *a/ b* of *Ulva*  
 2 *linza* cultured at different CO<sub>2</sub> levels and photoperiod regimes. CO<sub>2</sub>\*photoperiod means the interactive effects of CO<sub>2</sub> and photoperiod,  
 3 df mean degree of freedom and F means the value of F statistic, and Sig. means *p*-value.

| Source                       | Chl <i>a</i> |         |        | Chl <i>b</i> |         |        | Chl <i>a/ b</i> |       |       |
|------------------------------|--------------|---------|--------|--------------|---------|--------|-----------------|-------|-------|
|                              | df           | F       | Sig.   | df           | F       | Sig.   | df              | F     | Sig.  |
| Photoperiod                  | 2            | 121.283 | <0.001 | 2            | 111.035 | <0.001 | 2               | 2.625 | 0.113 |
| CO <sub>2</sub>              | 1            | 51.793  | <0.001 | 1            | 53.849  | <0.001 | 1               | 0.171 | 0.686 |
| Photoperiod* CO <sub>2</sub> | 2            | 26.932  | <0.001 | 2            | 21.719  | <0.001 | 2               | 8.389 | 0.005 |
| Error                        | 12           |         |        | 12           |         |        | 12              |       |       |

4