

Effects of prolonged continuous computer gaming on physical and ocular symptoms and binocular vision functions in young healthy individuals

Ji-Woo Lee ¹, Hyun Gug Cho ², Byeong-Yeon Moon ², Sang-Yeob Kim ², Dong-Sik Yu ^{Corresp. 2}

¹ Department of Optometry and Vision Science, Kyungwoon University, Gumi, South Korea

² Department of Optometry, Kangwon National University, Samcheok, South Korea

Corresponding Author: Dong-Sik Yu

Email address: yds@kangwon.ac.kr

Background and Objective. Addiction to computer gaming has become a social problem in Korea and elsewhere, and it has been enlisted as a mental health disorder by the World Health Organization. Most studies related to computer use and vision have individually assessed physical and ocular symptoms and binocular vision. Accordingly, the present study comprehensively assessed subjective physical and ocular symptoms and functions related to binocular vision after prolonged continuous computer gaming. This study aimed to investigate the effects of prolonged continuous computer gaming on physical and ocular health and visual functions in young healthy individuals.

Methods. Fifty healthy college students (35 male/ 15 female), aged 19-35 years old, were enrolled in this study. The inclusion criteria were no binocular vision problems and no reported history of ocular disease. Participants played continuously for 4 h from 6:00 to 10:00 p.m. Physical and ocular symptoms and visual functions such as convergence, accommodation, phoria, and the blink rate were assessed before and after continuous computer gaming for 4 h.

Results. Continuous computer gaming for 4 h resulted in convergence and accommodation disturbances and increased physical and ocular discomfort. Near phoria showed a divergent shift, whereas distance phoria showed no change. Moreover, the accommodative and vergence facilities and blink rate were significantly decreased. All visual functions recovered to the baseline levels by the following morning.

Discussion. Our findings suggest that excessive and continuous computer gaming impairs visual functions and causes ocular and physical fatigue. Our findings further the understanding of the adverse effects of excessive computer use on physical and ocular

health and highlight the need for measures such as visual exercises, and adequate rest periods for individuals performing computer-related tasks on a daily basis.

Effects of prolonged continuous computer gaming on physical and ocular symptoms and binocular vision functions in young healthy individuals

Ji-Woo Lee¹, Hyun Gug Cho², Byeong-Yeon Moon², Sang-Yeob Kim² and Dong-Sik Yu²

¹ Department of Optometry and Vision Science, Kyungwoon University, Gumi, South Korea

² Department of Optometry, Kangwon National University, Samcheok, South Korea

Corresponding Author:

Dong-Sik Yu

Hwangjogil 346, Dogye-up, Samcheok, Gangwondo 25949, Republic of Korea

Email address: yds@kangwon.ac.kr

ABSTRACT

Background and Objective. Addiction to computer gaming has become a social problem in Korea and elsewhere, and it has been enlisted as a mental health disorder by the World Health Organization. Most studies related to computer use and vision have individually assessed physical and ocular symptoms and binocular vision. Accordingly, the present study comprehensively assessed subjective physical and ocular symptoms and functions related to binocular vision after prolonged continuous computer gaming. This study aimed to investigate the effects of prolonged continuous computer gaming on physical and ocular health and visual functions in young healthy individuals.

Methods. Fifty healthy college students (35 male/ 15 female), aged 19-35 years old, were enrolled in this study. The inclusion criteria were no binocular vision problems and no reported history of ocular disease. Participants played continuously for 4 h from 6:00 to 10:00 p.m. Physical and ocular symptoms and visual functions such as convergence, accommodation, phoria, and the blink rate were assessed before and after continuous computer gaming for 4 h.

Results. Continuous computer gaming for 4 h resulted in convergence and accommodation disturbances and increased physical and ocular discomfort. Near phoria showed a divergent shift, whereas distance phoria showed no change. Moreover, the accommodative and vergence facilities and blink rate were significantly decreased. All visual functions recovered to the baseline levels by the following morning.

Discussion. Our findings suggest that excessive and continuous computer gaming impairs visual functions and causes ocular and physical fatigue. Our findings further the understanding of the adverse effects of excessive computer use on physical and ocular health and highlight the need for measures such as visual exercises, and adequate rest periods for individuals performing computer-related tasks on a daily basis.

INTRODUCTION

The ability to visualize near objects clearly and accurately is necessary for individuals using computers and other digital screen devices on a daily basis. In today's digital age, prolonged use of visual (or video) display terminals (VDTs) such as computers and smartphones has become very common and is one the main causes of ocular and physical discomfort (González-Pérez et al., 2014; Parihar et al., 2016). The majority of ocular and physical symptoms are associated with the effects of prolonged VDT usage on visual functions (Fenga et al., 2007; Tyrrell & Leibowitz, 1990; Yeow & Taylor, 1991). Computer vision syndrome (CVS), a condition associated with excessive computer use, is characterized by not only visual symptoms such as eyestrain, dry eyes, and blurred and double vision but also physical discomfort such as musculoskeletal shoulder disorders, neck pain, headache, and dizziness (Berqqvist et al., 1995; Hayes et al., 2007; Rosenfield, 2011). According to the Korea Games Users Survey Reports 2017, the rate of computer gaming in the general Korean population aged 10 to 65 years-old was increased from 67.9% in 2016 to 70.3% in 2017 (The Korea Creative Content Agency, 2017), which could lead to increased problems related to computer use.

Computer games have become a part of leisure activities in daily life, with young individuals spending excessive time indulging in such games (Wittek et al., 2016). The current most popular games require users to pay close attention to their VDTs, and the players show deep levels of emotional and physical addiction while gaming (Kim et al., 2016; Rikkers et al., 2016). Ubiquitous gaming cafes known as "PC bang" are very popular in Korea (Zastrow, 2017), where 24% of children are reportedly diagnosed with internet gaming addiction requiring hospitalization (Ahn, 2017). Computer gaming addiction can cause health and mental issues (Griffiths, Kussa & King, 2012; Saquib et al., 2017). Furthermore, excessive computer use has been reported to cause problems associated with poor computer (Toomingas, 2014). A recent study showed that the neck and shoulder are the most commonly reported areas affected by excessive computer use (James et al., 2018). Other commonly reported symptoms include headache, eyestrain, double vision, dry eyes, and ocular fatigue (Akinbinu & Mashalla, 2014). Some studies (Qu et al., 2005; Yekta, Pickwell, & Jenkins, 1989) have also reported that near work on VDTs may cause not only ocular and physical symptoms but also transient changes in refraction, visual acuity, accommodation, and convergence. Tosha et al. (2009) examined the relationship between visual discomfort and the accommodation response in a college population with visual discomfort and found that increased visual discomfort is characterized by accommodative fatigue, with a higher lag of accommodation developing at a near viewing distance over time.

Furthermore, a study of over 1,173 Korean school-going children reported that the participants spent approximately 1.35 h on computer-based academics and 2.03 h on computer gaming during a normal day (Yang et al., 2014). Computer overuse has led to progressive deterioration the visual environment yearly in the users' physical and mental health, and the "shutdown law" was recently introduced in Korea for the protection of youth from internet-based

computer games (Kim et al., 2016; Sung, 2014). Actually, most individuals addicted or immersed to computer games tend to play for prolonged hours without a break, and the effects of prolonged continuous viewing of VDTs at a near distance on physical and ocular health remain unclear. Moreover, most studies related to computer use and vision have individually evaluated physical and ocular symptoms and binocular vision (Parihar et al., 2016; Hamed et al., 2013). Therefore, we believe that a study focused on a comprehensive assessment of subjective physical and ocular symptoms and functions related to binocular vision after prolonged continuous computer use is necessary. Accordingly, the aim of this study was to investigate subjective physical and ocular symptoms and visual functions, as well as their interrelationships, after prolonged continuous computer gaming in young healthy individuals.

METHODS

Participants

This research complied with the tenets of the Declaration of Helsinki and was approved by the Institutional Review Board of Kangwon National University (KWNUIRB-2017-07-002-003). Informed consent was obtained from each participant.

Sample size was estimated at 0.05 for the alpha and 0.20 for the beta level using MedCalc software version 12.7.7.0 (MedCalc Software, Mariakerke, Belgium) based on the difference of computer-using time between a 3.6 ± 2.2 hours per day and a 2.4 ± 1.6 hours per day for internet-addicted and non-addicted Korea adolescents in the previous study, respectively (Yang et al., 2014). A sample size of 41 participants was required but fifty Korean healthy college students (age, 19-35 years, average age, 22.5 ± 3.4 years; male:female, 35:15) were randomly enrolled as subjects for our study. The inclusion criteria were as follows: no near or distance vision problems; a corrected or uncorrected distance visual acuity of 20/25 (0.8) or better in each eye; normal stereopsis; and no reported history of ocular disease. Twenty-nine subjects (58%) wore spectacles, and their mean spherical equivalent power was -1.99 ± 2.53 diopters (D).

This study was performed under immersive computer gaming that involved general tasks for a prolonged period (4 h) without a break versus a short period (≤ 2 h) of VDT exposure in order to determine the presence of physical and ocular discomfort.

Procedures

The subjects were asked to play a computer game (Diablo III, Blizzard Entertainment, USA) on a desktop computer screen (CX501N-KN/KOR with a 15-inch LCD monitor, Samsung, Korea) from a viewing distance of 50 cm under room illumination (approximately 50 lux). They played continuously for 4 h from 6:00 to 10:00 p.m. All subjects received a modified questionnaire (Ames et al., 2005) designed for assessing the effects of virtual reality viewing on monocular, binocular, and physical symptoms, as well as ophthalmological examinations for the assessment of visual functions, before and after the gaming session.

The questionnaire included 13 items pertaining to four physical and nine ocular symptoms. Each item was graded on a scale of 0 to 4 (0 = none, 1 = slight, 2 = moderate, 3 = severe, and 4 =

very severe). Visual functions were measured under habitual viewing conditions. Visual acuity was measured using visual charts (ACP-8, Topcon, Japan) and a phoropter (CV-3000, Topcon, Japan). The near point of convergence (NPC) is to evaluate the convergence amplitude, which represents a visual function to obtain single vision, was measured by bringing an accommodative target to the nose and recording the point time at which the subject could see double (Scheiman et al., 2003). The near point of accommodation (NPA) is the near point to measure the amplitude of accommodation under binocular conditions, which represents a visual function for maintaining a clear image (Abraham et al., 2005), was measured using an accommodative convergence rule (GR50, Bernell, USA) while the subject attempted to read small letters (near visual acuity, 20/32). Phoria is to assess the presence, direction, and amplitude of the eye alignment, which indicates latent misalignment of the eyes, was measured using Howell phoria cards (Wong, Fricke & Dinardo, 2002) (CDHP, Bernell, USA) at 3 m and 33 cm. Negative values indicate exophoria, whereas positive values indicate esophoria. Accommodative and vergence facilities is to evaluate the ability of the accommodative response and ability of the fusional vergence respectively (Weissberg, 2004; Gall, Wick & Bedell, 1998), which were measured using a ± 2.0 D binocular flipper lens and a prism flipper (3 Δ base-in + 12 Δ base-out; Δ : prism diopter) at 40 cm. Blink rate was measured using a camcorder combining a video camera (NV-GS400, Panasonic, Japan), complete and incomplete blinks were distinguished by the presence of the full and partial eye cycle of open-close-open. Binocular function was measured at time-points of before and after the computer game, and the next morning.

Statistical Analysis

All data were statistically analyzed using SPSS (ver. 18.0 for Windows, SPSS Inc., Chicago, IL, USA). Pearson's correlation analysis, paired t-tests, and repeated measures analysis of variance (ANOVA) were used for statistical analyses. The level of significance was set at $p = 0.05$. If significant differences were found, post hoc tests with Bonferroni corrections were used to identify the level of significance. Partial missing values in phoria and blink rate were excluded from descriptive statistics analyses.

RESULTS

Our results are not limited to gaming activities, but applicable under the environment of overuse of personal computers for the purpose of computer game after the college students' work day. The mean scores for the four physical (monocular) and nine ocular symptoms associated with computer game use are shown in Table 1. The mean overall scores for the physical and ocular discomfort domains increased from 0.39 ± 0.78 and 0.50 ± 0.83 , respectively, before the gaming session to 1.61 ± 1.24 ($t = 15.7$, $p < 0.001$) and 1.40 ± 1.22 ($t = 17.44$, $p < 0.001$), respectively, after the gaming session. The scores for all individual items also showed significant changes after the gaming session, and the mean difference was 1.00 ± 0.33 . In the physical discomfort domain, the score for neck discomfort after the gaming session was the highest, followed by the

scores for shoulder discomfort, headache, and back discomfort. In the ocular discomfort domain, the score for tired eyes was the highest (2.30 ± 1.09), followed by the scores for dry eyes, blurred vision, and eyestrain. The score for itchy eyes was the lowest (0.82 ± 1.00).

The effects of prolonged continuous computer gaming on visual functions are shown in [Table 2](#). The mean NPC values showed significant changes immediately after and the morning after the gaming session ($F = 47.83$, $p < 0.001$). A post hoc Bonferroni multiple comparisons test indicated that NPC significantly increased immediately after the gaming session and recovered to the baseline level by morning. The mean NPA value also showed significant changes ($F = 42.50$, $p < 0.001$). A multiple comparisons test with Bonferroni correction showed that the NPA value after the gaming session was significantly different from the next-day value and the baseline value ($p < 0.001$ for both); the next-day value and the baseline value showed no significant differences ($p = 1.000$).

The accommodative facility (AF) and vergence facility (VF) significantly decreased after the gaming session. Both facilities recovered to baseline levels by the following morning.

The blink rate before the gaming session was 16.24 times per minute; this significantly decreased to 8.27 and 9.51 times per minute at 1 and 4h after the start of the game, respectively ($F = 106.29$, $p < 0.001$).

The changes in phoria during the computer gaming session are shown in [Fig. 1](#). Near phoria showed a divergent shift (exophoria) during the session (repeated measures ANOVA; $F = 24.25$, $p < 0.001$) and recovered to baseline levels by the following morning. Distance phoria showed no significant differences during the game (repeated measures ANOVA; $F = 0.61$, $p = 0.610$). Horizontal phoria at distance showed no statistically significant differences among before, after, and the day after the gaming session ($F = 1.67$, $p = 0.193$), whereas horizontal phoria at near changed from -3.73 ± 3.93 before the gaming session to -5.75 ± 4.85 and -3.68 ± 4.86 immediately after and the day after the gaming session, respectively ($F = 18.39$, $p < 0.001$).

Pearson's correlation analysis showed a very weak negative correlation between itchy eyes and NPC after the gaming session ($r = -0.294$, $p = 0.039$) and a weak positive correlation between tired eyes and NPA ($r = 0.361$, $p = 0.01$) and between blurred vision and NPA ($r = 0.298$, $p = 0.036$) after the gaming session.

DISCUSSION

In the present study, we found that prolonged continuous computer use for gaming resulted in both physical and ocular discomfort as well as changes in binocular functions. In particular, the neck and shoulder, which remain in the same posture while playing, were affected. The major visual symptom was ocular fatigue (tired eyes), followed by dryness and blurred vision. These findings are consistent with those of other studies ([Hayes et al., 2007](#); [Klussmann et al., 2008](#)) reporting a high prevalence of neck and shoulder symptoms, ocular fatigue, and blurred vision after excessive computer use. The findings of a 13-item questionnaire used in our study revealed higher scores for physical symptoms than for ocular symptoms after 4 continuous hours of

gaming, although tired eyes was associated with the highest score. These results indicated that the physical problems related to the neck, shoulders, and backs were the most affected regions among the computer users, other ocular symptoms were the most frequently occurring health problems (Lanthers et al., 2016), and of the 59 included workers who use a computer for 3-6 h per day, the musculoskeletal problems were reported by 71.1% of subjects as compared to that for visual problems of 61.0% out (Talwar et al., 2009); thus, after physical discomfort, discomfort of the eye is the second most frequent problem reported by VDT operators. Tired eyes can be explained by the fact that high accommodation and convergence values are required for maintaining a clear vision at small distances from VDTs, and these values need to be maintained for the entire gaming duration. This invariably results in ocular fatigue. On the other hand, physical factors associated with discomfort had lower scores than did tired eyes due to frequent changes in the body posture to relieve physical stress.

We also found changes in visual functions after continuous computer gaming for 4 h in the present study. NPC, a visual function used to achieve single vision during near-distance work, showed a significant increase after the gaming session and recovered to the baseline level the next day. Qu et al. (2005) found that NPC was significantly increased after short-term VDT use. However, Yeow and Taylor (1991) found that NPC decreased with an increase in age, with no significant difference between VDT and non-VDT users. Hamed, David, and Marzieh (2013) reported that the evaluation of NPC helped in the differentiation of symptomatic and asymptomatic subjects, and that the average NPC value for break point was 11.7 cm. While NPC is a measurement of how close one can bring a fixation target to the nose while maintaining fusion, NPA is used to assess the amplitude of accommodation. Accommodation is connected with the function of vergence maintained by extracocular muscles. Because of the interaction that occur between accommodation and vergence, the accommodation and convergence disturbances observed after the gaming session in the present study can be attributed to intra- and extra ocular muscle fatigue resulting from the prolonged continuous computer gaming.

Accommodative and vergence facilities permit the ability to sustain clear and single binocular vision during near-distance work. A healthy individual's accommodative facility should be binocularly clear within 10.0 ± 5 cpm (Scheiman & Wick, 2002). In the present study, the value showed a significant change from 12.68 to 15.54 cpm. The vergence facility for single binocular vision also showed a significant change from 15.02 to 17.70 cpm; the norm is 15 cpm (Weissberg, 2004). These changes may be associated with symptoms such as ocular fatigue, eyestrain, and blurred vision, although NPC and NPA showed significant changes within normal values. Both accommodative and vergence facilities recovered by the next morning.

The results of the present study also showed that near phoria (dissociated phoria) increased from 3.73 Δ exophoria to 5.75 Δ exophoria with the development of symptoms after continuous computer gaming for 4 h, and it subsequently recovered to 3.68 Δ exophoria by the following morning. This result is consistent with those of other studies (Yekta et al., 1989; Tsubota & Nakamori, 1993) showing an exophoric shift in phoria due to near-distance tasks, with no significant change in distance phoria over time. According to other studies related to changes of

phoria, [Pickwell, Jenkins, and Yekta \(1987\)](#) reported that 30 min of reading in inadequate illumination or from an abnormally close distance resulted in an increase in phoria with the development of ocular symptoms. Also, [Yekta et al. \(1989\)](#) found that near-dissociated phoria increased from 4.70 Δ exophoria to 5.20 Δ exophoria after 30 min of reading and decreased to 4.87 Δ exophoria after 30 min of relaxation, with no accompanying symptoms. Collectively, these findings including our results suggest that near phoria for binocular vision changes to exophoria under the effect of visual stress caused by factors such as inappropriate illumination, abnormal working distance, and prolonged working hours.

The mean blinking rate at rest reportedly varies from 12 to 19 blinks per minute ([Karson et al., 1981](#)). [Tsubota and Nakamori \(1993\)](#) observed 22 blinks per minute under relaxed conditions and seven blinks per minute during VDT work. [Patel et al. \(1991\)](#) reported 18.4 blinks per minute before a computer task and 3.6 blinks per minute during the computer task. We found that the number of blinks per minute significantly decreased from 16.24 before the gaming session to 8.27 after 1 h of gaming and 9.51 after 4 h of continuous gaming. However, we do not know whether the decreased blink rate induces ocular fatigue, or vice versa. However, the blink rate may be an effective index for assessing visual fatigue during VDT tasks. Intentional blinking during VDT tasks is required to minimize visual fatigue. Blinking exercises have been shown to improve the glandular function and reduce the frequency of incomplete blinks ([Murakami et al., 2014](#); [Downie & Craig, 2017](#)). Therefore, we highly recommend blink training to increase the blinking rate during prolonged computer use.

Pearson's correlation analysis showed that ocular fatigue and blurred vision were weakly related to NPA, whereas itchiness was weakly related to NPC. These subjective symptoms may be provoked by the accommodation and convergence disturbances induced during prolonged computer gaming. Although these changes are usually temporary, visual fatigue caused by long-term VDT use tends to accumulate over time as reported by [Murata et al. \(1996\)](#).

This study is limited by the small study sample that included patients with healthy eyes without binocular vision problems so that there were minimum confounding variables. A larger sample of patients with binocular disorders is necessary for extensive estimation of the effects of prolonged continuous computer gaming. Also, the appeared points of physiological effects and the recovery were not exactly examined in this study.

The time-points of physiological effects and recovery were not exactly examined in this study. Nevertheless, our findings offer valuable insights and provide direct evidence of significant changes in physical and ocular symptoms and visual function after a 4-h period of computer game. The time-points of physiological effects could be variable due to influencing factors such as individual and ergonomic factors. The decreased accommodative functions in the subjects after performing VDT task for 2 h were recovered at the end of the 1-h lunch break ([Saito et al., 1994](#)). Accommodative functions after resting for 0.5 h following 1.5 h VDT task showed a tendency to return to the previous normal value ([Yoo et al., 1992](#)). These previous studies suggest that the recovery time of symptoms and signs may depended on the strength of VDT work.

CONCLUSIONS

In summary, the findings clearly indicate that changes in visual functions evoke ocular symptoms after prolonged continuous computer gaming, which also induces physical symptoms. Although these symptoms are usually temporary, the accumulation of fatigues could be potential risk factors for irreversible physical and visual disturbances associated with computer overuse. Users may need rest to alleviate fatigue due to computer use. Moreover, individuals should be advised regarding their working hours. Therefore, it seems reasonable that all individuals should be advised to take breaks and periodically gaze into the distance to minimize accommodation and vergence disturbances; longer rest periods will result in lesser physical and ocular fatigue.

ACKNOWLEDGEMENTS

Not applicable.

REFERENCES

- Abraham LM, Kuriakose T, Sivanandam V, Venkatesan N, Thomas R, Muliyl J. 2005. Amplitude of accommodation and its relation to refractive errors. *Indian J Ophthalmol* 53(2):105–108 DOI: 10.4103/0301-4738.16173.
- Ahn DH. 2007. Korean policy on treatment and rehabilitation for adolescents' internet addiction. In: *2007 International Symposium on the Counseling and Treatment of Youth Internet Addiction*. Seoul, Korea: National Youth Commission, 49.
- Akinbinu TR, Mashalla YJ. 2014. Impact of computer technology on health: computer vision syndrome (CVS). *Med Pract Rev* 5(3):20–30 DOI: 10.5897/MPR.2014.0121.
- Ames SL, Wolffsohn JS, McBrien NA. 2005. The development of a symptom questionnaire for assessing virtual reality viewing using a head-mounted display. *Optom Vis Sci* 82(3):168–176 DOI: 10.1097/01.OPX.0000156307.95086.6.
- Berqqvist U, Wolgast E, Nilsson B, Voss M. 1995. The influence of VDT work on musculoskeletal disorders. *Ergonomics* 38(4):754–762 DOI: 10.1080/00140139508925147.
- Downie LE, Craig JP, 2017. Tear film evaluation and management in soft contact lens wear: a systematic approach. *Clin Exp Optom* 100(5):438–458 DOI: 10.1111/cxo.12597.
- Fenga C, Di Pietro R, Fenga P, Di Nola C, Spinella R, Cacciola A, Germanò D, Aragona P. 2007. Asthenopia e lavoro al VDT: nostra esperienza. *G Ital Med Lav Ergon* 29(Suppl 3):500–501.
- Gall R, Wick B, Bedell H. 1998. Vergence facility: establishing clinical utility. *Optom Vis Sci* 75(10):731–742 DOI: 10.1097/00006324-199810000-00018.
- González-Pérez M, Susi R, Antona B, Barrio A, González E. 2014. The computer-vision symptom scale (CVSS17): development and initial validation. *Invest Ophthalmol Vis Sci* 55(7):4504–4511 DOI:10.1167/iops.13-13818.
- Griffiths MD, Kussa DJ, King DL. 2012. Video game addiction: past, present and future. *Curr Psychiatry Rev* 8(4):308–318 DOI: 10.2174/157340012803520414.
- Hamed MM, David AG, Marzieh E. 2013. The relationship between binocular vision

- symptoms and near point of convergence. *Indian J Ophthalmol* **61**(7):325–328
DOI: [10.4103/0301-4738.97553](https://doi.org/10.4103/0301-4738.97553).
- Hayes JR, Sheedy JE, Stelmack JA, Heaney CA. 2007.** Computer use, symptoms, and quality
of life. *Optom Vis Sci* **84**(8):738–744 DOI: [10.1097/OPX.0b013e31812f7546](https://doi.org/10.1097/OPX.0b013e31812f7546).
- James C, James D, Nie V, Schumacher T, Guest M, Tessier J, Marley J, Bohatko-Naismith
J, Snodgrass S. 2018.** Musculoskeletal discomfort and use of computers in the university
environment. *Appl Ergon* **69**:128–135 DOI: [10.1016/j.apergo.2018.01.013](https://doi.org/10.1016/j.apergo.2018.01.013).
- Karson CN, Berman KF, Donnelly EF, Mendelson WB, Kleinman JE, Wyatt RJ. 1981.**
Speaking, thinking and blinking. *Psychiatry Res* **5**(3):243–246 DOI: [10.1016/0165-1781\(81\)90070-6](https://doi.org/10.1016/0165-1781(81)90070-6).
- Kim T, Kang MY, Yoo MS, Lee D, Hong YC. 2016.** Computer use at work is associated with
self-reported depressive and anxiety disorder. *Ann Occup Environ Med* **28**:57 DOI:
[10.1186/s40557-016-0146-8](https://doi.org/10.1186/s40557-016-0146-8).
- Klussmann A, Gebhardt H, Liebers F, Rieger MA. 2008.** Musculoskeletal symptoms of the
upper extremities and the neck: a cross-sectional study on prevalence and symptom-predicting
factors at visual display terminal (VDT) workstations. *BMC Musculoskelet Disord* **9**:96 DOI:
[10.1186/1471-2474-9-96](https://doi.org/10.1186/1471-2474-9-96).
- Lanthers C, Pereira B, Garde G, Maublant C, Dutheil F, Coudeyre E. 2016.** Evaluation of ‘I-
Preventive’: a digital preventive tool for musculoskeletal disorders in computer workers—a
pilot cluster randomised trial. *BMJ Open* **6**:e011304 DOI: [10.1136/bmjopen-2016-011304](https://doi.org/10.1136/bmjopen-2016-011304).
- Murakami D, Blackie CA, Korb DR. 2014.** Blinking exercises can be used to decrease partial
blinking and improve gland function and symptoms in patients with evaporative dry eye.
Denver: American Academy of Optometry. Available at
<https://www.aaopt.org/detail/knowledge-base-article/> (accessed 9 December 2018).
- Murata K, Araki S, Yokoyama K, Yamashita K, Okumatsu T, Sakou S. 1996.** Accumulation
of VDT work-related visual fatigue assessed by visual evoked potential, near point distance
and critical flicker fusion. *Ind Health* **34**(2):61–69 DOI: [10.2486/indhealth.34.61](https://doi.org/10.2486/indhealth.34.61).
- Parihar JKS, Jain VK, Chaturvedi P, Kaushik J, Jain G, Parihar AK. 2016.** Computer and visual
display terminals (VDT) vision syndrome (CVDTs). *Med J Armed Forces India* **72**(3):270–
276 DOI: [10.1016/j.mjafi.2016.03.016](https://doi.org/10.1016/j.mjafi.2016.03.016).
- Patel S, Henderson R, Bradley L, Galloway B, Hunter L. 1991.** Effect of visual display unit
use on blink rate and tear stability. *Optom Vis Sci* **68**(11):888–892 DOI: [10.1097/00006324-199111000-00010](https://doi.org/10.1097/00006324-199111000-00010).
- Pickwell D, Jenkins T, Yekta AA. 1987.** The effect on fixation disparity and associated
heterophoria of reading at an abnormally close distance. *Ophthalmic Physiol Opt* **7**(4):345–
347 DOI: [10.1111/j.1475-1313.1987.tb00758.x](https://doi.org/10.1111/j.1475-1313.1987.tb00758.x).
- Qu XM, Chu RY, Wang L, Yao PJ, Liu JR. 2005.** Effects of short-term VDT usage on visual
functions. *Zhonghua Yan Ke Za Zhi* **41**(11):986–989 DOI: [10.3760/j.issn:0412-4081.2005.11.007](https://doi.org/10.3760/j.issn:0412-4081.2005.11.007).
- Rikkens W, Lawrence D, Hafekost J, Zubrick SR. 2016.** Internet use and electronic gaming

- by children and adolescents with emotional and behavioural problems in Australia - results from the second child and adolescent survey of mental health and wellbeing. *BMC Public Health* **16**:399 DOI: [10.1186/s12889-016-3058-1](https://doi.org/10.1186/s12889-016-3058-1).
- Rosenfield M. 2011.** Computer vision syndrome: a review of ocular causes and potential treatments. *Ophthalmic Physiol Opt* **31**(5):502–515 DOI: [10.1111/j.1475-1313.2011.00834.x](https://doi.org/10.1111/j.1475-1313.2011.00834.x).
- Saito S, Sotoyama M, Saito S, Taptagaporn S. 1994.** Physiological indices of visual fatigue due to VDT operation: pupillary reflexes and accommodative responses. *Ind Health* **32**(2):57-66.
- Saqui N, Saqui J, Wahid A, Ahmed AA, Dhuhayr HE, Zaghloul MS, Ewid M, Al-Mazrou A. 2017.** Video game addiction and psychological distress among expatriate adolescents in Saudi Arabia. *Addict Behav Rep* **6**:112–217 DOI: [10.1016/j.abrep.2017.09.003](https://doi.org/10.1016/j.abrep.2017.09.003).
- Scheiman M, Galloway M, Frantz KA, Peters RJ, Hatch S, Cuff M, Mitchell GL. 2003.** Nearpoint of convergence: test procedure, target selection, and normative data. *Optom Vis Sci* **80**(3):214–225 DOI: [10.1097/00006324-200303000-00011](https://doi.org/10.1097/00006324-200303000-00011).
- Scheiman M, Wick B. 2002.** *Clinical Management of Binocular Vision*. Philadelphia: Lippincott Williams and Wilkins, 20.
- Sung W. 2014.** A study on the effect of the policy of online game shutdown on the game time of youth. *Social Science Research Review* **30**(2):233–256.
- Talwar R, Kapoor R, Puri K, Bansal K, Singh S. 2009.** A study of visual and musculoskeletal health disorders among computer professionals in NCR Delhi. *Indian J Community Med* **34**(4):326–328 DOI: [10.4103/0970-0218.58392](https://doi.org/10.4103/0970-0218.58392).
- The Korea Creative Content Agency. 2017.** Game users survey report 2017. Available at <http://www.kocca.kr/cop/bbs/view/B0000147/1833674.do?> (accessed 25 June 2018).
- Toomingas A. 2014.** A definition of visual ergonomics. *Appl Ergon* **45**(4):1263–1264 DOI: [10.1016/j.apergo.2014.03.004](https://doi.org/10.1016/j.apergo.2014.03.004).
- Tosha C, Borsting E, Ridder WH, Chase C. 2009.** Accommodation response and visual discomfort. *Ophthalmic Physiol Opt* **29**(6):625–633 DOI: [10.1111/j.1475-1313.2009.00687.x](https://doi.org/10.1111/j.1475-1313.2009.00687.x).
- Tsubota K, Nakamori K. 1993.** Dry eyes and video display terminals. *N Engl J Med* **328**(8):584 DOI: [10.1056/NEJM199302253280817](https://doi.org/10.1056/NEJM199302253280817).
- Tyrrell RA, Leibowitz HW. 1990.** The relation of vergence effort to reports of visual fatigue following prolonged near work. *Hum Factors* **32**(3):341–357 DOI: [10.1177/001872089003200307](https://doi.org/10.1177/001872089003200307).
- Weissberg EM. 2004.** Diagnostic techniques. In: Weissberg EW, eds. *Essential of clinical binocular vision*. St.Louis: Butterworth-Heinemann, 158–164.
- Wittek CT, Finserås TR, Pallesen S, Mentzoni RA, Hanss D, Griffiths MD, Molde H. 2016.** Prevalence and predictors of video game addiction: a study based on a national representative sample of gamers. *Int J Ment Health Addict* **14**(5):672–686 DOI: [10.1007/s11469-015-9592-8](https://doi.org/10.1007/s11469-015-9592-8).
- Wong EP, Fricke TR, Dinardo C. 2002.** Interexaminer repeatability of a new, modified prentice card compared with established phoria tests. *Optom Vis Sci* **79**(6):370–375 DOI: [10.1097/00006324-200206000-00010](https://doi.org/10.1097/00006324-200206000-00010).

- 400 **Yang SJ, Stewart R, Lee JY, Kim JM, Kim SW, Shin IS, Yoon JS. 2014.** Prevalence and
- 401 correlates of problematic internet experiences and computer-using time: a two-year
- 402 longitudinal study in Korean school children. *Psychiatry Investig* **11(1)**:24–31. DOI:
- 403 [10.4306/pi.2014.11.1.24](https://doi.org/10.4306/pi.2014.11.1.24).
- 404 **Yekta AA, Pickwell LD, Jenkins TC. 1989.** Binocular vision without visual stress. *Optom Vis*
- 405 *Sci* **66(12)**:815–817 DOI: [10.1097/00006324-198912000-00002](https://doi.org/10.1097/00006324-198912000-00002).
- 406 **Yeow PT, Taylor SP. 1991.** Effects of long-term visual display terminal usage on visual
- 407 functions. *Optom Vis Sci* **68(12)**:930–941 DOI: [10.1097/00006324-199112000-00004](https://doi.org/10.1097/00006324-199112000-00004).
- 408 **Yoo JS, Yoon JW, Kim JD. 1992.** Influence of VDT work on accommodative function. *J*
- 409 *Korean Ophthalmol Soc* **33(7)**:693-397.
- 410 **Zastrow M. 2017.** News feature: is video game addiction really an addiction? *Proc Natl Acad*
- 411 *Sci U S A* **114(17)**:4268–4272 DOI: [10.1073/pnas.1705077114](https://doi.org/10.1073/pnas.1705077114).
- 412

Table 1(on next page)

Changes in physical and ocular symptom scores after continuous computer gaming for 4 h.

1

Symptoms	Score, mean \pm SD		Significance*
	Before gaming	After gaming	
Physical discomfort	0.39 \pm 0.78	1.61 \pm 1.24	t = 15.7, p < 0.001
Shoulder	0.48 \pm 0.81	1.88 \pm 1.19	t = 8.80, p < 0.001
Neck	0.44 \pm 0.81	1.92 \pm 1.21	t = 9.78, p < 0.001
Back	0.20 \pm 0.53	1.16 \pm 1.15	t = 6.60, p < 0.001
Headache	0.42 \pm 0.91	1.46 \pm 1.30	t = 6.76, p < 0.001
Ocular discomfort	0.50 \pm 0.83	1.40 \pm 1.22	t = 17.44, p < 0.001
Red eyes	0.48 \pm 0.81	1.38 \pm 1.09	t = 7.18, p < 0.001
Eyestrain	0.44 \pm 0.79	1.58 \pm 1.03	t = 8.32, p < 0.001
Itchy eyes	0.44 \pm 0.79	0.82 \pm 1.00	t = 2.57, p = 0.013
Tired eyes	0.94 \pm 0.98	2.30 \pm 1.09	t = 8.73, p < 0.001
Dry eyes	0.64 \pm 1.01	1.64 \pm 1.34	t = 6.86, p < 0.001
Tearful eyes	0.34 \pm 0.80	0.88 \pm 1.14	t = 3.62, p < 0.001
Irritated eyes	0.34 \pm 0.66	0.96 \pm 1.12	t = 3.73, p < 0.001
Blurred vision	0.46 \pm 0.73	1.60 \pm 1.20	t = 7.28, p < 0.001
Aching eyes	0.40 \pm 0.76	1.40 \pm 1.29	t = 6.29, p < 0.001

2 Symptoms were scored on a scale from 0 (none) to 4 (very severe).

3 SD: standard deviation

4 *The p-values were determined using paired t-tests. Differences between the means were statistically significant (p < 0.05).

5

Table 2 (on next page)

Changes in binocular function after continuous computer gaming for 4 h.

1

Binocular function	Before gaming	After gaming	Following morning	Significance
NPC (cm)	7.23 ± 1.64	8.77 ± 1.60	7.36 ± 1.29	F = 47.83, p < 0.001
NPA (cm)	7.79 ± 1.44	9.11 ± 1.93	7.78 ± 1.49	F = 42.50, p < 0.001
Phoria at distance (Δ)	-0.76 ± 1.71	-0.83 ± 2.25	-0.62 ± 1.69	F = 1.67, p = 0.193
Phoria at near (Δ)	-3.73 ± 3.93	-5.75 ± 4.85	-3.68 ± 4.86	F = 18.39, p < 0.001
Accommodative facility (cpm)	14.42 ± 3.20	12.68 ± 3.98	15.54 ± 2.95	F = 30.08, p < 0.001
Vergence facility (cpm)	16.94 ± 3.16	15.02 ± 3.67	17.70 ± 3.41	F = 38.97, p < 0.001
Blink rate (per minute)	16.24 ± 5.14	(8.27 ± 5.40) [†]	(9.51 ± 5.28) [‡]	F = 106.29, p < 0.001

2 Data are expressed as means ± standard deviations.

3 NPC: near point of convergence, NPA: near point of accommodation, Δ: prism diopter, cpm: cycles per minute. [†]Blink rate
4 after 1 h, [‡]Blink rate after 4 h.

5 The p-values were determined using repeated measures analysis of variance. Negative and positive values for phoria
6 represent exophoria and esophoria (horizontal), respectively. Missing data are 15 persons for phoria and 10 persons for
7 blink rate.

8

Figure 1

Changes in phoria during a continuous 4-h computer gaming session.

The error bars represent standard deviations. Negative values denote exophoria and positive values denote esophoria (horizontal).

