

# Random allogeneic blood transfusion in pigs: characterisation of a novel experimental model

Alexander Ziebart<sup>Corresp., 1</sup>, Moritz Schaefer<sup>1</sup>, Rainer Thomas<sup>1</sup>, Jens Kamuf<sup>1</sup>, Andreas Garcia-Bardon<sup>1</sup>, Christian Moellmann<sup>1</sup>, Robert Rueemmler<sup>1</sup>, Florian Heid<sup>1</sup>, Arno Schad<sup>2</sup>, Erik K Hartmann<sup>1</sup>

<sup>1</sup> Department of Anesthesiology, Medical Centre of the Johannes Gutenberg-University, Mainz, Germany

<sup>2</sup> Institute of Pathology, Medical Centre of the Johannes Gutenberg-University, Mainz, Germany

Corresponding Author: Alexander Ziebart

Email address: alexander.ziebart@unimedizin-mainz.de

## Abstract:

**Background:** Organ cross-talk describes interactions between a primary affected organ and a secondarily injured remote organ, particularly in lung-brain interactions. A common theory is the systemic distribution of inflammatory mediators that are released by the affected organ and transferred through the bloodstream. The present study characterises the baseline immunogenic effects of a novel experimental model of random allogeneic blood transfusion in pigs designed to analyse the role of the bloodstream in organ cross-talk.

**Methods:** After approval of the State and Institutional Animal Care Committee 20 anesthetized pig were randomized in a donor and an acceptor (each n=8): the acceptor animals each received high-volume whole blood transfusion from the donor (35-40 ml kg<sup>-1</sup>). Four animals received balanced electrolyte solution instead of blood transfusion (control group; n=4). Afterwards the animals underwent extended cardiorespiratory monitoring for eight hours. Post mortem assessment included pulmonary, cerebral and systemic mediators of early inflammatory response (IL-6, TNF-alpha, iNOS), wet to dry ratio, and lung histology.

**Results:** No adverse events or incompatibilities occurred during the blood transfusion procedures. No hints for systemic inflammatory response were found and the pulmonary function was not affected through blood transfusion. Lung histopathology scoring did not display relevant intergroup differences. Neither within the lung nor within the brain an up-regulation of inflammatory mediators was detected. High volume random allogeneic blood transfusion in pigs neither impaired pulmonary integrity nor induced systemic, lung, or brain inflammatory response.

**Conclusion:** This approach can represent a novel experimental model to characterize the blood-bound transmission in remote organ injury.

**Random allogeneic blood transfusion in pigs: characterisation of a novel  
experimental model**

Alexander Ziebart<sup>1\*</sup>, Moritz Schaefer<sup>1</sup>, Rainer Thomas<sup>1</sup>, Jens Kamuf<sup>1</sup>, Andreas Garcia-Bardon<sup>1</sup>,  
Christian Moellmann, Robert Ruemmler<sup>1</sup>, Florian Heid<sup>1</sup>, Arno Schad<sup>2</sup>, Erik K. Hartmann<sup>1</sup>

<sup>1</sup> Department of Anesthesiology, Medical Centre of the Johannes Gutenberg-University, Mainz,  
Germany

<sup>2</sup> Institute of Pathology, Medical Centre of the Johannes Gutenberg-University, Mainz, Germany

**\*Corresponding author:**

alexander.ziebart@unimedizin-mainz.de

24

25 **Abstract:**

26 **Background:** Organ cross-talk describes interactions between a primary affected organ and a  
 27 secondarily injured remote organ, particularly in lung-brain interactions. A common theory is the  
 28 systemic distribution of inflammatory mediators that are released by the affected organ and  
 29 transferred through the bloodstream. The present study characterises the baseline immunogenic  
 30 effects of a novel experimental model of random allogeneic blood transfusion in pigs designed to  
 31 analyse the role of the bloodstream in organ cross-talk.

32 **Methods:** After approval of the State and Institutional Animal Care Committee 20 anesthetized  
 33 pigs were randomized in a donor and an acceptor group (each n=8): the acceptor animals each  
 34 received high-volume whole blood transfusion from the donors (35-40 ml kg<sup>-1</sup>). Four animals  
 35 received balanced electrolyte solution instead of blood transfusion (control group; n=4).  
 36 Afterwards the animals underwent extended cardiorespiratory monitoring for eight hours. Post  
 37 mortem assessment included pulmonary, cerebral and systemic mediators of early inflammatory  
 38 response (IL-6, TNF-alpha, iNOS), wet to dry ratio, and lung histology. **Results:** No adverse  
 39 events or incompatibilities occurred during the blood transfusion procedures. Systemic cytokine  
 40 levels and pulmonary function were unaffected. Lung histopathology scoring did not display  
 41 relevant intergroup differences. Neither within the lung nor within the brain an up-regulation of  
 42 inflammatory mediators was detected. High volume random allogeneic blood transfusion in pigs  
 43 neither impaired pulmonary integrity nor induced systemic, lung, or brain inflammatory response.  
 44 **Conclusion:** This approach can represent a novel experimental model to characterize the blood-  
 45 bound transmission in remote organ injury.

46

**Introduction:**

Single organ injury or failure does not exert an isolated effect on the primary affected organ system. Remote organs consecutively develop secondary injuries that are not connected to the initial insult. This is referred to as deleterious organ cross-talk and described particularly for lung and brain interactions, where this pathomechanism is well known, but not fully understood. E.g. mechanical ventilation in the intensive care unit may cause inflammatory response and induce damage to distant organs. On the other hand, severe brain injury is associated with an increased risk of lung failure (Mascia et al. 2007; Pelosi & Rocco 2011).

Different theories for this phenomenon consider neuronal response, hormone release or circulating inflammatory mediators as causative (Han 2017; Mrozek et al. 2015; Pelosi & Rocco 2011). The hypothesis of inflammatory mediator distribution is promising as several studies show inflammatory response after a primary hit. The blood stream seems to be the distributor in remote organ injury and cross-talk (Hegeman et al. 2009; Klein et al. 2016; Mrozek et al. 2015). In theory, inflammatory cytokines that are released into the systemic circulation induce remote organ injury. Hence, for experimental purposes this effect might be transferable from one individual to another via whole blood transfusion. Porcine models, in this context, offer advantages in experimental assessment of blood stream-related distribution of inflammatory mediator transmission. Firstly, higher blood volumes can be transferred in comparison to common laboratory rodent species. Secondly, because major inter-individual blood incompatibilities are missing (Smith et al. 2006). Only few reports reveal bronchospasm, disseminated intravascular coagulation, bleeding, or progressive hypotension during blood transfusion in pigs (Hunfeld et al. 1984; Sheil et al. 1972;

Smith et al. 2006). This porcine system is based on blood groups A0, whereas the human group B is missing. Additional 16 different antigens are known to form the subgroups A-P (Smith et al. 2006). Unlike the human Rhesus antigen, which is highly immunogenic and prone for transfusion reactions or hemolysis, the porcine Rhesus antigen is coded by a single gen without any polymorphisms and less pathologic relevance (Avent & Reid 2000; Omi et al. 2003). However, the distinct effects of porcine blood transfusion with focus on end-organ inflammatory response are barely examined. In order to further elucidate the role of the blood stream as distributor in remote organ injury the present study aims to characterize the baseline immunogenic effects of random allogeneic blood transfusion in pigs. The approach may allow for experimental separation of blood-bound effects on organ integrity from an already injured subject: hence, we hypothesized that single, large-volume blood transfusion from a healthy donor neither exhibits inflammatory response nor impairs the pulmonary function in the acceptor. As secondary hypotheses, the systemic and cerebral inflammatory response is analyzed.

## **Materials & Methods**

The prospective-randomized animal study was approved by the State and Institutional Animal Care Committee (Landesuntersuchungsamt Rheinland-Pfalz, Koblenz, Germany; approval number: 23 177-07/G 14-1-084). The study was conducted in accordance with the ARRIVE guidelines and the international guidelines for the care and use of laboratory animals. The experiments were performed from 12/2014 till 3/2015. Twenty juvenile German Landrace pigs (*sus scrofa domestica*, weight  $28 \pm 2$  kg) were examined. Eight animals served as blood donors. Eight animals received allogenic blood transfusion. The control group consisted of four animals that received the same amount of balanced electrolyte solution instead of blood transfusion. One

acceptor animal was excluded due to complicated airway management and multiple intubation attempts with subsequent respiratory failure before the transfusion. Per protocol and approval, the donor animals that inevitably suffered from hemorrhagic shock were subsequently included in another study protocol focusing on fluid resuscitation. All experiments were conducted under continuous general anesthesia. This manuscript adheres to the applicable EQUATOR guidelines.

# **Anesthesia and instrumentation**

Anesthesia and instrumentation procedure was previously described in detail in our former studies (Ziebart et al. 2015; Ziebart et al. 2018): Briefly, in each case two animals received an intramuscular injection of ketamine (8mg kg<sup>-1</sup>) and midazolam (0.2 mg kg<sup>-1</sup>) followed by induction of general anesthesia via an ear vein cannula (propofol 4 mg kg<sup>-1</sup>, and fentanyl 4 µg kg<sup>-1</sup>, atracurium 0.5 mg kg<sup>-1</sup>), which was maintained by continuous infusion (propofol 8-12 mg kg<sup>-1</sup> h<sup>-1</sup>, fentanyl 0.1-0.2 mg h<sup>-1</sup>). After orotracheal intubation the animals were ventilated in volume-controlled mode (VCV; AVEA, CareFusion, USA): positive end-expiratory pressure (PEEP) 5 cmH<sub>2</sub>O, tidal volume 7 ml kg<sup>-1</sup>, inspiration to expiration ratio 1:2, fraction of inspired oxygen (FiO<sub>2</sub>) 0.4 and variable respiration rate to achieve an end-tidal CO<sub>2</sub> < 45 mmHg. In case of respiratory insufficiency, mechanical ventilation parameters escalated according to an adaption of the ARDS network tables. (Ziebart et al. 2014) Afterwards, the animals were randomized into an acceptor and a donor animal. Ultrasound guided femoral vascular access under sterile conditions included a central venous line, a pulse contour cardiac output catheter (PiCCO, Pulsion Medical, Germany), and a pulmonary arterial catheter in the acceptor animals. The donor animals received a central venous catheter, an arterial line, a pulse contour cardiac output system (PiCCO, Pulsion Medical Systems, Germany) and a large-bore venous introducer. Extended hemodynamics,

transpulmonary thermodilution-derived parameters, spirometry, and gas exchange were continuously monitored (Datex S/5 Monitor, GE Healthcare, Germany; PiCCO, Pulsion Medical Systems, Germany). Hematological and blood gas measurements were repetitively conducted. Electrical impedance tomography (EIT; Goe-MF II, CareFusion, Germany) was used to measure variations of the thoracic bioimpedance that are associated with pulmonary aeration. Sixteen adhesive electrodes were placed around the thorax and the ventilation distribution was measured for three lung compartments (Level 1-3; non-dependent, central, dependent) (Bodenstein et al. 2012; Ziebart et al. 2014). Multiple breath-nitrogen washout/washin technique was applied to measure the functional residual capacity (Kamuf et al. 2017). Body temperature was measured by a rectal probe and a surface warming device prevented hypothermia. All animals received a background infusion of 5 ml kg<sup>-1</sup> h<sup>-1</sup> balanced electrolyte solution (Sterofundin, B.Braun, Germany). After anesthesia, instrumentation, and 20-minute recovery period baseline values were measured and recorded.

### **Blood withdrawal procedure**

From the donor animals 35-40 ml kg<sup>-1</sup> blood was taken for allogenic blood transfusion through the arterial catheter. The whole blood was collected in specific collector bags system (Composelect, Fresenius-Kabi AG, Homburg Germany). These bags comprised 63 ml CPD/100 ml SAG-M - RCC (Citrate phosphate dextrose, Erythrocyte storage in hypertonic conservation medium). After collection in a primary bag. The blood was drained through a leucocyte-depleting filter into a second bag that contains the anticoagulants. Finally, the whole blood was collected into a bag with a connector for a transfusion system. Afterwards the whole blood was stored by room temperature

for about ten minutes before transfusion to avoid negative effects of long-time storing or cooling. Afterwards the donor animals were transferred to the fluid resuscitation protocol that was included in the same approval: they were euthanized by intravenous injection of propofol (200mg) and potassium chloride (40mval) within the scope of this protocol. (Ziebart et al. 2018)

# **Experimental protocol:**

The experimental protocol is summarized in figure 1. Immediately after blood withdrawal the whole blood was transfused into the recipients through the venous vascular access sheath over 15 minutes through a transfusion system with an integrated aggregate-filter. A control group (n=4) underwent the identical instrumentation and protocol, though received an equal amount of balanced electrolyte solution instead of allogeneic blood transfusion. Afterwards the animals were monitored for eight hours. In case of hemodynamic deterioration (mean arterial pressure < 60 mmHg) continuous noradrenaline infusion was administered. After finishing the protocol, the animals were killed in deep general anesthesia by injection of propofol (200 mg) and potassium chloride (40 mval).

# **Post mortem analysis**

Repetitively taken serum blood samples were used to determine the serum levels of TNF-alpha and IL-6 by enzyme-linked immunosorbent assays (Porcine Quantikine ELISA Kits, R/D Systems, Germany). After finishing the protocol, the lung and the brain each were removed en-bloc. Four representative tissue samples (dependent periphery, dependent central, non-dependent central, non-dependent periphery) from the right lung were extracted, formalin-fixed, and used for histological evaluation of alveolar damage in the upper and the lower lung. After paraffin



embedding and hematoxylin/eosin staining the samples were analyzed in investigator-blinded manner by means of standardized scoring scheme (Ziebart et al. 2015; Ziebart et al. 2014). Additional lung samples were cryopreserved for mRNA expression analysis of inflammatory mediators (IL-6, TNF-alpha, iNOS) using real-time polymerase chain reaction (rt-PCR; Lightcycler 480 PCR System, Roche Applied Science, Germany) as described in detail by Hartmann et al. (Hartmann et al. 2015). The mRNA expression was normalized to peptidylprolyl isomerase A (PPIA). The same technique and markers were used to quantify the cerebral mRNA expression from the right hippocampus and the right frontal cortex. Furthermore, the lung and brain water content were determined through the tissue wet to dry ratio. For this procedure the weight of the removed organs was determined directly and after two days of complete drying.

## Statistics

All parameters are presented mean and standard deviation ( $\pm$  SD) or displayed as plots. The analysis focusses on the relevant time points baseline, 0h, 4h 8h. Additional data are added as supplemental files. Group effects over time (Transfusion, Control) were compared by two-way analysis of variance (ANOVA) with post-hoc Student-Newman-Keuls-Test. Mann-Whitney-U-Test was used to analyze the post mortem parameters. A p-value lower than 0.05 was accepted as significant. The software package SigmaPlot 12.5 (Systat Software, USA) was used.

## Results

The estimated total blood volume of the pigs ( $74 \text{ ml kg}^{-1}$ ) was  $2084 \pm 166 \text{ ml}$  (Lyhs & Wachtel 1966). The transfused blood volume was  $48 \pm 4\%$ . No adverse events or major incompatibilities

occurred during the blood transfusion procedures. Beneath a temporary increase of the mean pulmonary arterial pressure following blood transfusion and a slight increase of the central venous pressure the cardiopulmonary monitoring and blood gas analyses showed no relevant differences between the transfusion and control animals over eight hours (table 1, 2 & 3). The lactate level was in a normal range and even decreased within eight hours. Through blood transfusion hemoglobin, hematocrit and erythrocyte counts significantly increased without relevant changes in leucocyte amounts (figure 2). Marker of systemic inflammatory response did not differ between the groups as well (figure 2 & 3). The pulmonary function as measured by blood gas analysis and ventilation distribution was not affected by blood transfusion (table 3 & figure 4). Only the alveolar-arterial oxygen difference increased. At baseline two animal started at a  $\text{PaO}_2/\text{FiO}_2 < 400$  mmHg without signs of local or systemic illness. A modification of the initial respiratory setting due to hypoxemia was not required in any animal. Lung histopathology scoring did not display relevant intergroup differences (control vs. transfusion;  $10 \pm 1$  vs.  $11 \pm 3$ ; each  $p > 0.05$ ). The tissue wet to dry ratio as surrogate of edema formation between both groups was not increased following transfusion (control vs. transfusion; lung:  $4.7 \pm 0.2$  vs.  $4.6 \pm 0.2$ ; brain:  $3.3 \pm 0.3$  vs.  $3.4 \pm 0.3$ ; each  $p > 0.05$ ). Neither within the lung nor within the brain the mRNA expression analysis revealed an up-regulation of inflammatory markers (figure 5 & 6).

## **Discussion**

The present study establishes a novel model for characterization of remote organ injury related to the distribution of inflammatory mediators through the systemic circulation. We found that one-time high-volume allogeneic blood transfusion in healthy pigs neither impairs pulmonary integrity nor mediates inflammatory response within the lung, brain, or systemic circulation. Accordingly, random allogeneic blood transfusion in pigs is feasible, and per itself does not trigger inflammatory

210 response in the acceptor animals. Hence, this model could be an appropriate approach to identify  
 211 the main mediators in injurious organ interactions, i.e. by transfusion of blood from severely brain  
 212 or lung injured subjects to healthy ones without co-founders. Other experimental protocols with  
 213 comparable porcine transfusion models investigate questions like acute kidney injury or effects of  
 214 blood storing and cooling. Several similar but, in some points, different protocols have been  
 215 described that make use of autologous instead homologous blood, heavier animals to minimize the  
 216 need for donor animals, or previously induce a hemorrhagic shock to reduce the rate of circulatory  
 217 overload (Biagini et al. 2017; Masuda et al. 2018; Patel et al. 2011; Wozniak et al. 2018).

218 In principle, blood transfusion can be associated with severe side effects. Beneath  
 219 incompatibilities, particularly pulmonary complications like transfusion-related acute lung injury  
 220 (TRALI) or transfusion-associated circulatory overload (TACO) are described. Other systemic  
 221 side effects include hypotension, anaphylaxis, hemolytic reactions and transfusion-transmitted  
 222 bacterial infections (Bux & Sachs 2008). Our study reveals a significant increase of mean arterial,  
 223 pulmonary arterial and central venous pressure immediately following transfusion, which was  
 224 underlined by increase of hemoglobin and hematocrit. These symptoms comply with subclinical  
 225 TACO induced by supraphysiological increase of intravascular volume (Roubinian et al. 2017).

226 Transfusion of longer stored but not freshly drawn blood was shown to considerably increase  
 227 pulmonary arterial pressure blood, which is presumably an effect of mediators released through  
 228 storage. (Baron et al. 2012; Solomon et al. 2013) In our study, similar but short-termed effects  
 229 occurred despite blood transfusion immediately after withdrawal. Furthermore, fluid overload,  
 230 which inevitably also occurs in our study, is also associated with pulmonary arterial hypertension.  
 231 (Yilmaz et al. 2016) (Wolsk et al. 2019) As we found no coincident pathologies that allow

distinguishing between fluid overload and mediator-based effects, identification the exact mechanism is beyond the scope of this study.

Relevant impairment of the pulmonary gas exchange or alterations in extravascular lung water content, functional residual capacity or ventilation distribution were not observed after blood transfusion. Over eight hours  $\text{PaO}_2/\text{FiO}_2$  decreases to a certain degree going along with a raised alveolar-arterial oxygen difference. In the absence of histopathologic lung injury and during low-grade invasive mechanical ventilation, this is most likely a consequence of atelectasis formation that inevitably occurs during general anesthesia, muscular paralysis and mechanical ventilation.(Hedenstierna & Edmark 2015) Two animals had a slightly reduced  $\text{PaO}_2/\text{FiO}_2$  ratio ( $< 400$  mmHg) already during baseline measurements without any signs for pulmonary or systemic illness: these two animals did neither respond differently to the transfusion nor showed signs of increased histopathologic lung damage despite formally exhibiting a  $\text{PaO}_2/\text{FiO}_2 < 300$  mmHg after eight hours. Increase and withdrawal of intravascular fluid volume in pigs was shown to alter the endexpiratory lung volume as measured by EIT, but did not affect tidal volume or ventilation distribution, which is in line with prior results (Bodenstein et al. 2012). TRALI represents another facet that directly affects the lung following blood transfusion. It is characterized by respiratory distress, hypoxemia, and non-cardiogenic lung edema that is induced by leukocyte-antibodies and neutrophil-priming substances (Bux & Sachs 2008). These interact with neutrophil granulocytes, which thereby are activated. The resulting clots impair pulmonary capillary perfusion and induce capillary leakage by releasing toxic enzymes and reactive oxygen species (Sachs et al. 2006; Toy et al. 2005). We did not find any TRALI surrogates such as edema formation or increasing counts of leucocytes or specifically neutrophil granulocytes in our porcine model. Furthermore, the histopathological analysis did not reveal relevant inflammatory cell immigration. Additionally, the

regional tidal volume distribution was unaltered, which is described as sign of severe pulmonary illnesses (Chiumello et al. 2013). Gender-specific forms of TRALI are particularly relevant in multiparous women as blood donors (Palfi et al. 2001) (Schmickl et al. 2015). Hence, the exclusive use of male pigs may represent an approach to minimize the TRALI risk. TACO on the other hand tends to appear primarily in patients suffering from serious pre-existing illnesses. In our study healthy animals with intact pulmonary vascular barrier and competent kidney function were used, which enables rapid compensation of the intravascular overload (Bux & Sachs 2008; Roubinian et al. 2017).

Single organ injury or failure can subsequently affect the integrity of distant remote organs leading to multiple organ failure. In particular, deleterious interactions of the lung and the brain have been described (Pelosi & Rocco 2011). Several models suggest that the lung may play a pivotal role as causative but also remotely injured organ: in mechanically ventilated rats that underwent an aggressive ventilator regime showed remarkable cerebral inflammatory response (Quilez et al. 2011). Cyclic variations of the  $\text{PaO}_2$  caused by non-protective ventilation in lung injured pigs are transmitted to the cerebral tissue and may trigger cerebral injury (Klein et al. 2013; Klein et al. 2016). Furthermore, also a cerebral insult is capable to induce or aggravate lung injury (Heuer et al. 2011). Even mechanical ventilation by itself can represent a trigger for cerebral cytokine expression. (Kamuf et al. 2018) Accordingly, we also assessed brain inflammatory response measured by cortical and hippocampal mRNA expression of key inflammatory mediators, which also were unaffected by the transfusion procedure. In comparison to human blood the use of porcine model may offer several advantages for studies concerning remote organ injury: Unlike human blood, where transfusion of incompatible blood products is deleterious, pig blood lacks the human blood group B, the secondary absorption of antigens from the plasma to the blood cells and

the Rh polymorphism (Omi et al. 2003). Additionally human blood is polygene coded, which results in a much higher incompatibility potential (Smith et al. 2006).

The study has several limitations. The study was not designed to assess long-term effects of blood transfusion but to exclude acute side effects of allogeneic blood transfusion itself with regard to a novel experimental approach to elucidate pathomechanisms of deleterious remote organ injury. Logistically it was not possible to investigate different pig subspecies or breeds. Hence, generalization of the reported findings lies beyond the scope of this study. Animal care reasons and the need for additional blood donor animal required a strict limitation of the animal numbers, whereas the chosen group numbers were in line with previous studies inflammatory acute effects in pigs (Hartmann et al. 2015; Ziebart et al. 2014). Exchange of the transfused blood by simultaneous blood withdrawal may be a possibility to avoid fluid overloading. However, given the risk of unpredictable removal of the donor blood, hemodynamic variations, and potential microcirculatory or inflammatory impairment during this complex procedure, we decided to decline this option. Small volume withdrawal prior to the transfusion procedure may represent a reasonable compromise, and may reduce the risk of TACO (Masuda et al. 2018). The use of plasma instead of whole blood is an interesting alternative approach to optimize this setting: theoretically, plasma administration will not increase the hemoglobin content, whereas full concentration of potentially inflammatory mediators should be present. This approach, however, was not feasible for the present study. Additionally, we did not measure cytokine contents in the transfused blood, as the donor animals were healthy and the blood was transfused within minutes. These measurements need to be included in studies focusing on severely injured donor animals.

## **Conclusion**

In conclusion, high volume random allogeneic blood transfusion in pigs neither impaired pulmonary integrity nor induced systemic, lung, or brain inflammatory response. No further side effects beyond expectable and temporary alterations of hemodynamics and blood count related to the rapid increase of blood volume were detected. Random allogeneic blood transfusion in pigs may represent an experimental model to characterize the blood-bound transmission in remote organ injury.

### **Acknowledgements:**

The authors thank Dagmar Dirvonskis and Dana Pieter for excellent technical support. Parts of this study are included in the doctoral thesis of Moritz Schaefer, and content of the professorial dissertation (habilitation) of Alexander Ziebart. Preliminary results of the study were presented in poster sessions at the German Anesthesia Congress (DAC), Nuernberg, Germany in 2017.

### **References**

- Avent ND, and Reid ME. 2000. The Rh blood group system: a review. *Blood* 95:375-387.
- Baron DM, Yu B, Lei C, Bagchi A, Beloiartsev A, Stowell CP, Steinbicker AU, Malhotra R, Bloch KD, and Zapol WM. 2012. Pulmonary hypertension in lambs transfused with stored blood is prevented by breathing nitric oxide. *Anesthesiology* 116:637-647. 10.1097/ALN.0b013e318246ef77
- Biagini S, Dale CS, Real JM, Moreira ES, Carvalho CRR, Schettino GPP, Wendel S, and Azevedo LCP. 2017. Short-term effects of stored homologous red blood cell transfusion on cardiorespiratory function and inflammation: an experimental study in a hypovolemia model. *Brazilian Journal of Medical and Biological Research* 51:e6258. 10.1590/1414-431x20176258
- Bodenstein M, Wang H, Boehme S, Vogt A, Kwiecien R, David M, and Markstaller K. 2012. Influence of crystalloid and colloid fluid infusion and blood withdrawal on pulmonary bioimpedance in an animal model of mechanical ventilation. *Physiol Meas* 33:1225-1236. 10.1088/0967-3334/33/7/1225
- Bux J, and Sachs UJ. 2008. Pulmonary transfusion reactions. *Transfus Med Hemother* 35:337-345. 10.1159/000151349
- Chiumello D, Froio S, Bouhemad B, Camporota L, and Coppola S. 2013. Clinical review: Lung imaging in acute respiratory distress syndrome patients--an update. *Crit Care* 17:243. 10.1186/cc13114
- Han DW. 2017. Brain and lung: dangerous crosstalk. *Korean J Anesthesiol* 70:116-117. 10.4097/kjae.2017.70.2.116

- Hartmann EK, Ziebart A, Thomas R, Liu T, Schad A, Tews M, Moosmann B, Kamuf J, Duenges B, Thal SC, and David M. 2015. Inhalation therapy with the synthetic TIP-like peptide AP318 attenuates pulmonary inflammation in a porcine sepsis model. *BMC Pulm Med* 15:7. 10.1186/s12890-015-0002-6
- Hedenstierna G, and Edmark L. 2015. Effects of anesthesia on the respiratory system. *Best Pract Res Clin Anaesthesiol* 29:273-284. 10.1016/j.bpa.2015.08.008
- Hegeman MA, Hennis MP, Heijnen CJ, Specht PA, Lachmann B, Jansen NJ, van Vught AJ, and Cobelens PM. 2009. Ventilator-induced endothelial activation and inflammation in the lung and distal organs. *Crit Care* 13:R182. 10.1186/cc8168
- Heuer JF, Pelosi P, Hermann P, Perske C, Crozier TA, Bruck W, and Quintel M. 2011. Acute effects of intracranial hypertension and ARDS on pulmonary and neuronal damage: a randomized experimental study in pigs. *Intensive Care Med* 37:1182-1191. 10.1007/s00134-011-2232-2
- Hunfeld MA, Hoitsma HF, Meijer S, van Haeringen H, and Rietveld FW. 1984. The role of A-O-incompatible blood transfusions in porcine orthotopic liver transplantations. *Eur Surg Res* 16:354-359.
- Kamuf J, Garcia-Bardon A, Duenges B, Liu T, Jahn-Eimermacher A, Heid F, David M, and Hartmann EK. 2017. Endexpiratory lung volume measurement correlates with the ventilation/perfusion mismatch in lung injured pigs. *Respir Res* 18:101. 10.1186/s12931-017-0585-y
- Kamuf J, Garcia-Bardon A, Ziebart A, Thomas R, Folkert K, Frauenknecht K, Thal SC, and Hartmann EK. 2018. Lung injury does not aggravate mechanical ventilation-induced early cerebral inflammation or apoptosis in an animal model. *PLoS One* 13:e0202131. 10.1371/journal.pone.0202131
- Klein KU, Boehme S, Hartmann EK, Szczyrba M, Heylen L, Liu T, David M, Werner C, Markstaller K, and Engelhard K. 2013. Transmission of arterial oxygen partial pressure oscillations to the cerebral microcirculation in a porcine model of acute lung injury caused by cyclic recruitment and derecruitment. *Br J Anaesth* 110:266-273. 10.1093/bja/aes376
- Klein KU, Johannes A, Bruckner M, Thomas R, Matthews S, Frauenknecht K, Leukel P, Mazur J, Poplawski A, Muellenbach R, Sommer CJ, Thal SC, and Engelhard K. 2016. Systemic PaO2 Oscillations Cause Mild Brain Injury in a Pig Model. *Crit Care Med* 44:e253-263. 10.1097/ccm.0000000000001399
- Lyhs L, and Wachtel W. 1966. [On the effect of physical stress on various circulatory functions in the pig]. *Arch Exp Veterinarmed* 20:1057-1063.
- Mascia L, Zavala E, Bosma K, Pasero D, Decaroli D, Andrews P, Isnardi D, Davi A, Arguis MJ, Berardino M, and Ducati A. 2007. High tidal volume is associated with the development of acute lung injury after severe brain injury: an international observational study. *Crit Care Med* 35:1815-1820. 10.1097/01.Ccm.0000275269.77467.Df
- Masuda R, Iijima T, Kondo R, Itoda Y, Matsuhashi M, Hashimoto S, Kohira T, Kobayashi N, and Okazaki H. 2018. Preceding haemorrhagic shock as a detrimental risk factor for respiratory distress after excessive allogeneic blood transfusion. *Vox Sang* 113:51-59. 10.1111/vox.12560
- Mrozek S, Constantin JM, and Geeraerts T. 2015. Brain-lung crosstalk: Implications for neurocritical care patients. *World J Crit Care Med* 4:163-178. 10.5492/wjccm.v4.i3.163
- Omi T, Vogeli P, Hagger C, Schelling C, Spilar S, Kajii E, Stranzinger G, and Neuenschwander S. 2003. cDNA cloning, mapping and polymorphism of the porcine Rhesus (RH) gene. *Anim Genet* 34:176-182.
- Palfi M, Berg S, Ernerudh J, and Berlin G. 2001. A randomized controlled trial of transfusion-related acute lung injury: is plasma from multiparous blood donors dangerous? *Transfusion* 41:317-322.
- Patel NN, Lin H, Toth T, Welsh GI, Jones C, Ray P, Satchell SC, Sleeman P, Angelini GD, and Murphy GJ. 2011. Reversal of anemia with allogenic RBC transfusion prevents post-cardiopulmonary bypass



acute kidney injury in swine. *American Journal of Physiology: Renal Physiology* 301:F605-614. 10.1152/ajprenal.00145.2011

Pelosi P, and Rocco PR. 2011. The lung and the brain: a dangerous cross-talk. *Crit Care* 15:168. 10.1186/cc10259

Quilez ME, Fuster G, Villar J, Flores C, Marti-Sistac O, Blanch L, and Lopez-Aguilar J. 2011. Injurious mechanical ventilation affects neuronal activation in ventilated rats. *Crit Care* 15:R124. 10.1186/cc10230

Roubinian NH, Hendrickson JE, Triulzi DJ, Gottschall JL, Chowdhury D, Kor DJ, Looney MR, Matthay MA, Kleinman SH, Brambilla D, and Murphy EL. 2017. Incidence and clinical characteristics of transfusion-associated circulatory overload using an active surveillance algorithm. *Vox Sang* 112:56-63. 10.1111/vox.12466

Sachs UJ, Hattar K, Weissmann N, Bohle RM, Weiss T, Sibelius U, and Bux J. 2006. Antibody-induced neutrophil activation as a trigger for transfusion-related acute lung injury in an ex vivo rat lung model. *Blood* 107:1217-1219. 10.1182/blood-2005-04-1744

Schmickl CN, Mastrobuoni S, Filippidis FT, Shah S, Radic J, Murad MH, Toy P, and Gajic O. 2015. Male-predominant plasma transfusion strategy for preventing transfusion-related acute lung injury: a systematic review. *Crit Care Med* 43:205-225. 10.1097/ccm.0000000000000675

Sheil AG, Halliday JP, Drummond JM, Bookallil MJ, Gaudry PL, and Yezerski SD. 1972. A modified technique for orthotopic liver transplantation. *Arch Surg* 104:720-724.

Smith DM, Newhouse M, Naziruddin B, and Kresie L. 2006. Blood groups and transfusions in pigs. *Xenotransplantation* 13:186-194. 10.1111/j.1399-3089.2006.00299.x

Solomon SB, Wang D, Sun J, Kanas T, Feng J, Helms CC, Solomon MA, Alimchandani M, Quezado M, Gladwin MT, Kim-Shapiro DB, Klein HG, and Natanson C. 2013. Mortality increases after massive exchange transfusion with older stored blood in canines with experimental pneumonia. *Blood* 121:1663-1672. 10.1182/blood-2012-10-462945

Toy P, Popovsky MA, Abraham E, Ambruso DR, Holness LG, Kopko PM, McFarland JG, Nathens AB, Silliman CC, and Stronck D. 2005. Transfusion-related acute lung injury: definition and review. *Crit Care Med* 33:721-726.

Wolsk E, Bakkestrom R, Kristensen CB, Aagaard Myhr K, Thomsen JH, Balling L, Andersen MJ, Dahl JS, Shah SJ, Gustafsson F, Hassager C, and Moller JE. 2019. Right Ventricular and Pulmonary Vascular Function are Influenced by Age and Volume Expansion in Healthy Humans. *J Card Fail* 25:51-59. 10.1016/j.cardfail.2018.11.013

Wozniak MJ, Qureshi S, Sullo N, Dott W, Cardigan R, Wiltshire M, Nath M, Patel NN, Kumar T, Goodall AH, and Murphy GJ. 2018. A Comparison of Red Cell Rejuvenation versus Mechanical Washing for the Prevention of Transfusion-associated Organ Injury in Swine. *Anesthesiology* 128:375-385. 10.1097/aln.0000000000001973

Yilmaz S, Yildirim Y, Taylan M, Demir M, Yilmaz Z, Kara AV, Aydin F, Sen HS, Karabulut A, and Topcu F. 2016. The Relationship of Fluid Overload as Assessed by Bioelectrical Impedance Analysis with Pulmonary Arterial Hypertension in Hemodialysis Patients. *Med Sci Monit* 22:488-494. 10.12659/msm.896305

Ziebart A, Garcia-Bardon A, Kamuf J, Thomas R, Liu T, Schad A, Duenges B, David M, and Hartmann EK. 2015. Pulmonary effects of expiratory-assisted small-lumen ventilation during upper airway obstruction in pigs. *Anaesthesia* 70:1171-1179. 10.1111/anae.13154

Ziebart A, Hartmann EK, Thomas R, Liu T, Duenges B, Schad A, Bodenstein M, Thal SC, and David M. 2014. Low tidal volume pressure support versus controlled ventilation in early experimental sepsis in pigs. *Respir Res* 15:101. 10.1186/s12931-014-0101-6

Ziebart A, Möllmann C, Garcia-Bardon A, Kamuf J, Schäfer M, Thomas R, and Hartmann EK. 2018. Effect of gelatin-polysuccinat on cerebral oxygenation and microcirculation in a porcine haemorrhagic


428 shock model. *Scandinavian Journal Trauma Resuscitation Emergency Medicin* 26:15.  
 429 10.1186/s13049-018-0477-2

430

# Figure 1

*Figure 1: Experimental flow chart*


After preparation and baseline measurement the animals were observed for eight hours. The hemodynamic and spirometric parameters were recorded continuously. Blood samples were taken every two hours. EIT (electrical impedance tomography) was measured hourly. Histopathological analysis of the lung was performed after the end of the experiment. Pulmonary and cerebral mRNA expression was quantified. (Transfusion n=7; Control n=4)



# Figure 2

*Figure 2: Hematological parameters.*


Parameters are displayed as percentage of baseline. A significant increase of hemoglobin, hematocrit and erythrocytes was observed after transfusion. The decrease of these parameters in the control group was not significant; \* =  $p < 0.05$ ; vs. baseline. (Transfusion  $n=7$ ; Control  $n=4$ )



# Figure 3

*Figure 3: Systemic inflammatory cytokines levels.*


Parameters are displayed are presented as pg/mL. The systemic inflammatory cytokines level did not differ between both study groups, but there is a significant decrease of TNF-alpha to baseline in both groups; \* =  $p < 0.05$ ; vs. baseline. (Transfusion n=7; Control n=4)



# Figure 4

*Figure 4: Regional distribution of the tidal volume.*

Non-dependent, central and dependent lung areas (% of the global tidal amplitude). No significant differences (each  $p > 0.05$ ). The regional distribution of the tidal volume did not differ between both study groups. (Transfusion  $n=7$ ; Control  $n=4$ )




# Figure 5

*Figure 5: Pulmonary mRNA expression of inflammatory mediators.*

Cryopreserved lung samples were collected after the experiment from upper and lower lung sections. mRNA expression of inflammatory parameters (IL-6, TNF-alpha, INOS) were analyzed via real-time polymerase chain reaction. mRNA expression is normalized to peptidylprolyl isomerase A (PPIA); no significant intergroups differences (each  $p > 0.05$ ). (Transfusion  $n=7$ ; Control  $n=4$ ).






# Figure 6

*Figure 6: Cerebral mRNA expression of inflammatory mediators.*

Cryopreserved cerebral samples were collected after the experiment from cortex and hippocampus regions. mRNA expression of inflammatory parameters (IL-6, TNF-alpha, INOS) were analyzed via real-time polymerase chain reaction. mRNA expression is normalized to peptidylprolyl isomerase A (PPIA); No significant intergroups differences (each  $p > 0.05$ ). (Transfusion  $n=7$ ; Control  $n=4$ ).



# **Table 1**(on next page)

## *Table 1: Hemodynamic parameters*

No significant intra- and intergroup changes of hemodynamic parameters were observed after blood transfusion or fluid infusion, with the exception of a temporary pulmonary arterial pressure increase and raise of the central venous and mean arterial pressure after transfusion / fluid administartion. (Transfusion n=7; Control n=4)

1

Parameter		BLH	0h	4h	8h
		MEAN (SD)	MEAN (SD)	MEAN (SD)	MEAN (SD)
MAP [mmHg]	Transfusion	70 (13)	98* (14)	82 (9)	81 (14)
	Control	77 (12)	93* (3)	73 (7)	71 (5)
HR [min <sup>-1</sup> ]	Transfusion	83 (19)	83 (13)	84 (20)	85 (13)
	Control	78 (9)	78 (10)	80 (7)	73 (6)
PAP [mmHg]	Transfusion	18 (5)	27 (6)*#	20 (2)	19 (3)
	Control	16 (4)	19 (3)	17 (3)	12 (7)
P <sub>es</sub> [mmHg]	Transfusion	5 (1)	7 (2)	4 (1)	3 (2)
	Control	6 (0)	6 (1)	4 (1)	4 (0)
CO [l min <sup>-1</sup> ]	Transfusion	3.6 (0.7)	3.7 (0.7)	3.2 (0.3)	3.2 (0.6)
	Control	3.4 (0.3)	4.3 (0.9)	3.4 (0.3)	2.9 (0.4)
GEDV [ml kg <sup>-1</sup> ]	Transfusion	820 (125)	850 (96)	778 (107)	812 (81)
	Control	820 (97)	916 (20)	808 (62)	803 (75)
EVLWI [ml kg <sup>-1</sup> ]	Transfusion	11 (2)	12 (2)	12 (2)	12 (3)
	Control	13 (5)	11 (0)	15 (3)	13 (2)
CVP [mmHg]	Transfusion	5 (2)	9 (2)*	7 (2)*	7 (2)*
	Control	7 (1)	8 (2)	7 (1)	7 (1)
S <sub>p</sub> O <sub>2</sub> [%]	Transfusion	100 (0)	100 (0)	99 (1)	99 (1)
	Control	100 (0)	100 (0)	100 (0)	100 (0)

\*indicates p<0.05 vs. Baseline value. # indicates p< 0.05 in intergroup comparison

MAP: mean arterial pressure; HR: heart rate;  
PAP: mean arterial pulmonary pressure; P<sub>es</sub>: esophageal pressure;  
CO: cardiac output; GEDV: global enddiastolic volume index;  
EVLWI: extravascular lung water index; CVP: central venous pressure;  
S<sub>p</sub>O<sub>2</sub>: oxygen saturation



## Table 2 (on next page)

### *Table 2: Spirometric parameters*

No significant intra- and intergroup changes of spirometric parameters were observed after blood transfusion or fluid infusion. (Transfusion n=7; Control n=4)

Parameter		BLH	0h	4h	8h
		MEAN (SD)	MEAN (SD)	MEAN (SD)	MEAN (SD)
F <sub>I</sub> O <sub>2</sub> [%]	Transfusion	39 (0)	39 (0)	39 (0)	39 (0)
	Control	39 (0)	39 (0)	39 (0)	39 (0)
MV [l/min]	Transfusion	6.2 (0.7)	6.1 (0.8)	6.2 (1)	6.6 (1.3)
	Control	6.4 (1.7)	5.8 (1.3)	6.5 (1.6)	7 (1)
TV [ml kg <sup>-1</sup> ]	Transfusion	8 (0)	8 (0)	8 (0)	8 (0)
	Control	8 (1)	8 (1)	8 (1)	8 (1)
P <sub>peak</sub> [cm H <sub>2</sub> O]	Transfusion	17 (2)	18 (2)	20 (2)	20 (2)
	Control	16 (2)	17 (1)	17 (3)	19 (3)
P <sub>mean</sub> [cm H <sub>2</sub> O]	Transfusion	9 (1)	9 (0)	10 (1)	10 (1)
	Control	9 (1)	8 (0)	9 (1)	10 (1)
PEEP [cm H <sub>2</sub> O]	Transfusion	4 (0)	4 (0)	4 (0)	4 (0)
	Control	4 (0)	4 (0)	4 (0)	4 (0)
C <sub>p</sub> [ml/cmH <sub>2</sub> O]	Transfusion	19 (3)	17 (2)	16 (2)	16 (3)
	Control	20 (3)	20 (2)	19 (3)	17 (3)
R <sub>AW</sub> [kPA/l/sec]	Transfusion	10 (1)	11 (1)	12 (1)	12 (1)
	Control	11 (1)	11 (1)	11 (1)	11 (2)
FRC [ml]	Transfusion	728 (206)	691 (260)	691 (240)	714 (235)
	Control	823 (369)	610 (87)	674 (77)	788 (351)

\*indicates p<0.05 vs. Baseline value. # indicates p< 0.05 in intergroup comparison

F<sub>I</sub>O<sub>2</sub>: fraction of inspired oxygen; MV: minute volume; TV: tidal volume;  
P<sub>peak</sub>: peak inspiratory pressure; P<sub>mean</sub>: mean airway pressure;  
PEEP: positive end-expiratory pressure C<sub>p</sub>: pulmonary compliance;  
R<sub>AW</sub>: airway resistance; FRC: functional residual capacity






# **Table 3**(on next page)

## *Table 3: Blood gas parameter*

No relevant intra- and intergroup changes of blood gas parameters were observed after blood transfusion or fluid infusion. Only the lactate levels decrease within the normal range and the AaDO<sub>2</sub> increases in both groups. (Transfusion n=7; Control n=4)

Parameter		BLH	0h	4h	8h
		MEAN (SD)	MEAN (SD)	MEAN (SD)	MEAN (SD)
S <sub>v</sub> O <sub>2</sub> [%]	Transfusion	60 (12)	66 (8)	58 (6)	56 (6)
	Control	63 (13)	64 (0)	63 (11)	53 (8)
pH	Transfusion	7.47 (0.028)	7.505 (0.027)	7.503 (0.017)	7.495 (0.019)
	Control	7.485 (0.02)	7.5 (0.033)	7.545 (0.054)	7.506 (0.069)
BE [mmol/ml]	Transfusion	4.5 (4)	7.3 (1.4)	8.5 (2.2)	6.6 (3.6)
	Control	5.9 (4.5)	6.1 (0)	8.9 (2.4)	6.4 (3.4)
a <sub>rt</sub> CO <sub>2</sub> [mmHg]	Transfusion	40 (4)	40 (4)	41 (3)	40 (5)
	Control	39 (5)	41 (4)	38 (6)	38 (4)
e <sub>t</sub> CO <sub>2</sub> [mmHg]	Transfusion	37 (3)	38 (3)	40 (1)	39 (3)
	Control	35 (5)	35 (4)	34 (5)	35 (4)
avCO <sub>2</sub> diff [mmHg]	Transfusion	9 (3)	8 (4)	8 (2)	5 (6)
	Control	10 (5)	-8 (21)	7 (1)	8 (1)
PaO <sub>2</sub> [mmHg]	Transfusion	185 (34)	181 (23)	154 (33)	141 (26)
	Control	206 (43)	230 (49)	186 (44)	171 (24)
PaO <sub>2</sub> /F <sub>i</sub> O <sub>2</sub> [mmHg]	Transfusion	464 (86)	452 (58)	386 (81)	352 (64)
	Control	514 (108)	576 (122)	464 (109)	428 (60)
AaDO <sub>2</sub> [mmHg]	Transfusion	53 (30)	58 (22)	82 (32)*	97 (27)*
	Control	33 (44)	7 (46)	55 (45)	69 (27)*
Temp. [°C]	Transfusion	37.1 (0.8)	36.9 (1.2)	38 (0.8)	38.5 (0.4)
	Control	35.8 (2.9)	35.4 (3.9)	37.6 (2.2)	38.3 (0.6)
Potassium [mmol/l]	Transfusion	4 (0.3)	3.6 (0.7)	4.2 (0.7)	3.9 (0.6)
	Control	4 (0.6)	4 (0.1)	4.5 (0.3)	4.2 (0.2)
Lactate [mmol/l]	Transfusion	1.4 (0.5)	1.1 (0.4)	0.7 (0.2)*	0.6 (0.2)*
	Control	1.4 (0.4)	1.3 (0.1)	0.9 (0.2)*	0.8 (0.3)*



$S_vO_2$ : central venous oxygen saturation, BE: base excess;  
 $art.CO_2$ : arterial carbon dioxide;  $ex.CO_2$ : expiratory carbon dioxide;  
 $avCO_2$  diff: arterial venous carbon dioxide difference;  
 $PaO_2$ : arterial oxygen;  $PaO_2/F_iO_2$ : Horovitz index;  
 $AaDO_2$ : alveolar-arterial oxygen difference; Temp.: Temperature