

The effect of elevational gradient on alpine gingers (*Roscoeia alpina* and *R. purpurea*) in the Himalayas

Babu Ram Paudel^{Corresp., 1, 2, 3}, Adrian G Dyer⁴, Jair E Garcia⁴, Mani Shrestha⁴

¹ Yunnan Key Laboratory of Plant Reproductive Adaption and Evolutionary Ecology, Yunnan University, Kunming, Yunnan, China

² Laboratory of Ecology and Evolutionary Biology, State Key Laboratory for Conservation and Utilization of Bio-Resources in Yunnan, Yunnan University, Kunming, Yunnan, China

³ Department of Botany, Prithvi Narayan Campus, Tribhuvan University, Pokhara, Gandaki, Nepal

⁴ School of Media and Communication, RMIT University, Melbourne, Victoria, Australia

Corresponding Author: Babu Ram Paudel
Email address: babu@xtbg.ac.cn

There is currently enormous interest in how morphological and physiological responses of herbaceous plants may be affected by changing elevational gradient. Mountain regions provide an excellent opportunity to understand how closely related species may adapt to the conditions that rapidly change with elevation. We investigated the morphological and physiological responses of two Himalayan alpine gingers (*Roscoeia alpina* and *R. purpurea*) along two different vertical transects of 400 m, *R. purpurea* between 2174- 2574 m a.s.l and *R. alpina* between 2675-3079 m a.s.l . We measured the variables of plant height, leaf length, leaf area, specific leaf area, and stomata density at five plots, along the vertical transect at an elevational gap of ca. 100 m. Results revealed that with increased elevation plant height, and leaf area decreased while stomata density increased, whereas changes in specific leaf area, were not correlated with the elevation. Our results reveal that these alpine gingers undergo local adaptation by modifying their plant height, leaf area and stomata density in response to the varying selection pressure associated with the elevational gradient. Thus, the findings of this research provide valuable information on how a narrow range of elevational gradient affects the herbaceous plants at the alpine habitat of the Himalayas.

RESEARCH ARTICLE

The effect of elevational gradient affect on alpine gingers (*Roscoea alpina* and *R. purpurea*) in the Himalayas

Babu Ram Paudel^{1,2,3*}, Adrian G. Dyer⁴, Jair E. Garcia⁴, and Mani Shrestha⁴

¹Yunnan Key Laboratory of Plant Reproductive Adaption and Evolutionary Ecology, Yunnan University, Kunming, China.

² Laboratory of Ecology and Evolutionary Biology, State Key Laboratory for Conservation and Utilization of Bio-Resources in Yunnan, Yunnan University, Kunming, 650091, Yunnan, P.R. China

³Department of Botany, Prithvi Narayan Campus, Tribhuvan University, Pokhara, Nepal

⁴ School of Media and Communication, RMIT University, Melbourne, Victoria 3001, Australia

*For correspondence: brp2033@gmail.com

Abstract

There is currently enormous interest in how morphological and physiological responses of herbaceous plants may be affected by changing elevational gradient. Mountain regions provide an excellent opportunity to understand how closely related species may adapt to the conditions that rapidly change with elevation. We investigated the morphological and physiological responses of two Himalayan alpine gingers (*Roscoea alpina* and *R. purpurea*) along two different vertical transects of 400 m, *R. purpurea* between 2174- 2574 m a.s.l and *R. alpina* between 2675-3079 m a.s.l. We measured the variables of plant height, leaf length, leaf area, specific leaf area, and stomata density at five plots, along the vertical transect at an elevational gap of ca. 100 m. Results revealed that with increased elevation plant height, and leaf area decreased while stomata density increased, whereas changes in specific leaf area, were not correlated with the elevation. Our results reveal that these alpine gingers undergo local adaptation by modifying their plant height, leaf area and stomata density in response to the varying selection pressure associated with the elevational gradient. Thus, the findings of this research provide valuable information on how a narrow range of elevational gradient affects the herbaceous plants at the alpine habitat of the Himalayas.

Short title: Response of alpine gingers with the elevational gradient in the Himalayas

47 Introduction

48 The elevational gradient is one of the key environmental factors that affect growth,
 49 morphology and physiology of plants (Cordell et al., 1998; Hultine & Marshall, 2000; Qiang et
 50 al., 2003). The elevational gradient in alpine regions provides a sharp environmental change across
 51 relatively short spatial distances because small changes in elevation can lead to a large shift in
 52 temperature, humidity, exposure, and concentration of atmospheric gases (Hovenden & Vander
 53 Schoor, 2004). Thus, alpine environments can provide useful natural avenues to investigate the
 54 response of plants to a suite of climatic conditions that are representative of the broader latitudinal
 55 range (Montesinos-Navarro et al., 2011). With the increase in elevation, there is typically an
 56 increase in both precipitation and light intensity including changes in distributions of short
 57 wavelength UV-A (315-400 nm) and UV-B (280-315 nm) radiation (Diffey, 1991; Rozema et al.,
 58 1997) whilst temperature and concentration of carbon dioxide and oxygen decrease (Friend &
 59 Woodward, 1990). These environmental variations may potentially alter the morphology and
 60 physiology of plants to endure the different stresses linked with changing elevation (Hovenden &
 61 Brodribb, 2000; Körner, 2007).

62 The alpine environment is potentially affected by climate change associated with global
 63 warming, and thus alpine plants may face rapidly changing environmental conditions that likely
 64 impose different stress levels on plants (Beniston, 2003; Byars, Papst & Hoffmann, 2007). Thus,
 65 based on the adaptative plasticity, the plant species exhibit local adaptation by altering the
 66 morphological and/or physiological traits over the range of elevational gradient (Hirano,
 67 Sakaguchi & Takahashi, 2017). For example, local adaptations of plants in response to variable
 68 climatic conditions at different elevations may result in variation of plant height and leaf length
 69 (Wang & Gao, 2004). Variation in carbon assimilation, energy balance and water relations along

the elevational gradient could result in variation of leaf morphological and physiological traits such as leaf area, specific leaf area (SLA) and stomata density (Ackerly et al., 2002). Therefore, the study of the variation in the growth forms, morphology and physiology of a plant species along an elevational gradient could provide valuable insights on how plants may respond to environmental stress imposed by rapid changes in climatic conditions (Premoli & Brewer, 2007; Körner, 2007; Bresson et al., 2011).

Although several previous studies have documented the effects of elevational gradient on the growth, morphology and physiology of the plants, most of the studies are focused on tree species (Cordell et al., 1998; Hultine & Marshall, 2000; Li et al., 2008). Recently a few studies have been conducted to understand how the elevational gradients affect the herbaceous plants (Gonzalo-Turpin & Hazard, 2009; Scheepens, Frei & Stöcklin, 2010; Hulshof et al., 2013; Bastida, Rey & Alcántara, 2015; Takahashi & Matsuki, 2017; Kiełtyk, 2018). These studies particularly focused on the variation of a specific trait, such as vegetative trait, reproductive trait or leaf trait. Currently however, there is a lack of empirical evidence on the adaptative potential of herbaceous plants along the elevational gradient in steep environments such as the Himalayas. As the alpine ecosystem in the Himalayas is likely to experience the adverse effects of the changing climate associated with global warming and anthropogenic disturbances (Beniston, 2003; Byars, Papst & Hoffmann, 2007), understanding the performance of herbaceous plants along the elevational gradient provides important insights for the enhanced prediction of the response of herbaceous plants under altered climatic conditions.

The genus *Roscoea*, with 22 known species, is a Himalayan endemic alpine perennial herb and the only alpine member of the predominately-tropical family Zingiberaceae (Cowley, 1982, 2007). The genus is distributed between the elevations of ca 1500 to 4500 m a.s.l (Cowley, 2007),

thus serving as a key model for how herbaceous plants respond to the potentially stressful environmental conditions associated with increasing elevation. All *Roscoea* species are small herbs with annual leafy shoots produced from a reduced erect rhizome (Cowley, 1982, 2007). Among the *Roscoea* species, *R. alpina* Royle and *R. purpurea* Smith are widely distributed in the Himalayan Mountains from Kashmir (Pakistan) in the west through Nepal, India, Bhutan and Tibet. As these two *Roscoea* species are widely distributed from low to high elevations, characterization of the variation in morphological and physiological traits along the elevational gradient will help to understand how these alpine gingers respond to changes in climatic conditions associated with elevation. In this study, we explore the changes in the morphological (plant height and leaf length) and physiological variables (leaf area, SLA and stomata density) of these alpine gingers along the well-defined elevational gradient in the Himalayan mountain range.

Materials and Methods

Study species

The two widespread *Roscoea* species used in this study were *R. alpina* and *R. purpurea* (Fig. 1). *Roscoea alpina* is a common species with a wide distribution between the elevations 2130-4270 m a.s.l in the Himalayan range from Kashmir (Pakistan) in the west through Bhutan in the east. The annual pseudostem may grow up to 12-20 cm high and presents flowers from the end of May to mid-August (Cowley, 2007). It has 2-3 obtuse sheathing leaves. Leaves are usually 1-2 in number and underdeveloped; occasionally the plant may bear up to four well-developed leaves. Leaves are linear, broadly elliptic or lanceolate. Only the first leaf is slightly auriculate and widest at the base while rest of the leaves are widest at the middle, with 17- 25 cm in length. Leaves are usually glabrous but young leaves are occasionally hairy at acute apex. Inflorescences are without exserted peduncle. Flowers are deep purple to white in appearance for a human observer (Fig. 1).

A single plant can develop up to five flowers, however only one flower blooms at a time. Obtuse to almost truncate bracts are shorter than the ovary. The calyx is much longer than the bract and bluntly bi-dentate. A long corolla tube is exerted from the calyx (Cowley, 2007).

Roscoea purpurea is also a widespread member of the Himalayan *Roscoea*, distributed between the elevations 1520-3100 m a.s.l. from Himachal Pradesh (India) in the west through Assam/Bhutan in the east. The annual erect pseudostem is most variable in habit and form and may grow up to 25-38 cm high, bearing 0-2 obtuse to truncate sheathing leaves. Leaves are usually 4-8 in number, lanceolate to oblong-ovate and 14-20 cm long with acuminate and sometimes with ciliated apex. Lower leaves are slightly auriculate at the base. The plant flowers from the end of June to early September (Cowley, 2007). The inflorescence is enclosed in upper leaf sheaths with only the upper part of bracts and flowers visible. Flowers are light purple or white with purple markings. Usually, 1-2 flowers open at a time. Bracts longer than calyx with acute apex which is pale green. The sharply bi-dentate and apiculate calyx is usually pale green and sometimes marked with pink. The corolla tube has a mauve or white colouration and is hardly exerted from the calyx (Cowley, 2007).

Study sites

The research was conducted along an elevational gradient at two sites, Daman and Ghorepani, Central Nepal (Fig. 2). Daman is located in Makawanpur district and forms a part of the Mahabharat mountain range (mountains lower than the Himalayas). This site lies about 70 km south-west of Kathmandu and is midway between Kathmandu and Hetauda. The vegetation type of this site typically comprises a mixed forest of *Pinus*, *Rhododendron* and *Quercus*. The site experiences cool temperate to subalpine climate with warm summers and cold winters that typically incur mild to heavy snowfall from November to February (Pers. Obs. BRP). Ghorepani,

located in Myagdi district, is about 270 km west of Kathmandu. The vegetation type of this site comprises a mixed forest of *Pinus* and *Rhododendron*. The site has a subalpine climate and cool weather throughout the year, and heavy snowfall from November to February (Pers. Obs. BRP). The geographical coordinates and the elevations of the study sites are presented in Table 1.

Measurement of traits

The field sampling was conducted from May to August 2014 and repeated the sampling again in 2017 (May to August). Five sampling plots were selected along a vertical transect from 2174 to 2574 m for *R. purpurea* and from 2675 to 3079 m for *R. alpina*. The sampling was done in a counterbalanced random fashion such that two adjacent sampling plots were at an elevation gap of ca 100 m. Plant height and leaf length were measured to examine the morphological variables. Physiological variables included leaf area, specific leaf area (SLA) and stomata density. At each sampling plot, a horizontal transect of 100 m length was laid down and twenty plants were randomly selected along the horizontal transect in such a way that the distance between the adjacent sampling plant was at least 5 m. A standard metric ruler was used to measure plant height (the distance from the ground to the topmost part of the stem). The largest leaf of every sampled plant was removed and leaf length was measured with a ruler. We used a graph paper to trace and quantify the area of each leaf, enabling robust repeatable measurements in remote locations. Specifically, two alternative methods were used to measure the area of the leaf. In 2014, the area of the leaf was measured after wet storage, while in 2017 the area was measured on the freshly plucked leaf. To prevent the leaves from possible shrinkage during wet storage, the leaves were first flattened if necessary and carefully placed in between the folds of a paper. Resulting samples were then placed in a sample box to avoid the external light and heat sources. For both respective leaf collection methods, the leaf (either wet stored or freshly plucked) was placed on a graph paper,

its outline was sketched and the number of squares enclosed within the leaf-outline were counted. Complete and greater than half squares were scored, whilst squares less than half a square were excluded. The measurements were repeated several times for each leaf to enable a robust field measurement of leaf area. Twenty leaves at each sampling plot were measured to assess variability. The area of leaf as measured by two alternative methods did not differ significantly (t test, $P>0.05$), thus data generated from the freshly removed leaf were used for further analysis. All collected leaves were gently pressed between the folds of an absorbent paper for five days to flatten the leaf surface and to absorb any excess moisture. The pressed leaves were subsequently oven dried at the university laboratory for 48 hours at 70⁰ C. Dry leaf weight was measured using a digital electronic balance (Fameway International (HK) Limited; accuracy 0.001g). Specific leaf area (SLA) of a leaf was calculated as the ratio of the area of a fresh leaf and its dry weight and expressed in cm²/g.

To determine the stomatal count, transparent nail polish was applied on the middle dorsal surface of a fresh leaf. After a few minutes, when nail polish had dried, a thin layer was peeled from the middle dorsal surface of a leaf. The peeled layers were separately preserved in a 10% glycerine solution for about 72 hours. In the laboratory, the temporary slide of each layer was prepared using safranin as a staining agent. The stained layers were individually mounted on microscope slides, and all stomata observed under a 10-x magnification microscopic field were counted. The stomata counts were repeated at three different microscopic fields to ensure the exact measurement of the stomata density. Area of the microscopic field was calculated using the formula $A = \pi r^2$ where r is the radius of microscopic field and density of stomata was calculated as the number of stomata under a microscopic field divided by the area of the microscopic field. The stomata density was expressed in terms of number per square millimetre.

Statistical Analyses

An independent sample t test was used to test the variation in measured traits between the years. Data from each of the measured morphological variables were summarized as Q-Q plots and tested for normality. Exploratory data analyses revealed that some of the response variables were not normally distributed and were better described by a Gamma distribution as most data consisted of positive values larger than zero (Zuur, Hilbe & Ieno, 2013). Consequently, non-parametric correlation analyses were performed among the five different traits measured for each species implementing Kendall's tau statistic (τ). This coefficient was chosen as it has a known standard error and provides a better estimate with low sample size. After the exploratory analyses, generalised linear regression models (GLM) were applied to test for the potential effects of elevation on the different traits measured for each species. For the five regression models, elevation was used as a predictor and it was assumed that the response variable followed a Gamma distribution. Link function for each model was selected based on a comparison of AIC scores obtained after fitting models implementing different link functions (Zuur, Hilbe & Ieno, 2013). Regression analyses were performed using the routine *glm* available as part of the base distribution of the R language and environment for statistical computing (version 3.3.1) (R Core Team, 2015).

Results

Correlation analyses

Our results indicated that all the measured variables did not differ significantly between years ($P > 0.05$), thus only 2017 data were used for further analyses. For *R. purpurea*, there was a significant correlation between stomata density and the variables of leaf area ($P < 0.001$) and specific leaf area (SLA) ($P = 0.020$) (Fig. 3). In *R. alpina*, leaf length was correlated with all

remaining variables (Fig. 4). Consequently, we separately performed the regression analyses for the two species for each of the measured response variables.

Variations of traits with elevation

Leaf length of *R. alpina* and plant height significantly decreased with increasing elevation ($P = 0.001$ for leaf length and $P = 0.017$ for plant height). The same trend was observed for leaf area ($P < 0.001$), while stomata density increased with elevation ($P = 0.005$). SLA values for this species were not significantly correlated with elevation ($P = 0.114$) (Fig. 5).

Plant height and leaf area significantly decreased, while stomata density increased, with increasing elevation in *R. purpurea* ($P = 0.044$, $P = 0.001$ and, $P = 0.002$ for plant height, leaf area and stomata density respectively). However, we did not find a significant relationship of elevation either with leaf length ($P = 0.471$) or with SLA ($P = 0.555$) (Fig. 5). Details on the regression analysis including coefficients and associated 95% confidence intervals are provided in Table 2 and Table 3.

Discussion

Variations of morphological traits with elevation

In the current study, we found a significant decrease in plant height of both species of *Roscoea* (*R. alpina* and *R. purpurea*) with increased elevation. Reduction of plant height in these alpine gingers with increased elevation is consistent with several previous findings reported for tree species (Körner, 1998; Cordell et al., 1998; Kronfus & Havranek, 1999; Paulsen, Weber & Körner, 2000; Kogami et al., 2001; Li, Yang & Kräuchi, 2003; Shi et al., 2006) and herbaceous species (Takahashi & Matsuki, 2017; Kiełtyk, 2018). Similarly, a decrease of leaf length of *R. alpina* with the increased elevation in the current study is consistent with the previous findings

(Hansen-Bristow, 1986; Schoettle, 1990; Kajimoto, 1993; Kao & Chang, 2001; Kieltyk, 2018). Based on the present result, we conclude that the elevational gradient has a significant effect on the growth form of these alpine gingers. At the lower elevation, environmental conditions are likely to be more favourable for optimum plant growth. The reduction of plant height and leaf length of these two alpine gingers with increasing elevation reflects the morphological adaptation to increased environmental stresses such as low concentration of carbon dioxide, decreased temperature, higher solar radiation and/or low water availability (Wang & Gao, 2004; Davis, Shaw & Etterson, 2005; Guerin, Wen & Lowe, 2012). The observed relatively smaller plants with shorter leaf characteristics of these gingers at higher elevation thus may reflect local adaptation at a higher altitude to enable reduction of transpiration and maintain efficient utilization of water (Ackerly et al., 2002; Royer et al., 2008; Peppe et al., 2011; Guerin, Wen & Lowe, 2012).

Variation of physiological traits with Elevation

Leaf traits variation with elevation:

Our findings revealed that variation in leaf area showed a significant but negative correlation with elevation, while the correlation between SLA and elevation was non-significant. Consistent with our result, Kouwenberg, Kurschner and McElwain (2007) found a decreasing trend in the leaf area of *Quercus kelloggii* with increasing elevation. Our result on the variation of leaf characters (leaf area and SLA) with the elevation is partially consistent with the previous findings reported by Hultine and Marshall (2000); Scheepens, Frei and Stöcklin (2010); Hulshof et al., (2013); Bastida, Rey and Alcántara, (2015), while the findings of Gonzalo-Turpin and Hazard (2009) indicate a different effect. Previous studies have suggested that the environment at higher elevations is characterized by higher solar radiation, lower water availability and lower stomatal conductance (Parkhurst & Loucks, 1972; Givnish & Vermeij, 1976; Ackerly et al., 2002).

253 Under such potentially stressful environmental conditions, small leaf size provides optimum
 254 adaptation to the plants by reducing boundary layer resistance and maintaining favorable leaf
 255 temperature and high photosynthetic water use efficiency (Renzhong et al., 2001). Thus, decreased
 256 leaf area of these alpine gingers with increased elevation may reflect an adaptation for the
 257 increased environmental stress and may be favourable to reduce water loss and maintain efficient
 258 use of absorbed water (Renzhong et al., 2001). In addition, some authors have implicated
 259 increasing UVB radiation levels as having a damaging effect on certain plant structures (Jansen,
 260 Gaba & Greenberg, 1998; Rozema, Aerts & Cornelissen, 2002); and there is some evidence of this
 261 affecting plant growth in some lowland terrestrial species (Rozema, Aerts & Cornelissen, 2002).
 262 These topics may be of high value to explore in alpine environments where there are likely large
 263 changes in UV levels. SLA is closely associated with leaf thickness, which mediates the trade-off
 264 between light capture, water loss and diffusion of carbon dioxide (Oberle & Schaal, 2011). Higher
 265 SLA leaves are thicker and contain more photosynthetic enzymes and there is more demand for
 266 carbon dioxide per unit area. Thus, stomata density increases to supply the higher demand for
 267 carbon dioxide. Consequently, the increase in SLA may be an advantage for carbon dioxide uptake.
 268 Non-linear change of SLA of both species along the elevational gradient may indicate that
 269 environmental factors associated with altitude alone cannot regulate the trade –off between light
 270 capture, water loss and diffusion of carbon dioxide in these alpine gingers. The smallest SLA of
 271 *R. alpina* at 2674 m a.s.l and *R. purpurea* at 2374 m a.s.l may indicate limited carbon gain and
 272 supply due to poor availability of resources and may be associated with the least productive zone
 273 of these species where retention of captured resources and protection from desiccation is of high
 274 priority (Wilson, Thompson & Hodgson, 1999).

Variation in Stomata density with Elevation

We found a significant increase in stomata density of both species (*R. alpina* and *R. purpurea*) with increased elevation. Many authors have made comprehensive efforts to relate the variation in stomata density along elevation gradients and have obtained different results. Consistent to our current results, Körner and Cochrane (1985); Friend and Woodward (1990); Hovenden and Brodribb (2000); and Kouwenberg, Kurschner and McElwain (2007) have found that stomata density increased linearly with elevation. Li et al. (2006) found that stomata density of *Quercus aquifolioides* increased linearly up to the height of 2800 m a.s.l., whilst above that height, it decreased linearly. Schoettle and Rochelle (2000) found that the stomata density of *Pinus flexilis* decreased linearly with altitude whilst Woodward (1986) did not observe any significant change in stomata density of *Vaccinium myrtillis* considering altitudes from 200 to 1100 m asl. The significant increase in stomata density with increasing elevation in our findings may be associated with lower availability of carbon dioxide, higher UV-B and long wave radiation, all reducing photosynthetic efficiency by decreasing stomatal absorption and conductance (Kouwenberg, Kurschner & McElwain, 2007; Körner, 2007). To adapt to such a harsh environmental conditions and maintain vitalities, stomata density of these gingers may have increased. The increase in stomata density provides compensation against the reduced stomatal conductance and carbon dioxide partial pressure to maintain photosynthetic efficiency (Kao & Chang, 2001; Kouwenberg, Kurschner & McElwain, 2007; Körner, 2007).

Our results indicate two major patterns in the vegetative traits of these alpine gingers with increased elevation: a significant decrease of leaf area and a significant increase of stomata density. These variations provide compensation to cope with the change in the concentration of atmospheric carbon dioxide, temperature, humidity and light at higher altitudes (Van de Water,

Leavitt & Betancourt, 1994; Hultine & Marshall, 2000; Qiang et al., 2003). A non-significant correlation between SLA and stomata density may suggest that leaf thickness have little role in regulating the carbon dioxide uptake and transpiration in these two alpine gingers. A negative correlation of stomata density with leaf area has previously indicated that with the increase of stomata density at a higher elevation, narrowing of leaves may reduce excess transpiration (Hermes & Mattson, 1992; Etterson & Shaw, 2001). The closely correlated variation in these two traits thus maintains a likely trade-off between photosynthesis and transpiration and provides local adaptation to the specific conditions, at different elevations.

Conclusions

Growth, morphology and physiology of *R. alpina* and *R. purpurea* were found to have a significant association with altitude. These alpine gingers exhibit optimum growth at their respective lowermost distribution range, and their growth response retards with increasing elevation. Based on the present result, it can be concluded that these alpine gingers favour shorter height, smaller leaf and higher stomata density at a higher elevation to adapt with the stressful factors associated with the change in elevational gradients. Variation in those traits at different elevations may reflect the response to the combined selection pressure of different abiotic and biotic factors that may generate different micro-environmental conditions at the respective elevation. Decreased growth forms and leaf area of these alpine gingers at a higher altitude may indicate a selection response to reduce water loss from the plant body during transpiration while increased stomata density may indicate the adaptation to cope with the decreased concentration of carbon dioxide. The closely correlated modification of these traits at different elevations may have played a significant role in providing local adaptation to these alpine gingers.

Acknowledgements

We are thankful to the Department of Plant Resources, Thapathali, Kathmandu for providing research permission. We thank Mr Kul Prasad Lamichhane, Kapil Paudel, and Dipesh Baral for assistance in the field. We thank Dr Subodh Adhikari for the constructive comments on the earlier version of the manuscript. We thank Dr Lalina Muir for proof reading the manuscript

References

- Ackerly DD, Knight CA, Weiss SB, Barton K, Starmer KP. 2002. Leaf size, specific leaf area and microhabitat distribution of chaparral woody plants: contrasting patterns in species level and community level analyses. *Oecologia* 130:449–457. DOI: 10.1007/s004420100805.
- Bastida JM, Rey PJ, Alcántara JM. 2015. Local adaptation to distinct elevational cores contributes to current elevational divergence of two *Aquilegia vulgaris* subspecies. *Journal of Plant Ecology* 8:273–283. DOI: 10.1093/jpe/rtu017.
- Beniston M. 2003. Climatic change in Mountain regions: a review of possible impacts. *Climate change* 59:5–31.
- Bresson CC, Vitasse Y, Kremer A, Delzon S. 2011. To what extent is altitudinal variation of functional traits driven by genetic adaptation in European oak and beech? *Tree physiology* 31:1164–74. DOI: 10.1093/treephys/tpr084.
- Byars SG, Papst W, Hoffmann A a. 2007. Local adaptation and cogradient selection in the alpine plant, *Poa hiemata*, along a narrow altitudinal gradient. *Evolution; international journal of organic evolution* 61:2925–41. DOI: 10.1111/j.1558-5646.2007.00248.x.
- Cordell S, Goldstein G, Mueller-Dombois D, Webb D, Vitousek PM. 1998. Physiological and morphological variation in *Metrosideros polymorpha*, a dominant Hawaiian tree species,

along an altitudinal gradient: the role of phenotypic plasticity. *Oecologia* 113:188–196.
DOI: 10.1007/s004420050367.

Cowley EJ. 1982. A revision of *Roscoe* (Zingiberaceae). *Kew Bulletin* 36:747–777.

Cowley EJ. 2007. *The genus Roscoe*. Royal Botanic Garden Kew, UK.

Davis MB, Shaw RG, Etterson JR. 2005. Evolutionary responses to changing climate. *Ecology* 86:1704–1714.

Diffey BL. 1991. Solar ultraviolet radiation effects on biological systems. *Physics in Medicine and Biology* 36:299–328.

Etterson JR, Shaw RG. 2001. Constraint to aAdaptive evolution in response to global warming. *Science* 294:151–154. DOI: 10.1126/science.1063656.

Friend AD, Woodward FI. 1990. Evolutionary and ecophysiological responses of mountain plants to the growing season environment. *Advances in Ecological Research* 20:59–124.

Givnish TJ, Vermeij GJ. 1976. Sizes and shapes of liane leaves. *The American naturalist* 110:743–778.

Gonzalo-Turpin H, Hazard L. 2009. Local adaptation occurs along altitudinal gradient despite the existence of gene flow in the alpine plant species *Festuca eskia*. *Journal of Ecology* 97:742–751. DOI: 10.1111/j.1365-2745.2009.01509.x.

Guerin GR, Wen H, Lowe AJ. 2012. Leaf morphology shift linked to climate change. *Biology Lett.* 1–5.

Hansen-Bristow K. 1986. Influence of increasing elevation on growth characteristics at timberline. *Canadian Journal of Botany* 64:2517–2523.

Herms DA, Mattson WJ. 1992. The dilemma of plants-to grow or defend. *Q. Rev. Biol.* 67:283–335.

- Hirano M, Sakaguchi S, Takahashi K. 2017. Phenotypic differentiation of the *Solidago virgaurea* complex along an elevational gradient: insights from a common garden experiment and population genetics. *Ecology and Evolution* 7:6949–6962. DOI: 10.1002/ece3.3252.
- Hovenden MJ, Brodribb T. 2000. Altitude of origin influences stomatal conductance and therefore maximum assimilation rate in Southern Beech, *Nothofagus cunninghamii*. *Aust. J. Plant Physiol.* 27:451–456.
- Hovenden MJ, Vander Schoor JK. 2004. Nature vs nurture in the leaf morphology of Southern beech, *Nothofagus cunninghamii* (Nothofagaceae). *New phytologist* 161:585–594.
- Hulshof CM, Violle C, Spasojevic MJ, McGill B, Damschen E, Harrison S, Enquist BJ. 2013. Intra-specific and inter-specific variation in specific leaf area reveals the importance of abiotic and biotic drivers of species diversity across elevation and latitude. *Journal of Vegetation Science* 24:921–931. DOI: 10.1111/jvs.12041.
- Hultine KR, Marshall JD. 2000. Altitude trends in conifer leaf morphology and stable carbon isotope composition. *Oecologia* 123:32–40. DOI: 10.1007/s004420050986.
- Jansen MAK, Gaba V, Greenberg B. 1998. Higher plants and UV-B radiation: balancing damage, repair and acclimation. *Trends in Plant Science* 3:131–135. DOI: [http://dx.doi.org/10.1016/S1360-1385\(98\)01215-1](http://dx.doi.org/10.1016/S1360-1385(98)01215-1).
- Kajimoto T. 1993. Shoot dynamics of *Pinus pumila* in relation to altitudinal and wind exposure gradients on the Kiso mountain range, central Japan. *Tree physiology* 13:41–53.
- Kao WY, Chang KW. 2001. Altitudinal trends in photosynthetic rate and leaf characteristics of *Miscanthus populations* from central Taiwan. *Aust. J. Bot.* 49:509–514.
- Kieltyk P. 2018. Variation of vegetative and floral traits in the alpine plant *Solidago minuta*: evidence for local optimum along an elevational gradient. *Alpine Botany* 128:47–57. DOI:

10.1007/s00035-017-0197-7.

Kogami H, Hanba YT, Kibe T, Terashima I, Masuzawa T. 2001. CO₂ transfer conductance, leaf structure and carbon isotope composition of *Polygonum cuspidatum* leaves from low and high altitudes. *Plant, Cell and Environment* 24:529–538. DOI: 10.1046/j.1365-3040.2001.00696.x.

Körner C. 1998. A re-assessment of high elevation treeline positions and their explanation. *Oecologia* 115:445–459.

Körner C. 2007. The use of “altitude” in ecological research. *Trends in ecology & evolution* 22:569–74. DOI: 10.1016/j.tree.2007.09.006.

Körner C, Cochrane PM. 1985. Stomatal responses and water relations of *Eucalyptus pauciflora* in summer along an elevational gradient. *Oecologia* 66:443–455.

Kouwenberg LLR, Kurschner WM, McElwain JC. 2007. Stomatal frequency change over altitudinal gradients: prospects for paleoaltimetry. *Reviews in Mineralogy and Geochemistry* 66:215–241. DOI: 10.2138/rmg.2007.66.9.

Kronfus H, Havranek WM. 1999. Effects of elevation and wind on the growth of *Pinus cembra* L. in a subalpine afforestation. *Phyton(Austria)* 39:99–106.

Li M-H, Xiao W-F, Shi P, Wang S-G, Zhong Y-D, Liu X-L, Wang X-D, Cai X-H, Shi Z-M. 2008. Nitrogen and carbon source-sink relationships in trees at the Himalayan treelines compared with lower elevations. *Plant, cell & environment* 31:1377–1387. DOI: 10.1111/j.1365-3040.2008.01848.x.

Li MH, Yang J, Kräuchi N. 2003. Growth responses of *Picea abies* and *Larix decidua* to elevation in subalpine areas of Tyrol, Austria. *Canadian Journal of Forest Research* 33:653–662. DOI: 10.1139/X02-202.

- Li C, Zhang X, Liu X, Luukkanen O, Berninger F. 2006. Leaf morphological and physiological responses of *Quercus aquifolioides* along an Altitudinal Gradient. *Silva Fennica* 40:5–13.
- Montesinos-Navarro A, Wig J, Pico FX, Tonsor SJ. 2011. *Arabidopsis thaliana* populations show clinal variation in a climatic gradient associated with altitude. *New phytologist* 189:282–294. DOI: 10.1111/j.1469-8137.2010.03479.x.
- Oberle B, Schaal BA. 2011. Responses to historical climate change identify contemporary threats to diversity in *Dodecatheon*. *Proceedings of the National Academy of Sciences of the United States of America* 108:5655–60. DOI: 10.1073/pnas.1012302108.
- Parkhurst D, Loucks O. 1972. Optimal leaf size in relation to environment. *Journal of Ecology* 60:505–537.
- Paulsen J, Weber UM, Korner C. 2000. Tree growth near treeline : abrupt or gradual reduction with altitude ? *Arct. Antarc. Alp. Res.* 32:14–20.
- Peppe DJ, Royer DL, Cariglino B, Oliver SY, Newman S, Leight E, Enikolopov G, Fernandez-Burgos M, Herrera F, Adams JM, Correa E, Currano ED, Erickson JM, Hinojosa LF, Hoganson JW, Iglesias A, Jaramillo C a, Johnson KR, Jordan GJ, Kraft NJB, Lovelock EC, Lusk CH, Niinemets U, Peñuelas J, Rapson G, Wing SL, Wright IJ. 2011. Sensitivity of leaf size and shape to climate: global patterns and paleoclimatic applications. *New phytologist* 190:724–39. DOI: 10.1111/j.1469-8137.2010.03615.x.
- Premoli AC, Brewer CA. 2007. Environmental vs. genetically driven variation in ecophysiological traits of *Nothofagus pumilio* from contrasting elevations. *Australian Journal of Botany* 55:585. DOI: 10.1071/BT06026.
- Qiang W, Xun-ling W, Chen T, Hu-yuan F, Li-zhe A, Yuan-qin H, Gang W. 2003. Variations of stomatal density and carbon isotope values of *Picea crassifolia* at different altitudes in the

- Qilian Mountains. *Trees* 17:258–262. DOI: 10.1007/s00468-002-0235-x.
- R Core Team. 2015. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.
- Renzhong W, Ripley EA, Yuangang Z, Shaoquan N. 2001. Demography of reproductive and biomass allocation of grassland and dune *Leymus chinensis* on the Songnen Plain, north-eastern China. *Journal of Arid Environments* 49:289–299. DOI: 10.1006/jare.2000.0787.
- Royer DL, McElwain JC, Adams JM, Wilf P. 2008. Sensitivity of leaf size and shape to climate within *Acer rubrum* and *Quercus kelloggii*. *New Phytologist* 179:808–817. DOI: 10.1111/j.1469-8137.2008.02496.x.
- Rozema J, Aerts R, Cornelissen H (eds.). 2002. *Plants and Climate Change*. Vrije Universiteit, Amsterdam, The Netherlands.
- Rozema J, Staaïj van de J, Björn L., Caldwell M. 1997. UV-B as an environmental factor in plant life: stress and regulation. *Trends in Ecology & Evolution* 12:22–28.
- Scheepens JF, Frei ES, Stöcklin J. 2010. Genotypic and environmental variation in specific leaf area in a widespread alpine plant after transplantation to different altitudes. *Oecologia* 164:141–150. DOI: 10.1007/s00442-010-1650-0.
- Schoettle AW. 1990. The interaction between leaf longevity and shoot growth and foliar biomass per shoot in *Pinus contorta* at two elevations. *Tree physiology* 7:209–214.
- Schoettle AW, Rochelle SG. 2000. Morphological variation of *Pinus flexilis* (Pinaceae), a bird-dispersed pine, across a range of elevations. *American Journal of Botany* 87:1797–1806. DOI: 10.2307/2656832.
- Shi Z, Liu S, Liu X, Centritto M. 2006. Altitudinal variation in photosynthetic capacity, diffusional conductance and $\delta^{13}\text{C}$ of butterfly bush (*Buddleja davidii*) plants growing at

high elevations. *Physiologia Plantarum* 128:722–731. DOI: 10.1111/j.1399-3054.2006.00805.x.

Takahashi K, Matsuki S. 2017. Morphological variations of the *Solidago virgaurea* L. complex along an elevational gradient on Mt Norikura, central Japan. *Plant Species Biology* 32:238–246. DOI: 10.1111/1442-1984.12148.

Wang R, Gao Q. 2004. Morphological responses of *Leymus chinensis* (Poaceae) to the large-scale climatic gradient along the North-east China Transect (NECT). *Diversity and Distributions* 10:65–73. DOI: 10.1111/j.1472-4642.2004.00056.x.

Van de Water PK, Leavitt SW, Betancourt JL. 1994. Trends in stomatal density and $^{13}\text{C}/^{12}\text{C}$ ratios of *Pinus flexilis* needles during last Glacial-Interglacial Cycle. *Science* 264:239–243. DOI: 10.1126/science.264.5156.239.

Wilson PJ, Thompson K, Hodgson JG. 1999. Specific leaf area and leaf dry matter content as alternative predictors of plant strategies. *New phytologist* 143:155–162.

Woodward FI. 1986. Ecophysiological studies on the shrub *s* L. taken from a wide altitudinal range. *Oecologia* 70:580–586.

Zuur AF, Hilbe JM, Ieno EN. 2013. *A Beginner's Guide to GLM and GLMM with R*. Highland Statistics, Newburgh, UK.

Figure 1

Figure 1

Study species *Roscoea purpurea* (A) and *R. alpina* (B) in their natural habitat.



Figure 2

Figure 2

Map of the study area: Yellow hexagons represent the study site (Ghorepani) of *Roscoea alpina* whereas Light-Oliventine triangles represent the study site (Daman) of *R. purpurea* (See Table 1 for detail). The top map represents the elevational gradient of study locations.




Figure 3

Figure 3

Correlations between the different traits of *R. purpurea* measured at five different elevations.




Figure 4

Figure 4

Correlations between the different traits of *R. alpina* measured at five different elevations.




Figure 5

Figure 5

Generalised linear regression models showing the effect of elevation on plant height (fig. a), leaf length (fig. b), leaf area (fig. c), specific leaf area (fig. d) and, stomata density (fig. e) for *R. purpurea* (blue line with filled markers) and *R. alpina* (purple line with empty markers). Markers indicate the mean value of the corresponding trait at each elevation and error bars indicate standard deviation. Solid lines represent the regression function for each trait and species. A significant correlation of elevation on the value for each trait is indicated by two asterisks (**) while a non-significant correlation of elevation is indicated by “NS” above the corresponding regression line.




Table 1 (on next page)

Table 1

Geographical details of the study sites.

Table 1: Geographical details of study sites.

<i>Roscoea purpurea</i>			<i>R. alpina</i>		
Latitude	Longitude	Elevation (a.s.l.)	Latitude	Longitude	Elevation (a.s.l.)
27°36'45.7"	85°5'32"	2174 m	28°23'21.9"	83°42'22.1"	2675 m
27°36'44.7"	85°5'37.6"	2274 m	28°23'42.2"	83°42'9.2"	2770 m
27°36'2.1"	85°5'13.4"	2374 m	28°24'4.9"	83°41'58.9"	2874 m
27°35'37.1"	85°5'57.3"	2474 m	28°24'5.2"	83°41'46.8"	2968 m
27°35'37.4"	85°5'17.3"	2574 m	28°24'0.9"	83°41'34.7"	3079 m

Table 2(on next page)

Table 2

Results of regression analysis between various traits of *R. alpina* and elevation.

Table 2: Results of regression analysis between various traits of *R. alpina* and elevation.

Traits	Parameters	Coefficients and 95 % Cis			Distribution	link	P
		2.5	50	97.5			
Plant height	m	3.85E-05	6.48E-05	9.12E-05	Gamma	Inverse	0.017
	b	-1.52E-01	-7.71E-02	-1.85E-05			
Leaf length	m	6.99E-05	8.37E-05	9.75E-05	Gamma	Inverse	<0.001
	b	-1.87E-01	-1.47E-01	-1.08E-01			
Leaf area	m	-4.00E-02	-3.60E-02	-3.30E-02	Gamma	Identity	<0.001
	b	1.09E+02	1.21E+02	1.33E+02			
SLA	m	3.91E-07	3.33E-06	6.31E-06	Gamma	Inverse	0.114
	b	-1.43E-02	-5.85E-03	2.55E-03			
Stomata Density	m	9.10E-02	1.24E-01	1.58E-01	Gaussian	Identity	0.005
	b	-2.75E+02	-1.78E+02	-8.15E+01			

Table 3(on next page)

Table 3

Results of regression analysis between various traits of *R. purpurea* and elevation.

1 **Table 3:** Results of regression analysis between various traits of *R. purpurea* and elevation.

Traits	Param eters	Coefficients and 95 % Cis			Distrib ution	link	P
		2.5	50	97.5			
Plant height	m	-7.90E-03	-5.00E-03	-2.10E-03	Gaussian	Identity	0.044
	b	2.59E+01	3.29E+01	3.99E+01			
Leaf length	m	-7.00E-03	-2.00E-03	3.00E-03	Gamma	Identity	0.47
	b	1.15E+01	2.32E+01	3.49E+01			
Leaf area	m	-6.70E-02	-5.80E-02	1.55E+02	Gamma	Identity	<0.001
	b	-4.80E-02	1.79E+02	2.03E+02			
SLA	m	-7.55E-06	-1.90E-06	3.71E-06	Gamma	Inverse	0.55
	b	-4.40E-03	8.91E-03	2.26E-02			
Stomata Density	m	-7.31E-02	-6.18E-06	-5.05E-06	Gamma	Inverse	0.002
	b	2.04E-02	2.32E-02	2.59E-02			

2

3

4

5