

Efficient virus-induced gene silencing in *Hibiscus hamabo* Sieb. et Zucc. using tobacco rattle virus

Zhiquan Wang¹, Xiaoyang Xu¹, Longjie Ni^{1,2}, Jinbo Guo¹, Chunsun Gu^{1*}

¹Jiangsu Key Laboratory for the Research and Utilization of Plant Resources, Institute of Botany, Jiangsu Province and Chinese Academy of Sciences, Nanjing, Jiangsu, China.

²College of Forest Sciences, Nanjing Forestry University, Nanjing, Jiangsu, China.

Corresponding Author:

Chunsun Gu¹

1 Qianhuhoucun Road, Nanjing, Jiangsu, 210014, China

Email address: chunsungu@cnbg.net

Abstract

Background. *Hibiscus hamabo* Sieb. et Zucc. is a semi-mangrove plant used for the ecological restoration of saline-alkali land, coastal afforestation and urban landscaping. Because the genetic transformation of *H. hamabo* is difficult, molecular breeding and gene functional studies have been severely restricted. In plants, virus-induced gene silencing provides a pathway to rapidly and effectively create targeted gene knockouts for gene functional studies.

Methods. In this study, we tested the efficiency of a tobacco rattle virus vector in silencing the chloroplasts alterados 1 (*CLAI*) gene through agroinfiltration.

Results. The leaves of *H. hamabo* showed white streaks typical of *CLAI* gene silencing three weeks after agroinfiltration. In agroinfiltrated *H. hamabo* plants, the *CLAI* expression levels in leaves with white streaks were all significantly lower than in those of mock-infected and control plants.

Conclusions. Thus, this virus-induced gene silencing system was efficient in *H. hamabo* and may be a powerful tool for large-scale reverse-genetic analyses of gene functions in *H. hamabo*.

Introduction

Hibiscus hamabo Sieb. et Zucc., which is a shrub plant in the genus *Hibiscus*, family Malvaceae, is an important semi-mangrove plant (Nakanishi 1979). Because of its excellent salt tolerance and morphological characteristics, *H. hamabo* is widely used in public parks, waysides and coastal sands near sea level (Fowler 2017; Li et al. 2012; Yang et al. 2008). In addition, *H. hamabo* is a good plant material for exploring the salt-stress response mechanisms of woody plants (Li et al. 2012). The desire to determine gene functions and regulatory mechanisms in *H. hamabo* has led to the urgent need to manipulate genes. However, to date, the inefficient and laborious transformation procedures used have impeded this research. Additionally, transcriptome

Commented [XX-A1]: Molecular breeding is not necessarily dependent on genetic transformation. Instead, it is associated with molecular mapping.

Formatted: Highlight

Commented [XX-A2]: Please rephrase this sentence.

Formatted: Highlight

Commented [XX-A3]:

analyses have mined many excellent genes that are awaiting functional identification. Appropriate techniques need to be applied successfully to allow the study of gene functions in this plant.

Commented [XX-A4]: Revisions are needed.

Virus-induced gene silencing (VIGS) is a powerful technology that uses engineered viruses to specifically silence host gene expression through post-transcriptional gene silencing (Becker & Lange 2010; Krishnan et al. 2015; Purkayastha & Dasgupta 2009). VIGS is an effective method for the large-scale analysis of genes and their functions, and it has been successfully performed in many plants, including tobacco, *Arabidopsis*, tomato, cotton, wheat, and many woody plants (Burch-Smith et al. 2006; Jiang et al. 2014; Kumagai et al. 1995; Orzaez et al. 2009; Scofield et al. 2005). VIGS works via a mechanism that is similar to that of RNA interference (Baulcombe 1999; Baulcombe 2004; Burch-Smith et al. 2004; Lu et al. 2003; Waterhouse et al. 2001). Double-stranded (ds) RNA is the key to the VIGS process; the dsRNA could be cleaved into short interfering (si) RNAs of 21 to 25 nucleotides (Burch-Smith et al. 2004; Jiang et al. 2014; Lu et al. 2003). Two strands can be obtained from the siRNAs, the guide and passenger strands. The RNA-induced silencing complex incorporated the guide strand to degrade the specific single-stranded RNA that was complementary to the guide RNA, and then, the passenger strand was degraded (Mustafa et al. 2016). As a result, the target gene was silenced and large amounts of siRNAs were produced (Fuchs et al. 2004). Previous studies had reported the detailed protocol of agrobacterium-mediated VIGS system in cotton and had optimized the conditions for VIGS in three cultivated cotton species, indicated that tobacco rattle virus (TRV) was a useful vector for VIGS in *Gossypium* species (Gao et al. 2011; Ge et al. 2016).

Formatted: Highlight

Tobacco rattle virus (TRV), belonging to genus *Tobravirus* (family *Virgaviridae*), is a suitable virus vector system for VIGS (Jiang et al. 2014). A positive sense single-stranded RNA genome exists in TRV, consisting of two components, RNA 1 and RNA 2 (Mustafa et al. 2016). RNA 1 encodes genes with viral replication and movement functions, while RNA 2 encodes the coat protein and some nonessential structural proteins that can be replaced by foreign sequences (Hayward et al. 2011). The TRV vector has been used in *G. spp.*, *Arabidopsis* and *Vernicia fordii* in silencing the chloroplasts *alterados 1 (CLAI)* gene, which is involved in chloroplast development (Jiang et al. 2014; Manhães et al. 2015; Mustafa et al. 2016). The *CLAI* gene is highly conserved in various plant species (Jiang et al. 2014). The silencing phenotypes of albino leaves were observed in *Vernicia fordii* two weeks after inoculation using a heterologous TRV-based VIGS system, which *CLAI* was isolated from *Populus tomentosa* Carr. (Jiang et al. 2014). Silencing *CLAI* is a useful marker for determining the silencing efficiency because of the bleached phenotype (Mustafa et al. 2016). In this study, we tested the feasibility of TRV-VIGS system in *H. hamabo* using *HhCLAI* gene as a reporter. The agroinfiltrated leaves of *H. hamabo* showed white streaks typical at three weeks after infection, and the expression levels of the *HhCLAI* gene in leaves with white streaks were significantly lower than that in leaves from mock-infected and control plants. Thus, the TRV-VIGS system was efficient in *H. hamabo*. To our knowledge, this is the first report of the successful application of VIGS in *H. hamabo*.

Commented [XX-A5]: What protocol? VIGS system or TVR-based VIGS system protocol? Revisions are needed for this paragraph.

Commented [XX-A6]: Revisions are needed.

Commented [XX-A7]: Revisions are needed

Commented [XX-A8]:

Commented [XX-A9]: Revisions are needed

Materials & Methods

Plant materials and growth conditions.

Seeds of *H. hamabo* were collected from Nanjing's Sun Yat-Sen Memorial Botanical Garden. The seeds were then treated with concentrated sulfuric acid for 15 min and rinsed thoroughly with sterile water. Next, the pretreated seeds were sown into flowerpots containing a mixture of peat and vermiculite (1: 1, v: v) in an illuminated incubator with day/night temperatures of 26°C /22°C and a 16/8 h photoperiod.

Sequence analysis.

Based on the *HhCLA1* sequence (GenBank accession no. MK229167), the deduced protein sequence was analyzed with CLA1 proteins of other species by ClustalX (Liu et al. 2015). The amino acid sequences were obtained from NCBI (<https://www.ncbi.nlm.nih.gov/>). Then, the sequences were used to construct a phylogenetic tree, which was drawn with MEGA 7.0 using the Neighbor-Joining (NJ) method and 1,000 bootstrap replicates.

VIGS vector construction.

Total RNA was extracted from the leaves of *H. hamabo* using a Plant RNeasy Mini Kit (Qiagen, Hilden, Germany). The first-strand cDNA was synthesized using a SuperScript II reverse transcriptase kit (TaKaRa, Dalian, China). The primer pair *HhCLA1*-F and *HhCLA1*-R (Table 1) was designed using Oligo 6.0 software (Molecular Biology Insights, Inc., Cascade, CO, USA) based on the conserved domain of *HhCLA1*. To amplify partial fragments of *HhCLA1*, the primer pair, cDNA and PrimeSTARTM HS DNA polymerase (TaKaRa) were used. *Eco*RI enzyme cleavage sites were added to the upstream primers and *Sac*I enzyme cleavage sites were added to the downstream primers. The *pTRV1* and *pTRV2* vectors were used in this study as described previously (Gao et al. 2011; Liu et al. 2002). The PCR products were ligated into pTRV2 (Supplementary Fig. S1) (double-digested with *Eco*RI and *Sac*I enzymes) using a ClonExpress[®] II One Step Cloning Kit (Vazyme, Nanjing, China). The resulting vector was designated as pTRV2-*HhCLA1*.

Agroinfiltration.

pTRV2-*HhCLA1* was transformed into *Agrobacterium tumefaciens* strain 'GV3101' using the freeze-thawing method (Höfgen & Willmitzer 1988). PCR-confirmed single colonies were then selected and independently inoculated into 3 mL of Luria-Bertani medium containing 25 mg/L rifampicin and 50 mg/L kanamycin and grown overnight in a 28°C shaker. For the VIGS assay, 3-mL cultures of *A. tumefaciens* strain GV3101 independently containing pTRV1 and pTRV2 were grown overnight in the same culture conditions. These overnight starter cultures were subsequently used to inoculate 50-mL cultures that were grown overnight at 28°C. *Agrobacterium* cultures were harvested by centrifugation at 4,000 × g for 10 min, and the pellets were resuspended in an infiltration buffer (10 mM MES (2- (N-morpholino) ethanesulfonic acid), 10 mM MgCl₂ and 200 μM acetosyringone, pH 5.6) at an optical density of 2.0 at 600 nm and incubated at room temperature for 3 h without shaking. *Agrobacterium* cultures containing mixtures of pTRV1 and pTRV2-*HhCLA1* (1: 1 ratio) were infiltrated with needleless 1-mL syringes into the backs of cotyledons of 2-week-old *H. hamabo* seedlings, following a protocol described previously (Gao et al. 2011). To determine whether the TRV vector can directly infect *H. hamabo*, a mixture of

Commented [XX-A10]: PLEASE REPHRASE THIS SENTENCE.

Formatted: Highlight

Commented [XX-A11]: You may provide detailed PCR protocol here.

Commented [XX-A12]: You mean "in a shaker at 28°C"?

Formatted: Highlight

Formatted: Highlight

Commented [XX-A13]: There are two cultures, right? Please rephrase this sentence.

Commented [XX-A14]: You have two cultures, right? Please rephrase the sentence.

Formatted: Highlight

Agrobacterium cultures containing pTRV1 and pTRV2 constructs in a 1: 1 ratio was infiltrated into the backs of cotyledons of eight 2-week-old *H. hamabo* to serve as the mock. Experimental and non-injected control plants were transferred to a growth chamber and maintained under set conditions.

Quantitative real-time PCR (qPCR).

To determine the relative levels of the endogenous *HhCLA1* transcripts in infected leaves exhibiting visible silencing phenotypes, qPCR was performed using the primer pair q*HhCLA1*-F/q*HhCLA1*-R (Table 1). For the experiments, leaves from plants with significant white streak symptoms were analyzed in comparison with leaves of the mock and control plants after three weeks of agroinfiltration. Total RNA was extracted from these leaves using a Plant RNeasy Mini Kit (Qiagen) and treated with DNase I to remove residual DNA. The first-strand cDNA was synthesized using a SuperScript II reverse transcriptase kit (TaKaRa). The qPCR assays were performed using the SYBR Green PCR Master Mix (Bimake, Houston, TX, USA) and a StepOne™ System (ABI, USA). The transcript level of *18S rRNA* was served as internal standards. All experiments were repeated three times. The relative gene expression level was calculated using the $2^{-\Delta\Delta Ct}$ method (Gu et al. 2018; Liao et al. 2016).

Statistical analysis.

Results were expressed as means \pm standard errors. Statistical procedures were conducted with IBM SPSS (Version 21). Data of qPCR was analyzed with one-way analysis of variance (ANOVA) followed by Duncan's multiple range test ($P < 0.05$).

Results

Characterization of the *HhCLA1* gene in *H. hamabo*.

The amino acid sequence alignment indicated that the HhCLA1 protein showed high homology to known CLA1 proteins from *G. barbadense*, *G. hirsutum* and other species (Fig. 1 and Supplementary Table S1). To understand the evolutionary relationship of CLA1 proteins from different species, the amino acid sequences of HhCLA1 was aligned with other CLA1 proteins, and an un-rooted neighbor-joining (NJ) phylogenetic tree was constructed using MEGA 7.0 software. The phylogenetic analysis showed that HhCLA1 clustered with *G. barbadense* and *G. hirsutum* in a clade (Fig. 2).

Silencing efficiency of the *HhCLA1* gene in *H. hamabo* using the VIGS system.

In total, fifty-two *H. hamabo* plants were inoculated with *A. tumefaciens* 'GV3101' harboring pTRV2-*HhCLA1*. Two weeks after agroinfiltration, white streaks began to appear in the emerging leaves of partially agroinoculated plants. At three weeks post infiltration, 87% of the *H. hamabo* plants showed white-streak leaf symptoms (Table 2; Fig. 3A and Supplementary Fig. S2). At three weeks after agroinfiltration, plants inoculated with pTRV1 and pTRV2 (Mock) showed no obvious differences in leaf morphology compared with the control (Fig. 3B, 3C and Supplementary Fig. S2). The three leaves on the left in Fig. 3D were picked from plants infiltrated with pTRV2-*HhCLA1* (*CLA1*), the leaf in the middle was got from empty vector infiltrated plants (Mock) and the right one was from control plants (CK). Compared with the plants of mock and CK, the leaf

Formatted: Highlight

Commented [XX-A15]: Rephrase this paragraph

Formatted: Highlight

Commented [XX-A16]: Please give detailed results: plants showing symptoms.

Commented [XX-A17]: This part should be included in the legend of Fig. 3D.

phenotypic characteristics suggested that *HhCLA1* gene expression maybe silenced in these treated plants infiltrated with pTRV2-*HhCLA1*.

q-PCR analysis of the knockdown levels of *HhCLA1*.

The gene silencing efficiency was analyzed at the molecular level by monitoring expression levels of *HhCLA1* in plants showing white-streak leaf symptoms. Four groups of plants with significant white streak symptoms, one group of control and one group of mock, in order to analyze the test results more accurately, were analyzed in this experiment. Each group contained three biological replicates. The results showed that *HhCLA1* gene expression levels were unchanged in mock-injected plants, while the *HhCLA1* expression levels were 62.6%-76.4% lower in the pTRV2-*HhCLA1* agroinfiltrated plants than that in the non-infiltrated plants (control) (Fig. 4 and Table S2). The phenotypic characteristics were consistent with the expression characteristics of *HhCLA1*, clearly indicated that the expression of *HhCLA1* was significantly down-regulated through TRV-VIGS in *H. hamabo*, and led to an albino phenotype on true leaves.

Formatted: Highlight

Formatted: Highlight

Commented [XX-A18]: Please rephrase.

Discussion

In this study, we demonstrated for the first time that TRV-VIGS could effectively down-regulate endogenous gene expression levels in the salt-tolerant species *H. hamabo*. The genetic transformation of this species is laborious, time-consuming and technically challenging. To resolve these problems, effective and low-cost techniques need to be developed to enable the rapid validation of gene functions. In future studies, stress-responsive genes isolated in *H. hamabo* could be silenced in loss-of-function screens using the TRV-VIGS system.

The *CLA1* gene is involved in chloroplast development and is a useful marker in the TRV-VIGS system (Mustafa et al. 2016). In this research, multiple sequence alignments indicated that *HhCLA1* was similar to those *CLA1* proteins of other species. Additionally, the phylogenetic analysis indicated that *HhCLA1* was highly similar to *CLA1* proteins in Malvaceae, including *G. barbadense* and *G. hirsutum*.

The most cost-efficient and effective method of inoculating plants with virus-based vectors is agroinfection (Grimsley et al. 1986), but its efficiency varies among plants (Zhang et al. 2016). In turf grass, the silencing efficiency of the RTBV-VIGS system in *Cynodon dactylon* was such that 65.8- 72.5% of the agroinfected plants developed symptoms typical for phytoene desaturase gene silencing, while the silencing efficiency in *Zoysia japonica* was much weaker, with only 52.7%- 55% of agroinfected plants developing the phenotype (Zhang et al. 2016). The ability of the TRV vector to directly infect woody plant species has been tested, and TRV-mediated VIGS was effective in *Vernicia fordii*, weak in *Populus tomentosa* Carr., and ineffective in *Camellia oleifera* (Jiang et al. 2014). In this study, the silencing efficiency of the TRV-VIGS system in *H. hamabo* was high, with 87% of agroinfected plants developing a white-streak leaf phenotype, indicating *HhCLA1* gene was silenced. The *HhCLA1* mRNA level was also down-regulated by TRV-VIGS in *H. hamabo*.

Conclusions

In conclusion, we demonstrated that TRV-mediated VIGS was effective in *H. hamabo*, which added to the increasing list of wood species for which VIGS-mediated studies can be used. The loss-of-function assay using TRV-mediated VIGS developed in this study provided an alternative tool for functional genes studies of *H. hamabo*.

Acknowledgements

We thank International Science Editing (<http://www.internationalscienceediting.com>) for editing this manuscript.

References

- Baulcombe D. 1999. Viruses and gene silencing in plants. *100 Years of Virology*: Springer, 189-201.
- Baulcombe D. 2004. RNA silencing in plants. *Nature* 431(7006): 356.
- Becker A, and Lange M. 2010. VIGS—genomics goes functional. *Trends in Plant Science* 15: 1-4.
- Burch-Smith TM, Schiff M, Liu Y, and Dinesh-Kumar SP. 2006. Efficient virus-induced gene silencing in Arabidopsis. *Plant Physiology* 142: 21-27.
- Burch-Smith TM, Anderson JC, Martin GB, and Dinesh-Kumar S. P. 2004. Applications and advantages of virus - induced gene silencing for gene function studies in plants. *Plant Journal* 39(5): 734-746.
- Fowler E. 2017. Assessment and characterization of microbial communities in salt affected soils on Galveston Island. SFA scholarworks, Electronic Theses and Dissertations. 76, Stephen F Austin State University.
- Fuchs U, Damm-Welk C, and Borkhardt A. 2004. Silencing of disease-related genes by small interfering RNAs. *Current Molecular Medicine* 4: 507-517.
- Gao X, Britt Jr RC, Shan L, and He P. 2011. Agrobacterium-mediated virus-induced gene silencing assay in cotton. *Journal of Visualized Experiments: JoVE* 54: e2938.
- Ge X, Wu J, Zhang C, Wang Q, Hou Y, Yang Z, Yang Z, Xu Z, Wang Y, Lu L, Zhang X, Hua J, and Li F. 2016. Prediction of VIGS efficiency by the Sfold program and its reliability analysis in *Gossypium hirsutum*. *Science Bulletin* 61(7): 543-551.
- Grimsley N, Hohn B, Hohn T, and Walden R. 1986. "Agroinfection," an alternative route for viral infection of plants by using the Ti plasmid. *Proceedings of the National Academy of Sciences* 83:3282-3286.
- Gu C, Liu L, Song A, Liu Z, Zhang Y, and Huang S. 2018. *Iris lactea* var. *chinensis* (Fisch.) cysteine-rich gene *lICDT1* enhances cadmium tolerance in yeast cells and *Arabidopsis thaliana*. *Ecotoxicology and Environmental Safety* 157:67-72.
- Höfgen R, and Willmitzer L. 1988. Storage of competent cells for Agrobacterium transformation. *Nucleic Acids Research* 16:9877.
- Hayward A, Padmanabhan M, and Dinesh-Kumar S. 2011. Virus-induced gene silencing in *Nicotiana benthamiana* and other plant species. *Plant Reverse Genetics: Springer* 55-63.
- Jiang Y, Ye S, Wang L, Duan Y, Lu W, Liu H, Fan D, Zhang F, and Luo K. 2014. Heterologous gene silencing induced by tobacco rattle virus (TRV) is efficient for pursuing functional genomics studies in woody plants. *Plant Cell Tissue & Organ Culture* 116:163-174.
- Krishnan A, Mahadevan C, Mani T, and Sakuntala M. 2015. Virus-induced gene silencing (VIGS)

- for elucidation of pathogen defense role of serine/threonine protein kinase in the non-model plant *Piper colubrinum* Link. *Plant Cell Tissue & Organ Culture* 122:269-283.
- Kumagai M, Donson J, Della-Cioppa G, Harvey D, Hanley K, and Grill L. 1995. Cytoplasmic inhibition of carotenoid biosynthesis with virus-derived RNA. *Proceedings of the National Academy of Sciences* 92:1679-1683.
- Li J, Liao J, Guan M, Wang E, and Zhang J. 2012. Salt tolerance of *Hibiscus hamabo* seedlings: a candidate halophyte for reclamation areas. *Acta Physiologiae Plantarum* 34:1747-1755.
- Liao YL, Xu F, Zang WW, Cheng SY, Shen YB, and Chang J. 2016. Cloning, characterization and expression analysis of GbWRKY11, a novel transcription factor gene in *Ginkgo biloba*. *International Journal of Agriculture and Biology* 18(1): 117-124.
- Liu C, Wang G, Wang H, Xia T, Zhang S, Wang Q, and Fang Y. 2015. Phylogenetic relationships in the genus *Rosa* revisited based on rpl16, trnL-F, and atpB-rbcL sequences. *HortScience* 50(11): 1618-1624.
- Liu Y, Schiff M, and Dinesh-Kumar S. 2002. Virus-induced gene silencing in tomato. *The Plant Journal* 31: 777-786.
- Lu R, Martin-Hernandez AM, Peart J R, Malcuit I, and Baulcombe DC. 2003. Virus-induced gene silencing in plants. *Methods* 30(4): 296-303.
- Manhães AMEdA, de Oliveira MV, and Shan L. 2015. Establishment of an Efficient Virus-Induced Gene Silencing (VIGS) Assay in *Arabidopsis* by Agrobacterium-Mediated Rubbing Infection. *Plant Gene Silencing: Springer* 235-241.
- Mustafa R, Shafiq M, Mansoor S, Briddon RW, Scheffler BE, Scheffler J, and Amin I. 2016. Virus-induced gene silencing in cultivated cotton (*Gossypium* spp.) using tobacco rattle virus. *Molecular Biotechnology* 58:65-72.
- Nakanishi H. 1979. Distribution and ecology of a semi-mangrove plant, *Hibiscus hamabo* SIEB. et ZUCC. and its community. *Acta Phytotaxonomica Et Geobotanica* 30:169-179.
- Orzaez D, Medina A, Torre S, Fernández-Moreno JP, Rambla JL, Fernández-del-Carmen A, Butelli E, Martin C, and Granell A. 2009. A visual reporter system for virus-induced gene silencing in tomato fruit based on anthocyanin accumulation. *Plant Physiology* 150:1122-1134.
- Purkayastha A, and Dasgupta I. 2009. Virus-induced gene silencing: a versatile tool for discovery of gene functions in plants. *Plant Physiology and Biochemistry* 47:967-976.
- Scofield SR, Huang L, Brandt AS, and Gill BS. 2005. Development of a virus-induced gene-silencing system for hexaploid wheat and its use in functional analysis of the Lr21-mediated leaf rust resistance pathway. *Plant Physiology* 138:2165-2173.
- Waterhouse PM, Wang MB, and Lough T. 2001. Gene silencing as an adaptive defence against viruses. *Nature* 411:834.
- Yang H, Du GJ, and Wang KH. 2008. Study on the Physiological Characteristics of *Hibiscus hamabo* under Stress. *Journal of Zhejiang Forestry Science and Technology* 3:011.
- Zhang B, Shi J, Chen J, Li D, Li J, Guo H, Zong J, Wang Y, Guo A, and Liu J. 2016. Efficient virus-induced gene silencing in *Cynodon dactylon* and *Zoysia japonica* using rice tungro bacilliform virus vectors. *Scientia Horticulturae* 207:97-103.