

Gradual, temperature-induced change of secondary sexual characteristics in *Trichogramma pretiosum* infected with parthenogenesis-inducing *Wolbachia*

Su-fang Ning^{Equal first author, 1}, Jin-cheng Zhou^{Equal first author, 1}, Quan-quan Liu¹, Qian Zhao¹, Hui Dong^{Corresp. 1}

¹ College of Plant Protection, Shenyang Agricultural University, Shenyang, Liaoning, China

Corresponding Author: Hui Dong
Email address: biocontrol@163.com

Intersex is an intermediate stage of sexual differentiation in insects. Determining intersex morphology and the cause of its production will aid in the understanding of the mechanism of sexual differentiation in insects. In this paper, *Wolbachia*-infected *Trichogramma pretiosum* (*T. preW*⁺) that shows thelytokous parthenogenesis were used as subjects. In order to determine the causes of the *T. preW*⁺ intersex and the influence of parental generation temperature on gradual changes in secondary masculinization in intersex offspring, we examined the occurrence of intersex offspring (F1 and F2 generation) after the parental generations were treated with high temperature (27°C, 29°C, 31°C, and 33°C) and described the external morphology of the intersexes. The results showed that the *T. preW*⁺ parental generation temperature is positively correlated with the probability of intersex offspring. The probability of F1 intersex is significantly higher than that of F2 intersex in different high temperature. The degree of secondary masculinization in *T. preW*⁺ intersexes increases as parental generation temperature increases. In addition, our study first identified 11 intersex types in *T. preW*⁺ and found that the primary and secondary sexual characteristics showed a regular distribution. And we found that the D type and H type of intersex have the highest frequency of appearance. The external genitalia of most intersexes were female, and only three intersex types have male external genitalia. Conclusions were ultimately obtained: *Wolbachia* is a direct factor that causes the occurrence of intersexes, while high temperature is an indirect factor that determines the external morphology of intersexes. The effects of high temperature on *T. preW*⁺ intersexes is passed through the parental generation to offspring, and this maternal effect weakens as the number of generations increases. In *T. preW*⁺ intersex individuals, most exhibit female primary sexual characteristics, and secondary sexual characteristics exhibit

signs of masculinization.

Gradual, temperature-induced change of secondary sexual characteristics in *Trichogramma pretiosum* infected with parthenogenesis-inducing *Wolbachia*

Su-fang Ning[†], Jin-cheng Zhou[†], Quan-quan Liu, qian Zhao, hui Dong*

* Correspondence to: Hui Dong, College of Plant Protection, Shenyang Agricultural University, Shenyang, Liaoning, 110866, China. E-mail: biocontrol@163.com

[†] These authors contributed equally to the work.

College of Plant Protection, Shenyang Agricultural University, Shenyang, Liaoning, China

Key words: *Trichogramma pretiosum*; temperature; *Wolbachia*; intersex; secondary sexual characteristics

Abstract

Intersex is an intermediate stage of sexual differentiation in insects. Determining intersex morphology and the cause of its production will aid in the understanding of the mechanism of sexual differentiation in insects. In this paper, *Wolbachia*-infected *Trichogramma pretiosum* (*T. preW*⁺) that shows thelytokous parthenogenesis were used as subjects. In order to determine the causes of the *T. preW*⁺ intersex and the influence of parental generation temperature on gradual changes in secondary masculinization in intersex offspring, we examined the occurrence of

intersex offspring (F1 and F2 generation) after the parental generations were treated with high temperature (27°C, 29°C, 31°C, and 33°C) and described the external morphology of the intersexes. The results showed that the *T. preW*⁺ parental generation temperature is positively correlated with the probability of intersex offspring. The probability of F1 intersex is significantly higher than that of F2 intersex in different high temperature. The degree of secondary masculinization in *T. preW*⁺ intersexes increases as parental generation temperature increases. In addition, our study first identified 11 intersex types in *T. preW*⁺ and found that the primary and secondary sexual characteristics showed a regular distribution. And we found that the D type and H type of intersex have the highest frequency of appearance. The external genitalia of most intersexes were female, and only three intersex types have male external genitalia. Conclusions were ultimately obtained: *Wolbachia* is a direct factor that causes the occurrence of intersexes, while high temperature is an indirect factor that determines the external morphology of intersexes. The effects of high temperature on *T. preW*⁺ intersexes is passed through the parental generation to offspring, and this maternal effect weakens as the number of generations increases. In *T. preW*⁺ intersex individuals, most exhibit female primary sexual characteristics, and secondary sexual characteristics exhibit signs of masculinization.

Introduction

The difference between the sexes of the same species is called sexual dimorphism in nature. Generally, two types of sexual characteristics are used to describe sexual dimorphism. Primary sexual characteristics include male and female reproductive organs that are directly related to reproduction (such as external genitalia and internal genitalia). Whereas the Secondary sexual characteristics include body size, external morphology, antenna, etc., and are used to describe

characteristics that aid in reproduction (Schärer *et al.*, 2012; Tulgetske & Stouthamer *et al.*, 2012; Bear *et al.*, 2013). Usually, sexually aberrant individuals with conflicting primary and secondary sexual characteristics are known as intersexes, gynandromorphs and sexual mosaics (Bowen & Monteiro, 1966; Pintureauand & Bolland, 2001; Pintureau *et al.*, 2002; Tulgetske *et al.*, 2012). Of these terms, intersexes refer to individuals with uniform genetic constitution, but some tissues show sexual phenotypes that conflict with the associated genotype (Goldschmidt *et al.*, 1949). However, this concept is frequently confused or wrongly used with gynandromorphs (Tulgetske *et al.*, 2012). In actual fact, the latter refers to individuals that are composed of genetically distinct tissues that correspond to the sexual phenotypes observed (Brust *et al.*, 1966). An example is the genus *Trichogramma* in the order Hymenoptera, sexually aberrant individuals were often described as gynandromorphs and intersexes. However, Tulgetske et al.(2012) showed that these individuals are intersexes and not gynandromorphs. The intersexes can be falls into two categories: female intersex and male intersex in Hymenoptera (Whiting *et al.*, 1934; Tulgetske *et al.*, 2012). Female intersexes begin their development as females but organs developed later are masculinized, and male intersexes begin their development as males but organs developed later are feminized (Whiting *et al.*, 1934). As intersex is an intermediate stage of sexual differentiation in insects (Bowen *et al.*, 1966), in-depth research on these rare sexually aberrant individuals is necessary and crucial. Therefore, determining intersex morphology and the cause of its production will aid in the understanding of the mechanism of sexual differentiation in insects.

Many studies showed that intersex formation may be affected by single or multiple regulatory factors that participate in sex determination. These regulatory factors include mutations (Cline *et al.*, 1984; Schutt & Nöthiger, 2000), chromosomal aberrations (Seiler *et al* 1958; Beukeboom &

69 Kamping, 2006), and epigenetic factors (eg. environment, sex hormones and symbionts) (Rigaud
70 & Juchault, 1998; Narita *et al.*, 2010; Tulgetske *et al.*, 2012; Martin & Scholtz, 2012). In
71 *Trichogramma*, intersex formation is associated with *Wolbachia* regulation. The intracellular
72 symbiotic bacteria, *Wolbachia*, is a type of sex regulatory element that is ubiquitous in
73 arthropods. *Wolbachia* can induce insects to undergo parthenogenesis (PI), cytoplasmic
74 incompatibility(CI), male-killing(MK), and feminization (Werren & Windsor, 2000; Almeida &
75 Stouthamer, 2018). The sex determination mechanism for *Trichogramma* is haplodiploidy,
76 where fertilized eggs develop into diploid females and unfertilized eggs develop into haploid
77 males (Perlman *et al.*, 2008). However, PI *Wolbachia* can induce parthenogenesis in
78 *Trichogramma* by causing unfertilized eggs to undergo diploidization and feminization. Thus,
79 females could produce nearly 100% female offsprings even without mating (Ma *et al.*, 2015).
80 According to our knowledge of *Trichogramma*, sexually aberrant individuals are only present in
81 parthenogenetic species. (Tulgetske *et al.*, 2010). The *Wolbachia*-mediated phenotype are closely
82 related with *Wolbahcia* titers. Many studies revealed *Wolbachia* were sensitive to mutiple factors
83 (e.g. temperature and antibiotic treatment) (Tulgetske *et al.*, 2010; Tulgetske *et al.*, 2012;
84 Almeida & Stouthamer, 2018). The reduction of *Wolbachia* titers was often reported in high
85 temperature (Zchori-Fein *et al.*, 2000; Ma & Schwander, 2017). Almost no intersexes appear
86 under normal temperature. However, the number of intersexes significantly increases after high
87 temperature or antibiotic treatment (Pintureau & Bolland, 2001; Tulgetske *et al.*, 2010). The
88 specific reason for this may be that *Wolbachia* regulation of *Trichogramma* wasps dominate at
89 normal temperatures but high temperature and antibiotics eliminated the dominance of
90 *Wolbachia*, thereby decreasing *Wolbachia* titers in infected parthenogenetic insects (Schilthuizen
91 & Stouthamer, 1997; Kent *et al.*, 2011; Fukui *et al.*, 2015). This disrupts or inhibits *Wolbachia*-

induced feminization (Rigaud *et al.*, 1998), ultimately resulting in a higher proportion of intersex offspring (Pintureau & Bolland, 2001; Pintureau *et al.*, 1999). Intersexes were reported at frequencies of parental generation after continuous high temperature treatment (Pintureau *et al.*, 1999; Pintureau & Bolland, 2001). However, whether parental generation treatment will affect intersex offspring and whether parental generation effects can be transmitted to the next generation are still unknown. In addition, there is a lack of objective and accurate estimation of gradual changes in secondary sexual characteristics as influenced by temperature in intersexes. In view of the aforementioned phenomena, we carried out the following study.

Trichogramma is an extremely effective natural enemy of egg parasites and has an indispensable role in biological pest control (Sigsgaard *et al.*, 2017). In this study, *T. preW*⁺ populations were used as subjects to study the effects of different feeding temperature in the parental generation (27°C, 29°C, 31°C, 33°C) on the probability of intersexes in the F1 and F2 generation. The levels of masculinization, which were determined based on previous descriptions (Tulgetske *et al.*, 2012), were introduced to reveal the effector pattern of high temperature on gradual changes in secondary sexual characteristics in *T. preW*⁺. Our study systematically described and identified the external morphological characteristics of intersexes. The results of this study may aid in the understanding of sex determination and sexual differentiation in *T. preW*⁺.

Materials & Methods

Insect breeding

Trichogramma pretiosum Riley (*T. preW*⁺) infected with *Wolbachia* were provided by the Pest Biological Control Laboratory of Shenyang agricultural university. And the uninfected with *Wolbachia* of *Trichogramma pretiosum* Riley (*T. preW*) were obtained by feeding starved one-

day-old wasps a 2mg/ml solution of the antibiotic, rifampicin, mixed with honey. The population has been maintained in our laboratory for a long time over 20 generations (ca. 200 days) after collection from the field. The presence of *Wolbachia* in the females was detected by using forward primer 5'-TGGTCCAATAAGTGATGAAGAAAC-3' and reverse primer 5'-AAAAATTAAACGCTACTCCA-3' for the *wsp* gene of *T. pretiosum* strains. The wasps were reared on fresh eggs of *Corcyra cephalonica* Stainton (< 24h) that were scattered on paper cards coated with Arabic gum and irradiated with ultraviolet (UV) lamp (30W, TUV30W; Philips; Amsterdam, the Netherlands) for 30 min. A group of 30 females were introduced in a finger tubes (1.8cm×5cm) containing a host egg card with ca. 600 *C. cephalonica* eggs for 2 h. Thereafter, the females were removed and the host eggs were reared for ca.10 days until the emergence of wasps. The *T. preW*⁺ have been cultured in laboratory conditions (25 ± 1 °C, 70 ± 5% relative humidity (RH) with a 16 : 8 h light : dark photoperiod) in growth chamber (MLR-352H-PC, Panasonic, Osaka, Japan) before use.

Maternal temperature treatment

Four elevated temperatures: 27°C, 29°C, 31°C, 33°C, were used in this study. And 3 repetitions were performed for each temperature treatment. In addition, no intersex were found at 25°C in *T. preW*, so they are not described in this paper. Take *T. preW* as the control group. *T. preW*⁺ is the experimental group.

We firstly selected 25 newly emerged females for each temperature and evenly separated them into 5 finger tubes (1.8cm×5cm) with surplus *C. cephalonica* eggs glued on paper cards. These wasps were allowed to lay eggs within 2 hours at room temperature. After that, the finger tubes were placed at 27°C, 29°C, 31°C, and 33°C in the growth chamber (MLR-352H-PC, Panasonic, Osaka, Japan). The subsequently emerged wasps (F0 generation) were maintained under

laboratory conditions and supplied with superfluous *C. cephalonica* eggs and 50% honey-water. After 3 days, the wasps were gently removed. Then the emerged F1 generation wasps were allowed to reproduce to get F2 generation. The 1-day-old wasps of F1 or F2 were killed under -80°C conditions for observation. The intersexes were discriminated under the stereomicroscope lens (Motic SMZ-161; Motic; China). The external characteristics of intersex individuals from F1 and F2 generations were observed and described, and the intersex number in each generation was counted. The intersex individuals were determined based on external characteristics with Tulgetsk's method (Tulgetsk *et al.*, 2012). Based on the characteristics of antennae and external genitals, we were able to classify the intersexes into 11 morphological categories (from level 1 to level 11 which was labeled by A to K respectively) with increasing masculinization (Fig. 1). The number of intersex in each level was counted as well.

Sample preparation and photograph

In order to obtain the pictures of all masculinization levels of intersex and make the size of the sample remains unchanged for a long time, the intersex samples were fixed 2 hours in Carnoy's solution (methanol : glacial acetic acid = 3 : 1). A step by step dehydration was performed with different gradient ethanol, 10-20 min each. Subsequently, the solution in different proportions of ethanol and glycerol was treated step by step for only 1 time with a span of 30 min. Finally, the samples were treated with pure glycerol twice for 30 min each time. Then, samples were stored in the refrigerator (HYC-940; Haier; Qingdao, China) at 4°C.

The intersex samples were softened at room temperature. To observe the characteristics of antennae and external genitals, we first removed the wings and feet from the body and then the head and abdomen were cut off by a scalpel under the stereomicroscope lens (Motic SMZ-161;

Motic; China). Thereafter, the samples were photographed with the camera (Nikon SMZ25; Nikon; Japan).

Data analysis

The probability of the intersex (binomial distribution) and the level of masculinization of the intersex (poisson distribution) as influenced by generations and temperature were analyzed by the logistic model and generalized log-linear model respectively. The multiple comparisons of all treatment pairs were carried by Tukey-Kramer tests. The homogeneity of all models was tested by studentized Breusch-Pagan test and Shapiro test respectively (Hall, 1992; Koenker, 1981). All analyses were carried out with R ver. 3.5.1 (R Core Team, 2017).

Results

General characterization of intersex individuals produced by *T. preW*⁺

In this study, we found 235 intersexes from 83699 offspring wasps produced by female of *T. preW*⁺ infected *Wolbachia* under different temperatures (Supplemental Information S1). Based on external characters, we were able to classify the intersex into 11 morphological categories (A-K) with increasing masculinization (Fig. 1). The external genitalia of most intersexes were female, and only I, J, K intersex types have male external genitalia(Supplemental Information S2). In addition, the intersexes were not found in the population of *T. preW* (Supplemental Information S1).

The probability of intersex as influenced by temperature

The probability of intersex was significantly influenced by generations ($\chi^2 = 17.16$, $P < 0.001$) or temperature ($\chi^2 = 143.13$, $P < 0.001$), but it was not influenced by the interaction of generations

and temperature ($\chi^2 = 0.68$, $P = 0.41$) (Fig. 2). The probability of intersex increased with temperature (Coefficient \pm SE = 0.40 ± 0.036 , $z = 11.10$, $P < 0.001$). The average probability of intersex in F1 (27°C: 0.17 ± 0.12 % (Mean \pm SE); 29°C: 0.51 ± 0.25 %; 31°C: 1.43 ± 0.40 %; 33°C: 1.41 ± 0.52 %) was significantly higher ($z = 4.70$, $P < 0.001$) than it was in F2 (27°C: 0.07 ± 0.016 %; 29°C: 0.26 ± 0.040 %; 31°C: 0.25 ± 0.029 %; 33°C: 1.06 ± 0.12 %).

The degree of secondary masculinization level of intersex individuals in *T. preW*⁺

The masculinization level of intersex individuals were significantly influenced by temperature ($\chi^2 = 8.25$, $P < 0.01$), but they were not influenced by generations ($\chi^2 = 0.37$, $P = 0.54$) and the interaction of generations and temperature ($\chi^2 = 0.56$, $P = 0.45$) (Fig. 3). The masculinization level of intersex individuals significant increased with temperature (Coefficient \pm SE = 0.039 ± 0.014 , $z = 2.85$, $P < 0.01$).

Discussion

In order to determine the causes of intersex formation in *T. preW*⁺ and the effects of parental generation temperature of gradual changes in the masculinization of secondary sexual characteristics in intersex offspring, we focused on sexually aberrant intersexes and found that intersex offspring occurs after the *T. preW*⁺ parental generation experiences high temperature. The parental generation temperature affects the probability of occurrence of intersex offspring and the degree of masculinization in secondary sexual characteristics in intersexes. A large volume of studies has showed that high temperature will decrease *Wolbachia* titers inside *Trichogramma* wasps and produce males and intersexes, thereby changing the original reproduction method and offspring sex ratio in *Trichogramma* wasps (Pintureau *et al.*, 1999;

206 Zchori-Fein *et al.*, 2000; Ma & Schwander, 2017). In addition, it is found that there is no intersex
 207 in the offspring in *Wolbachia*-uninfected *T. pretiosum* in this paper. The following conclusions
 208 can be obtained: *Wolbachia* is a direct factor that causes the occurrence of intersexes, while high
 209 temperature is an indirect factor that determines the external morphology of intersexes.

210 All intersexes are caused by sex determination and aberrant sexual differentiation. Therefore, a
 211 strict classification of intersex types has low significance (Gusmao & McKinnon, 2009; Narita *et*
 212 *al.*, 2010). Hence, our study did not compare the external morphology of different types of
 213 intersexes but examined the relationship between parental generation temperature and different
 214 offspring with intersexes through calculating the probability of intersex occurrence. Results
 215 showed that *T. preW*⁺ parental generation temperature is positively correlated with the
 216 probability of intersex occurrence. The probability of intersex occurrence in the F1 generation is
 217 significantly higher than that of the F2 generation. This shows that the effects of high
 218 temperature on *T. preW*⁺ intersexes is passed through the parental generation to offspring, and
 219 this maternal effect weakens as the number of generations increases. A previous study found that
 220 high temperature can affect *Wolbachia* titers to affect the sex of offspring (Sakamoto *et al.*,
 221 2008; Narita *et al.*, 2007; Sugimoto *et al.*, 2015). If this is true, it is not difficult to imagine that
 222 the effect of high temperature on the first generation is stronger, causing lower *Wolbachia* titers.

223 At this point, the effect of sex determination mechanisms of the host is relatively stronger than
 224 feminization induced by *Wolbachia*, and it can causes stronger genetic conflict between
 225 *Wolbachia* and host genes, and the number of intersex individuals increases. In contrast, the
 226 weakening of the effects of high temperature on the second generation combined with self-
 227 recovery of *Wolbachia* increases *Wolbachia* titers. This causes *Wolbachia* to gradually restore
 228 the ability that control host reproductive patterns, and the number of intersex individuals

decreases. However, another potential explanation is that intersexes may be induced in some Hymenoptera species with complementary sex determination (CSD) mechanism (Tuljatske & Stouthamer, 2012). CSD is a sex-determining mechanism of alleles at sex locus determine the sex of offspring (Vorburger C, 2014). However, our unpublished results has showed that the sex determination mechanisms for *Trichogramma* wasps does not conform to complementary sex determination (CSD) but may conform to a form of genome imprinting (QQ Liu, 2018-2019, personal observations). The genome imprinting model has been described in *Nasonia vitripennis* (Heimpel & de Boer, 2008). In this model, a maternally active sex-determination gene imprints a sex-determination gene (potentially the same gene) of fertilized eggs. Haploid eggs contain only a maternally derived, and thus imprinted, sex-determination gene and develop into males. Diploid eggs contain a paternally derived nonimprinted copy of the sex determination gene in addition and develop as diploid females. Genomic imprinting is often caused by DNA methylation in organisms, but not yet been found in hymenopteran insects(Heimpel & de Boer, 2008). With regards to *T. preW*⁺ in this study, if we assume that genome imprinting is retained in the offspring by the parental generation and the presence of *Wolbachia* inhibits the effects of genome imprinting, when temperature reduces *Wolbachia* titers, the effects of parental generation genome imprinting will become stronger. This may be the cause of intersex formation.

In this study, the degree of masculinization in primary and secondary sexual characteristics in intersexes was used to identify 11 intersex types. In addition, we found that the frequency of the D (one male and one female antenna, male antennae contain long setae, female antennae do not contain bristles, external genitalia present in females) and H types (two male antennae that possess long setae, female external genitalia present) of intersex was the highest. The external

genitalia of most intersex types were female, and only 3 intersex types have male external genitalia. In *T. preW⁺* intersex individuals, most exhibit female primary sexual characteristics, and secondary sexual characteristics exhibit signs of masculinization. Tulgetske et al. (2012) found that the degree of masculinization in secondary sexual characteristics of intersexes is randomly distributed and not associated with high temperature. Another paper has showed that high temperature will cause feminization of secondary sexual characteristics in *Culex stimulans* intersexes with a male genotype (Anderson & Horsfall, 1965). In contrast, high temperature can also induce masculinization of secondary sexual characteristics in *Trichogramma deion* (Bowen & Stern, 1966), *S. triquetrella* and *Carausius morosus* intersexes (Narita et al., 2010). In our study, we found that the degree of masculinization of secondary sexual characteristics in *T. preW⁺* intersexes increases as parental generation temperature increases. The differences in results may be due to the differences in sex determination mechanisms in different hosts and the presence of *Wolbachia*. The genetic conflict between host genes and *Wolbachia* may also be the cause of intersex development. (Rigaud & Juchault, 1993; Rigaud & Juchault, 1998; Stouthamer et al., 2010).

In summary, high temperature affects *Wolbachia* in the parental generation to affect sex changes in the host offspring, and the maternal effect that causes this phenomenon decreases as the number of generations increases. Many papers on insect intersexes in the order Hymenoptera have focused on external morphology and behavior and less on sex-determining genes. As the frequency of intersex occurrence in *T. preW⁺* is low, and *T. preW⁺* individuals are small, it is relatively difficult to collect or dissect these intersexes and determine whether all intersex can reproduce. More studies should be carried in future. Therefore, we only studied the external genitalia of intersexes in this paper. In the future, in-depth research on internal genitalia and the

fertility of *T. preW*⁺ can be carried out to facilitate the understanding of intersex formation mechanisms and a deeper understanding of sex determination and differentiation in the order Hymenoptera.

Conclusions

we focused on sexually aberrant intersexes and found that intersex offspring occurs after the *T. preW*⁺ parental generation experiences high temperature. The following conclusions can be obtained: *Wolbachia* is a direct factor that causes the occurrence of intersexes, while high temperature is an indirect factor that determines the external morphology of intersexes. This change of sex may be associated with the changes in the expression levels of sex-determining genes. The changes in the expression levels of sex-determining genes in *T. preW*⁺ that are jointly caused by temperature-*Wolbachia* is the root cause for intersex formation. In view of the aforementioned situation, we recommend that more in-depth research on relevant sex-determining gene regulation mechanisms for intersexes be carried out in the future to determine whether changes in the expression levels of sex-determining genes results in intersex formation to elucidate the mechanisms for intersex formation.

Acknowledgements

We thank Su-ying Bao for supplying *Corcyra cephalonica* strains. We thank Stephen Johnson and two anonymous reviewers for constructive comments on a previous version of this manuscript.

298

299 References

- 300 **Almeida RP, Stouthamer R. 2018.** Phylogeny of the *Trichogramma* endosymbiont *Wolbachia*,
301 an alpha-proteobacteria (Rickettsiae). *Brazilian Journal of Biology* **78**:421–428 DOI
302 10.1590/1519-6984.166671.
- 303 **Anderson JF, Horsfall WR. 1965.** Dimorphic development of transplanted juvenile gonads of
304 mosquitoes. *Science* **147**:642–645.
- 305 **Bear A, Monteiro A. 2013.** Both cell-autonomous mechanisms and hormones contribute to
306 sexual development in vertebrates and insects. *Bioessays* **35**:725–732 DOI
307 10.1002/bies.201300009.
- 308 **Beukeboom LW, Kamping A. 2006.** No patrigenes required for femaleness in the haplodiploid
309 wasp *Nasonia vitripennis*. *Genetics* **172**:981–989 DOI 10.1534/genetics.105.044743.
- 310 **Bowen WR, Stern VM. 1966.** Effect of temperature on the production of males and sexual
311 mosaics in a uniparental race of *Trichogramma*. *Annals of the Entomological Society of America*
312 **59**:823–834 DOI 10.1093/aesa/59.4.823.
- 313 **Brust RA. 1966.** Gynandromorphs and intersexes in mosquitoes (Diptera:Culicidae). *Canadian*
314 *Journal of Zoology* **44**:911–921 DOI 10.1139/z66-093.
- 315 **Cline TW. 1984.** Autoregulatory functioning of a *Drosophila* gene product that establishes and
316 maintains the sexually determined state. *Genetics* **107**:231–277 DOI 10.1007/BF00056440.
- 317 **Fukui T, Kawamoto M, Shoji K, Kiuchi T, Sugano S, Shimada T, Suzuki Y, Katsuma S.**
318 **2015.** The Endosymbiotic Bacterium *Wolbachia* Selectively Kills Male Hosts by Targeting the
319 Masculinizing Gene. *PLoS Pathog* **11**:e1005048 DOI 10.1371/journal.ppat.1005048.

- 320 **Goldschmidt RB. 1949.** The Intersexual Males of the Beaded Minute Combination in
321 *Drosophila Melanogaster*. *Proceedings of the National Academy of Sciences of the United States*
322 *of America* **35**:314–316 DOI 10.1073/pnas.35.6.314.
- 323 **Gusmao LFM, McKinnon AD. 2009.** Sex ratios, intersexuality and sex change in copepods.
324 *Journal of Plankton Research* **31**:1101–1117 DOI 10.1093/plankt/fbp059.
- 325 **Hall AD. 1992.** A study of various score test statistics for heteroscedasticity in the general linear
326 model. *Mathematics & Computers in Simulation* **33**:563–568 DOI 10.1016/0378-
327 4754(92)90153-8.
- 328 **Heimpel GE, de Boer JG. 2008.** Sex determination in the Hymenoptera. *Annu Rev Entomol*
329 **53**:209–230 DOI 10.1146/annurev.ento.53.103106.093441.
- 330 **Kent BN, Funkhouser LJ, Setia S, Bordenstein SR. 2011.** Evolutionary genomics of a
331 temperate bacteriophage in an obligate intracellular bacteria (*Wolbachia*). *PLoS One* **6**:e24984
332 DOI 10.1371/journal.pone.0024984.
- 333 **Koenker R. 1981.** A note on studentizing a test for heteroscedasticity. *Journal of Econometrics*
334 **17**:107–112 DOI 10.1016/0304-4076(81)90062-2.
- 335 **Ma WJ, Pannebakker BA, van de Zande L, Schwander T, Wertheim B, Beukeboom LW.**
336 **2015.** Diploid males support a two-step mechanism of endosymbiont-induced thelytoky in a
337 parasitoid wasp. *BMC Evolutionary Biology* **15**:84–96 DOI 10.1186/s12862-015-0370-9.
- 338 **Ma WJ, Schwander T. 2017.** Patterns and mechanisms in instances of endosymbiont-induced
339 parthenogenesis. *J Evol Biol* **30**:868–888 DOI 10.1111/jeb.13069.
- 340 **Martin P, and Scholtz G. 2012.** A case of Intersexuality in the Parthenogenetic Marmorkrebs
341 (Decapoda: Astacida: Cambaridae). *J Biomater Sci Polym Ed* DOI 10.1163/193724012X629031.

342 **Narita S, Kageyama D, Nomura M, Fukatsu T. 2007.** Unexpected mechanism of symbiont-
 343 induced reversal of insect sex: feminizing *Wolbachia* continuously acts on the butterfly *Eurema*
 344 *hecabe* during larval development. *Appl Environ Microbiol* **73**:4332–4341 DOI
 345 10.1128/AEM.00145-07.

346 **Narita S, Pereira R, Kjellberg F, Kageyama D. 2010.** Gynandromorphs and intersexes:
 347 potential to understand the mechanism of sex determination in arthropods. *Terrestrial Arthropod*
 348 *Reviews* **3**:63–96 DOI 10.1163/187498310x496190.

349 **Perlman SJ, Kelly SE, Hunter MS. 2008.** Population biology of cytoplasmic incompatibility:
 350 maintenance and spread of *Cardinium* symbionts in a parasitic wasp. *Genetics* **178**:1003–1011
 351 DOI 10.1534/genetics.107.083071.

352 **Pintureau B, Chapelle L, Delobel B. 1999.** Effects of repeated thermic and antibiotic
 353 treatments on a *Trichogramma* (Hym., *Trichogrammatidae*) symbiont. *Journal of Applied*
 354 *Entomology* **123**:473–483 DOI 10.1046/j.1439-0418.1999.00412.x.

355 **Pintureau B, Lassabliere F, Daumal J, Grenier S. 2002.** Does a cyclic natural thermal cure
 356 occur in *Wolbachia*-infected *Trichogramma* species. *Ecological Entomology* **27**:366–372 DOI
 357 10.1046/j.1365-2311.2002.00416.x.

358 **Pintureauand B, Bolland P. 2001.** A *Trichogramma* Species Showing a Better Adaptation to
 359 High Temperature than its Symbionts. *Biocontrol Science and Technology* **11**:13–20 DOI
 360 10.1080/09583150020029691.

361 **R Core Team. 2017.** R: A language and environment for statistical computing. R Foundation for
 362 Statistical Computing, Vienna, Austria.<http://www.R-project.org/>.

363 **Rigaud T, Juchault P. 1993.** Conflict Between Feminizing Sex Ratio Distorters and an
 364 Autosomal Masculinizing Gene in the Terrestrial Isopod *Armadillidium vulgare* Latr. *Genetics*
 365 **133**:247–252 DOI 10.1016/1050-3862(93)90021-A.

366 **Rigaud T, Juchault P. 1998.** Sterile intersexuality in an isopod induced by the interaction
 367 between a bacterium (*Wolbachia*) and the environment. *Canadian Journal of Zoology* **76**:493–
 368 499 DOI 10.1139/z97-216.

369 **Sakamoto H, Kageyama D, Hoshizaki S, Ishikawa Y. 2008.** Heat Treatment of the Adzuki
 370 Bean Borer, *Ostrinia scapularis* Infected with *Wolbachia* gives Rise to Sexually Mosaic
 371 Offspring. *Journal of Insect Science* **8**:1–5 DOI 10.1673/031.008.6701.

372 **Schärer L, Rowe L, Arnqvist G. 2012.** Anisogamy, chance and the evolution of sex roles.
 373 *Trends in Ecology & Evolution* **27**:260–264 DOI 10.1016/j.tree.2011.12.006.

374 **Schilthuizen MO, Stouthamer R. 1997.** Horizontal transmission of parthenogenesis-inducing
 375 microbes in *Trichogramma* wasps. *Proc Biol Sci* **264**: 361–366 DOI 10.1098/rspb.1997.0052.

376 **Schutt C, Nöthiger R . 2000.** Structure, function and evolution of sex determining systems in
 377 Dipteran insects. *Development* **127**: 667–677 DOI 10.1007/PL00008234.

378 **Seiler J, Puchta O, Brunold E. 1958.** Die Entwicklung des Genitalapparates bei triploiden
 379 Intersexen von *Solenobia triquetrella* F.R. (Lepid. Psychidae). Deutung des
 380 Intersexualitätsphänomens. Wilhelm Roux' Archiv für Entwicklungsmechanik der Organismen
 381 **150**: 190–372.

382 **Sigsgaard L, Herz A, Korsgaard M, Wührer B. 2017.** Mass Release of *Trichogramma*
 383 evanescens and *T. cacoeciae* Can Reduce Damage by the Apple Codling Moth *Cydia pomonella*
 384 in Organic Orchards under Pheromone Disruption. *Insects* **8**:41–49 DOI
 385 10.3390/insects8020041.

386 **Stouthamer R, Russell JE, Vavre F, Nunney L. 2010.** Intra-genomic conflict in populations
 387 infected by Parthenogenesis Inducing *Wolbachia* ends with irreversible loss of sexual
 388 reproduction. *BMC Evolutionary Biology* **10**:229 DOI 10.1186/1471-2148-10-229.

389 **Sugimoto TN, Kayukawa T, Matsuo T, Tsuchida T, Ishikawa Y. 2015.** A short, high-
 390 temperature treatment of host larvae to analyze *Wolbachia*-host interactions in the moth *Ostrinia*
 391 *scapularis*. *Journal of Insect Physiology* **81**:48–51 DOI 10.1016/j.jinsphys.2015.06.016.

392 **Tulgettske GM, Stouthamer R. 2012.** Characterization of intersex production in *Trichogramma*
 393 *kaykai* infected with parthenogenesis-inducing *Wolbachia*. *Naturwissenschaften* **99**:143–152
 394 DOI 10.1007/s00114-011-0880-2.

395 **Tulgettske, Michelle G. 2010.** Investigations into the Mechanisms of *Wolbachia*-Induced
 396 Parthenogenesis and Sex Determination in the Parasitoid Wasp, *Trichogramma*. Ph.D. Thesis,
 397 University of California.

398 **Vorburger C. 2014.** Thelytoky and Sex Determination in the
 399 *Hymenoptera*: Mutual Constraints. *Sexual Development* **8**:50–58 DOI 10.1159/000356508.

400 **Werren JH, Windsor DM. 2000.** *Wolbachia* infection frequencies in insects: evidence of a
 401 global equilibrium? Proceedings. *Biological Sciences* **267**:1277–1285 DOI
 402 10.1098/rspb.2000.1139.

403 **Whiting PW, Greb RJ, Speicher BR. 1934.** A New Type of Sex-Intergrade. *Biological Bulletin*
 404 **66**:152–165 DOI 10.2307/1537329.

405 **Zchori-Fein E, Gottlieb Y, Coll M. 2000.** *Wolbachia* density and host fitness components in
 406 *Muscidifurax uniraptor* (Hymenoptera: Pteromalidae). *Journal of Invertebrate Pathology* **75**:
 407 267–272 DOI 10.1006/jipa.2000.4927.

Figure 1

Description of the external morphology of 11 types of intersexes in *T. preW*⁺.

(1) Picture A to X: Female intersex, the external genitalia are female and the antennae tend to masculinized. The picture A, D, G, J, M, P, S, V represents the ventral (Secondary sexual characteristics), the picture B, E, H, K, N, Q, T, W represents the antennae (Secondary sexual characteristics) and C, F, I, L, O, R, U, X represents the external genitalia (Primary sexual characteristics) respectively. Type A of intersex: Picture A to C (A: the ventral of intersex; B: 2 female antennae, 1 with few short setae; C: female genitalia). Type B of intersex: Picture D to F (D: the ventral of intersex; E: 2 female antennae, both with few short setae; F: female genitalia). Type C of intersex: Picture G to I (G: the ventral of intersex; H: 1 male antenna with few short setae, 1 female antenna; I: female genitalia). Type D of intersex: Picture J to L (J: the ventral of intersex; K: 1 male antenna with few long setae, 1 female antenna; L: female genitalia). Type E of intersex: Picture M to O (M: the ventral of intersex; N: 1 male antenna with few long setae, 1 female antenna with few short setae; O: female genitalia). Type F of intersex: Picture P to R (P: the ventral of intersex; Q: 2 male antennae, both with short setae; R: female genitalia). Type G of intersex: Picture S to U (S: the ventral of intersex; T: 2 male antennae, 1 with few long setae, 1 with few short setae; U: female genitalia). Type H of intersex: Picture V to X (V: the ventral of intersex; W: 2 male antennae, both with long setae; X: female genitalia).

(2) Picture Y to GG: Male intersex, the external genitalia are male and the antennae tend to feminized. The picture Y, BB, EE represents the ventral (Secondary sexual characteristics), the picture Z, CC, FF represents the antennae (Secondary sexual characteristics) and AA, DD, GG represents the external genitalia (Primary sexual characteristics) respectively. Type I of intersex: Picture Y to AA (Y: the ventral of intersex; Z: 2 female antennae; AA: male genitalia). Type J of intersex: Picture BB to DD (BB: the ventral of intersex; CC: 1 male antenna with long setae, 1 female antenna; DD: male genitalia). Type K of intersex: Picture EE to GG (EE: the ventral of intersex; FF: 1 male antenna with long setae, 1 female antenna with few short setae; GG: male genitalia).

(3) Picture HH to KK: As control group, is the normal female in *T. preW* and *T. preW*⁺, normal male in *T. preW*. The picture HH is the lateral of female and II is the lateral of male. The picture JJ and KK represents ventral of female and male respectively. The external morphology of female is 2 female antennae, none of setae, female genitalia. The external morphology of male is 2 male antennae, both with long setae, male genitalia.




Figure 2 (on next page)

The effects of different temperature in the parental generation (27°C, 29°C, 31°C, and 33°C) on the probability of intersex in the F1 and F2 generation in *T. preW*⁺.

The blue line indicates the F2 generation and the red line indicates the F1 generation.




Figure 3(on next page)

The degree of masculinization level of intersexes in *T. preW*⁺.

The left ordinate of the graph represents the masculinization level of intersex individuals. The right ordinate of the graph represents the types of intersexes. The shaded area said confidence interval.

The masculinization level of intersex individuals

