

# Comparative transcriptome analysis of panicle development under heat stress in two rice (*Oryza sativa* L.) cultivars differing in heat tolerance

Yaliang Wang<sup>Equal first author, 1</sup>, Yikai Zhang<sup>Equal first author, 1</sup>, Qiang Zhang<sup>1</sup>, Yongtao Cui<sup>1</sup>, Jing Xiang<sup>1</sup>, Huizhe Chen<sup>1</sup>, Guohui Hu<sup>1</sup>, Yanhua Chen<sup>1</sup>, Xiaodan Wang<sup>1</sup>, Defeng Zhu<sup>Corresp., 1</sup>, Yuping Zhang<sup>Corresp. 1</sup>

<sup>1</sup> State Key Laboratory of Rice Biology, China National Rice Research Institute, Hangzhou, Zhejiang, China

Corresponding Authors: Defeng Zhu, Yuping Zhang  
Email address: cnrice@qq.com, cnrrizyp@163.com

Heat stress inhibits rice panicle development and reduces the spikelet number per panicle. This study investigated the mechanism involved in heat-induced damage to panicle development and spikelet formation in rice cultivars that differ in heat tolerance. Transcriptome data from developing panicles grown at 40°C or 32°C were compared for two rice cultivars: heat-tolerant Huanghuazhan and heat-susceptible IR36. Of the differentially expressed genes (DEGs), 4,070 heat stress-responsive genes were identified, including 1,688 heat-resistant-cultivar-related genes (RHR), 707 heat-susceptible-cultivar-related genes (SHR), and 1,675 common heat stress-responsive genes. A Gene ontology (GO) analysis showed that the DEGs in the RHR category were significantly enriched in 54 gene ontology terms, some of which improved heat tolerance, including those in the WRKY, HD-ZIP, ERF, and MADS transcription factor families. A Kyoto Encyclopedia of Genes and Genomes (KEGG) analysis showed that the DEGs in the RHR and SHR categories were enriched in 15 and 11 significant metabolic pathways, respectively. Improved signal transduction capabilities of endogenous hormones under high temperature seemed to promote heat tolerance, while impaired starch and sucrose metabolism under high temperature might have inhibited young panicle development. Our transcriptome analysis provides insights into the different molecular mechanisms of heat stress tolerance in developing rice.

# Comparative transcriptome analysis of panicle development under heat stress in two rice (*Oryza sativa* L.) cultivars differing in heat tolerance

Yaliang Wang<sup>#</sup>, Yikai Zhang<sup>#</sup>, Qiang Zhang, Yongtao Cui, Jing Xiang, Huizhe Chen, Guohui Hu, Yanhua Chen, Xiaodan Wang, Defeng Zhu\*, Yuping Zhang\*

State Key Laboratory of Rice Biology, China National Rice Research Institute, Hangzhou, Zhejiang, China

<sup>#</sup>The first two authors contributed equally to this paper

\*Corresponding authors:

Defeng Zhu, E-mail: [cnrice@qq.com](mailto:cnrice@qq.com);

Yuping Zhang, E-mail: [cnrrizyp@163.com](mailto:cnrrizyp@163.com)

## Abstract

Heat stress inhibits rice panicle development and reduces the spikelet number per panicle. This study investigated the mechanism involved in heat-induced damage to panicle development and spikelet formation in rice cultivars that differ in heat tolerance. Transcriptome data from developing panicles grown at 40°C or 32°C were compared for two rice cultivars: heat-tolerant Huanghuazhan and heat-susceptible IR36. Of the differentially expressed genes (DEGs), 4,070 heat stress-responsive genes were identified, including 1,688 heat-resistant-cultivar-related genes (RHR), 707 heat-susceptible-cultivar-related genes (SHR), and 1,675 common heat stress-responsive genes. A Gene ontology (GO) analysis showed that the DEGs in the RHR category were significantly enriched in 54 gene ontology terms, some of which improved heat tolerance, including those in the WRKY, HD-ZIP, ERF, and MADS transcription factor families. A Kyoto Encyclopedia of Genes and Genomes (KEGG) analysis showed that the DEGs in the RHR and SHR categories were enriched in 15 and 11 significant metabolic pathways, respectively. Improved signal transduction capabilities of endogenous hormones under high temperature seemed to promote heat tolerance, while impaired starch and sucrose metabolism under high temperature might have inhibited young panicle development. Our transcriptome analysis provides insights into the different molecular mechanisms of heat stress tolerance in developing rice.

Key words: Rice; Panicle development; Heat stress; Transcriptome analysis

## Introduction

Climate change is predicted to increase the average global temperatures by 0.3-4.8°C by the end of the 21st century (Stocher *et al.* 2013). Unusually high temperatures occur frequently during the rice growing season (Dwivedi *et al.* 2015; Tao *et al.* 2013), and cause reductions in the yield and quality in several rice producing regions, including China, India, and Japan (Anand *et al.* 2018; Morita *et al.* 2016; Wang *et al.* 2019). The primary cause of rice yield reductions is a reduction in spikelet fertility due to high temperatures during the flowering period (Espe *et al.* 2017). Rice quality is also influenced by high temperature, which causes carbohydrate metabolism disorders (Yamakawa & Hakata 2010). As climate change has intensified, extremely high temperatures above 40°C have become more frequent. Such high temperatures inhibit rice panicle development, reduce the spikelet number by 5%-15%, and aggravate rice yield losses (Wang *et al.* 2017).

High temperatures adversely affect floral development by reducing the antioxidant capacity, inhibiting nutrient accumulation, and degenerating tapetal cells (Prasad *et al.* 2017). A previous study showed that high temperature (39°C) conditions downregulated certain genes related to tapetum function, pollen adhesion, and germination, including *OsINV4* and *OsMST8*, which influenced spikelet fertilization (Endo *et al.* 2009). In addition, sugar and endogenous hormone metabolism under high temperature reportedly plays an important role in pollen formation in both rice and cotton (Islam *et al.* 2018; Min *et al.* 2014). At the rice ripening stage, high temperature induces early termination of grain filling (Kim *et al.* 2011). Grain chalkiness increases under a mean temperature greater than 32°C, resulting in the deterioration of eating and cooking quality, which are both closely linked to starch and sucrose metabolism (Zhong *et al.* 2010). Transcriptome analysis has shown that high temperatures influence the expression of genes involved in the inhibition of sucrose degradation and starch biosynthesis while promoting starch degradation and storage proteins synthesis (Yamakawa & Hakata 2010; Yamakawa *et al.* 2007). Takehara *et al.* (2018) reported that the upregulation of *OsSUS3*, which encodes sucrose synthase, improved high-temperature tolerance.

The panicle initiation stage is an important period of spikelet proliferation. Dry matter accumulation is essential for panicle development; however, the pathway for carbohydrate accumulation during spikelet formation under heat stress remains vague. The reduction in spikelet number that occurs under high temperature conditions has been associated with heat-induced phytohormone changes, especially enhanced cytokinin degradation (Wu *et al.* 2017; Wu *et al.*

2016). The number of spikelets per panicle is determined by spikelet differentiation and degeneration. Spikelet differentiation is correlated with dry matter accumulation and influenced by environmental factors (Liu *et al.* 2005). Ding *et al.* (2016) reported that hormone metabolism, the stress response, carbohydrate metabolism and transport, and protein degradation were regulated to influence panicle initiation. Additionally, certain genes, such as MADS-box genes, are related to panicle initiation (Kang *et al.* 2013; Kobayashi *et al.* 2012). Quantitative trait loci for spikelet degeneration have been identified (Yamagishi *et al.* 2004), and the genes *SPI*, *ASPI*, *TUT1*, *PAA2*, and *OsALMT7* have been found to control spikelet degeneration (Bai *et al.* 2015; Heng *et al.* 2018; Li *et al.* 2010). However, the mechanism of panicle development under high temperature conditions is still unclear. In this study, an RNA-Seq analysis was used to explore the mechanism of heat tolerance during panicle development. Huanghuazhan (HHZ) is a heat-tolerant rice cultivar widely grown in the middle and lower reaches of the Yangtze River in China (Cao *et al.* 2009; Zhou *et al.* 2012). IR36 is a heat-susceptible cultivar (Fang *et al.* 2006) and a parental line of HHZ. In the current study, we investigated transcriptome differences between these two cultivars exposed to different temperatures 40°C and 32°C, during the spikelet differentiation stage. We identified differentially expressed genes (DEGs) in young panicles of the two cultivars under the two temperature treatments and performed Gene Ontology (GO) enrichment and the Kyoto Encyclopedia of Genes and Genomes (KEGG) analysis. This work improves our understanding of the molecular mechanism underlying the heat-induced inhibition of spikelet development and provides important insights into rice breeding.

## Materials and methods

### Plant materials and heat stress treatments

We used the rice cultivars HHZ and IR36 in this study. Pre-germinated seeds were sown in seed trays filled with a mixture of vermiculite (20%), charcoal (30%), soil (40%), and slow-release fertilizer (10%). After 20 days, the seedlings were transplanted into pots with four seedlings per pot. Each pot (24 cm length × 22.5 cm width × 21.5 cm height) contained 10 kg air-dried paddy soil. Pots were kept under natural environmental conditions (the average temperature was 30–35°C).

Before seedlings were transplanted into pots, fertilizer was applied to each pot based on a field application rate of 14 kg nitrogen per 666.7 m<sup>2</sup>. Before transplanting into the pots, 3.5 g compound fertilizer (nitrogen: phosphorus: potassium = 15%: 15%: 15%) was applied to each pot. At the tillering stage, 0.6 g urea was supplemented in each pot. At panicle initiation, 0.6 g urea and 0.5 g potassium chloride were also applied to each pot. Pests, diseases, and weeds were intensively controlled.

Automatic growth chambers (Qiushi Environment Corporation, Hangzhou, China) were used to conduct the temperature treatments. Plants were moved to the growth chambers on the approximate date of spikelet differentiation when the panicle length was approximately 0.2 cm (around 60-70 d after seed sowing). The high-temperature (40°C) and control temperature (32°C) treatments were implemented for eight hours each day from 9:30 to 17:30 h (the setting details are shown in Table S1) for nine days. The humidity in the chambers was maintained at 75-80%. Rice plants were grown under natural ambient conditions during all growth stages before and after the temperature treatments. Each treatment contained three replicates with 20 pots per replicate.

### **Panicle and spikelet morphology**

Ten main tillers were sampled per replicate on day 9 of treatment at 40°C or 32°C to investigate the development of young panicles at high temperature.

Spikelet differentiation or degeneration of the main tiller panicles was determined at the heading stage. The number of degenerated spikelets was calculated by counting the vestiges present on the panicles. The number of differentiated spikelets was the sum of the surviving and degenerated spikelets. The proportion of degenerated spikelets was then calculated.

Spikelet morphology was observed under a stereomicroscope (Olympus SZX7, Olympus Corporation, Tokyo, Japan) and the glume length and width (mm) were measured at 0.63x and 2.5x using the microscale in the Image Pro-Plus 5.1 image processing software (Olympus SZX7, Olympus Corporation, Tokyo, Japan). Fifteen spikelets were collected from the upper, middle and lower parts of each panicle, with five panicles sampled for each replicate.

### **RNA extraction, transcriptome sequencing, and mapping**

After nine days at the 40 °C or 32 °C treatment, young panicles from 20 main tillers were collected for each replicate at 12:00-13:00 and immediately frozen in liquid nitrogen. In quick succession, TRIzol reagent (Invitrogen, Carlsbad, CA, USA) was used to extract total RNA from the young panicles according to the manufacturer's instructions. A TruSeq RNA Sample Preparation Kit (Illumina Inc., CA, USA) was used to generate 12 sequencing libraries according to the manufacturer's instructions. The sequencing libraries were then sequenced on a HiSeq platform (Illumina, Inc., CA, USA). High-quality sequence reads were obtained by filtering the raw data and then compared to the 9311-reference genome (*Oryza indica*.ASM465v1.dna.toplevel. fa) obtained from <http://www.ensembl.org/>. The raw RNA sequence data were submitted to the NCBI Sequence Read Archive with accession number PRJNA508820.

### **Gene expression level and differential expression analysis**

We used HTSeq (0.9.1) to statistically compare the read count values of each gene, which represent the original expression of each gene. Fragments per kilobase of transcript per million mapped reads (FPKM) was used to standardize the expression. Next, we used DESeq (1.30.0) to analyze the differential expression of genes with the following screening conditions: an expression difference of  $|\log_2\text{foldChange}| > 1$  and a significant  $P\text{-value} < 0.05$ .

## GO and KEGG enrichment analysis of DEGs

For the GO enrichment analysis of DEGs, we used the Singular Enrichment Analysis tool in AgriGO (<http://bioinfo.cau.edu.cn/agriGO/analysis.php>) with the default parameters, and a  $P\text{-value} \leq 0.05$  indicated significant enrichment. The KEGG enrichment analysis of DEGs was performed using KOBAS software with default parameters and a  $P\text{-value} \leq 0.05$  indicated significant pathway enrichment.

## Verification of RNA-Seq by quantitative real-time PCR (qRT-PCR)

First-strand cDNA was synthesized using ReverTra Ace qPCR RT Master Mix with gDNA Remover (Toyobo, Osaka, Japan) according to the manufacturer's instructions. The qRT-PCR analyses were performed using an Applied Biosystems 7500 Real-Time PCR system with Power SYBR Green PCR Master Mix (Applied Biosystems, Carlsbad, CA, USA). The primers used for qRT-PCR are listed in Supplementary Table S2. The *OsUBQ* gene was used as an internal control. Relative gene expression levels were determined from the equation  $2^{-\Delta\Delta CT}$  (Czechowski *et al.* 2004), where  $\Delta\Delta CT$  represents  $\Delta CT$  (target gene of interest)  $-\Delta CT$  (control gene).

## Statistical analyses

Microsoft Excel 2016 (Microsoft Inc., Redmond, WA, USA) was employed for data collection. The panicle and spikelet morphological data collected for the 40°C and 32°C treatments (mean of three replicates) were statistically analyzed by Student's t-test ( $P < 0.05$ ). Graphs were created using Origin 9.1 (Ver. 9.1, OriginLab, Northampton, MA, USA).

# Results

## Spikelet development at high temperature

A preliminary experiment showed a significant difference in panicle development, which was measured as spike differentiation, after nine days of the high-temperature treatment. The results reported in the current study are consistent with these preliminary findings. Spikelet differentiation inhibited young panicle growth after nine days of the high temperature treatment (Fig. 1). After

the temperature treatments, the panicles required an additional 15-20 days to complete growth.

Compared to the control temperature treatment, the high temperature treatment reduced spikelet survival by 22.3% ( $P < 0.05$ ) for HHZ and 53.6% ( $P < 0.05$ ) for IR36. With high temperature, the number of differentiated spikelets decreased by 9.6% and 33.2% ( $P < 0.05$ ) for HHZ and IR36, respectively, and the proportion of degenerated spikelets significantly increased by 32.3% ( $P < 0.05$ ) and 67.4% ( $P < 0.05$ ), respectively. In addition, the heat treatment reduced the glume length by 10.3% ( $P < 0.05$ ) for HHZ and by 16.0% ( $P < 0.05$ ) for IR36 and reduced the glume width by 12.0% ( $P < 0.05$ ) and 8.0% ( $P < 0.05$ ), respectively. The reductions in spikelet number and size led to reductions in panicle weight of 33.2% ( $P < 0.05$ ) for HHZ and 67.7% ( $P < 0.05$ ) for IR36. The larger reduction in panicle weight in IR36 suggests that high temperature has a greater effect on young panicle development in heat susceptible cultivars (Table 1).

## Transcriptome analysis

Under the 32°C control temperature, a total of 44.2 million and 48.9 million raw reads were obtained from HHZ (referred to as HHZ\_32) and IR36 (referred to as IR36\_32), respectively. Under the 40°C treatment, a total of 45.5 million raw reads were obtained from both HHZ (HHZ\_40) and IR36 (IR36\_40) (Table 2 and Table S3). More than 99.0% clean reads were obtained for the downstream analysis. The results of RNA sequence mapping indicated that 85.8-88.0% of the clean reads could be mapped onto the reference genome and most were uniquely mapped (Table 2).

## Identification of DEGs

To compare the differences between the two cultivars at 40°C and 32°C, we used four comparison groups: HHZ\_32 vs HHZ\_40, IR36\_32 vs IR36\_40, IR36\_40 vs HHZ\_40, and IR36\_32 vs HHZ\_32. DEGs for the four groups were restricted to those with a  $|\log_2\text{fold change}| > 1$  and a  $P\text{-value} < 0.05$ . With these criteria, 3,342, 2,469, 2,949, and 2,461 DEGs were detected for HHZ\_32 vs HHZ\_40, IR36\_32 vs IR36\_40, IR36\_40 vs HHZ\_40, and IR36\_32 vs HHZ\_32, respectively (Fig. 2). Significantly different gene expression was observed both between cultivars and between treatments. For HHZ, 1,794 genes were upregulated and 1,548 genes were downregulated in the 40°C treatment compared with the 32°C treatment (Fig. 2). Furthermore, 1,140 genes were upregulated and 1,329 genes were downregulated in IR36 under the 40°C treatment compared with the 32 °C treatment (Fig. 2). For comparisons within treatments, 1,408 genes were upregulated and 1,541 were downregulated in the IR36\_40 vs HHZ\_40 group and 893 genes were upregulated and 1,751 genes were downregulated in the IR36\_32 vs HHZ\_32 group (Fig.2c and 2d).

# Classification of DEGs

In all four groups, a total of 5,533 unique DEGs were identified, and they could be divided into 15 disjointed subgroups (Fig. 3). Among the 15 subgroups, eight from the IR36\_32 vs HHZ\_32 group were excluded from the analysis because they were not influenced by high temperature. In addition, 1,157, 603, 524, and 402 DEGs were uniquely identified in the HHZ\_32 vs HHZ\_40, IR36\_32 vs IR36\_40, IR36\_40 vs HHZ\_40, and IR36\_32 vs HHZ\_32 groups, respectively. The DEGs in groups that were responsive to high temperature could be further sorted into three categories: heat-tolerance-cultivar-related genes (RHR, 1,688 genes), heat-susceptible-cultivar-related genes (SHR, 707 genes), and common heat stress-response genes (CHR, 1,675 genes) (Table 3 and Table S4). The DEGs in the RHR category might have played an important role in heat tolerance, whereas the DEGs in the SHR category might be associated with heat injuries in the heat-susceptible cultivar.

# Analysis of GO annotation

The purpose of the GO enrichment analysis was to obtain GO functional terms with significant enrichment of DEGs and thus reveal the possible functions of the DEGs. Of all DEGs, 2,307 (69.0%), 1,680 (68.0%), 1,832 (62.1%), and 1,472 (59.8%) DEGs were enriched in GO terms in HHZ\_32 vs HHZ\_40, IR36\_32 vs IR36\_40, IR36\_40 vs HHZ\_40, and IR36\_32 vs HHZ\_32 groups, respectively. There were 75, 11, 13, and 31 significant GO terms observed in HHZ\_32 vs HHZ\_40, IR36\_32 vs IR36\_40, IR36\_40 vs HHZ\_40, and IR36\_32 vs HHZ\_32, respectively (Fig. 4). The maximum number of DEGs was observed for the heterocycle biosynthetic process in the IR36\_40 vs HHZ\_40 group. In IR36\_32 vs IR36\_40 and HHZ\_32 vs HHZ\_40, the DEGs were enriched in the terms response to stimulus, response to temperature stimulus, and response to heat in the biological process category. Within the cellular component category, the DEGs were commonly enriched in the terms chromatin, DNA packaging complex, and nucleosome in the IR36\_32 vs IR36\_40 and HHZ\_32 vs HHZ\_40 groups. However, there were no common GO terms in the category of molecular function in the IR36\_32 vs IR36\_40 and HHZ\_32 vs HHZ\_40 groups.

We further identified GO term categories for DEGs in the RHR, SHR, and CHR categories (Fig. 5 and Table S5). Among the 1,689 DEGs in RHR, 54 significant GO terms were detected. However, no significant GO terms were observed among the 707 DEGs in SHR. In CHR, 30 significant GO terms were detected. In the CHR group, eight significant GO terms were observed in the biological process category, including response to stimulus, response to temperature stimulus, and response to heat; 17 GO terms were in the cellular component category; and two significant GO terms were in the molecular function category. In the RHR group, 30, 14 and 10 significant GO terms were in the biological process, cellular component, and molecular function



categories, respectively. The most significant GO terms, in decreasing order, were RNA biosynthetic process, nucleus, and DNA binding. In the molecular function category, 50 DEGs were specifically assigned to DNA-binding transcription factor activity, which may play an important role in heat stress tolerance.

The 50 DEGs of DNA-binding transcription factor activity could be divided into 11 transcription factor (TF) families, including HSF (1), WRKY (6), MADS (12), HD-ZIP (7), GATA (3), ERF (12), ABAI (1), b-ZIP (4), ARR-B (2), E2F (1), and NF-YA (1). Expression of the genes *BGIOSGA006348* of HSF, *BGIOSGA010835* of ABAI, *BGIOSGA010142* of HAP, and *BGIOSGA000303* and *BGIOSGA000304* of ARR-B was significantly upregulated. In addition, five genes in WRKY, eight genes in MADS, two genes in HD-ZIP, two genes in GATA, six genes in ERF, and two genes in b-ZIP were also upregulated (Table S6). These results suggest that, these 30 TF genes may play important roles in heat stress resistance.

### Analysis of KEGG pathway enrichment

In the KEGG analysis, 1,158 DEGs were classified into 225, 191, 239, and 211 functional pathways in HHZ\_32 vs HHZ\_40; 838 DEGs in IR36\_32 vs IR36\_40; 732 DEGs in IR36\_40 vs HHZ\_40; and 539 DEGs in IR36\_32 vs HHZ\_32. A total of 79 pathways were significant ( $P$ -value  $< 0.05$ ) (Fig. 6). Among these pathways, the phenylpropanoid biosynthesis pathway was common in HHZ\_32 vs HHZ\_40, IR36\_32 vs IR36\_40, and IR36\_40 vs HHZ\_40, which suggests that heat stress impaired phenylpropanoid biosynthesis.

Based on further analysis of the three categories with different heat-stress responses, 146 DEGs in RHR were involved in 15 overrepresented pathways, including purine metabolism, pyrimidine metabolism, and amino sugar and nucleotide sugar metabolism; 45 DEGs in SHR were involved in 11 overrepresented pathways, including arginine biosynthesis, starch and sucrose metabolism, and polyketide sugar unit biosynthesis; and 184 DEGs in CHR were involved in 29 overrepresented pathways (Fig. 7 and Table S7).

A previous study showed that plant hormones are important for panicle development. Among the 15 KEGG pathways in RHR, 21 DEGs were involved in plant hormone signal transduction, of which 14 DEGs were upregulated in HHZ; three DEGs were involved in cytochrome P450 metabolism, which plays a role in brassinosteroid (BR) biosynthesis; and two were upregulated (Table 4).

In SHR and CHR, there were three common pathways: the starch and sucrose metabolism pathway, the NOD-like receptor signaling pathway, and the estrogen signaling pathway. Carbohydrate accumulation was essential for panicle development. In the KEGG analysis, seven DEGs involved in starch and sucrose metabolism were observed in SHR and 18 DEGs involved in starch and sucrose metabolism were observed in CHR. In SHR, the genes in HHZ were not

different between HHZ\_40 and HHZ\_32. However, genes *BGIOGA010570* and *BGIOGA026140* encoding sucrose synthase (EC 2.4.1.13), genes *BGIOGA026976*, *BGIOGA009181*, and *BGIOGA030796* encoding trehalose-6-phosphate synthase (EC 2.4.1.15), and gene *BGIOGA000509* encoding trehalose-6-phosphate phosphatase (EC 3.1.3.12) were significantly downregulated in IR36\_40 compared with IR36\_32. However, gene *BGIOGA031385* encoding beta-amylase (EC 3.2.1.2) was significantly upregulated in IR36\_40 compared with IR36\_32 (Table 5).

## qRT-PCR verification

To confirm the accuracy of the RNA-Seq results, ten representative DEGs each from the HHZ\_32 vs HHZ\_40 (a) and IR36\_32 vs IR36\_40 (b) groups, as well as five DEGs each from the IR36\_40 vs HHZ\_40 (c) and IR36\_32 vs HHZ\_32 (d) groups were chosen to determine relative expression. Of the ten DEGs from the HHZ\_32 vs HHZ\_40 group, five were in RHR: *BGIOGA022020* is related to BR synthesis, *BGIOGA006348* encodes a heat shock factor (Hsf), *BGIOGA017088* is involved in the ETH TF family, *BGIOGA006285* participates in ethylene responsive regulation, and *BGIOGA024710* is an auxin-responsive gene involved in plant hormone transduction. Among the ten DEGs from the IR36\_32 vs IR36\_40 group, five were in SHR and encoded cytokinin oxidase/dehydrogenase (*BGIOGA005140*), sucrose synthase (*BGIOGA026140*), trehalose-6-phosphate synthase (*BGIOGA026976*), trehalose-6-phosphate phosphatase (*BGIOGA000509*), and catalase (*BGIOGA007252*). Four DEGs were in CHR from the HHZ\_32 vs HHZ\_40 and IR36\_32 vs IR36\_40 groups and two common genes, *BGIOGA032653* and *BGIOGA015767*, were validated. *BGIOGA032653* is involved in phenylpropanoid biosynthesis and *BGIOGA015767* encodes a heat shock protein (HSP). The qRT-PCR results for the DEGs were all consistent with the RNA-Seq data (Fig.8).

## Discussion

The exposure of rice plants to high temperature growing conditions during spikelet differentiation inhibited panicle initiation and reduced spikelet number per panicle (Fig. 1). Previous studies have shown that the genes *SPI*, *ASPI*, *TUT1*, *PAA2*, and *OsALMT7* are closely related to branch and spikelet development in rice (Bai *et al.* 2015; Heng *et al.* 2018; Li *et al.* 2010). However, in the current study, we observed no significant difference in the expression of these genes between the 40°C treatment and the 32°C control treatment in either rice cultivar, which indicates that the expression of these genes might not be inhibited in young panicles exposed to high temperature.

In general, the upregulation of HSPs contributes to the heat stress response in plants (Guan *et*

*al.* 2010; Jagadish *et al.* 2010; Jung *et al.* 2013). Moon *et al.* (2014) reported that heterologous overexpression of *OsHSP1* (*BGIOSGA015767*, encoding a HSP) increased heat tolerance in *Arabidopsis*. However, in the current study, *BGIOSGA015767* expression was upregulated in both HHZ ( $\log_2(\text{HHZ}_{40}/\text{HHZ}_{32}) = 5.7$ ,  $P\text{-value} = 0$ ) and IR36 ( $\log_2(\text{IR36}_{40}/\text{IR36}_{32}) = 5.0$ ,  $P\text{-value} = 0$ ). In addition, there was no gene expression difference in the GO term of HSPs between cultivars, which demonstrates that the heat stress reaction is common to both rice cultivars when exposed to high temperature. The GO enrichment analysis revealed that the DEGs for the CHR group were commonly enriched in response to GO terms representing heat, stress, and temperature stimuli in the biological process category (Fig. 5). These results demonstrate that the heat stress response did not directly inhibit panicle development but rather may disrupt physiological processes related to panicle development.

An important factor determining heat tolerance is antioxidant capacity (Lan *et al.* 2016). Buer *et al.* (2010) reported that flavonoids can positively regulate reactive oxygen species (ROS), which can affect the transport of plant hormones and influence pollen development. The flavonoid synthesis pathway was overrepresented in the IR36<sub>32</sub> vs IR36<sub>40</sub> group. Specifically, five genes involved in flavonoid synthesis were downregulated at 40°C, which might indicate a reduction in the antioxidant capacity of IR36 under heat stress. In addition, 14 DEGs in the IR36<sub>32</sub> vs IR36<sub>40</sub> group were enriched in the peroxisome pathway. Among these, 10 DEGs were significantly downregulated and four DEGs were significantly upregulated. However, the peroxisome pathway was not significant in the KEGG analysis of HHZ<sub>32</sub> vs HHZ<sub>40</sub> (Fig. 6). *BGIOSGA007252* and *BGIOSGA011520*, which encode catalase (EC:1.11.1.6), were significantly downregulated in IR36 at 40°C compared with 32°C, whereas no expression differences were observed in HHZ<sub>32</sub> vs HHZ<sub>40</sub>. This finding suggests that high temperature had a greater negative effect on the antioxidant capacity of IR36 than of HHZ, which provides a primary explanation for the greater heat injury observed in the young IR36 panicles than in those of HHZ.

Regulation of endogenous hormones affects the development of young panicles. Wu *et al.* (2017) reported that a lower spikelet number under high temperature growing conditions was associated with cytokinin degradation. In the current study, *BGIOSGA001314*, which encodes a cytokinin-activity enzyme, did not differ between the 40°C and 32°C treatments in HHZ ( $\log_2(\text{HHZ}_{40}/\text{HHZ}_{32}) = -0.41$ ) or IR36 ( $\log_2(\text{IR36}_{40}/\text{IR36}_{32}) = -0.38$ ). However, the gene *BGIOSGA005140*, which encodes cytokinin oxidase/dehydrogenase, was significantly upregulated in the IR36<sub>32</sub> vs IR36<sub>40</sub> group ( $\log_2$  fold change = 1.67,  $P\text{-value} = 0.004$ ), but was not different in the HHZ<sub>32</sub> vs HHZ<sub>40</sub> group ( $\log_2$  fold change = 0.86,  $P\text{-value} = 0.088$ ). These results are consistent with those of Wu *et al.* (2016) and suggest that spikelet formation is associated with cytokinin degradation and greater degradation occurs at high temperatures in the heat-susceptible cultivar than in the heat-tolerant cultivar.

The DEGs in RHR were enriched in 54 GO terms (Fig. 5). The GO term analysis revealed biological processes promoting resistance to heat stress in the heat-tolerant cultivar HHZ. Downregulation of *BGIOSGA022020* in the heterocycle biosynthetic process induces GRAS protein reduction, which promotes BR synthesis to enhance heat tolerance (Vriet *et al.* 2012). In the molecular function category for RHR, 50 DEGs were involved in DNA-binding transcription factor activity. *BGIOSGA006348* encoded an HSF TF and was upregulated in the HHZ\_32 vs HHZ\_40 group, although differences were not observed in the IR36\_32 vs IR36\_40 group. Wang *et al.* (2009) reported that the higher expression of heat shock TFs contributed to high temperature tolerance. WRKY genes encode TFs that play important roles in abiotic stress responses (Chen *et al.* 2010), especially to abscisic acid (ABA) (Zhen *et al.* 2005). In this study, six DEGs were WRKY TFs, namely, *BGIOSGA003134*, *BGIOSGA017063*, *BGIOSGA029574*, *BGIOSGA005924*, *BGIOSGA024948*, and *BGIOSGA033505*, which might promote young panicle development associated with sucrose consumption mediated by ABA under high temperature (Feng *et al.* 2018). However, few studies have reported the relationship between the WRKY family and heat resistance, which should be further studied. *BGIOSGA029574* is a general stress-response gene, which has putative functions in distinct cellular processes, such as transcription regulation, stress response, and sugar metabolism under Fe-excess-induced, dark-induced, and drought-induced stress (Ricachenevsky *et al.* 2010). Among the six WRKY genes, *BGIOSGA017063* was downregulated while the other five genes were upregulated, although the gene has not been cloned for the gene function analysis and therefore requires further study. Of the 10 DEGs in the ETH family, five genes were downregulated and the downregulation of *BGIOSGA017088* reduced the ABA content and promoted gibberellin (GA) signal transduction, which is beneficial for rice plant growth (Yaish *et al.* 2010). The upregulation of *BGIOSGA006285*, *BGIOSGA010867*, *BGIOSGA030019*, *BGIOSGA005915*, and *BGIOSGA012535* plays an important role in ethylene response regulation. Cao *et al.* (2006) reported that the upregulation of *BGIOSGA005915* enhanced tolerance to salt, cold, drought, and wounding, and the current study reveals that this gene might also contribute to the improvement of high-temperature stress resistance. *BGIOSGA000303* and *BGIOSGA000304* are genes in the cytokinin receptor family, and the upregulation of these two genes promotes cytokinin activation (Ito & Kurata 2006). The MADs box gene is related to flower development (Kobayashi *et al.* 2012) and the upregulation of the MAD genes in RHR indicated that the MAD family might enhance heat stress tolerance. The HZ-ZIP TF family might have a similar function.

In the RHR category, the DEGs enriched in the KEGG pathways appear beneficial for heat-stress tolerance, including plant hormone signal transduction and BR biosynthesis. Twenty-one DEGs were involved in plant hormone signal transduction, of which 14 DEGs were upregulated, including the auxin-responsive genes *BGIOSGA024710*, *BGIOSGA001585*, *BGIOSGA019301*,

and *BGIOSGA037837*, which facilitate rice plant growth (Hagen & Guilfoyle 2002). In BR biosynthesis, *BGIOSGA002945*, which encodes *D2/CYP90D2*, a gene that catalyzes the steps from 6-deoxoteasterone to 3-dehydro-6-deoxoteasterone and from teasterone to 3-dehydroteasterone, was upregulated to promote BR synthesis in the latter pathway (Hong *et al.* 2003), and *BGIOSGA001585* was downregulated to promote BR activity (Sakamoto *et al.* 2011). The genes related to hormone signal transduction and BR biosynthesis might contribute to young panicle development under high temperature. Seven DEGs involved in plant hormone signal transduction were downregulated, and among these, *BGIOSGA036617*, *BGIOSGA034767*, and *BGIOSGA010559* have not been cloned for functional analysis while *BGIOSGA034772* plays a more important role in organismal development. The genes *BGIOSGA024374*, *BGIOSGA023368*, *BGIOSGA000304* and *BGIOSGA005312* are A-type response regulated genes (Jain *et al.* 2006). However, it is unclear whether the downregulation of *BGIOSGA024374* and *BGIOSGA023368* contributes to improved heat tolerance in rice varieties. In addition, the downregulated gene, *BGIOSGA010919*, is an ABA receptor. Tian *et al.* (2015) reported that ABA accumulation upregulates gene expression. In the current study, downregulation of *BGIOSGA010919* may contribute to excessive ABA accumulation. The role of ABA in panicle development requires further study. *BGIOSGA014915*, which participates in BR synthesis, was downregulated in RHR. Previous reports have found that BRs can modulate the metabolic responses of plants to abiotic environmental stresses (Vriet *et al.* 2012; Wang *et al.* 2018). BR accumulation reportedly reduces spikelet degeneration under nitrogen application (Zhang, 2018). *BGIOSGA002945* and *BGIOSGA014915* participate in different BR biosynthesis pathways (Shi, 2015), but *BGIOSGA002945* may play a more important role in modulating spikelet development under high temperature.

Carbohydrate storage and utilization are essential for panicle initiation (Tian *et al.* 2016). The KEGG analysis showed that the phenylpropanoid biosynthesis pathway was commonly overrepresented in HHZ\_32 vs HHZ\_40, IR36\_32 vs IR36\_40, and IR36\_40 vs HHZ\_40. The phenylpropanoid biosynthesis pathway is involved in lignin synthesis, which suggests that high temperature inhibits lignin synthesis; however, phenylpropanoid biosynthesis was not associated with heat tolerance in our heat resistant cultivar (Fig. 6). In the SHR category, seven DEGs were enriched in the starch and sucrose metabolism pathway (Fig.7b). The gene *BGIOSGA031385*, which encodes beta-amylase, was significantly upregulated in IR36\_32 vs IR36\_40, suggesting that it promoted starch hydrolysis and reduced carbohydrate storage. The genes *BGIOSGA010570* and *BGIOSGA026140*, which encode sucrose synthesis, were significantly downregulated in the IR36\_32 vs IR36\_40 group, whereas no difference in expression was observed in the HHZ\_32 vs HHZ\_40 group. Sucrose degrades into uridine 5'-diphosphoglucose and fructose, which are major forms of carbon used for energy. Impairment of sucrose synthase activity reportedly reduced

resistance to heat stress (Hirose *et al.* 2008; Takehara *et al.* 2018). The results of the current study suggest that impaired carbohydrate metabolism in the heat-susceptible cultivar aggravated spikelet reduction. The starch and sucrose pathway genes were also highly represented in the CHR group (Fig.7c). Such genes are involved in the downregulation of genes encoding beta-fructofuranosidase, fructokinase, beta-glucosidase, trehalose-6-phosphate phosphatase, alpha-trehalase, and others. Trehalose-6-phosphate synthase, trehalose-6-phosphate phosphatase, and alpha-trehalase are involved in trehalose synthesis. Trehalose plays an important role in abiotic stress resistance, and trehalose-6-phosphate, an intermediate product of trehalose synthesis participates in sucrose signal transduction (Lunn *et al.* 2006; Ruan 2014). Nunes *et al.* (2013) reported that trehalose-6-phosphate served as a sugar signal that could induce the expression of genes associated with the alleviation of abiotic stress injury. In this study, certain DEGs in the CHR group were also upregulated to promote trehalose-6-phosphate synthesis, and the upregulation of *BGIOSGA026976*, *BGIOSGA009181*, and *BGIOSGA030796* promoted trehalose-6-phosphate synthesis in SHR. These findings indicate that trehalose-6-phosphate synthesis may be a normal response of young rice panicles to high temperature and that the heat-sensitive rice cultivar synthesizes trehalose-6-phosphate more readily than the heat-tolerant cultivar in response to heat stress. However, the gene encoding trehalose-6-phosphate phosphatase, *BGIOSGA000509*, was significantly downregulated in IR36 at 40°C compared with that at 32°C, which might cause a decrease in trehalose content and in turn disrupt carbohydrate distribution. Our results suggest that trehalose-6-phosphate metabolism was disordered under the high temperature condition and that the effects were more severe in the heat-susceptible cultivar than in the heat-tolerant cultivar.

A close relationship is observed between endogenous hormones and carbohydrate accumulation, which may suggest that the regulation of endogenous hormones in heat-tolerant varieties promotes carbohydrate utilization. The identification of DEGs in this study could improve understanding of the molecular mechanisms of heat resistance in young panicles. In the practice of rice production and breeding, DEGs associated with hormone metabolism in the RHR category and DEGs associated with starch and metabolism in the SHR category under high temperature could be used to quickly identify heat tolerant cultivars.

## Conclusions

In summary, heat stress-responsive DEGs in young panicles were identified by a transcriptome analysis of a heat-tolerant rice cultivar and a heat-susceptible rice cultivar grown at high temperature (40°C) and a control temperature (32°C). The statistical analysis of 5,533 DEGs revealed three categories of genes (RHR, SHR, and CHR) containing a total of 4,070 DEGs. We highlighted the differential expression of a group of DNA-binding TFs that was significantly

enriched in the RHR category as well as the differential expression of genes involved in the starch and sucrose metabolism pathway that were overrepresented in the SHR category. Overall, DEGs related to plant hormones and signal transduction might be specifically beneficial for young panicle development at high temperature. Heat-tolerant cultivars seem to increase endogenous hormones and maintain a stable carbohydrate metabolism pathway under high temperature. However, certain metabolic pathways, including starch and sucrose metabolism, were much more damaged in the heat susceptible cultivars under high temperatures, and this damage might have inhibited the of panicle development.

## References

- Anand RY, Kumar SD, and Narayan TK. 2018. Effect of irrigation method on adaptation capacity of rice to climate change in subtropical India. *International Journal of Plant Production*, 2018, 12: 203-217.
- Bai JT, Zhu XD, Wang Q, Zhang J, Chen HQ, Dong GJ, Zhu L, Zheng HK, Xie QJ, Nian JQ, Chen F, Fu Y, Qian Q, and Zuo JR. 2015. Rice *TUTOUI* encodes a SCAR-like protein that is important for actin organization and panicle development. *Plant Physiology*, 169:1179-1191.
- Buer CS, Imin N, and Djordjevic MA. 2010. Flavonoids: new roles for old molecules. *Journal of Integrative Plant Biology*, 52:98-111.
- Cao YY, Duan H, Yang LN, Wang ZQ, Zhou SC, and Yang JC. 2009. Effect of heat-stress during meiosis on grain yield of rice cultivars differing in heat-tolerance and its physiological mechanism. *Acta Agronomica Sinica*, 34:2134-2142.
- Cao Y, Song F, Goodman RM, and Zheng Z. 2006. Molecular characterization of four rice genes encoding ethylene-responsive transcriptional factors and their expressions in response to biotic and abiotic stress. *Journal of Plant Physiology*, 163:1167-1178.
- Chen H, Lai ZB, Shi JW, Xiao Y, Chen ZX, and Xu XP. 2010. Roles of *Arabidopsis* WRKY18, WRKY40 and WRKY60 transcription factors in plant responses to abscisic acid and abiotic stress. *BMC Plant Biology*, 10:1-15.
- Czechowski T, Bari RP, Stitt M, Scheible WR, and Udvardi MK. 2004. Real-time RT-PCR profiling of over 1400 *Arabidopsis* transcription factors: unprecedented sensitivity reveals novel root- and shoot-specific genes. *Plant Journal* 38:366-379.
- Ding C, Wang S, and Ding Y. 2016. Functional genes of panicle development in response to nitrogen fertilizer in rice. *Chinese Bullrtin of Botany*, 51:488-498.
- Dwivedi SK, Kumar S, Prakash V, Mondal S, and Mishra JS. 2015. Influence of rising atmospheric CO<sub>2</sub> concentrations and temperature on morpho-physiological traits and yield of rice genotypes in sub humid climate of eastern India. *American Journal of Plant Sciences*, 6:2239-2249.
- Endo M, Tsuchiya T, Hamada K, Kawamura S, Yano K, Ohshima M, Higashitani A, Watanabe M, and Kawagishi-Kobayashi M. 2009. High temperatures cause male sterility in rice plants with transcriptional alterations during pollen development. *Plant Cell Physiology*, 50:1911-1922.
- Espe MB, Hill JE, Hijmans RJ, McKenzie K, Mutters R, Espino LA, Leinfelder-Miles M, van Kessel C, and Linquist BA. 2017. Point stresses during reproductive stage rather than warming seasonal temperature determine yield

- in temperate rice. *Global Change Biology*, 23:4386-4395.
- Fang XW, Tang LH, and Wang YP. 2006. Selection on rice germplasm tolerant to high temperature. *Journal of Plant Genetic Resources*, 7:342-344.
- Feng BH, Zhang CX, Chen TT, Zhang XF, Tao LX, and Fu GF. 2018. Salicylic acid reverses pollen abortion of rice caused by heat stress. *BMC Plant Biology*, 18: 245.
- Guan JC, Yeh CH, Lin YP, Ke YT, Chen MT, You JW, Liu YH, Lu CA, Wu SJ, and Lin CY. 2010. A 9 bp cis-element in the promoters of class I small heat shock protein genes on chromosome 3 in rice mediates L-azetidine-2-carboxylic acid and heat shock responses. *Journal of Experimental Botany*, 61:4249-4261.
- Hagen G, and Guilfoyle T. 2002. Auxin-responsive gene expression: genes, promoters and regulatory factors. *Plant Molecular Biology*, 49:373-385.
- Heng YQ, Wu CY, Long Y, Luo S, Ma J, Chen J, Liu JF, Zhang H, Ren YL, Wang M, Tan JJ, Zhu SS, Wang JL, Lei CL, Zhang X, Guo XP, Wang HY, Cheng ZJ, and Wan JM. 2018. *OsALMT7* maintains panicle size and grain yield in rice by mediating malate transport. *The Plant Cell*, 30:889-906.
- Hirose T, Scofield GN, and Terao T. 2008. An expression analysis profile for the entire sucrose synthase gene family in rice. *Plant Science*, 174:534-543.
- Hong Z, Ueguchitanaka M, Umemura K, Uozu S, Fujioka S, Takatsuto S, Yoshida S, Ashikari M, Kitano H, and Matsuoka M. 2003. A rice brassinosteroid-deficient mutant, *ebisu dwarf* (d2), is caused by a loss of function of a new member of cytochrome P450. *The Plant Cell*, 15:2900-2910.
- Islam MR, Feng BH, Chen TT, Fu WM, Zhang CX, Tao LX, and Fu GF. 2018. Absciscic acid prevents pollen abortion under high temperature stress by mediating sugar metabolism in rice spikelets. *Physiologia Plantarum*, 2019, 165(3): 644-663.
- Ito Y, and Kurata N. 2006. Identification and characterization of cytokinin-signalling gene families in rice. *Gene*, 382:57-65.
- Jagadish SV, Muthurajan R, Oane R, Wheeler TR, Heuer S, Bennett J, and Craufurd PQ. 2010. Physiological and proteomic approaches to address heat tolerance during anthesis in rice (*Oryza sativa* L.). *Journal of Experimental Botany*, 61:143-156.
- Jain M, Tyagi AK, and Khurana JP. 2006. Molecular characterization and differential expression of cytokinin-responsive type-A response regulators in rice (*Oryza sativa*). *BMC Plant Biology*, 6:1-11.
- Jung KH, Ghosh HJ, Nguyen MX, Kim SR, and An G. 2013. Genome-wide expression analysis of *HSP70* family genes in rice and identification of a cytosolic *HSP70* gene highly induced under heat stress. *Functional & Integrative Genomics*, 13:391-402.
- Kang HG, Jang S, Chung JE, Cho YG, and An G. 2013. Characterization of two rice MADS box genes that control flowering time. *Molecular Cells*, 7:559-566.
- Kim J, Shon J, Lee CK, Yang W, Yoon Y, Yang WH, Kim YG, and Lee BW. 2011. Relationship between grain filling duration and leaf senescence of temperate rice under high temperature. *Field Crops Research*, 122:207-213.
- Kobayashi K, Yasuno N, Sato Y, Yoda M, Yamazaki R, Kimizu M, Yoshida H, Nagamura Y, and Kozuka J. 2012. Inflorescence meristem identity in rice is specified by overlapping functions of three *API1/FUL*-Like MADS Box Genes and *PAP2*, a *SEPALLATA* MADS Box Gene. *The Plant Cell*, 24:1848-1859.
- Lan X, Gu ZD, Ding YF, Wang K, Jiang Q, and Zhu C. 2016. Effect of high temperature stress on physiological characteristics of spikelet of rice during flowering stage. *Chinese Journal of Rice Science*, 30:637-646.
- Li SB, Qian Q, Fu ZM, Zeng DL, Meng XB, Kozuka J, Maekawa M, Zhu XD, Zhang J, and Li JY. 2010. Short



- panicle1 encodes a putative *PTR* family transporter and determines rice panicle size. *Plant Journal*, 58:592-605.
- Liu XW, Meng YL, Zhou ZG, and Cao WX. 2005. Dynamic characteristics of floret differentiation and degeneration in rice. *Acta Agronomica Sinica*, 31: 451-455.
- Lunn JE, Feil R, Hendriks JHM, Gibon Y, Morcuende R, Osuna D, Scheible W, Carillo P, Hajirezaei MR, and Stitt M. 2006. Sugar-induced increases in trehalose 6-phosphate are correlated with redox activation of ADPglucose pyrophosphorylase and higher rates of starch synthesis in *Arabidopsis thaliana*. *Biochemical Journal*, 397:139-148.
- Min L, Li Y, Hu Q, Zhu L, Gao W, Wu Y, Ding Y, Liu S, Yang X, and Zhang X. 2014. Sugar and auxin signaling pathways respond to high-temperature stress during anther development as revealed by transcript profiling analysis in cotton. *Plant Physiology*, 164:1293-1308.
- Moon JC, Ham DJ, Hwang SG, Yong CP, Lee C, and Jang CS. 2014. Molecular characterization of a heat inducible rice gene, *OsHSP1*, and implications for rice thermotolerance. *Genes & Genomics*, 36:151-161.
- Morita S, Wada H, and Matsue Y. 2016. Countermeasures for heat damage in rice grain quality under climate change. *Plant Production Science*, 19(1):1-11.
- Nunes C, and Paul MJ. 2013. The trehalose 6-Phosphate/*SnRK1* signaling pathway primes growth recovery following relief of sink limitation. *Plant Physiology*, 162:1720-1732.
- Prasad PVV, Bheemanahalli R, and Jagadish SVK. 2017. Field crops and the fear of heat stress-Opportunities, challenges and future directions. *Field Crops Research*, 200:114-121.
- Ricachenevsky FK, Sperotto RA, Menguer PK, Fett JP. 2010. Identification of Fe-excess-induced genes in rice shoots reveals a WRKY transcription factor responsive to Fe, drought and senescence. *Molecular Biology Reports*, 37:3735-3745.
- Ruan YL. 2014. Sucrose metabolism: gateway to diverse carbon use and sugar signaling. *Annual Review Plant Biology*, 65:33-67.
- Sakamoto T, Kawabe A, Tokidasegawa A, Shimizu B, Takatsuto S, Shimada Y, Fujioka S, and Mizutani M. 2011. Rice *CYP734As* function as multisubstrate and multifunctional enzymes in brassinosteroid catabolism. *Plant Journal*, 67:1-12.
- Shi ZY. 2015. Fine mapping of grain width locus GW12 and functional verification of a grain length gene SG4 in Rice. Ph. D. dissertation, Chinese Academy of Agricultural Sciences, Beijing.
- Stocher TF, Plattner GK, Tignor MMB, Allen SK, Boschung J, Nauel A, Xia Y, Bex V, and Midgley P. 2013. CLIMATE CHANGE 2013: The Physical Science Basis, working group I contribution to the fifth assessment report of the intergovernmental panel on climate change. Cambridge University Press, Cambridge, UK.
- Takehara K, Murata K, Yamaguchi T, Yamaguchi K, Chaya G, Kido S, Iwasaki Y, Ogiwara H, Ebitani T, and Miura K. 2018. Thermo-responsive allele of *sucrose synthase 3 (Sus3)* provides high-temperature tolerance during the ripening stage in rice (*Oryza sativa* L.). *Breeding Science*, 63:336-342.
- Tao FL, Zhang Z, Shi WJ, Liu YJ, Xiao DP, Zhang S, Zhu Z, Wang M, and Liu FS. 2013. Single rice growth period was prolonged by cultivars shifts, but yield was damaged by climate change during 1981-2009 in China, and late rice was just opposite. *Global Change Biology*, 19:3200-3209.
- Tian QL, Liu B, Zhong XY, Zhao M, Sun H, and Ren WJ. 2016. Relationship of NSC with the formation of branches and spikelets and the yield traits of Indica hybrid rice in different planting methods. *Scientia Agricultura Sinica* 49:35-53.

- 581 Tian XJ, Wang ZY, LiXF, Lv TX, Liu HZ, Wang LZ, Niu HB, Bu QY. Characterization and functional analysis of  
582 pyrabactin resistance-like abscisic acid receptor family in rice. *Rice*, 2015,8:28.
- 583 Vriet C, Russinova E, and Reuzeau C. 2012. Boosting crop yields with plant steroids. *The Plant Cell*, 24:842-857.
- 584 Wang C, Qian Z, and Shou HX. 2009. Identification and expression analysis of OsHsfs in rice. *Journal of Zhejiang*  
585 *University-Science B(Biomedicine & Biotechnology)*, 10:291-300.
- 586 Wang YL, Wang L, Zhou JX, Hu SB, Chen HZ, Xiang J, Zhang YK, Zeng YJ, Shi QH, Zhu DF, and Zhang YP.  
587 Research progress on heat stress of rice at flowering stage. *Rice Science*, 2019, 26(1): 1-10.
- 588 Wang YL, Zhang YP, Xiang J, Wang L, Chen HZ, Zhang YK, Zhang WQ, and Zhu F. 2017. Response of indica rice  
589 spikelet differentiation and degeneration to air temperature and solar radiation of different sowing dates.  
590 *Chinese Journal of Applied Ecology*, 28:3571-3580.
- 591 Wang ZQ, Zhang WY, and Yang JC. 2018. Physiological mechanism underlying spikelet degeneration in rice. *Journal*  
592 *of Integrative Agriculture*, 17:1475-1481.
- 593 Wu C, Cui KH, Wang WC, Li Q, Fahad S, Hu QQ, Huang JL, Nie LX, Mohapatra PK, and Peng SB. 2017. Heat-  
594 induced cytokinin transportation and degradation are associated with reduced panicle cytokinin expression  
595 and fewer spikelets per panicle in rice. *Frontiers in Plant Science*, 8:371.
- 596 Wu C, Cui KH, Wang WC, Li Q, Fahad S, Hu QQ, Huang JL, Nie LX, and Peng SB. 2016. Heat-induced  
597 phytohormone changes are associated with disrupted early reproductive development and reduced yield in  
598 rice. *Scientific Reports*, 6:34978.
- 599 Yaish MW, Elkereamy A, Zhu T, Beatty PH, Good AG, Bi YM, and Rothstein SJ. 2010. The *APETALA-2-like*  
600 transcription factor *OsAP2-39* controls key interactions between abscisic acid and gibberellin in rice. *Plos*  
601 *Genetics*, 6:e1001098.
- 602 Yamagishi J, Miyamoto N, Hirotsu S, Laza RC, and Nemoto K. 2004. QTLs for branching, floret formation, and pre-  
603 flowering floret abortion of rice panicle in a temperate japonica x tropical japonica cross. *Theoretical &*  
604 *Applied Genetics*, 109:1555-1561.
- 605 Yamakawa H, and Hakata M. 2010. Atlas of rice grain filling-related metabolism under high temperature: joint  
606 analysis of metabolome and transcriptome demonstrated inhibition of starch accumulation and induction of  
607 amino acid accumulation. *Plant & Cell Physiology*, 51:795-809.
- 608 Yamakawa H, Hirose T, Kuroda M, and Yamaguchi T. 2007. Comprehensive expression profiling of rice grain filling-  
609 related genes under high temperature using DNA microarray. *Plant Physiology*, 144:258-277.
- 610 Zhen X, Zhang ZL, Zou XL, Huang J, Ruas P, Thompson D, and Shen QJ. 2005. Annotations and Functional Analyses  
611 of the Rice WRKY Gene Superfamily Reveal Positive and Negative Regulators of Abscisic Acid Signaling  
612 in Aleurone Cells[J]. *Plant Physiology*, 137: 176-189.
- 613 Zhong LJ, Cheng FM, Wen X, Sun ZX, and Zhang GP. 2010. The deterioration of eating and cooking quality caused  
614 by high temperature during grain filling in early-season indica rice cultivars. *Journal of Agronomy & Crop*  
615 *Science*, 191:218-225.
- 616 Zhou SC, Li H, Huang D, Lu D, Lai Z, Zhou D, Li K, Wang C, and Li H. 2012. Breeding and application of  
617 Huanghuazhan-A new variety with 1st class rice quality of mational standard. *Hubei Agricultural Sciences*,  
618 51:1960-1964.
- 619 Zhang WY. 2018. Mechanism underlying water and nitrogen regulation spikelet development and grain filling of rice.  
620 Ph.D. dissertation, Yangzhou university, Yangzhou.
- 621

**Figure legends:**

**Figure 1.** Effects of high temperature on panicle development. (a) young panicle morphologies after 9 d of high-temperature treatment; and (b) panicle morphologies at the heading stage after high-temperature treatment. Bars = 0.5 cm in (a) and 3 cm in (b).

**Figure 2.** Gene expression in the four comparison groups. (a) HHZ\_32 vs HHZ\_40, (b) IR36\_32 vs IR36\_40, (c) IR36\_40 vs HHZ\_40, and (d) IR36\_32 vs HHZ\_32. Red (upregulated) and blue (downregulated) dots indicate significant differences in gene expression, whereas gray dots represent genes with no significant difference in expression.

**Figure 3.** Venn diagrams for DEGs in the four comparison groups. (a) HHZ\_32 vs HHZ\_40, (b) IR36\_32 vs IR36\_40, (c) IR36\_40 vs HHZ\_40, and (d) IR36\_32 vs HHZ\_32.

**Figure 4.** Enriched GO terms ( $P < 0.05$ ) of all DEGs. (a) biological process, (b) cellular component, and (c) molecular function.

**Figure 5.** Enriched GO terms ( $P < 0.05$ ) of the DEGs in the RHR and CHR groups. (a) biological process, (b) cellular component, and (c) molecular function.

**Figure 6.** KEGG enrichment analysis of all DEGs. (a) HHZ\_32 vs HHZ\_40, (b) IR36\_32 vs IR36\_40, (c) IR36\_40 vs HHZ\_40, and (d) IR36\_32 vs HHZ\_32.

**Figure 7.** KEGG enrichment analysis for heat stress responsive genes from the three categories. (a) RHR, (b) SHR, and (c) CHR.

**Figure 8.** Gene expression levels determined by RNA-Seq and qRT-PCR. (a) HHZ\_32 vs HHZ\_40, (b) IR36\_32 vs IR36\_40, (c) IR36\_40 vs HHZ\_40, and (d) IR36\_32 vs HHZ\_32.

**Table 1** (on next page)

Panicle characters after high temperature treatment

1 **Table 1 Panicle characters after high temperature treatment**

Cultivar s	Treatme nt	Panicle weight(g)	Spikelet number	The number of differentiated spikelet	The proportion of degenerated spikelet (%)	Spikelet fertility (%)	Grain weight(mg)	Glume length (mm)	Glume width (mm)
HHZ	32°C	3.6±0.4	235.0±20.0	335.3±20.5	30.0±1.7	83.0±2.0	18.5±0.3	8.7±0.2	2.5±0.0
	40°C	2.4±0.3**	182.7±11.2*	303.3±10.1	39.7±5.3	78.5±1.3**	17.0±0.3**	7.8±0.1**	2.2±0.1**
IR36	32°C	3.1±0.4	183.3±7.6	264.7±13.8	30.7±1.5	81.5±1.8	20.5±0.3	8.1±0.3	2.5±0.0
	40°C	1.0±0.1**	85.0±13.5**	176.7±17.6**	51.4±9.3**	73.5±1.1**	16.0±0.1**	6.8±0.2**	2.3±0.1**

2 \* and \*\* indicate significance differences between the control (32°C) and high (°C) temperature treatments (one-tailed Student's t-test): \* $P<0.05$ ; \*\* $P<0.01$ .

3

# **Table 2**(on next page)

Table 2. RNA sequencing results

**Table 2.** Statistics of RNA sequencing results

Sample	HHZ_32	HHZ_40	IR36_32	IR36_40
Raw reads	44231722	45513241	45877838	46465046
Clean reads	44032896	45256701	45580821	46252929
	(99.6%)	(99.4%)	(99.4%)	(99.5%)
Total mapped	38834391	39148950	39541858	40418126
	(87.8%)	(86.0%)	(86.2%)	(87.0%)
Uniquely mapped	37502957	37759013	38120438	38853775
	(84.8%)	(83.0%)	(83.1%)	(83.1%)
Multiply mapped	1331434	1389937	1421421	1561018
	(3.0%)	(3.1%)	(3.1%)	(3.6%)

Note: HHZ\_32: The sample of HHZ treated with 32°C; HHZ\_40: The sample of HHZ treated with 40°C; IR36\_32: The sample of IR36 treated with 32°C; IR36\_40: The sample of IR36 treated with 40°C.





# **Table 3**(on next page)

Table 3. Classification of three categories of DEGs

**Table 3.** Classification of three categories of DEGs.

Categories	Subgroups	Number of DEGs
RHR	Only HHZ_32 vs HHZ_40	1157
	HHZ_32 vs HHZ_40 $\cap$ IR36_40 vs HHZ_40	531
SHR	Only IR36_32 vs IR36_40	603
	IR36_32 vs IR36_40 $\cap$ IR36_40 vs HHZ_40	104
CHR	Only IR36_40 vs HHZ_40	524
	HHZ_32 vs HHZ_40 $\cap$ IR36_32 vs IR36_40,	
	HHZ_32 vs HHZ_40 $\cap$ IR36_32 vs IR36_40 $\cap$ IR36_40 vs HHZ_40	1151

Note: RHR, heat-resistant-cultivar-related genes; SHR, heat-susceptible-cultivar-related genes; CHR, common heat stress-response genes.

**Table 4**(on next page)

Table 4. Gene expression of DEGs in plant hormone signal transduction and BR biosynthesis of RHR

**Table 4.** Gene expression of DEGs in Plant hormone signal transduction and BR biosynthesis of RHR

ID	Gene annotation	Cultiva r	baseMean	32°C	40°C	log2FoldChange	pval
BGIOSGA018672	Pseudo histidine-containing phosphotransfer protein 2	HHZ	64.7	36.9	92.5	1.33	0.00
		IR36	59.4	49.2	69.7	0.50	0.06
BGIOSGA004140	Probable protein phosphatase 2C 8	HHZ	665.2	219.8	1110.7	2.34	0.00
		IR36	465.6	355.3	575.9	0.70	0.00
BGIOSGA005312	Two-component response regulator ORR3	HHZ	50.6	28.6	72.6	1.35	0.00
		IR36	26.3	24.2	28.5	0.24	0.58
BGIOSGA024710	Auxin-responsive protein IAA24	HHZ	807.3	458.2	1156.3	1.34	0.00
		IR36	828.1	653.4	1002.7	0.62	0.00
BGIOSGA010835	ABSCISIC ACID-INSENSITIVE 5-like protein 2	HHZ	146.5	85.8	207.2	1.27	0.00
		IR36	83.4	74.0	92.8	0.33	0.26
BGIOSGA011032	Probable protein phosphatase 2C 30	HHZ	102.9	53.4	152.4	1.51	0.00
		IR36	113.6	108.0	119.3	0.14	0.69
BGIOSGA015611	Probable protein phosphatase 2C 37	HHZ	86.1	44.3	127.8	1.53	0.00
		IR36	75.8	52.0	99.6	0.94	0.00
BGIOSGA019301	Auxin-responsive protein IAA16	HHZ	97.4	56.6	138.3	1.29	0.00
		IR36	79.7	76.6	82.8	0.11	0.61
BGIOSGA008704	Auxin-responsive protein SAUR36	HHZ	36.6	22.9	50.3	1.14	0.00
		IR36	23.1	22.5	23.6	0.07	0.91
BGIOSGA012535	ARATH Protein ETHYLENE INSENSITIVE 3	HHZ	2890.1	1268.4	4511.8	1.83	0.00
		IR36	2293.2	1543.8	3042.6	0.98	0.00
BGIOSGA037772	ARATH Transcription factor PIF1	HHZ	27.5	14.2	40.8	1.52	0.00
		IR36	17.3	13.3	21.4	0.69	0.24
BGIOSGA000304	Two-component response regulator ORR26	HHZ	143.8	92.5	195.0	1.08	0.00
		IR36	110.4	85.9	134.9	0.65	0.00

BGIOGA004789	Probable protein phosphatase 2C	HHZ	522.2	301.9	742.5	1.30	0.00
		IR36	623.2	501.8	744.5	0.57	0.02
BGIOGA037837	Auxin-responsive protein SAUR72	HHZ	3.5	1.0	6.0	2.55	0.04
		IR36	0.8	1.3	0.3	-2.07	0.67
BGIOGA024374	Two-component response regulator ORR7	HHZ	17.8	29.5	6.2	-2.26	0.00
		IR36	49.3	58.2	40.4	-0.53	0.10
BGIOGA036617	Transcription factor TGAL11	HHZ	308.0	423.3	192.8	-1.13	0.00
		IR36	572.4	700.8	444.0	-0.66	0.00
BGIOGA034772	BTB/POZ domain and ankyrin repeat-containing protein NH5.1	HHZ	1148.1	1629.2	667.0	-1.29	0.00
		IR36	1335.2	1698.8	971.6	-0.81	0.00
BGIOGA010559	Protein TIFY 10a	HHZ	339.2	465.1	213.4	-1.12	0.00
		IR36	374.0	492.0	256.0	-0.94	0.00
BGIOGA010919	Absciscic acid receptor PYL5	HHZ	34.8	55.0	14.5	-1.92	0.00
		IR36	52.9	58.0	47.8	-0.28	0.33
BGIOGA023368	Two-component response regulator ORR25	HHZ	4.4	8.8	0.0	-Inf	0.00
		IR36	3.2	5.5	1.0	-2.49	0.15
BGIOGA034767	BTB/POZ domain and ankyrin repeat-containing protein NH5.2	HHZ	1147.8	1623.6	672.0	-1.27	0.00
		IR36	1304.2	1737.3	871.0	-1.00	0.00
BGIOGA002945	Cytochrome P450 90D2	HHZ	178.2	118.0	238.4	1.01	0.00
		IR36	184.9	164.3	205.5	0.32	0.05
BGIOGA014915	Cytochrome P450 724B1	HHZ	1872.9	2570.7	1175.2	-1.13	0.00
		IR36	1251.4	1482.6	1020.2	-0.54	0.00
BGIOGA001585	Cytochrome P450 734A6	HHZ	123.1	178.3	67.9	-1.39	0.00
		IR36	202.3	267.6	136.9	-0.97	0.00

# **Table 5**(on next page)

Table 5. Gene expression of DEGs in starch and sucrose metabolism in SHR

**Table 5.** Gene expression of DEGs in starch and sucrose metabolism in SHR


ID	Gene annotation	Cultivar	baseMean	IR36_32	IR36_40	log2FoldChange	P-value
BGIOGA01057	Sucrose synthase	HHZ	14318.1	18545.9	10090.4	-0.88	0.00
0		IR36	13352.6	18616.3	8088.8	-1.20	0.00
BGIOGA02614	Sucrose synthase	HHZ	13.5	16.5	10.4	-0.67	0.24
0		IR36	16.8	24.7	8.9	-1.47	0.01
BGIOGA02697	trehalose-6-phosphate synthase, putative, expressed	HHZ	651.9	572.0	731.7	0.36	0.12
6		IR36	691.3	399.7	982.9	1.30	0.00
BGIOGA00918	trehalose-6-phosphate synthase, putative, expressed	HHZ	731.2	562.7	899.8	0.68	0.01
1		IR36	988.8	578.3	1399.3	1.28	0.00
BGIOGA03079	trehalose-6-phosphate synthase, putative, expressed	HHZ	2.2	1.9	2.4	0.38	0.98
6		IR36	4.3	0.0	8.6	Inf	0.00
BGIOGA00050	Trehalose-6-phosphate phosphatase	HHZ	175.0	229.9	120.2	-0.94	0.00
9		IR36	179.2	268.6	89.9	-1.58	0.00
BGIOGA03138	beta-amylase, putative, expressed	HHZ	19.3	17.7	20.8	0.23	0.65
5		IR36	28.0	17.4	38.5	1.15	0.01

# Figure 1

Figure 1. Effects of high temperature on panicle development.

(a) young panicle morphologies after 9 d of high-temperature treatment; and (b) panicle morphologies at the heading stage after high-temperature treatment. Bars = 0.5 cm in (a) and 3 cm in (b).






# Figure 2

Figure 2. Gene expression in the four comparison groups.


(a) HHZ\_32 vs HHZ\_40, (b) IR36\_32 vs IR36\_40, (c) IR36\_40 vs HHZ\_40, and (d) IR36\_32 vs HHZ\_32. Red (upregulated) and blue (downregulated) dots indicate significant differences in gene expression, whereas gray dots represent genes with no significant difference in expression.



# Figure 3

Figure 3. Venn diagrams for DEGs in the four comparison groups.


(a) HHZ\_32 vs HHZ\_40, (b) IR36\_32 vs IR36\_40, (c) IR36\_40 vs HHZ\_40, and (d) IR36\_32 vs HHZ\_32. 2.



# Figure 4

Figure 4. Enriched GO terms ( $P < 0.05$ ) of all DEGs.

(a) biological process, (b) cellular component, and (c) molecular function.




# Figure 5

Figure 5. Enriched GO terms ( $P < 0.05$ ) of DEGs in RHR and CHR.

(a) biological process, (b) cellular component, and (c) molecular function.










# Figure 7

Figure 7. KEGG enrichment analysis for heat stress responsive genes from the three categories.

(a) RHR, (b) SHR, and (c) CHR.



# Figure 8

Figure 8. Gene expression levels determined by RNA-Seq and qRT-PCR.

(a) HHZ\_32 vs HHZ\_40, (b) IR36\_32 vs IR36\_40, (c) IR36\_40 vs HHZ\_40, and (d) IR36\_32 vs HHZ\_32.

