

Histological changes of female reproductive organs subjected to different jumping exercise intensities and honey supplementation in rats

Maryam Mosavat ¹, Mahaneem Mohamed ², Foong Kiew Ooi ^{Corresp., 3}, Mitra Mirsanjari ⁴, Anani Aila Mat Zin ⁵, Aminah Che Romli ²

¹ Sports Science Unit, School of Medical Sciences,, Universiti Sains Malaysia, Kota Bharu, Kelantan, Malaysia

² Department of Physiology, School of Medical Sciences, Universiti Sains Malaysia, Kota Bharu, Kelantan, Malaysia

³ Exercise and Sports Science Programme, School of Health Sciences, Universiti Sains Malaysia, Kota Bharu, Kelantan, Malaysia

⁴ Nutrition Programme, School of Health Sciences, Universiti Sains Malaysia, Kota Bharu, Kelantan, Malaysia

⁵ Department of Pathology, School of Medical Sciences, Universiti Sains Malaysia, Kota Bharu, Kelantan, Malaysia

Corresponding Author: Foong Kiew Ooi

Email address: fkooi@usm.my

Background: We assessed histopathological changes of ovaries and uterus in female rats subjected to different jumping exercise intensities combined with honey supplementation at 1 g/kg body weight/day. **Methods:** Seventy-two rats were divided into 6 groups, 12 rats in each: control(C), 20 and 80 jumps (20E, 80E), honey (H), and 20 and 80 jump with honey (20EH, 80EH). **Results:** The endometrium was significantly thicker in the rats in H, 20EH and 80EH groups compared to C, 20E and 80E. The myometrium thickness was significantly lower in 80E and significantly higher in 80EH compared to C, respectively. There was significantly higher myometrium thickness in 20EH and 80EH compared to 20E and 80E and H. The number of glands of the uterus in 20E and 80E was significantly lower than C. However, there was a significantly higher number of glands in H, 20EH and 80EH compared to 20E and 80E. The numbers of uterus vessels were significantly lower in 80E compared to 20E. However, the numbers of vessels were significantly higher in H, 20EH, and 80EH compared to 80E. The number of ovarian haemorrhagia was significantly lower in 20E, 80E, H, 20EH and 80EH compared to C. The number of corpora lutea was significantly lower in 80EH, H, 80E and 20E compared to C. However, the number of corpora lutea was significantly higher in 20EH compared to J20 and H. **Conclusion:** This study suggested that jumping exercises in particularly high-intensity exercise may induce histopathological changes in uterus and ovary in rats, and honey supplementation may ameliorate these effects.

Histological changes of female reproductive organs subjected to different jumping exercise intensities and honey supplementation in rats

Maryam Mosavat¹, Mahaneem Mohamed², Foong Kiew Ooi^{1, 5*}, Mitra Mirsanjari³, Anani Aila Mat Zin⁴, Aminah Che Romli².

¹ Sport Science Unit, School of Medical Sciences, Universiti Sains Malaysia, Kubang Kerian, Malaysia

² Department of Physiology, School of Medical Sciences, Universiti Sains Malaysia, Kubang Kerian, Malaysia

³ Nutrition Programme, School of Health Sciences, Universiti Sains Malaysia, Kubang Kerian, Malaysia

⁴ Department of Pathology, School of Medical Sciences, Universiti Sains Malaysia, Kubang Kerian, 16150, Malaysia

⁵ Exercise and Sports Science Programme, School of Health Sciences, Universiti Sains Malaysia, Kubang Kerian, Malaysia

Corresponding Author:

Foong Kiew Ooi*

Exercise and Sports Science Programme, School of Health Sciences, Universiti Sains Malaysia, Kubang Kerian, Kota Bharu, Kelantan, 16150 Malaysia

Email address: fkooi@usm.my

Abstract

Background: We assessed histopathological changes of ovaries and uterus in female rats subjected to different jumping exercise intensities combined with honey supplementation at 1 g/kg body weight/day.

Methods: Seventy-two rats were divided into 6 groups, 12 rats in each: control(C), 20 and 80 jumps (20E, 80E), honey (H), and 20 and 80 jump with honey (20EH, 80EH).

Results: The endometrium was significantly thicker in the rats in H, 20EH and 80EH groups compared to C, 20E and 80E. The myometrium thickness was significantly lower in 80E and significantly higher in 80EH compared to C, respectively. There was significantly higher myometrium thickness in 20EH and 80EH compared to 20E and 80E and H. The number of glands of the uterus in 20E and 80E was significantly lower than C. However, there was a significantly higher number of glands in H, 20EH and 80EH compared to 20E and 80E. The numbers of uterus blood vessels were significantly lower in 80E compared to 20E. However, the numbers of blood vessels were significantly higher in H, 20EH, and 80EH compared to 80E. The number of ovarian haemorrhage was significantly lower in 20E, 80E, H, 20EH and 80EH compared to C. The number of corpora lutea was significantly lower in 80EH, H, 80E and 20E compared to C. However, the number of corpora lutea was significantly higher in 20EH compared to 20E and H.

Conclusion: This study suggested that jumping exercises in particularly high-intensity exercise may induce histopathological changes in uterus and ovary in rats, and honey supplementation may ameliorate these effects.

Introduction

Apart from the important health benefits earned by exercise and physical activity, high-intensity physical activity may accompany with female reproductive disorders. Low energy availability and stress induced by intense exercise may cause hypothalamic dysfunction such as disturbance of gonadotropin releasing hormone (GnRH) pulsatility and hypoestrogenism which in turn may result in menstrual disorder, infertility and osteoporosis (1–3). On the other side, high intensity and prolonged physical activity increase production of reactive oxygen species (ROS) by metabolic and physiological processes and cause cellular damage such as lipid damage, depletion of adenosine triphosphate and inhibition of protein synthesis (4). Furthermore, ROS may influence the biology of the female reproduction system at the levels of the ovary, follicle, and oocyte (5). The increased stress hormone concentrations such as cortisol reduce estradiol production by reducing the function of granulosa cell within the follicle, which lead to deteriorating oocyte quality (5). Recently, we have reported that jumping exercise at high-intensity level could cause an increment in serum cortisol levels, and it was accompanied with lower levels of luteinizing hormone (LH), follicle stimulating hormone (FSH) and progesterone in adult female rats (6). It also has been found that an intense and exhaustive exercise programme is accompanied by the reduction in uterus thickness in female rats (7).

Several non-pharmacological therapies such as increased caloric intake reduced exercise energy expenditure, and pharmacological treatment has been described to prevent and/ or treatment of

disorders caused by intense exercises (8). Honey is a natural complex of sugars contains carbohydrates such as fructose, glucose, raffinose and sucrose, and flavonoids, enzymes, antioxidants, minerals, organic acids, proteins, phenolic acids, phytochemicals and vitamins such as vitamins C and E (9). Tualang honey is a wild multi-floral honey found in the Malaysian Rain Forest with intermediate glycemic index as well as phenolic compounds and antioxidant activity (10), and the flavonoids present in honey particularly kaempferol and quercetin have been revealed to have estrogenic activity which may be beneficial for female reproductive health (11,12). It has been reported that administration of 0.2g/ kg Tualang honey elicited beneficial effects in the enhancement of uterus weight and thickness of uterus endometrium and vaginal epithelium in ovariectomized rats (13). Furthermore, we have shown that administration of Tualang honey at 1g/kg body weight could reduce the adverse effect induced by exercise on female reproductive hormones in rats (6,14). To date, however, whether different intensities of exercise may induce changes on histology of uterus and the possible protective effect of honey supplementation have not yet been reported.

Therefore, we aimed to investigate the effect of different jumping exercise intensities on uterus histology and the possible protective effect of honey supplementation in rats.

Materials & Methods

Seventy-two, 9-week old female Sprague- Dawley rats with no significant mean difference in initial body weights (nearest 0.1 g) were entered in this study (7, 2). The experimental protocol was approved by the Animal Ethics Committee, Universiti Sains Malaysia [16 August 2011, Reference: USM/Animal Ethics Approval/ 2011/ (71) (325)], and has been described previously (6,14).

The estrous cycle is the main reproductive cycle of rodents that lasts four days and is categorized as: proestrus, estrus, metestrus (diestrus I) and diestrus (diestrus II), which can be determined according to the cell types observed in the vaginal smear (15). Diestrus phase, which lasts 55–57 h and more than half of the cycle (16), suits to observe the population of different types of cells. To aim of standardization in hormonal phase, the rats were scrutinized to detect diestrus phase two times; beginning and end of the experiment. Vaginal secretion of each rat was collected by flushing the vagina with 1 µL of normal saline (0.9%) using a clean pipette. The unstained wet smear was observed under a light microscope at 100x magnifications, similar pattern was used as standard for determining diestrus phase of the rats. The rats were assigned into 6 experimental groups (12 rats in each group) by block-randomization; control group with free cage activity and no intervention (C), low intensity; 20 jumps per day at 5 days per week for 8 weeks (20E), high intensity; 80 jumps per day for 5 days per week for 8 weeks (80E), honey supplementation for 7 days per week for 8 weeks (H), 20 jumps per day for 5 days per week combined with honey supplementation for 8 weeks (20EH), and 80 jumps per day for 5 days per week combined with honey supplementation for 8 weeks (80EH). Each rat in the training groups was positioned at the

bottom of a specially designed wooden box and jumping was commenced by applying an electrical grid and after a few days of training, the rats jumped without electrical stimulation. Each jump took 4 seconds. The rats in the control group (C) were also handled during the duration of the study to imitate the stress induced by handling. At the end of the experiment, the rats were anaesthetized using chloroform by lying for 2-3 minutes in a dried jar containing chloroform-soaked gauze pad and then they were decapitated in diestrus phase (Scientific Research Instrument, U. K)(6,14).

After decapitation, the uterus horns and ovaries were carefully removed and cleaned. The required organs were fixed in 10% neutral buffered formalin. These specimens were stored in a screw-capped specimen container prior to the next step procedures within a reasonable time. Each tissue was subjected to tissue fixation, processing, paraffin wax embedding, section cutting, and staining according to the previous study (17). Two histological slices were made from each organ for each rat. The standard hematoxylin and eosin (H& E) stained tissues of ovaries and uterine horns were observed for any changes under a light microscope attached with an image analysis system. Primary follicle consists of a central oocyte surrounded by a single layer of cuboidal granulosa cells. Graafian follicles is a large and mature follicle that has an antrum containing follicular fluid. Corpus hemorrhagicum is a temporary structure formed immediately after ovulation from the ovarian follicle as it collapses and is filled with blood clot. Corpus luteum is formed after ovulation which consists of luteinized granulosa and theca cells. Ovarian luteal cyst is corpus luteum that has large cystic centre. Endometrium is the tissue lining the inner cavity of the uterus and is bordered with simple epithelium line and comprises plentiful tubular glands. The myometrium is the middle layer of the uterine wall and comprised of a three smooth muscle layers, which are microscopically difficult to distinct. Uterine blood vessels is the myometrium that has strong and well-perfused vascular layer that is quite running around the uterus. The uterine gland or endometrial gland is a simple tubular gland shaped by invagination of the uterine endometrium. Qualitative and quantitative parameters of ovaries and uterine horns including the thickness of endometrium and myometrium, as well as the number of uterine glands and blood vessels were observed and measured. The mean of uterus thickness, including endometrium and myometrium of each rat was measured based on the maximum and minimum sagittal section thickness in μm . The number of uterine glands and blood vessels was counted once per slide for each rat and then the mean was calculated (Figure 1). Regarding ovaries, the number of Graafian follicles, primary follicle, cyst, haemorregia, corpora lutea, and blood vessels were counted twice for each rat and then the mean was calculated (Figure 2).

Honey supplementation

The rats were fed with Malaysian Tualang honey at a dosage of 1 g/ kg body weight/ day by oral gavage for 7 days per week for 8 weeks. The rats in combined honey with jumping exercise groups (20EH & 80EH) were fed with honey, 30 minutes prior to the exercise session (6). The honey dosage was prescribed based on the rat biweekly body weight.

Statistical analysis

The numerical data were studied using SPSS version 18.0. All variables were assessed for normality using the Normality Test as well as the Levene's Test to check for homogeneity of variance at 5% level of significance. One-way analysis of variance (ANOVA) with post-hoc test were performed to determine the significance of the differences between groups. The results are presented as mean \pm standard error (SE). The p -value of < 0.05 was defined as statistically significant and used for all the comparisons.

Results

The data of body weights and reproductive organ weights presented in Table 1. The rats were weight matched at the beginning of the study, and there was no significant difference in weight gain percentage among all the experimental groups. There were also no significant differences in weight of ovarian among all the groups. Mean uterine weight in 20EH group was significantly greater compared to control, and there were no significant differences in uterine weight among all the experimental groups compared to control.

The quantitative histopathological findings and cross sections of uterine glands and blood vessels, the thickness of myometrium and endometrium are presented in Table 2 and Figure 1, respectively. Mean number of primary and Graafian follicles, corpora lutea, cysts, haemorrhagia, blood vessels and cross sections of the ovaries are presented in Table 3 and Figure 2, respectively.

The present data show that the 80 jumps/per day were associated with lower endometrium thickness in comparison with the control group, however, results were not significant at 0.05 level. The endometrium was significantly thicker in the rats in H, 20EH and 80EH groups compared to the rats in C, 20E and 80E. The myometrium thicknesses of the rats in 80E were significantly lower compared to controls. However, the myometrium thickness was significantly higher in 80EH compared to C. There was significantly higher myometrium thickness in 20EH and 80EH compared to 20E and 80E. Additionally, the myometrium thickness was significantly higher in 20EH and 80EH compared to H.

Regarding numbers of glands and blood vessels of the uterus, it was found that the numbers of glands of the uterus in 20E and 80E were significantly lower compared to C. However, there was a significantly high level of glands number in H, 20EH, and 80EH compared to 20E and 80E. The numbers of uterus blood vessels were significantly lower in 80E compared to 20E. However, the numbers of blood vessels were significantly higher in H, 20EH, and 80EH compared to 80E.

The histopathological finding of ovaries was revealed that the number of Graafian follicles did not differ between all experimental groups with exception of 20EH with significantly ($P<0.05$) greater count of follicles compared to C, J20 and J80. The number of primary follicles ($P<0.05$) were

higher in 80E, 20EH and 80EH compared to C. There were significantly ($P<0.05$) higher of primary follicles in 20EH and 80EH compared to H. There was no significant difference in the number of ovarian cysts among the experimental groups. There was no count of haemorrhagia in 20EH group. The number of haemorrhagia were significantly ($p<0.05$) lower in 20E, 80E, H and 80EH compared to C. The number of haemorrhagia were significantly ($p<0.05$) lower in 20E, H and 80EH compared to J80. The number of corpora lutea were significantly ($p<0.05$) lower in 80EH compared to H. However, the number of corpora lutea were significantly ($p<0.05$) higher in 20EH compared to J20 and H. The number of ovary blood vessels was significantly ($p<0.05$) higher in H compared to 80E. However, the levels of ovary blood vessels in HJ20 were significantly higher compared to C, J20, J80, and 80EH.

Discussion

This study firstly evaluated the probable histomorphometric changes of female reproductive organs influenced by different jumping exercise intensities. We observed that jumping exercise mainly 80 jumps per day (high-intensity exercise) induced negative effects on the measured histopathological parameters with significantly decreased in the number of uterus glands and myometrium thickness. Consequently, we observed that honey supplementation has played an effective role in diminishing these adverse effects induced by jumping exercise on uterus as well as improvement in ovary parameters characteristics.

It has been shown that intense exercise is commonly correlated with the boosted generation of ROS which may damage tissue and organ redox homeostasis (18). Furthermore, it has also been shown that intense exercise-induced ischemic/reperfusion actions can be related to increased ROS production due to the deviation of the cardiac output to muscle mass inactivity and skin, consequently causing ischemia in the pelvis section (19). Our finding showed that honey supplementation may have potential in increasing the endometrium thickness in the rats feeding with honey alone or combined with 20 and 80 jumps/ day compared to the rats in control and exercise alone groups. Similar effects of honey supplementation on reproductive organs have been reported in the study done by Zaid et al. (15) on ovariectomized rats, in which administration of honey supplementation for 2 weeks significantly increased the weight of the uterus and the thickness of vaginal epithelium. This previous study showed that improvement of uterus and vagina atrophy which might be attributed to the biologically active estrogen-like molecules or phytoestrogens exists in honey supplementation. In the present study, the myometrium thickness was significantly lower in 80 jumps/ day group compared to the control group suggesting that there were negative effects of high exercise intensities on the female reproductive system. Previously, we have shown that high intensity of jumping exercises elicited negative effects on follicle stimulating hormone (FSH), luteinizing hormone (LH) and progesterone concentrations in female rats which can possibly explain the changes in the thickness of endometrium and myometrium (6,14). It has been suggested by Warren and Perlroth (1) that stress induced by exercise can arrest

the gonadal function, through the increased glucocorticoids and catecholamines levels with activation of the corticotropin-releasing hormone neurons. Meanwhile, low caloric intake and high caloric expenditure, which can occur among athletes, suppress reproductive function (20), and this could be a suppressor factor for the gonadotropin-releasing hormone. In the present study, we observed that the myometrium was thicker in 20 and 80 jumps/ day combined with honey supplementation groups compared to the control group and the rats with jumping alone, i.e. 20 and 80 jumps/ day groups. Additionally, the myometrium thickness was significantly higher in 20EH and 80EH compared to H. Regarding the number of glands and blood vessels of the uterus, it was found that the numbers of glands of the uterus in 20E and 80E were significantly lower than C, and the numbers of uterus blood vessels were significantly lower in 80E compared to 20E. These results implied that jumping exercise mainly high intensity with 80E /day may elicit negative effects on these measured parameters. We also found that there were increases in these measured parameters in jumping groups with honey supplementation compared to exercise groups without honey supplementation. Flavonoids contained in the honey with their antioxidant compounds may able to retard biologically damaging chemical reactions in living organisms through their ability to scavenge oxidants and free radicals (21). This positive effects of honey on endometrium may due to its high nutritional contents, particularly flavonoids which have antioxidant property (22) in scavenging ROS that may occur during exercise (16, 17). Furthermore, based on Oh et al (12) and Jaganathan (11) findings, flavonoids specially kaempferol and quercetin are natural phytoestrogens as they show the estrogenic property. It also was reported that phytoestrogens were associated with an increased incidence of endometrial proliferation (23). Furthermore, these positive effects of honey particularly in combination with jumping exercise on myometrium thickness, number of gland and blood vessels may also be due to increase in supply of honey's components, i.e. sugars, phenolic acid and flavonoids through the increased of blood flow to the muscle (24) induced by jumping exercise and these vital elements contained in honey may have beneficial effects on uterus.

Our finding showed that high and low intensity jumping combined with honey supplementation were most effective in primary follicle generation in female rats. This observation is similar to the findings of the study conducted by Zaid et al. (23) in which the researcher showed that animals treated with Bisphenol A together with Tualang honey exhibited an improvement in the percentage of normal oestrous cycle compared to those treated with Bisphenol A. This finding may be attributed to the effect of honey on improving the normal oestrous cycle. The noteworthy histological findings of ovaries showed that low-intensity jumping exercise containing 20 jumps per day combined with 1 g per kg body weight of daily honey supplementation caused more obvious beneficial effects on the number of Graafian follicles, corpora lutea, and ovarian blood vessels formation compared to other groups in female rats.

The appearance of corpora lutea is considered the occurrence of ovulation. These results could be explained with this fact that exercise might promote follicular maturation and ovulation by

decreasing sympathetic activity. The possible justification for the improvement of cyclicity is that reduced sympathetic activity caused by exercise may have a direct impact on the ovaries and sex steroid synthesis pathways (26). It has been shown that physical exercise decreased sympathetic nerve activity, improve menstrual frequency, and improved hyperandrogenism in women with the polycystic ovarian syndrome (27).

The findings of this study showed that honey supplementation induced by different jumping exercise intensities caused no significant difference in the number of ovarian cysts in rats. The cystic follicle is formed from anovulatory follicle encircled by thin layers of granulosa cells with non-detectable theca cell layers (25). These findings suggest that exercise promotes follicular maturation and ovulation. The possible clarification for the improvement of cyclicity is that reduced sympathetic action may have a direct effect on the ovaries and female steroid production pathways. It has been reported that physical activity improved menstrual cycle through a reduction in sympathetic nerve activity and the levels of several sex steroids in women diagnosed with the polycystic ovarian syndrome (27).

Conclusions

In summary, this study demonstrated that honey has a protective effect against the histological and structural changes induced by jumping exercise on uterus and ovary in rats.

Acknowledgements

The authors would like to acknowledge Universiti Sains Malaysia for providing the research grant (No: 304 /PPSP/61312031). We would also like to thank the staff of Physiology and Sports Science Laboratories, School of Medical Sciences, Universiti Sains Malaysia.

References

- Warren MP, Perlroth NE. The effects of intense exercise on the female reproductive system. *J Endocrinol*. 2001;170(1):3–11.
- Mountjoy M, Sundgot-Borgen J, Burke L, Carter S, Constantini N, Lebrun C, et al. The IOC consensus statement: beyond the female athlete triad—Relative Energy Deficiency in Sport (RED-S). *Br J Sports Med*. 2014;48(7):491–497.
- De Souza MJ, Williams NI. Physiological aspects and clinical sequelae of energy deficiency and hypoestrogenism in exercising women. *Hum Reprod Update*. 2004;10(5):433–448.
- Powers SK, Radak Z, Ji LL. Exercise-induced oxidative stress: past, present and future. *J Physiol*. 2016;594(18):5081–5092.
- Prasad S, Tiwari M, Pandey AN, Shrivastav TG, Chaube SK. Impact of stress on oocyte quality and reproductive outcome. *J Biomed Sci*. 2016;23(1):36.

6. Mosavat M, Ooi FK, Mohamed M. Stress hormone and reproductive system in response to honey supplementation combined with different jumping exercise intensities in female rats. *BioMed Res Int.* 2014;2014.
7. Costa AE, Silva JL, Simões MJ, Nouailhetas VL. Morphofunctional alterations of the nonpregnant murine uterus in response to intense and exhaustive exercise are not related to oxidative stress. *J Appl Physiol.* 2014;116(6):604–610.
8. De Souza MJ, Nattiv A, Joy E, Misra M, Williams NI, Mallinson RJ, et al. 2014 Female Athlete Triad Coalition Consensus Statement on treatment and return to play of the female athlete triad: 1st International Conference held in San Francisco, California, May 2012 and 2nd International Conference held in Indianapolis, Indiana, May 2013. *Br J Sports Med.* 2014;48(4):289–289.
9. Aljadi AM, Kamaruddin MY. Evaluation of the phenolic contents and antioxidant capacities of two Malaysian floral honeys. *Food Chem.* 2004;85(4):513–518.
10. Mohamed M, Sirajudeen KNS, Swamy M, Yaacob M, Sulaiman S. Studies on the antioxidant properties of Tualang honey of Malaysia. *Afr J Tradit Complement Altern Med.* 2010;7(1).
11. Jaganathan SK, Mandal M. Antiproliferative effects of honey and of its polyphenols: a review. *BioMed Res Int.* 2009;2009.
12. Oh SM, Chung KH. Antiestrogenic activities of Ginkgo biloba extracts. *J Steroid Biochem Mol Biol.* 2006;100(4–5):167–176.
13. Zaid SS, Sulaiman SA, Sirajudeen KN, Othman NH. The effects of Tualang honey on female reproductive organs, tibia bone and hormonal profile in ovariectomised rats-animal model for menopause. *BMC Complement Altern Med.* 2010;10(1):82.
14. Mosavat M, Ooi FK, Mohamed M. Effects of honey supplementation combined with different jumping exercise intensities on bone mass, serum bone metabolism markers and gonadotropins in female rats. *BMC Complement Altern Med.* 2014;14(1):126.
15. Marcondes FK, Bianchi FJ, Tanno AP. Determination of the estrous cycle phases of rats: some helpful considerations. *Braz J Biol.* 2002;62(4A):609–614.
16. Hubscher CH, Brooks DL, Johnson JR. A quantitative method for assessing stages of the rat estrous cycle. *Biotech Histochem.* 2005;80(2):79–87.
17. Suvarna KS, Layton C, Bancroft JD. Bancroft's Theory and Practice of Histological Techniques E-Book. Elsevier Health Sciences; 2012.
18. Quindry JC, Kavazis AN, Powers SK. Exercise-Induced Oxidative Stress: Are Supplemental Antioxidants Warranted? *Encycl Sports Med IOC Med Comm Publ Vol* 19. 2013;263–276.
19. Vollaard NB, Shearman JP, Cooper CE. Exercise-induced oxidative stress. *Sports Med.* 2005;35(12):1045–1062.
20. Gibbs JC, Williams NI, Scheid JL, Toombs RJ, De Souza MJ. The association of a high drive for thinness with energy deficiency and severe menstrual disturbances: confirmation in a large population of exercising women. *Int J Sport Nutr Exerc Metab.* 2011;21(4):280–290.
21. Bertoncelj J, Doberšek U, Jamnik M, Golob T. Evaluation of the phenolic content, antioxidant activity and colour of Slovenian honey. *Food Chem.* 2007;105(2):822–828.
22. Buratti S, Benedetti S, Cosio MS. Evaluation of the antioxidant power of honey, propolis and royal jelly by amperometric flow injection analysis. *Talanta.* 2007;71(3):1387–1392.

23. Unfer V, Casini ML, Costabile L, Mignosa M, Gerli S, Di Renzo GC. Endometrial effects of long-term treatment with phytoestrogens: a randomized, double-blind, placebo-controlled study. *Fertil Steril*. 2004;82(1):145–148.
24. Laughlin MH, Roseguini B. Mechanisms for exercise training-induced increases in skeletal muscle blood flow capacity: differences with interval sprint training versus aerobic endurance training. *J Physiol Pharmacol Off J Pol Physiol Soc*. 2008;59(Suppl 7):71.
25. Zaid SSM, Othman S, Kassim NM. Potential protective effect of Tualang honey on BPA-induced ovarian toxicity in prepubertal rat. *BMC Complement Altern Med*. 2014;14(1):509.
26. Wu C, Lin F, Qiu S, Jiang Z. The characterization of obese polycystic ovary syndrome rat model suitable for exercise intervention. *PloS One*. 2014;9(6):e99155.
27. Benrick A, Maliqueo M, Miao S, Villanueva JA, Feng Y, Ohlsson C, et al. Resveratrol is not as effective as physical exercise for improving reproductive and metabolic functions in rats with dihydrotestosterone-induced polycystic ovary syndrome. *Evid Based Complement Alternat Med*. 2013;2013.

Table 1(on next page)

Body weights and reproductive organ weights (mean \pm SE)

*significant from C ($p < 0.05$)

Title

Table 1. Body weights and reproductive organ weights (mean ± SE)

Group	Initial body weight (g)	Weight gain% (g)	Ovarian weight (mg)	Uterine weight (mg)
C	186.7(16.5)	76.2 (2.0)	0.10(0.02)	0.41(0.1)
H	192.5(16.3)	76.6 (1.5)	0.10(0.03)	0.46(0.1)
20E	188.9 (10.5)	75.4 (1.1)	0.10(0.02)	0.48(0.1)
80E	189.1(9.3)	75.8(1.9)	0.11(0.02)	0.48(0.1)
20EH	191.6 (13.5)	78.1(2.3)	0.10(0.02)	0.53(0.3) *
80EH	193.3(8.6)	76.3(1.8)	0.10(0.01)	0.45(0.1)

*significant from C (p<0.05)

Table 2 (on next page)

Uterus quantitative histopathological findings and presented as mean \pm SE

^a significant from C (P<0.05), ^b significant form 20E (P<0.05), ^c significant from 80E(P<0.05), ^d significant from H (P<0.05)

Title

Table 2. Uterus quantitative histopathological findings and presented as mean \pm SE

Groups	No. uterus glands	No. uterus vessels	Endometrium thickness (μ m)	Myometrium thickness (μ m)
C	22.2(1.9)	8.4(2.4)	3.6(0.5)	107.1(7.9)
20E	13.9(1.2) ^a	13.2(3.9)	3.6(0.6)	89.2 (6.6) ^a
80E	12.6(2.1) ^a	7.3(1.7) ^b	2.2(0.4)	79.9(7.7) ^a
H	21.9(2.3) ^{b,c}	18.5(3.9) ^c	7.5(0.9) ^{a,b,c}	96.3(6.6)
20EH	21.6(2.1) ^{b,c}	19.3(3.8) ^c	6.3(0.9) ^{a,b,c}	121.9(9.2) ^{b,c,d}
80EH	21.6(3.6) ^{b,c}	21.5(2.8) ^c	6.1(0.5) ^{a,b,c}	140.6(8.2) ^{a,b,c,d}

^a significant from C (P<0.05), ^bsignificant form 20E (P<0.05), ^c significant from 80E(P<0.05),

^d significant from H (P<0.05)

Table 3(on next page)

Ovary quantitative histopathological findings and presented as mean \pm SE

^a significant from C (P<0.05), ^b significant form 20E (P<0.05), ^c significant from 80E(P<0.05), ^d significant from H (P<0.05), ^e significant from 20EH (P<0.05)

Title

Table 3. Ovary quantitative histopathological findings and presented as mean \pm SE

Groups	Primary follicle	Graafian follicle	Corpora lutea	Ovarian luteal cysts	Corpus hemorrhagicum	Blood vessels
C	4.1(1.4)	7.7(1.4)	8.4(1.0)	7.3(1.3)	2.2(0.9)	7.0(1.3)
20E	2.2(0.3)	6.0(0.6)	6.7(0.9)	7.9(2.1)	0.3(0.2) ^a	7.2(1.8)
80E	5.9(1.5) ^b	7.4(1.4)	8.8(1.7)	6.1(2.0)	1.8(0.5) ^{a,b}	6.0(1.3)
H	2.3(0.4) ^c	9.7(1.0)	8.1(1.6)	6.5(1.5)	0.2(0.2) ^{a,c}	11.9(2.8) ^c
20EH	6.4(1.1) ^{b,d}	12.4(1.6) ^{a,b,c}	12.5(2.3) ^{b,d}	6.0(1.9)	0.0	15.1(2.4) ^{a,b,c}
80EH	7.3(1.6) ^{b,d}	8.7(1.7)	7.0(1.2) ^e	7.6(2.1)	0.2(0.2) ^{a,c}	7.8(2.5) ^e

^a significant from C (P<0.05), ^b significant form 20E (P<0.05),

^c significant from 80E(P<0.05), ^d significant from H (P<0.05), ^e significant from 20EH (P<0.05)

Figure 1

Cross sections of the uterus (Magnification 4X with scale bar: 500µm and 10X with scale bar: 200µm) of A: Control (C), B: H (honey), C: 20E (20 jumps/day), D: 80E (80 jumps/day), E: 20EH (20 jumps and honey supplementation), F: 80EH (80 jumps and honey

supplementation). G: Gland; M: Myometrium; E: Endometrium; BV: Blood vessels.




Figure 2

Cross sections of the ovaries (Magnification 4X with scale bar: 500µm and 10X with scale bar: 200µm). A: Control (C), B: H (honey), C: 20E (20 jumps/day), D: 80E (80 jumps/day), E: 20EH (20 jumps and honey supplementation), F: 80EH (80 jumps and honey supplementation). PF: Primary follicle; CL: Corpus Lutea; GF: Graafian follicle; BV: Blood vessels; C: Ovarian luteal cyst; H: Corpus hemorrhagicum.

