

***Nurhachius luei*, a new istiodactylid pterosaur (Pterosauria, Pterodactyloidea) from the Early Cretaceous Jiufotang Formation of Chaoyang City, Liaoning Province (China); and comments on the Istiodactylidae**

Xuanyu Zhou^{1,2}, Rodrigo Vargas Pêgas³, Maria Eduarda de Castro Leal^{4,5}, Niels Bonde^{5,6}

1. Institute of Geology, Chinese Academy of Geological Sciences, Beijing, China
2. China University of Geosciences, Beijing, China
3. Laboratory of Vertebrate Paleontology and Animal Behavior, Universidade Federal do ABC, São Bernardo, São Paulo, Brazil
4. Departamento de Geologia, Universidade Federal do Ceará, Fortaleza, Ceará, Brazil
5. Zoological Museum (SNM), Copenhagen University, Copenhagen, Denmark
6. Fur Museum (Museum Saling), Fur, Denmark

Corresponding Author:

Xuanyu Zhou

No.26 Baiwanzhuang Street, Beijing, Beijing, 100037, China

Email address: zhouxy2017@yeah.net

Rodrigo Vargas Pêgas

Alameda da Universidade, s/n - Anchieta, São Bernardo do Campo - SP, 09606-045, Brazil

Email address: rodrigo.pegas@hotmail.com

Abstract

A new istiodactylid pterosaur, *Nurhachius luei* sp. nov., is here reported based on a complete skull with mandible and some cervical vertebrae from the lower part of the Jiufotang Formation of western Liaoning (China). This is the second species of *Nurhachius*, the type-species being *N. ignaciobrito* from the upper part of the Jiufotang Formation. A revised diagnosis of the genus *Nurhachius* is presented. In this genus, a slight dorsal deflection of the palatal anterior tip is present, which is homoplastic with

Comentario [GP1]: Replace by “provided, being this taxon characterized by the presence of a slight dorsal deflection of the palatal anterior tip, ”

Con formato: Diseño: Claro (Amarillo), Resaltar

Con formato: Resaltar

Con formato: Resaltar

the Anhangueria and *Cimoliopterus*. *Nurhachius luei* sp. nov. show an unusual pattern of tooth replacement. The relationships within the Istiodactylidae and with their closest taxa are investigated through a phylogenetic analysis by parsimony.

Eliminado: s

Comentario [GP2]: Insert after “replacement “: “, with respect to other dactyloid species.”

Introduction

Istiodactylid pterosaurs are characterized by rhombic teeth with lancet-shaped crowns, long skulls with short pre-antorbital portions of the rostrum, and nasoantorbital fenestrae representing over 50 percent of the total skull length and height (Howse *et al.*, 2001; Andres & Ji, 2006; Lü *et al.*, 2013). The group was originally named by Howse *et al.* (2001) in order to accommodate, then, only *Istiodactylus latidens*. Later, the Istiodactylidae were phylogenetically defined by Andres *et al.* (2014) as the least inclusive clade containing *Nurhachius* and *Istiodactylus*.

Four pterosaur genera and five species (all represented by a single specimen) have been referred to the Istiodactylidae sensu Andres *et al.* (2014) in the literature, namely *Istiodactylus latidens*, *I. sinensis*, *Liaoxipterus brachyognathus*, *Nurhachius ignaciobritoi* and *Longchengpterus zhaoi*. However, *Longchengpterus zhaoi* has been considered a junior synonym of *N. ignaciobritoi* by Lü *et al.* (2008), a view that is followed here (see Discussion). Therefore, *N. ignaciobritoi* is the only Chinese istiodactylid species to be represented by two specimens so far. *Haopterus gracilis*, *Hongshanopterus lacustris* and *Archaeoistiodactylus linglongtaensis* have been reported in literature as taxa that are close to the Istiodactylidae (Wang XL & Lü, 2001; Wang XL *et al.*, 2008a; Lü & Fucha, 2010). However, the affinity of *Archaeoistiodactylus linglongtaensis* has been questioned by Sullivan *et al.* (2014).

All istiodactylid pterosaurs are from the Early Cretaceous Jiufotang Formation of northeastern China with the exception of *Istiodactylus latidens*, which is from the Early Cretaceous Vectis Formation of the Isle of Wight, Southern England. Also, the three taxa that are reported as close to istiodactylids come from northeastern China and surrounding areas: *Haopterus gracilis* is from the Early Cretaceous Yixian Formation, *Hongshanopterus lacustris* from the Jiufotang Formation, and *Archaeoistiodactylus linglongtaensis* from the Middle Jurassic Tiaojishan Formation. Apart from the latter, these Chinese pterosaurs belong to the Jehol Biota (see Chang *et al.*, 2003).

By the end of 2016, 23 species of pterosaurs from the Jiufotang Formation have been

reported (Andres & Ji, 2006; Dong & Lü, 2005; Dong *et al.*, 2003; Jiang *et al.*, 2016; Rodrigues *et al.*, 2015; Li *et al.*, 2003; Lü & Ji, 2005; Lü & Yuan, 2005; Lü *et al.*, 2006, 2007, 2008, 2016a, 2016b; Wang L *et al.*, 2006; Wang XL & Zhou, 2003a, 2003b; Wang XL *et al.*, 2005, 2008a, 2008b, 2012, 2014).

In this paper, we describe a second species of *Nurhachius* from the Jiufotang Formation and investigate the phylogenetic relationships of the istiodactylids and purported close taxa.

Geological, paleontological and geochronological information

The Jiufotang Formation is known worldwide for its paleontological richness and the exquisite preservation of its fossils, which include plants, insects, fishes, mammals, birds, non-avian dinosaurs and pterosaurs (Wang X, 2018; Meng *et al.*, 2011; Wang M and Zhou, 2019; Yao *et al.*, 2019). Fossils occur mainly in the lower part of the formation, known as Boluochi Beds or Boluochi Member (see Chang *et al.*, 2003), which is characterized by the *Jinanichthys* – *Cathayornis* Fauna that includes small feathered dinosaurs like the four-winged *Microraptor* (Xu *et al.*, 2003) and several pterosaurs (Chang *et al.*, 2003; Zhou *et al.*, 2003).

The Jiufotang Formation is 206–2685 m thick according to Chang *et al.* (2009) and is mainly composed of mudstone, siltstone, shale, sandstone and tuff. A tuff from the basal part of formation (two meters above the boundary between the Yixian and Jiufotang formations) in western Liaoning was dated to 122.1 ± 0.3 Ma by Chang *et al.* (2009). A basalt in the upper part of the formation in Inner Mongolia was dated to 110.59 ± 0.52 Ma by Eberth *et al.* (1993), but Chang *et al.* (2009) objected that the correlation between the Jiufotang Formation in Liaoning and Inner Mongolia is unclear and the age of the uppermost Jiufotang Formation remains unknown. The Aptian age of the Early Cretaceous ranges ~125–113 Ma according to the International Chronostratigraphic Chart 2018/08. Therefore, the Jiufotang Formation is Aptian in age, but might reach the Albian.

The Jiufotang Formation and the underlying Yixian Formation traditionally constitute the Jehol Group. The Yixian Formation is 225–4000 m thick, varying in thickness and lithology in different areas according to Chang *et al.* (2009), but only a fraction is made of sedimentary rocks because basalts and lavas represents a substantial part of the

Comentario [GP3]: Not in references

Con formato: Resaltar

Con formato: Fuente: Cursiva

section. Chang *et al.* (2009) dated the basal part of the Yixian Formation in Western Liaoning to 129.7 ± 0.5 and the uppermost part of the underlying Tuchengzi Formation to 139.5 ± 1.0 Ma. The upper part of the Yixian Formation (the Jingangshan Beds) was dated to 126.5 Ma (Chang *et al.*, 2003). Therefore, the Yixian Formation represents an interval of ~ 7 Ma from early Barremian to early Aptian and the Jiufotang Formation might represents an interval of over 11 Ma from early Aptian to early Albian.

The Jehol Group has yielded the famous Jehol biota. Four fossil-bearing levels with partly different fossil associations have been distinct within the Yixian Formation and only one (corresponding to the Boluochi Beds) in the Jiufotang Formation (Chang *et al.* 2003).

Both the holotype of *N. ignaciobritoi* and its referred specimen (the holotype of *Longchengpterus zhaoi*) come from the upper part of the Jiufotang Formation (see Wu *et al.*, 2018), whereas the new species comes from the Boluochi Beds (lower part of the Jiufotang Formation).

Material & Methods

The holotype and only specimen of the new species consists of a skull with mandible and seven articulated cervical vertebrae. It was previously figured in Lü *et al.* (2013, figures at pp. 81-82) and reported as an unnamed istiodactylid. The specimen was found near the village of Huanghuatan (Dapingfang town, Chaoyang City, western Liaoning).

For the comparisons we present, the following taxa/specimens were analyzed first-hand (by XZ): *Nurhachius ignaciobritoi* (both specimens, LPM 00023 and IVPP V-13288), *Liaoxipterus brachyognathus* holotype (CAR-0018), *Hongshanopterus lacustris* holotype (IVPP V14582) and *Haopterus gracilis* holotype (IVPP V11726). Data from other taxa was gathered from the literature.

A phylogenetic analysis was performed based on the data matrix by Holgado *et al.* (2019) modified with the inclusion of characters by Lü *et al.* (2008), Witton (2012), Andres *et al.* (2014), and new characters; and the addition of the following taxa:

Archaeoistiodactylus linglongtaensis, *Kunpengopterus sinensis*, *Liaoxipterus brachyognathus* and *Nurhachius luei* sp. nov. (see SI). The analysis was performed by TNT (Goloboff *et al.*, 2008) using the Traditional Search option, 10000 replicates, random seed = 0 and collapsing trees after search. The character and character states list and the TNT file with the data matrix are available in the SI.

The electronic version of this article in Portable Document Format (PDF) will represent a published work according to the International Commission on Zoological Nomenclature (ICZN), and hence the new names contained in the electronic version are effectively published under that Code from the electronic edition alone. This published work and the nomenclatural acts it contains have been registered in ZooBank, the online registration system for the ICZN. The ZooBank LSIDs (Life Science Identifiers) can be resolved and the associated information viewed through any standard web browser by appending the LSID to the prefix <http://zoobank.org/>. The LSID for this publication is:

urn:lsid:zoobank.org:pub:03EF173E-4AB5-4C74-B80C-A6AAFA65E61C. The online version of this work is archived and available from the following digital repositories: PeerJ, PubMed Central and CLOCKSS.

Results

Systematic Paleontology

Pterosauria Kaup, 1834

Pterodactyloidea Plieninger, 1901

Istiodactylidae Howse *et al.*, 2001 (sensu Andres *et al.*, 2014)

Nurhachius Wang XL *et al.*, 2005

Type species. *Nurhachius ignaciobritoi* Wang XL *et al.*, 2005

Synonym. *Longchengpterus zhaoi* Wang L *et al.*, 2006

Emended Diagnosis. Istiodactylids that share the following features: slight dorsal deflection of the palatal anterior tip; orbit piriform; craniomandibular joint located

under the anterior margin of the orbit; dentary symphysis about one third the length of the mandible; dentary symphysis with gradual taper of the lateral margins; triangular, laterally compressed teeth lacking carinae; crowns with both labial and lingual slight concavities; slight constriction between tooth crown and root.

Nurhachius luei sp. nov.

ZooBank **LSID** **for** **species.**
urn:lsid:zoobank.org:act:6F93DC7F-20A7-4CBC-8A38-1D6C802A1906.

Etymology. The specific name *luei* (/ lyi/) honors the late Prof. Junchang Lü, who has made great contributions to the study of Chinese pterosaurs.

Holotype. Skull, mandible and seven cervical vertebrae (BPMC-0204). The specimen is permanently deposited and available for researchers at a public repository, the Beipiao Pterosaur Museum of China, Beipiao, Liaoning Province, China (Fig. 1).

Type Locality and Horizon. Huanghuatan village, Dapingfang town, Chaoyang City, Liaoning Province, China (Fig. 2); lower part of the Jiufotang Formation, Early Cretaceous (Aptian).

Differential diagnosis. The new species is diagnosed based on the following features: quadrate inclined at 150°; medial process of the pterygoid broad and plate-like; dorsal median sulcus of the mandibular symphysis extending up to the first pair of mandibular teeth; dorsally directed odontoid (pseudotooth) of the mandibular symphysis, lacking a foramen on the lateral side and with a blunt occlusal surface; ceratobranchial I of the hyoids accounting for 60% of mandibular length; mandibular teeth extending distally beyond the symphysis.

Description

Skull and mandible. The skull is exposed in right lateral view, with some palatal elements that are visible in dorsal view. The mandible is exposed in right dorsolateral view. The skull is 300 mm long from the squamosal to the premaxillary tip (total skull length), and 74 mm high at its greatest height, which is at the level of the occiput. The nasoantorbital fenestra is long, corresponding to 45% of the total skull length

(premaxilla to squamosal) and 55% of the length from the craniomandibular joint to the premaxilla. Anterior to the nasoantorbital fenestra, the long axis of the rostrum is slightly deflected dorsally, as in other istiodactylids (Wang XL *et al.*, 2005; Andres & Ji, 2006; Lü *et al.*, 2008; Witton, 2012), as well as *Ikrandraco avatar* and anhanguerians (e.g. Kellner & Tomida, 2000; Wang XL *et al.*, 2014; 2015; Holgado *et al.*, 2019), but unlike boreopterids (Lü & Ji, 2005; Lü, 2010; Jiang *et al.*, 2014). There is a strong palatal keel extending from pre-narial part of the rostrum to the anterior third of the nasoantorbital fenestra. The craniomandibular joint levels with the anterior margin of the orbit, similarly to both specimens of *N. ignaciobrito* (see Fig. 3; both specimens, LPM 00023 and IVPP V-13288; see Wang XL *et al.*, 2005; Wang L *et al.*, 2006; Lü *et al.*, 2008), *Anhanguera* spp. (see Kellner & Tomida, 2000) and *Linlongopterus jennya* (see Rodrigues *et al.*, 2015), but unlike *Istiodactylus* spp., in which the joint is located anterior to the orbit (see Andres & Ji, 2006; Witton, 2012), and *Ikrandraco avatar* (see Wang XL *et al.*, 2015), *Hamipterus tianshanensis* (see Wang XL *et al.*, 2014) and *Ludodactylus sibbicki* (Frey *et al.*, 2003), in which the joint is located under the middle of the orbit. The orbit is piriform, with the narrowest part being ventral, and without a suborbital vacuity. This is similar to the condition seen in the referred specimen of *N. ignaciobrito* and unlike the rounded orbit of *Istiodactylus*, which has also a suborbital vacuity (see Andres & Ji, 2006; Lü *et al.*, 2008; Witton, 2012). The infratemporal fenestra is elliptical and much smaller than the orbit. The supratemporal fenestra is poorly preserved.

Premaxilla and Maxilla. The premaxilla is fused with the maxilla and the suture is obliterated, thus the boundary between the two bones cannot be traced. Consequently, the premaxillary and maxillary teeth count is unknown. There is no premaxillary crest, as in all other istiodactylids and *Haopterus gracilis*. The rostral tip of the premaxilla exhibits a slight dorsal deflection of palatal anterior tip (Fig. 4), as evidenced from the uplifted positions of the two anteriormost teeth. This is similar to what has already been reported for *Cimoliopterus* and anhanguerians (see Rodrigues & Kellner, 2013).

Nasal and Lacrimal. The nasal and lacrimal form the anterodorsal margin of the orbit and the posterodorsal margin of the nasoantorbital fenestra. The anterior end of the nasolacrimal coincides with the highest point of the nasoantorbital fenestra, as in both specimens of *N. ignaciobrito* and also *Ikrandraco avatar* (Wang XL *et al.*, 2005; Wang L *et al.*, 2006; Andres & Ji, 2006; Lü *et al.*, 2008; Wang XL *et al.*, 2015), but unlike

Istiodactylus latidens and most anhanguerians (e.g., *Anhanguera*, *Tropeognathus* and *Hamipterus*), except for *Ludodactylus sibbicki*, in which the highest point is posterior to the anterior end of the nasolacrimal (Campos & Kellner, 1985; Wellnhofer, 1987; Kellner & Tomida, 2000; Wang XL *et al.*, 2014; Frey *et al.*, 2003). A nasal descending process cannot be seen in BPMC-0204, possibly because it is still covered by rock. There are no traces of an orbital process of the lacrimal invading the orbit, but the posterior margin of the lacrimal is slightly damaged and a small process similar to the one seen in the holotype of *N. ignaciobrito* may have been present and got lost (see Wang XL *et al.*, 2005; Wang L *et al.*, 2006; Andres & Ji, 2006; Lü *et al.*, 2008). The lacrimal contacts the lacrimal process of the jugal at about the mid-height of the posterior margin of the nasoantorbital fenestra. The nasal is bordered dorsally by the premaxilla and by the prefrontal posteroventrally.

Jugal and Quadratojugal. The jugal is partially preserved, missing part of the maxillary process and the base of the lacrimal process. The jugal sends a postorbital process to contact the postorbital, separating the orbit and the infratemporal fenestra. Posteriorly, the jugal contacts the quadratojugal, which forms the anteroventral of the infratemporal fenestra.

Quadrate. Sutures at the lateral surface of the quadrate are unclear. It is unclear whether the articulation with the mandible is helical or not. The mid-region of the quadrate is lost. The dorsal portion of the quadrate contacts the quadratojugal anteriorly and the squamosal dorsally. The quadrate is inclined backwards at an angle of 150°, unlike both specimens of *N. ignaciobrito* (Fig. 3), in which it slopes at ~160° (160°4' in the holotype; Wang XL *et al.*, 2005; 163° in the referred specimen; Wang L *et al.*, 2006).

Prefrontal. The prefrontal is a small bone that forms the anterodorsal margin of the orbit, contacting the nasolacrimal. A suture between these two bones can be seen anteroventrally. The dorsoposterior tip of this bone contacts the frontal.

Frontal. The frontal seems to be fused with the premaxilla and parietal, with no visible sutures. It is unclear whether the posterodorsal extension of the frontal forms a blunt and low frontoparietal crest as in *Anhanguera* (see Kellner & Tomida, 2000) or not.

Parietal and Squamosal. The parietal and squamosal are poorly preserved, especially the latter. The squamosal outline cannot be properly identified. The parietal preserves a shallow depression in its surface that corresponds to the medial wall of the

supratemporal fenestra. The dorsal limits of this fossa level with the orbit and extend ventrally to the region of contact between the squamosal and the postorbital.

Postorbital. The postorbital is slender and does not exhibit a triangular shape, oppositely to the triangular condition that is seen in anhanguerids (e.g. Kellner & Tomida, 2000). Instead, it is like a three-pointed star (= concave equilateral hexagon) as in *Haopterus gracilis* (see Wang XL & Lü, 2001); that is, is essentially composed of three connected processes. The anterior region of the postorbital which is formed by the frontal and jugal processes, is arched and forms the posterior margin of the orbit. The squamosal process is shorter than the other processes and separates the supra and infratemporal fenestrae. There is no orbital process of the postorbital invading the orbit, unlike *Istiodactylus* (Andres & Ji, 2006; Witton, 2012).

Palatal elements. Due to crushing, some palatal elements are visible in dorsal view, though few details can be observed. The vomers form a long, slender bony bar that separates the choanae, as in *Hongshanopterus lacustris* (see Wang XL *et al.*, 2008a). Of the pterygoid, only the medial process can be seen. It is large and plate-like as that of *Hongshanopterus lacustris* (see Wang XL *et al.*, 2008a) and, to a lesser extent, the anhanguerids, in which the process is also broad but less medially expanded (e.g., Campos & Kellner, 1985; Frey *et al.*, 2003). This differs from the slender medial processes of the pterygoid of azhdarchoids (e.g., Pinheiro & Schultz, 2012; Kellner, 2013; Pêgas *et al.*, 2018) or those of the referred specimen of *N. ignaciobrito* (see Wang L *et al.*, 2006; Lü *et al.*, 2008) and *Ikrandraco avatar* (see Wang XL *et al.*, 2015).

Dentary. The dentaries are fused rostrally forming a symphysis that accounts for 36% of total mandibular length, which is 240 mm long. The dorsal surface of the symphysis presents a deep and broad median sulcus that extends anteriorly up to the level of the first pair of teeth (Fig. 5). The rostral tip of the dentary symphysis has an odontoid (pseudotooth), that is located between the first pair of teeth, is smaller than the adjacent tooth crowns and is dorsally directed. The odontoid lacks the neurovascular foramen piercing its surface in the referred specimen of *N. ignaciobrito* (see Wang L *et al.*, 2006). The odontoid has the same orientation as that of *Istiodactylus latidens* (see Witton, 2012; Martill, 2014) and *Lonchodraco giganteus* (see Rodrigues & Kellner, 2013, fig. 4E-F), unlike the sub-horizontal odontoids of both specimens of *N. ignaciobrito* (see Wang XL *et al.*, 2005; Wang L *et al.*, 2006) and *Ikrandraco avatar*

(see Wang XL *et al.*, 2015). The symphysis presents 11 tooth positions per side.

Surangular, Articular and Angular. The lateral surface of the posterior region of the right mandibular ramus is composed by the surangular, angular and articular. A suture separates the long anterior process of the surangular from the dentary dorsally. Posteriorly, the surangular becomes deeper and is sutured with the angular. The boundary between the angular and the dentary, however, cannot be distinguished, nor the boundary between the angular and the articular. The articular forms the posterior part of the mandible, including the articular surface for the quadrate and the retroarticular process, which is pointed, dorsoventrally low and distally tapering.

Hyoid. Only the right ceratobranchial I is exposed along the ventral margin of the right mandibular ramus (Fig. 1). A small portion of the posterior part is missing. The ceratobranchial I is a long rod-like bone extending along the whole length of the mandibular rami.

Dentition. There are 12 tooth positions along the right side of the upper jaw and 11 tooth positions along each side of the lower jaw, with an inferred total count of 46 tooth positions. The first two teeth of the upper jaw (which are presumably premaxillary teeth) are procumbent. The first tooth forms an angle of 130° with the main axis of the rostrum, while the second forms and angle of 123°. The third tooth is also slightly procumbent, forming an angle of 100° with the palatal plane. All subsequent teeth are perpendicular to the main axis of the rostrum. The first two dentary teeth are also slightly procumbent. The last two alveoli of the right maxilla are empty, and the last one is placed just anterior to the level of the rostral end of the nasoantorbital fenestra. All of the crowns are triangular in labiolingual view and labiolingually compressed, as typical of the Istiodactylidae. The base of the crowns is mesiodistally inflated. The lingual surface of the crown is concave with a well-marked basoapical depression and a low transversal convexity at the base, that forms a lingual cingulum. The labial surface is mostly convex with a shallow concavity in the middle of the basal part of the crown. No carinae are present along the mesial and distal cutting margins of the crowns. The same features occur in the crowns of the holotype of *N. ignaciobritoi*. The teeth exhibit a slight constriction between crown and root (Fig. 5). This feature is also shared with *N. ignaciobritoi* (see Wang XL *et al.*, 2005).

The first nine pairs of teeth of the upper jaw are large and subequal in size. Their

Comentario [GP4]: Replace by “an”

Con formato: Diseño: Claro (Amarillo)

apicobasal total length of their crowns is about 12 mm and the crown apicobasal length is about 7 mm, and the mesiodistal width of the socket is 4 mm. The smallest crown is 6 mm in apicobasal length and 2 mm in mesiodistal width.

N. luei sp. nov. also presents an interesting pattern of tooth replacement. Two teeth occur in the tenth alveolus of the right dentary: a large functional one and a not yet fully erupted replacement tooth. The replacement tooth was erupting anterolabially to the functional tooth, instead of posterolingually as reported in other pterodactyloid pterosaurs like *Anhanguera* (see Kellner & Tomida, 2000; Fastnacht, 2001) and '*Cearadactylus*' *ligabuei* (see Dalla Vecchia, 1993).

Cervical vertebrae. Seven cervical vertebrae are preserved, including the atlas-axis complex (although the atlas itself cannot be identified). They are articulated, except for the 7th vertebra, which is disarticulated but still contacting the 6th vertebra. The 3rd cervical, with 38.4 mm, is longer than the 4th through 7th cervicals which are of similar length. The neural spine is damaged in most cervicals, except that of the 4th vertebra, which is high and with a peculiar shape (its anterior margin is anteriorly inclined). The apex of the neural spine is gently rounded. The postzygapophyses are posterodorsally oriented. The centrum extends posterior to the postzygapophyses. In the 3rd to the 7th cervicals, a large pneumatic foramen can be seen on the posterior half of the centrum below the neural arch.

Phylogenetic analysis results

The phylogenetic analysis by parsimony produced 51 most parsimonious trees with a minimum length of 360 steps, with minimum consistency index of 0.642 and retention index of 0.862. As suspected by Witton (2012), *Hongshanopterus lacustris* and *Haopterus gracilis* were found to be closely related to the Istiodactylidae in the strict consensus tree (Fig. 6), but they fall outside them. The Istiodactylidae have been defined by Andres *et al.* (2014) as the least inclusive clade containing *Istiodactylus latidens* Seeley 1901 and *Nurhachius ignaciobrito* Wang *et al.* 2005. In the strict consensus tree (Fig. 6), the Istiodactylidae contain *Nurhachius*, *Liaoxipterus brachyognathus* and *Istiodactylus*. *N. ignaciobrito* and *N. luei* sp. nov. were recovered as sister taxa. *Istiodactylus latidens* and *I. sinensis* were also recovered as sister taxa and *Liaoxipterus brachyognathus* is the sister-taxon of *Istiodactylus*.

Comentario [GP5]: Should be included as a reference

Con formato: Resaltar

Istiodactylus has the following five synapomorphies: presence of a suborbital opening (character 11, state 1); prenasal portion of the rostrum less than 20% the skull length (character 25, state 0); presence of an orbital process in the jugal (character 56, state 1); and sharp carinae in the teeth (character 96, state 1). *Istiodactylus* shares with *Liaoxipterus brachyognathus* the following synapomorphies: subparallel lateral margins of the jaws (character 24, state 1); mandibular symphysis shorter than 33% of the mandible length (character 78, state 1); and rounded outline of the rostral end of the mandible (character 79, state 0).

The genus *Nurhachius* is characterized by the following five synapomorphies: piriform orbit (character 7, state 2); cranio-mandibular articulation under the anterior margin of orbit (character 58, state 2); dorsal deflection of the palatal anterior tip (character 71, state 1); teeth crowns with labial and lingual depressions (character 100, state 1); teeth with a mesiodistal constriction between crown and root (character 101, state 1). The presence of a dorsal deflection of the palatal anterior tip represents a homoplasy with *Anhangueria* + *Cimoliopterus*.

The Istiodactylidae share the following seven synapomorphies: ventral margin of the nasoantorbital fenestra longer than 40% of skull length (character 4, state 1); orbit reaching high in the skull, with the dorsal margin surpassing the dorsal margin of the nasoantorbital fenestra (character 10, state 1); skull height (exclusive of cranial crests) over 25% of the jaw length (character 23, state 1); lacrimal process of jugal inclined posteriorly (character 54, state 2); helical jaw-joint absent (character 59, state 0); palatal occlusal surface: strong palatal ridge confined to the posterior portion of the palate (character 71, state 3); teeth confined to about the anterior third of the jaws (character 86, state 3).

Hongshanopterus lacustris results to be the sister-taxon of the Istiodactylidae. The clade *Hongshanopterus lacustris* + Istiodactylidae presents two synapomorphies: tooth crowns strongly compressed laterally (character 95, state 2); and mesial crowns under twice as long as wide (character 97, state 0).

Haopterus gracilis results as the sister-taxon of *Hongshanopterus lacustris* + Istiodactylidae, sharing teeth that are confined to the anterior half of the jaws (character 86, state 2).

The Istiodactylidae, *Hongshanopterus lacustris* and *Haopterus gracilis* share with *Ikrandraco avatar* four synapomorphies: narrow lacrimal process of jugal (character 53, state 1); quadrate inclination relative to the ventral margin of the skull is 150° or more (character 57, state 3); tooth crowns slightly compressed laterally (character 95, state 2); lingual cingulum present at the base of tooth crown (character 102, state 1). The last two character states are also shared with *Lonchodraco giganteus*.

Bremer support values were 1 for the genus *Istiodactylus*, 3 for *Istiodactylus* + *Liaoxipterus*, 5 for the genus *Nurhachius*, 3 for Istiodactylidae, 3 for *Hongshanopterus* + Istiodactylidae, 1 for *Haopterus* + (*Hongshanopterus* + Istiodactylidae), 1 for *Ikrandraco* + *Lonchodraco* and 2 for the clade that joins all of these taxa.

Discussion

For over a century, *Istiodactylus latidens* was the only known istiodactylid (Witton, 2012). In the last 15 years, three new istiodactylids have been reported from the Jiufotang Formation of China: *N. ignaciobritoi* (described in 2005); *Istiodactylus sinensis* and *Longchengpterus zhaoi* (both described in 2006); and *Liaoxipterus brachyognathus* (originally described in 2005 as a purported ctenochasmatid and referred to the Istiodactylidae in 2008; Dong & Lü, 2005; Wang XL *et al.*, 2005; Wang L *et al.*, 2006; Andres & Ji, 2006; Lü *et al.*, 2008). However, the validity of some of them is debated. According to Lü *et al.* (2008), the holotypes of *Longchengpterus zhaoi* and *N. ignaciobritoi* are indistinguishable, sharing general skull shape and tooth morphology. Therefore, *Longchengpterus zhaoi* was considered a junior synonym of *N. ignaciobritoi* by Lü *et al.* (2008). Witton (2012) provisionally considered both of them as valid and distinct taxa, coding them separately in his phylogenetic analysis though without discussing it further. They were coded differently as for tooth count and spacing, with *Nurhachius* that was considered to have more numerous and more spaced teeth. However, both specimens exhibit a similar number of upper teeth (13 pairs in the holotype of *N. ignaciobritoi* and 12 pairs in *L. zhaoi*) and similar spacing (Wang XL *et al.*, 2005; Lü *et al.*, 2008). Furthermore, the holotypes and only specimens of *Longchengpterus zhaoi* and *N. ignaciobritoi* share further features that are unique within istiodactylids (Fig. 3): the high quadrate inclination ($\sim 160^{\circ}$), the reduced medial

process of the pterygoid, the upper dentition ending at the level of the nasoantorbital fenestra, and the sub-horizontal odontoid in the mandibular symphysis. Therefore, we follow Lü *et al.* (2008) in considering *Longchengpterus zhaoi* as a junior synonym of *N. ignaciobritoi*.

N. luei sp. nov. is an istiodactylid based on the following features: nasoantorbital fenestra longer than 40% of skull length, dentary symphysis less than 33% of mandible length; and triangular, labiolingually compressed tooth crowns. It shares with *N. ignaciobritoi* a piriform orbit, a dorsally deflected palatal anterior tip, a cranio-mandibular articulation positioned under the anterior margin of the orbit (Fig. 3), tooth crowns with labial and lingual depressions, and teeth with mesiodistal constrictions between crown and root (Fig. 5; 6). All these features are present in the holotype of *N. ignaciobritoi*, while in the referred specimen the dorsal deflection of the palatal anterior tip cannot be assessed as the rostrum tip is missing. Concerning the crown, with both a labial and a lingual depression, we note that this morphology is reflected in the shape of the alveoli in *N. ignaciobritoi*, with concave labial and lingual margins (Fig. 5). The dorsal deflection of the palatal anterior tip is present in the rostral tip of the skull of the holotype of *N. ignaciobritoi*, as evidenced from the dorsal position of the first pair of alveoli, (see Fig. 4C-D), although it was not mentioned in the original description (Wang XL *et al.*, 2005), as well as in the holotype of *N. luei*. This character was utilized in a data matrix for the first time by Rodrigues & Kellner (2013), and resulted to be a synapomorphy of Anhangueria + *Cimoliopterus*. According to our phylogenetic analysis (Fig. 7), this feature was independently acquired by *Nurhachius* and Anhangueria + *Cimoliopterus*.

N. luei sp. nov. differs from *N. ignaciobritoi* in the following features: the quadrate is inclined at 150° instead of the ~160° of *N. ignaciobritoi*; the medial process of the pterygoid is broad and plate-like, whereas it is reduced in *N. ignaciobritoi* (see Fig. 3); the dorsal median sulcus of the mandibular symphysis extends up to the first pair of teeth, whereas it reaches the sixth pair of teeth in *N. ignaciobritoi* (see Fig. 5); the odontoid (pseudotooth) lacks a lateral foramen, whereas a foramen is present in the referred specimen of *N. ignaciobritoi* (see Martill, 2014, fig. 7C-D); the odontoid has a blunt occlusal surface, whereas the surface is sharp in *N. ignaciobritoi* (Fig. 5); the odontoid is dorsally directed, whereas it is anterodorsally directed in *N. ignaciobritoi*.

(but see Martill, 2014, p. 57, right column, lines 21-23); and the ceratobranchial I of the hyoid apparatus accounts for 60% of the mandibular length, whereas it accounts for 35% of the mandibular length in *N. ignaciobritoi* (Fig. 5).

Concerning quadrate inclination, we do not regard this variation as intraspecific as the variation does not surpass 3° in the archaeopterodactyloid *Pterodactylus antiquus* (specimens BSPG AS I 739, BSPG 1929 I 18, BMMS 7; see Bennett, 2012; Vidovic & Martill, 2014), 3° in the anhanguerian *Hamipterus tianshanensis* (IVPP V18931.1, holotype, and IVPP V18935.1, paratype; see Wang et al., 2014), 5° in *Aerodactylus scolopaciceps* (BSPG 1883 XVI 1, BSPG 1937 I 18, BSPG AS V 29 a/b; see Vidovic & Martill, 2014), and is less than 6° in *Pteranodon longiceps* (specimens YPM 1177, USNM 13656, KUPV 2212, KUPV 27821) and also in *Pteranodon sternbergi* (specimens FHSM VP 339, YPM 1179, UALVP 24238; see Bennett, 1994; 2001). Quadrate inclination has indeed been regarded as diagnostic before for tapejarids (Kellner, 2013) and *Pteranodon* (Bennett, 1994). We regard this feature as potentially diagnostic at least until further specimens of each species of *Nurhachius* are found.

Both specimens of *N. ignaciobritoi* come from the upper part of the Jiufotang Formation (see Wu et al., 2018), while the holotype of *N. luei* comes from the lowermost part of the Jiufotang Formation. This stratigraphic distribution might be suggestive of an anagenetic link between the two species, similar to the case of *Pteranodon longiceps* (from the upper Smoky Hill Chalk) and *Pteranodon sternbergi* (from the lower Smoky Hill Chalk) according to Bennett (1994), but see taxonomic controversies (Kellner, 2010; 2017; Martin-Silverstone et al., 2017; Acorn et al., 2017). The same has been speculated as a possible explanation for the occurrence of multiple species of *Anhanguera* (Pinheiro & Rodrigues, 2017), *Thalassodromeus* and *Tupuxuara* (Pêgas et al., 2018) in the Romualdo Formation. However, these cases still lack stratigraphic control for support, and the time resolution of the Romualdo Formation is still in question (see Pinheiro & Rodrigues, 2017).

Wang XL et al. (2008) were unable to differentiate *Liaoxipterus brachyognathus* from *Longchengpterus zhaoi*, and suggested that *Longchengpterus zhaoi* could be a junior synonym of *Liaoxipterus brachyognathus*. However, as observed by Lü et al. (2008), the rostral end of the mandibular symphysis is rounded in *Liaoxipterus brachyognathus* (in dorsal view, as it is in *Istiodactylus latidens*), whereas it is triangular in

Comentario [GP6]: Not in the references list

Con formato: Resaltar

Comentario [GP7]: Wang XL?

Con formato: Resaltar

Con formato: Resaltar

Comentario [GP8]: Not in references

Con formato: Resaltar

Comentario [GP9]: Bennett, 2001 was not included in the references

Con formato: Resaltar

Comentario [GP10]: It is as Martin-Silverstone, 2017 in the references. Please check.

Comentario [GP11]: A or b?

“*Longchengpterus zhaoi*”. Furthermore, as coded by Andres *et al.* (2014), both jaws show an attenuated taper (in occlusal view) in *Longchengpterus zhaoi*, while the lateral margins of the lower jaw are sub-parallel in *Liaoxipterus brachyognathus* (Fig. 8A) as in *Istiodactylus latidens*. Additionally, the mandibular symphysis of *Liaoxipterus brachyognathus* is relatively stouter than that of *Longchengpterus zhaoi*: their length/width ratios are 0.43 and 0.27, respectively. It is worthy of being noticed that the mandibular symphysis of *Longchengpterus zhaoi* is incorrectly drawn in Martill (2014, fig. 7B), as resulting to be much shorter than it is. The actual configuration can be clearly assessed in the description by Lü *et al.* (2008) and in Fig. 5A. We thus follow Lü *et al.* (2008) and Witton (2012) in considering *Liaoxipterus brachyognathus* as distinct from *Longchengpterus zhaoi*, which we consider as a junior synonym of *N. ignaciobritoi*.

Lü *et al.* (2008) and Witton (2012) noticed that comparison between *Liaoxipterus brachyognathus* and *Istiodactylus sinensis* is very limited because the former is represented by a mandible exposed in occlusal view, while the latter is a partial skeleton including a mandible exposed in lateral view. However, *Liaoxipterus brachyognathus* differs from *Istiodactylus* in the lack of mesial carinae, according to Lü *et al.* (2008) and the dataset of Andres *et al.* (2014). We thus follow these authors in considering *Liaoxipterus brachyognathus* as a valid taxon.

According to our phylogenetic analysis, *Istiodactylus* is monophyletic, comprising *I. latidens* and *I. sinensis*. *Liaoxipterus brachyognathus* is the sister-taxon of *Istiodactylus*, and *Nurhachius* is the sister-taxon of *Liaoxipterus brachyognathus* + *Istiodactylus*, in agreement with the results of the phylogenetic hypothesis published by Longrich *et al.* (2018). In our analysis, *N. luei* results to be the sister-taxon of *N. ignaciobritoi*, supporting their congeneric status. The relationships within the Istiodactylidae obtained in our analysis are similar to those found by Andres *et al.* (2014), but *Longchengpterus zhaoi* is not the sister-taxon of *N. ignaciobritoi* in the cladogram of figure S2 of Andres *et al.* (2014).

Haopterus gracilis was first described by Wang XL & Lü (2001) and referred to the Pterodactylidae. However, it resulted to be close to *Istiodactylus latidens* in the 50% majority-rule tree by Lü *et al.* (2008) and formed a polytomy with *Nurhachius* and *Istiodactylus* in the strict consensus tree by Lü *et al.* (2009). *Hongshanopterus lacustris*

was described by Wang XL *et al.* (2008) and interpreted as a primitive istiodactylid. In the strict consensus tree by Witton (2012), the Istiodactylidae include *Nurhachius igniaciobritoi*, *Longchengpterus zhaoi*, *Istiodactylus latidens*, *Istiodactylus sinensis*, and *Liaoxipterus brachyognathus*. *Haopterus gracilis* and *Hongshanopterus lacustris* form a polytomy with *Pteranodon longiceps*, *Coloborhynchus spielbergi* and the Istiodactylidae (Witton, 2012).

Comentario [GP12]: A or b?

In the phylogenetic analysis of Andres *et al.* (2014; fig. S2), *Haopterus gracilis* results to be a basal eupterodactyloidean and *Hongshanopterus* a basal ornithocheiromorph. Recently, Holgado *et al.* (2019) have published a phylogenetic hypothesis in which *Hongshanopterus lacustris* would be the sister-group of the Istiodactylidae (although it is erroneously reported within this clade as the basal member in their fig. 5A), while *Haopterus gracilis* would be closer to anhanguerians than to istiodactylids (see Holgado *et al.*, 2019).

In our analysis, *Haopterus gracilis* and *Hongshanopterus lacustris* are closely related to the Istiodactylidae, as found by Lü *et al.* (2008) and Wang XL *et al.* (2008), respectively. *Hongshanopterus lacustris* results to be the sister-taxon of the Istiodactylidae, as in the analysis by Holgado *et al.* (2019). *Hongshanopterus lacustris* shares with the istiodactylids the presence of labiolingually compressed teeth with triangular crowns. *Haopterus gracilis* results to be the sister-taxon of *Hongshanopterus lacustris* + Istiodactylidae, a relationship that is supported by the possession of a dentition restricted to the anterior half of the jaws.

Haopterus, *Hongshanopterus* and istiodactylids share also the presence of a lingual cingulum in the tooth crown, a feature that occurs also in *Ikrandraco avatar*. A lingual cingulum can be seen in *Nurhachius luei* and *N. igniaciobritoi* (Fig. 6). The same feature has been previously reported for *Liaoxipterus brachyognathus* (see Lü *et al.*, 2008) and depicted for *Ikrandraco avatar* (see the second figure of Wang XL *et al.*, 2015). In *Haopterus gracilis*, the labiodistal view of the third right upper tooth presents a lingually oriented convexity that also suggests the presence of this feature (Fig. 8).

Ikrandraco avatar shares with istiodactylids also a narrow lacrimal process of the jugal and a quadrate inclined at 150° or over (the inclination of the quadrate is unknown in *Haopterus gracilis* and *Hongshanopterus lacustris*). *Ikrandraco avatar* and *Haopterus*

gracilis also exhibit a certain degree of labiolingual compression of the teeth, at least in the distal part of the dentition (Wang XL & Lü, 2001; Wang XL *et al.*, 2015), though not to the same degree seen in the istiodactylids and *Hongshanopterus*. The last two mandibular alveoli preserved in the holotype of *Lonchodraco giganteus* (the sister-taxon of *Ikrandraco avatar* in our analysis) are also labiolingually narrow (see Martill, 2011; Rodrigues & Kellner, 2013).

Furthermore, *Ikrandraco avatar* and *Lonchodraco giganteus* also share with istiodactylids the presence of an odontoid, which is anterodorsally oriented in the former and dorsally oriented in the latter (see Rodrigues & Kellner, 2013; Wang XL *et al.*, 2015).

A close relationship among *Ikrandraco avatar*, *Lonchodraco giganteus* and istiodactylids is found here for the first time. *Ikrandraco avatar* formed a polytomy with the Istiodactylidae, *Cimoliopterus* and the Anhangueria in the phylogenetic analysis by Wang XL *et al.* (2015) and *Lonchodraco giganteus* is outside the Lanceodontia in the phylogenetic analysis by Longrich *et al.* (2018).

Archaeoistiodactylus linglongtaensis is based on the sole holotype (JPM04-0008), including fragments of skull and one displaced maxillary tooth, a partial lower jaw in occlusal view with two teeth in place, an almost complete forelimb, a femur and a tibia. It is from the Middle Jurassic (Bathonian-Oxfordian) Tiaojishan Formation. *A. linglongtaensis* was described by Lü & Fucha (2010) who interpreted it as the "ancestor form of the known istiodactylid pterosaur [sic]" (Lü & Fucha, 2010, p. 113). Lü & Fucha (2010) observed that JPM04-0008 and the istiodactylids share teeth with triangular crowns and an odontoid (pseudotooth) on the mandibular symphysis. That odontoid was mistaken for a mid-line, unpaired tooth by Sullivan *et al.* (2014), but it had been explicitly described as a bony process by Lü & Fucha (2010, p. 116). Lü & Fucha (2010) also observed that the single maxillary tooth is recurved as in *Hongshanopterus lacustris*, and reported the presence of a warped deltopectoral crest in the humerus, which is a diagnostic feature of the Pteranodontoidea (Kellner, 2003). They noted that *A. linglongtaensis* differs from istiodactylids and all other pterodactyloids in the relatively short fourth metacarpal and in the presence of tibia, and second and third phalanges of the wing digit with subequal lengths. If actually a

pterodactyloid, it would represent one of the oldest occurrences of the Pterodactyloidea, being coeval or even older than the Callovian-Oxfordian basalmost pterodactyloid *Kryptodrakon progenitor* (see Andres *et al.*, 2014).

Its identification as a pterodactyloid was disputed by Martill & Etches (2013), who affirmed that JPM04-0008 is probably a badly preserved specimen of *Darwinopterus*, though they did not present any evidence to support this statement. According to Sullivan *et al.* (2014), the short fourth-metacarpal, the long humerus and short first wing phalanx are typical of non-pterodactyloid pterosaurs (see Kellner, 2003; Unwin, 2003; Andres *et al.*, 2010), thus JPM04-0008 is not a pterodactyloid. These features united to the presence of a confluent nasoantorbital fenestra in JPM04-0008, led Sullivan *et al.* (2014) to interpret *A. linglongtaensis* as a basal monofenestratan.

However, *A. linglongtaensis* has never been included in any phylogenetic analysis to test its basal monofenestratan affinity, thus it was included in the analysis performed in this paper. Our results (Fig. 6) confirm the interpretation by Sullivan *et al.* (2014). *Archaeoistiodactylus linglongtaensis* lacks the following pterodactyloid features: humerus length under 1.5 times metacarpal IV length; ulna under double the length of metacarpal IV; and femur subequal to or shorter than metacarpal IV. The humerus of JPM04-0008 is crushed and the original orientation of the deltopectoral crest cannot be assessed. Differently from pterodactyloids, the deltopectoral crest of JPM04-0008 is confined to the proximal region of the humerus (Wang XL *et al.*, 2009). *A. linglongtaensis* also lacks pneumatic foramina on the centra of the mid-cervical vertebrae, which is a diagnostic feature of the Dsungaripteroidea (the least inclusive clade containing *Nyctosaurus* and *Quetzalcoatlus*, which includes also the Istiodactylidae; Kellner, 2003; Andres *et al.*, 2014). Furthermore, *A. linglongtaensis* exhibits low neural spines, like wukongopterids (see Wang XL *et al.* 2009; 2010; Lü *et al.*, 2009; 2011; Cheng *et al.*, 2017) and unlike istiodactylids (see Wang L *et al.*, 2006; Lü *et al.*, 2008).

The dentition of *A. linglongtaensis* is indeed reminiscent of that of the Istiodactylidae due to the short triangular aspect of the crowns in labiolingual view. However, this feature is also present in the wukongopterids *Wukongopterus lüi*, *Darwinopterus robustodens*, *Darwinopterus linglongtaensis* and *Kunpengopterus sinensis*, though not in *Darwinopterus modularis* (see Wang XL *et al.* 2009; 2010; Lü *et al.*, 2009; 2011;

Comentario [GP13]: Not in the references

Con formato: Resaltar

Comentario [GP14]: This reference was not included in the list

Con formato: Resaltar

Cheng *et al.*, 2017). Furthermore, in *A. linglongtaensis* the alveoli are circular (Lü & Fucha, 2010), as in wukongopterids, not labiolingually compressed triangular teeth as in istiodactylids. The presence or absence of an odontoid in the lower jaw cannot be confidently assessed in *Wukongopterus* and *Darwinopterus*, but can be seen in a specimen referred to *Kunpengopterus sinensis* (see Cheng *et al.*, 2017), in convergence with the istiodactylids. Finally, *A. linglongtaensis* shares with *Darwinopterus* and *Kunpengopterus*, but not with *Wukongopterus*, the subequal in length second and third phalanges of the wing digit. Thus, *A. linglongtaensis* may be closely related to *Darwinopterus* or *Kunpengopterus*. In our analysis, *A. linglongtaensis* falls in a polytomy with *Darwinopterus linglongtaensis*, *D. robustodens* and *Kunpengopterus sinensis* (Fig. 6). However, we were unable to access the specimen first-hand and further scrutiny is desirable in order to confirm or deny this affinity.

Conclusions

The new specimen here described represents the second species for the genus *Nurhachius*, previously restricted to its type-species *N. ignaciobritoi*. A slight dorsal deflection of the palatal anterior tip revealed to be a synapomorphy of *N. ignaciobritoi* and *N. luei*. That feature was previously thought to be restricted to the Anhangueria and *Cimoliopterus*. Unlike other pterodactyloids, the holotype of *N. luei* sp. nov. shows an anterolabial tooth replacement. The position of *Hongshanopterus lacustris* and *Haopterus gracilis* as close taxa to the Istiodactylids is supported by the performed phylogenetic analysis. *Ikrandraco avatar* and *Lonchodraco giganteus* resulted to be sister taxa, and closer to istiodactylids than to other lanceodontians. The phylogenetic analysis supports the reinterpretation of *Archaeoistiodactylus linglongtaensis* as a non-pterodactyloid monofenestratan, probably a wukongopterid.

Acknowledgements

We thank Shu'an Ji and Xuefang Wei (IG-CAGS, Institute of Geology, Chinese Academy of Geological Sciences) for the help all along. Thanks to Cunyu Liu (BPMC, Beipiao Pterosaur Museum of China), Dongyu Hu (SNU, Shenyang Normal University), Xiaolin Wang & Shunxing Jiang (IVPP, Institute of Vertebrate

Paleontology and Paleoanthropology) for access to specimens under their care. RVP thanks Kamila Bandeira, Lucy Souza and Natan Brilhante (Museu Nacional/UFRJ) for technical help with image software. We thank Zoological Museum (SNM), Copenhagen University for hospitality during X. Y. Zhou's and R. V. Pêgas' stay in Copenhagen with access to important pterosaur specimens, and for M. E. C. Leal's status as guest researcher, and N. Bonde's work space as emeritus (and Senior Scientist, Fur Museum). Thanks to Fabio M. Dalla Vecchia, Felipe Pinheiro, Chris Bennett and an anonymous reviewer for their thoughtful and constructive critiques; and to editor Graciela Piñeiro for her kind attention.

References

- Acorn JH, Martin-Silverstone E, Glasier JRN, Mohr S, Currie PJ. 2017.** Response to Kellner (2017) 'Rebuttal of Martin-Silverstone, E., JRN Glasier, JH Acorn, S. Mohr, and PJ Currie, 2017'. *Vertebrate Anatomy Morphology Palaeontology* **3**: 90-92.
- Andres B, Ji Q. 2006.** A new species of *Istiodactylus* (Pterosauria, Pterodactyloidea) from the Lower Cretaceous of Liaoning, China. *Journal of Vertebrate Paleontology* **26(1)**: 70-78.
- Andres B, Clark JM, Xu X. 2010.** A new rhamphorhynchid pterosaur from the Upper Jurassic of Xinjiang, China, and the phylogenetic relationships of basal pterosaurs. *Journal of Vertebrate Paleontology* **30(1)**: 163-187.
- Andres B, Clark JM, Xu X. 2014.** The earliest pterodactyloid and the origin of the group. *Current Biology* **24(9)**: 1011-1016.
- Bennett SC. 1994.** Taxonomy and systematics of the Late Cretaceous pterosaur *Pteranodon* (Pterosauria, Pterodactyloidea). *Occasional papers of the Natural History Museum/The University of Kansas* (169):1-70.
- Campos DA, Kellner AWA. 1985.** Panorama of the flying reptiles study in Brazil and South America. *Anais da Academia Brasileira de Ciências* **57**: 453-466.
- Chang MM, Chen PJ, Wang YQ, Wang Y, Miao DS. eds. 2003.** *The Jehol Biota*. Shanghai Sci. & Tech. Publ. 208 pp.
- Chang SC, Zhang HC, Renne PR, Fang Y. 2009.** High-precision $^{40}\text{Ar}/^{39}\text{Ar}$ age for the Jehol Biota. *Palaeogeography, Palaeoclimatology, Palaeoecology* **280**: 94-104.
- Cheng X, Jiang SX, Wang XL, Kellner AWA. 2017.** Premaxillary crest variation within the Wukongopteridae (Reptilia, Pterosauria) and comments on cranial

structures in pterosaurs. Anais da Academia Brasileira de Ciências **89(1)**: 119-130.

Dalla Vecchia FM. 1993. *Cearadactylus? ligabuei*, nov. sp., a new Early Cretaceous (Aptian) pterosaur from Chapada do Araripe (Northeastern Brazil). Bolletino della Societa Paleontologica Italiana **32**: 401–409.

Con formato: Inglés (Estados Unidos)

Dong ZM, Lü JC. 2005. A new ctenochasmatid pterosaur from the Early Cretaceous of Liaoning Province. Acta Geologica Sinica **79(2)**: 164-167.

Dong ZM, Sun YW, Wu SY. 2003. On a new pterosaur from the Lower Cretaceous of Chaoyang Basin, Western Liaoning, China. Global Geology **22(1)**: 1-8.

Con formato: Inglés (Estados Unidos)

Eberth DA, Russell DA, Braman DR, Deino AL. 1993. The age of the dinosaur bearing sediments at Tebch, Inner Mongolia, People's Republic of China. Canadian Journal of Earth Science **30**: 2101–2106.

Fastnacht M. 2001. First record of *Coloborhynchus* (Pterosauria) from the Santana Formation (Lower Cretaceous) of the Chapada do Araripe, Brazil. Paläontologische Zeitschrift **75(1)**: 23-26.

Comentario [GP15]: Not cited

Frey E, Martill DM, Buchy MC. 2003. A new crested ornithocheirid from the Lower Cretaceous of northeastern Brazil and the unusual death of an unusual pterosaur. Geological Society, London, Special Publications **217(1)**: 55–63.

Goloboff PA, Farris JS, Nixon KC. 2008. TNT, a free program for phylogenetic analysis. Cladistics **24(5)**: 774–786.

Holgado B, Pêgas RV, Canudo JI, Fortuny J, Rodrigues T, Company J, Kellner AWA. 2019. On a new crested pterodactyloid from the Early Cretaceous of the Iberian Peninsula and the radiation of the clade Anhangueria. Scientific reports **9(1)**: 4940.

Howse SCB, Milner AR, Martill DM. 2001. "Pterosaurs". In Martill D M, Naish D. *Dinosaurs of the Isle of Wight*. The Palaeontological Association. Pp:324–335.

Jiang SX, Wang XL, Meng X, Cheng X. 2014. A new boreopterid pterosaur from the Lower Cretaceous of western Liaoning, China, with a reassessment of the phylogenetic relationships of the Boreopteridae. Journal of Paleontology **88(4)**: 823-828.

Jiang SX, Cheng X, Ma YX, Wang XL. 2016. A new archaeopterodactyloid pterosaur from the Jiufotang Formation of western Liaoning, China, with a comparison of sterna in Pterodactylomorpha. Journal of Vertebrate Paleontology **36 (6)**: e1212058.

Kaup SS. 1834. Versuch einer Eintheilung der Säugethiere in 6 Stämme und der Amphibien in 6 Ordnungen. Isis von Oken, 1834, cols. 311–315.

Kellner AWA. 2003. Pterosaur phylogeny and comments on the evolutionary history of the group. Geological Society, London, Special Publications **217(1)**: 105–137.

- | **Kellner AWA. 2010.** Comments on the Pteranodontidae (Pterosauria, Pterodactyloidea) with the description of two new species. *Anais da Academia Brasileira de Ciências* **82(4)**: 1063-1084.
- | **Kellner AWA. 2013.** A new unusual tapejarid (Pterosauria, Pterodactyloidea) from the Early Cretaceous Romualdo Formation, Araripe Basin, Brazil. *Earth and Environmental Science Transactions of the Royal Society of Edinburgh* **103(3-4)**: 409-421.
- | **Kellner AWA. 2017.** Rebuttal of Martin-Silverstone *et al.* 2017, 'Reassessment of *Dawndraco kanzai* Kellner 2010 and reassignment of the type specimen to *Pteranodon sternbergi* Harksen, 1966'. *Vertebrate Anatomy Morphology Palaeontology* **3**: 81-89.
- | **Kellner AWA, Tomida Y. 2000.** Description of a new species of Anhangueridae (Pterodactyloidea) with comments on the pterosaur fauna from the Santana Formation (Aptian–Albian), northeastern Brazil. *National Science Museum Monographs* **17**: ix-137.
- | **Li JJ, Lü JC, Zhang BK. 2003.** A new Lower Cretaceous sinopterid pterosaur from the western Liaoning, China. *Acta Palaeontologica Sinica* **42(3)**: 442-447.
- | **Longrich NR, Martill DM, Andres B. 2018.** Late Maastrichtian pterosaurs from North Africa and mass extinction of Pterosauria at the Cretaceous-Paleogene boundary. *PLoS Biology* **16(3)**: e2001663.
- | **Lü JC. 2010.** A new boreopterid pterodactyloid pterosaur from the Early Cretaceous Yixian Formation of Liaoning Province, northeastern China. *Acta Geologica Sinica* **84(2)**: 241-246.
- | **Lü JC, Fucha XH. 2010.** A new pterosaur (Pterosauria) from Middle Jurassic Tiaojishan Formation of western Liaoning, China. *Global Geology* **13 (3/4)**: 113–118.
- | **Lü JC, Ji Q. 2005.** New azhdarchid pterosaur from the Early Cretaceous of western Liaoning. *Acta Geologica Sinica* **79 (3)**: 301–307.
- | **Lü JC, Yuan CX. 2005.** New tapejarid pterosaur from western Liaoning, china. *Acta Geologica Sinica* **79(4)**: 453-458.
- | **Lü JC, Jin XS, Unwin DM, Zhao LJ, Azuma Y, Ji Q. 2006.** A new species of *Huaxiapterus* (Pterosauria: Pterodactyloidea) from the Lower Cretaceous of western Liaoning, China with comments on the systematics of tapejarid pterosaurs. *Acta Geologica Sinica* **80(3)**: 315–326.
- | **Lü JC, Gao YB, Xing LD, Li ZX, Sun ZY. 2007.** A new species of *Huaxiapterus* from the Early Cretaceous of western Liaoning,China. *Acta Geologica Sinica* **81 (5)**: 683–687.
- | **Lü JC, Xu L, Ji Q. 2008.** Restudy of *Liaoxiapterus* (Istiodactylidae:Pterosauria), with

comments on the Chinese istiodactylid pterosaurs. *Zitteliana* **B28**: 229-241.

- | **Lü JC, Unwin DM, Jin XS, Liu YQ, Ji Q. 2009.** Evidence for modular evolution in a long-tailed pterosaur with a pterodactyloid skull. *Proceedings of the Royal Society B: Biological Sciences* **277(1680)**: 383-389.
- | **Lü JC, Xu L, Chang HL, Zhang XL. 2011.** A new darwinopterid pterosaur from the Middle Jurassic of western Liaoning, northeastern China and its ecological implications. *Acta Geologica Sinica* **85(3)**: 507-514.
- | **Lü JC, Jin XS, Gao CL, Du TM, Ding M, Sheng YM, Wei XF. 2013.** *Dragons of the Skies (Recent advances on the study of pterosaurs from China)*. Zhejiang Science & Technology Press . 127 Pp.
- | **Lü J C, Liu C Y, Xu L, Pan L J, Shen C Z. 2016a.** A new pterodactyloid pterosaur from the Early Cretaceous of the Western Part of Liaoning Province, Northeastern China. *Acta Geologica Sinica*, **90 (3)**: 777–782.
- | **Lü J C, Teng F F, Sun D Y, Shen C Z, Li G Q, Gao X, Liu H F. 2016b.** The toothless pterosaurs from China. *Acta Geologica Sinica*, **90 (9)**: 2513–2525.
- | **Martill DM. 2011.** A new pterodactyloid pterosaur from the Santana Formation (Cretaceous) of Brazil. *Cretaceous Research* **32 (2)**: 236-243.
- | **Martill DM. 2014.** A functional odontoid in the dentary of the Early Cretaceous pterosaur *Istiodactylus latidens*: Implications for feeding. *Cretaceous Research* **47**: 56-65.
- | **Martill DM, Etches S. 2013.** A new monofenestratan pterosaur from the Kimmeridge Clay Formation (Kimmeridgian, Upper Jurassic) of Dorset, England. *Acta Palaeontologica Polonica* **58(2)**: 285-295.
- | **Martin-Silverstone E. 2017.** Redescription of *Dawndraco kanzai* Kellner, 2010 and reassignment of the type specimen to *Pteranodon sternbergi* Harksen, 1966. *Vertebrate Anatomy Morphology Palaeontology* **3**: 47-59.
- | **Meng J, Wang YQ, Li CK. 2011.** Transitional mammalian middle ear from a new Cretaceous Jehol eutriconodont. *Nature* **472(7342)**: 181–185.
- | **Pêgas RV, Costa FR, Kellner AWA. 2018.** New information on the osteology and a taxonomic revision of the genus *Thalassodromeus* (Pterodactyloidea, Tapejaridae, Thalassodrominae). *Journal of Vertebrate Paleontology* **38(2)**: e1443273.
- | **Pinheiro FL, Rodrigues T. 2017.** *Anhanguera* taxonomy revisited: is our understanding of Santana Group pterosaur diversity biased by poor biological and stratigraphic control?. *PeerJ* **5**: e3285.
- | **Pinheiro FL, Schultz CL. 2012.** An unusual pterosaur specimen (Pterodactyloidea,? Azhdarchoidea) from the Early Cretaceous Romualdo Formation of Brazil, and the evolution of the pterodactyloid palate. *PLOS ONE* **7(11)**: e50088.

- Plieninger F. 1901.** Beiträge zur Kenntnis der Flugsaurier. *Palaeontographica* **48**: 65–90.
- Rodrigues T, Kellner AWA. 2013.** Taxonomic review of the *Ornithocheirus* complex (Pterosauria) from the Cretaceous of England. *ZooKeys* **308**: 1–112.
- Rodrigues T, Jiang SX, Cheng X, Wang X, Kellner AWA. 2015.** A new toothed pteranodontoid (Pterosauria: Pterodactyloidea) from the Jiufotang Formation (Lower Cretaceous, Aptian) of China and comments on *Liaoningopterus gui* Wang and Zhou, 2003. *Historical Biology* **27**(6): 782–795.
- Sullivan C, Wang Y, Hone DW, Wang YQ, Xu X, Zhang FC. 2014.** The vertebrates of the Jurassic Daohugou Biota of northeastern China. *Journal of Vertebrate Paleontology* **34**(2): 243–280.
- Unwin DM. 2003.** On the phylogeny and evolutionary history of pterosaurs. Geological Society, London, Special Publications **217**(1): 139–190.
- Wang L, Li L, Duan Y, Cheng SL. 2006.** A new istiodactylid pterosaur from western Liaoning, China. *Geological Bulletin of China* **25**(6): 737–740.
- Wang M, Zhou ZH. 2019.** A new enantiornithine (Aves: Ornithothoraces) with completely fused premaxillae from the Early Cretaceous of China. *Journal of Systematic Palaeontology* **Online edition**: 1–14.
- Wang X. 2018.** Background for the Plant Fossils. Pp. 47–59 in: *The Dawn Angiosperms*. Springer Geology. Springer, Cham. 334 Pp.
- Wang XL, Lü JC. 2001.** Discovery of a pterodactylid pterosaur from the Yixian Formation of western Liaoning, China. *Chinese Science Bulletin* **46**(13): 1112–1117.
- Wang XL, Zhou ZH. 2003a.** A new pterosaur (Pterodactyloidea, Tapejaridae) from the Early Cretaceous Jiufotang Formation of Western Liaoning, China and its implications for biostratigraphy. *Chinese Science Bulletin* **48**(1): 16–23.
- Wang XL, Zhou ZH. 2003b.** Two new pterodactyloid pterosaurs from the Early Cretaceous Jiufotang Formation of Western Liaoning, China. *Vertebrata Palasiatica* **41** (1): 34–41.
- Wang XL, Zhou ZH. 2003c.** Pterosaurs. – In: M. M. Chang, *et al.* (Eds), *The Jehol Biota*; Shanghai (Shanghai Scientific and Technical Publishers): 99–108.
- Wang XL, Kellner AWA, Zhou ZH, Campos DA. 2005.** Pterosaur diversity and faunal turnover in Cretaceous terrestrial ecosystems in China. *Nature* **437**: 875–879.
- Wang XL, Campos DA, Zhou ZH, Kellner AWA. 2008a.** A primitive istiodactylid pterosaur (Pterodactyloidea) from the Jiufotang Formation (Early Cretaceous), northeast China. *Zootaxa* **1813**: 1–18.

Comentario [GP16]: Not cited

- | **Wang XL, Kellner AWA, Zhou ZH, Campos DA. 2008b.** Discovery of a rare arboreal forest-dwelling flying reptile (Pterosauria: Pterodactyloidea) from China. *Proceedings of the National Academy of Sciences* **105(6)**: 1983-1987
- | **Wang XL, Kellner AWA, Jiang SX, Cheng X, Meng X, Rodrigues T. 2010.** New long-tailed pterosaurs (Wukongopteridae) from western Liaoning, China. *Anais da Academia Brasileira de Ciências* **82(4)**: 1045-1062.
- | **Wang XL, Kellner AWA, Jiang SX, Cheng X. 2012.** New toothed flying reptile from Asia; close similarities between early Cretaceous pterosaurs faunas from China and Brazil. *Naturwissenschaften* **99(4)**: 249-257.
- | **Wang XL, Kellner AWA, Jiang SX, Wang Q, Ma YX, Paidoula Y, Cheng X, Rodrigues T, Meng X, Zhang JL, Li N, Zhou ZH. 2014.** Sexually dimorphic tridimensionally preserved pterosaurs and their eggs from China. *Current Biology* **24(12)**: 1323-1330.
- | **Wang XL, Rodrigues T, Jiang SX, Cheng X, Kellner AWA. 2015.** An Early Cretaceous pterosaur with an unusual mandibular crest from China and a potential novel feeding strategy. *Scientific Reports* **4**: 6329.
- | **Wellnhofer P. 1987.** New crested pterosaurs from the Lower Cretaceous of Brazil. *Mitteilungen der Bayerischen Staatssammlung für Paläontologie und Historische Geologie* **27**: 17–186.
- | **Witton MP. 2012.** New insights into the skull of *Istiodactylus latidens* (Ornithocheiroidea, Pterodactyloidea). *PloS one* **7(3)**: e33170.
- | **Wu ZJ, Gao FL, Pan YQ, Wang X. 2018.** Division and comparison of Jiufotang Formation of western Liaoning area and its rare fossil bed. *Geoscience* **32(4)**: 758-765.
- | **Yao X, Liao CC, Sullivan C, Xu X. 2019.** A new transitional therizinosaurian theropod from the Early Cretaceous Jehol Biota of China. *Scientific Reports* **9**: 5026
- | **Zhou ZH, Barrett PM, Hilton J. 2003.** An exceptionally preserved Lower Cretaceous ecosystem. *Nature* **421(6925)**: 807–814.

Figures

Figure 1. *Nurhachius luei* sp. nov., BPMC-0204, holotype, photograph and line drawing. The scale bar in the line drawing equals 50 mm. Abbreviations: alv, alveolus; an, angular; art, articular; ax, axis; cel, ceratobranchial I; ch, choana; cv, cervical vertebra; d, dentary; f, frontal; j, jugal; la, lacrimal; m, maxilla; n, nasal; naof, nasoantorbital fenestra; odp, odontoid process; or, orbit; pa, parietal; pf, prefrontal;

pmax, premaxilla; po, postorbital; prid, palatal ridge; pty, pterygoid; q, quadrate; vo, vomer. Isolated numbers indicate tooth positions. Note: the visible region of the pterygoid corresponds to the medial process of the bone. Photo by Xuanyu Zhou. Drawing by Maria Eduarda Leal.

Figure 2. Location of the site where BPMC-0204 was found.

Figure 3. *Nurhachius ignaciobritoi* specimens, photographs and line drawings. (A) LPM 00023, referred specimen (holotype of *Longchengpterus zhaoi*), skull and mandible in right lateral view. (B) IVPP V-13288, holotype, skull (mirrored) and mandible in right lateral view. Scale bars equal 50 mm. Abbreviations: art, articular; ch, choana; d, dentary; f, frontal; ios, interorbital septum; j, jugal; ls, laterosphenoid; m, maxilla; n, nasal; op, opisthotic; or, orbit; pa, parietal; pf, prefrontal; pm, premaxilla; prid, palatal ridge; pty, pterygoid; q, quadrate; san, surangular; sq, squamosal. Photographs by Xuanyu Zhou. Drawings by Rodrigo V. Pêgas.

Figure 4. Close view of the rostral tip of *Nurhachius* species in right lateral view. (A) *Nurhachius luei* sp. nov., holotype and (C) *Nurhachius ignaciobritoi*, IVPP V-13288, holotype, mirrored. (B) and (D), respective schematic drawings of (A) and (C), showing the slight dorsal deflection of the palatal anterior tip (notice the positions of the first and second alveoli in both specimens). Numbers indicate tooth positions. Scale bars equal 20 mm. Photos by Xuanyu Zhou. Drawings by Rodrigo V. Pêgas.

Figure 5. Close view of the dentary symphysis of *Nurhachius* species. (A) *Nurhachius luei* sp. nov. holotype in dorsolateral view, and (B) line drawing. (C) *Nurhachius ignaciobritoi*, LPM 00023, referred specimen, occlusal view, and (D), line drawing. Abbreviations: mg, median groove; odp, odontoid process. Numbers indicate tooth positions. Red arrows indicate the mesiodistal constriction between crown and root. Photos by Xuanyu Zhou. Drawings by Maria Eduarda Leal and Rodrigo V. Pêgas.

Figure 6. Close view of the dentition in *Nurhachius* species. (A) *Nurhachius ignaciobritoi*, LPM 00023, referred specimen, isolated tooth in lingual view, and (B) line drawing. (C) *Nurhachius luei* sp. nov. holotype, ninth left mandibular tooth in lingual view, and (D) line drawing. Red arrows indicate the mesiodistal constrictions

between crown and root. Blue arrows indicate the horizontal elevation at the base of the crown (cingulum). Photos by Xuanyu Zhou. Drawings by Rodrigo V. Pêgas.

Figure 7. *Nurhachius luei* sp. nov. phylogenetic relationships. Strict consensus tree of 51 most parsimonious trees. Tree length is 358, consistency index 0.644 and retention index 0.867. The red rectangle indicates the Istiodactylidae and its two closest taxa.

Figure 8. Other istiodactylids and close taxa. (A) *Liaoxipterus brachyognathus*, CAR-0018, holotype, lower jaw in dorsal view. (B) *Haopterus gracilis*, IVPP V11726, holotype, skull in right lateral view. (C) *Hongshanopterus lacustris*, IVPP V14582, holotype, skull in ventral view. All scale bars equal 50 mm. (A) and (C) by Xuanyu Zhou; (B) by Shunxing Jiang (courtesy of IVPP).