

# Combined effect of the entomopathogenic fungus *Metarhizium robertsii* and avermectins on the survival and immune response of *Aedes aegypti* larvae

Yuriy A Noskov<sup>Corresp., 1, 2</sup>, Olga V Polenogova<sup>1</sup>, Olga N Yaroslavtseva<sup>1</sup>, Olga E Belevich<sup>1</sup>, Yuriy Y Yurchenko<sup>1</sup>, Ekaterina A Chertkova<sup>1</sup>, Natalya A Kryukova<sup>1</sup>, Vadim Y Kryukov<sup>1</sup>, Viktor V Glupov<sup>1</sup>

<sup>1</sup> Institute of Systematics and Ecology of Animals, Novosibirsk, Russian Federation

<sup>2</sup> Tomsk State University, Tomsk, Russian Federation

Corresponding Author: Yuriy A Noskov  
Email address: yunoskov@gmail.com

Combination of insect pathogenic fungi and microbial metabolites is a prospective method for mosquito control. The effect of the entomopathogenic fungus *Metarhizium robertsii* and avermectins on the survival and physiological parameters of *Aedes aegypti* larvae (dopamine concentration (DA), glutathione S-transferase (GST), nonspecific esterases (EST), acid proteases, lysozyme-like, phenoloxidase (PO) activities) was studied. It is shown that the combination of these agents leads to a synergistic effect on mosquito mortality. Systemic colonization of the larvae of *Ae. aegypti* by hyphal bodies following water inoculation with conidia is shown for the first time. The larvae affected by fungi are characterized by a decrease in PO and dopamine levels. In the initial stages of toxicosis and/or fungal infection (12 h posttreatment), increases in the activity of insect detoxifying enzymes (GST and EST) and acid proteases are observed after monotreatments, and these increases are suppressed after combined treatment with the fungus and avermectins. Lysozyme-like activity is also most strongly suppressed under combined treatment with the fungus and avermectins in the early stages posttreatment (12 h). Forty-eight hours posttreatment, we observe increases in GST, EST, acid proteases, and lysozyme-like activity under the influence of the fungus and/or avermectins. The larvae affected by avermectins accumulate lower levels of conidia than avermectin-free larvae. On the other hand, a burst of bacterial CFUs is observed under treatment with both the fungus and avermectins. We assume that disturbance of the responses of the immune and detoxifying systems under the combined treatment and the development of opportunistic bacteria may be among the causes of the synergistic effect.

**Combined effect of the entomopathogenic fungus *Metarhizium robertsii* and avermectins on the survival and immune response of *Aedes aegypti* larvae**

Yuriy A. Noskov<sup>1,2</sup>, Olga V. Polenogova<sup>1</sup>, Olga N. Yaroslavtseva<sup>1</sup>, Olga E. Belevich<sup>1</sup>, Yuriy A. Yurchenko<sup>1</sup>, Ekaterina A. Chertkova<sup>1</sup>, Natalya A. Kryukova<sup>1</sup>, Vadim Y. Kryukov<sup>1</sup>, Viktor V. Glupov<sup>1</sup>

<sup>1</sup> Institute of Systematics and Ecology of Animals SB RAS, Novosibirsk, Russia

<sup>2</sup> National Research Tomsk State University, Tomsk, Russia

Corresponding author:

Yuriy Noskov <sup>1,2</sup>

Frunze str., 11, Novosibirsk, 630091, Russia

Email address: yunoskov@gmail.com

# Abstract

Combination of insect pathogenic fungi and microbial metabolites is a prospective method for mosquito control. The effect of the entomopathogenic fungus *Metarhizium robertsii* and avermectins on the survival and physiological parameters of *Aedes aegypti* larvae (dopamine concentration (DA), glutathione S-transferase (GST), nonspecific esterases (EST), acid proteases, lysozyme-like, phenoloxidase (PO) activities) was studied. It is shown that the combination of these agents leads to a synergistic effect on mosquito mortality. Systemic colonization of the larvae of *Ae. aegypti* by hyphal bodies following water inoculation with conidia is shown for the first time. The larvae affected by fungi are characterized by a decrease in PO and dopamine levels. In the initial stages of toxicosis and/or fungal infection (12 h posttreatment), increases in the activity of insect detoxifying enzymes (GST and EST) and acid proteases are observed after monotreatments, and these increases are suppressed after combined treatment with the fungus and avermectins. Lysozyme-like activity is also most strongly suppressed under combined treatment with the fungus and avermectins in the early stages posttreatment (12 h). Forty-eight hours posttreatment, we observe increases in GST, EST, acid proteases, and lysozyme-like activity under the influence of the fungus and/or avermectins. The larvae affected by avermectins accumulate lower levels of conidia than avermectin-free larvae. On the other hand, a burst of bacterial CFUs is observed under treatment with both the fungus and avermectins. We assume that disturbance of the responses of the immune and detoxifying systems under the combined treatment and the development of opportunistic bacteria may be among the causes of the synergistic effect.

# Introduction

Mosquitoes are obligate intermediate hosts for a variety of pathogens that cause human mortality and morbidity worldwide. *Aedes aegypti* is considered to be an important vector of human diseases such as dengue and yellow fever, chikungunya, and Zika infections (Tolle, 2009; Bhatt et al., 2013), and its control is therefore an objective to prevent the transmission of these diseases. Chemical insecticides are still the most important element in mosquito control programs, despite direct and indirect toxic effects on nontarget organisms, including humans. In addition, chemicals induce resistance in a number of vector species (Hemingway & Ranson, 2000; Vontas, Ranson & Alphey, 2010). Therefore, there is a need for alternative nonchemical vector control approaches. Classical biological control based on using various microorganisms such as entomopathogenic fungi and bacteria is a frequent tool for addressing this issue. Among the biological agents employed for mosquito larvae control, bacteria from the genus *Bacillus* are the most widely used. In addition, products of the entomopathogenic fungi (EPF) *Metarhizium* and *Beauveria* are actively being developed for use against mosquito adults and larvae (Butt et al., 2013; Greenfield et al., 2015; Ortiz-Urquiza et al., 2015). It should be noted that mosquitoes and other insects can develop resistance to the *Bacillus thuringiensis* biological larvicide (Tilquin et al., 2008; Paris et al., 2011; Boyer et al., 2012). However, the resistance of insects to entomopathogenic fungi develops very slowly (Dubovskiy et al., 2013). Various species of mosquito larvae present different susceptibilities to *Metarhizium*, among which *Ae. aegypti* is the least susceptible (Greenfield et al., 2015; Garrido-Jurado et al., 2016). Thus, a concentration of conidia that is effective for *Ae. aegypti* control would affect a range of nontarget aquatic invertebrates. Recent studies have found that some nontarget aquatic species are more sensitive to fungal metabolites (Garrido-Jurado et al., 2016) and conidia (Belevich et al., 2017) than target mosquito species. To reduce toxic effects on the aquatic environment and increase

efficacy against mosquitoes, entomopathogenic fungi may be combined with other biocontrol agents or low doses of natural insecticides. For example, combined treatment with *Metarhizium* and mosquito predator species (*Toxorhynchites*) has shown additive or synergistic effects on the mortality of *Ae. aegypti* (Alkhaibari et al., 2018). However, few studies have been carried out to determine the effect of combined treatment with EPF and other insecticides or plant or microbial metabolites as a potential tool for improving mosquito larvae control. Synergistic effects between EPF and some chemical insecticides (temephos, spinosad) (Shoukat et al., 2018; Vivekanandhan et al., 2018) or biological agents (*Azadirachta indica*) (Badiane et al., 2017) on the mortality of mosquito larvae have been found. However, the physiological and biochemical aspects of this synergism were not considered.

One type of promising insecticide that can be effectively used for mosquito vector control is the avermectins. Avermectins are a class of macrocyclic lactones isolated from the soil actinomycete *Streptomyces avermitilis* (Drinyaev et al., 1999) and include several commercial derivatives (ivermectin, abamectin, doramectin and eprinomectin) with the same mode of action – activation of glutamate-gated chloride channels, followed by uncontrolled influx of chloride ions into the cells, which leads to paralysis and death of the organism (Campbell et al., 1983). At the same time, avermectins are relatively safe for humans (Crump, Omura, 2011). Previous studies have shown that avermectins are efficient for the control of *Culex quinquefasciatus* (Freitas et al., 1996; Alves et al., 2004), *Anopheles albimanus* and *An. stephensi* (Dreyer, Morin & Vaughan, 2018), and *An. gambiae* (Alout et al., 2014; Chaccour et al., 2017). However, most of these studies have been carried out with adult mosquitoes feeding on blood containing ivermectin. Avermectins exhibit a relatively short half-life period, which limits their ability to kill mosquitoes but may be compensated by the application of multiple treatments or use of higher concentrations. However, these approaches may contribute to the development of mosquito larvae resistance to avermectins (Su et al., 2017). We hypothesize that the interaction of EPF with avermectins can have a stable insecticidal effect at relatively low concentrations and is a promising combination for safe and effective mosquito control.

It is important that EPF such as *Metarhizium* are adapted to terrestrial hosts and that in mosquito larvae, the fungi do not adhere to the cuticle surface and do not germinate through integuments into the body cavity. Conidia ingested by mosquito larvae do not penetrate the gut wall (Butt et al., 2013). Thus, a "classic" host-pathogen interaction does not occur, and larval mortality is associated with stress induced by spore-bound proteases on the surface of ingested conidia (Butt et al., 2013). These authors suggest that fungal proteases cause an increase in the activity of caspases in mosquitoes, which leads to apoptosis, autolysis of tissues and death of the larvae. The activation of detoxifying enzymes and antimicrobial peptides (AMPs) occurs in larvae infected with the fungus but is not sufficient to protect the larvae from death. As a rule, mosquito larvae die showing symptoms of bacterial decomposition after treatment with *Metarhizium* and *Beauveria* (Scholte et al., 2004). Therefore, this pathogenesis can be considered mixed (both bacterial and fungal).

During this process, particular superfamilies of enzymes such as glutathione-S-transferases (GST) and nonspecific esterases (EST) are usually involved in the biochemical transformation of xenobiotics. Various hormones such as biogenic amines are involved in insect stress reactions. Among these hormones, the role of the neurohormone dopamine (DA) in this process remains poorly understood. It is known that DA mediates phagocytosis and is involved in the activation of the pro-phenoloxidase (proPO) cascade, thus playing an important role in fungal and bacterial pathogenesis as well as in the development of toxicoses caused by insecticides (Delpuech, Frey

& Carton, 1996; Gorman, An & Kanost, 2007; Wu et al., 2015). In addition, both pathogens and toxicants can lead to changes in the antimicrobial activity of insects and the bacterial load that can affect the sensitivity of insects to pathogens (Wei et al., 2017; Ramirez et al., 2018; Polenogova et al., 2019). It should be noted that the above mentioned physiological reactions in mosquito larvae under the combined action of entomopathogenic fungi and insecticides have not yet been studied.

The aims of this study were (1) to determine the susceptibility of *Aedes aegypti* larvae to combined treatment with avermectins and *Metarhizium robertsii* J.F. Bischoff, S.A. Rehner & Humber and (2) estimated their immune and detoxificative responses to *M. robertsii* and avermectins either alone or their combination.

## Materials & methods

### Insecticides and fungi

The entomopathogenic fungus *Metarhizium robertsii* (strain P-72; GenBank No. KP172147) from the collection of microorganisms of the Institute of Systematics and Ecology of Animals SB RAS was used in this work. The conidia of the fungus were grown on autoclaved millet for 10 days at 26°C in the dark, followed by drying and sifting (Belevich et al., 2017). The industrial product “Phytoverm” 0.2 % (SPC “Pharmbiomed”, Russia) was used in these experiments and includes a complex of natural avermectins (A1a (9 %), A2a (18 %), B1a (46 %), B2a (27 %)) produced by *Streptomyces avermitilis*.

### Insect maintenance and toxicity tests

*Aedes aegypti* larvae from the collection of the Institute of Systematics and Ecology of Animals SB RAS were maintained in tap water in the laboratory at 24°C (±1°C) under a natural photoperiod (approximately 16:8 light:dark). The larvae were fed Tetramin Junior fish food (Tetra, Germany). The susceptibility of *Ae. aegypti* to both avermectins and the conidia of *M. robertsii* was tested in 200 ml plastic containers containing 100 ml of water with 15 larvae. 3rd-4th-instar larvae were used in the experiment. The experiment involved four treatments: control, fungus, avermectins, and fungus + avermectins. The fungal conidia and avermectins were suspended (or dissolved) in distilled water, vortexed and applied separately or together to the containers with mosquito larvae at a volume of 2 ml per container. The final conidial concentration for infection was  $1 \times 10^6$  conidia / ml ( $\approx$  LC50). The final concentration of avermectins was 0.00001 % / ml ( $\approx$  LC50). The control was treated with the same amount of distilled water. Mortality was assessed daily for 6 days. Ten replicates with 15 larvae were performed for each treatment.

### Light microscopy and colonization assessment

Forty-eight hours posttreatment (pt), mosquito larvae (n = 3 for each treatment) were collected in 2% glutaraldehyde containing 0.1 M Na-cacodylate buffer (pH 7.2) and were maintained at 4 °C for 1–24 h. Semithin sections were stained with crystal violet and basic fuchsin and were observed with a phase-contrast microscope (Axioskop 40, Carl Zeiss, Germany). To assess fungal colonization, 48 h pt and newly dead larvae (4–6 days pt) were cut open, and their internal contents were squeezed onto a glass slide. The contents were examined for the presence / absence of hyphal bodies using light microscopy (n = 30 for each treatment). Newly dead larvae were placed on moistened filter paper in Petri dishes (n = 30) to determine the

germination and surface sporulation of *Metarhizium*.

### **Total larval body supernatant**

Mosquito larvae bodies of individual 4th-instar larvae of *Ae. aegypti* were collected in 50 µl of cool (+4°C) 0.01M PBS (50 mM, pH 7.4, 150 mM NaCl) with 0.1 mM N-phenylthiourea (PTU) to measure GST or without PTU to measure phenoloxidase (PO) activity. Then, the samples were sonicated in an ice bath with three 10 s bursts using a Bandelin Sonopuls sonicator. The sample solution was centrifuged at 20.000 g for 5 min at +4°C. The obtained supernatant was directly used to determine enzyme activities.

### **Detection of phenoloxidase, glutathione-S-transferase and esterase activity**

The activities of PO, GST and EST were measured at 12 and 48 h after exposure (n = 20 per treatment for each enzyme).

PO activity was assayed by using a method modified from that described by Ashida and Söderhäll (1984). The PO activity of the larval homogenates was determined spectrophotometrically on the basis of the formation of dopachrome at a wavelength of 490 nm. Aliquots of the samples (10 µl) were added to microplate wells containing 200 µl of 10 mM 3,4-dihydroxyphenylalanine (DOPA) and incubated at 28 °C for 45 min.

The activity of EST was measured using the method of Prabhakaran and Kamble (1995) with some modifications. Aliquots of the samples (3 µl) were added to microplate wells containing 200 µl of 0.01 % p-nitrophenylacetate and incubated for 5 min at 28 °C. The activity of EST was determined spectrophotometrically at a wavelength of 410 nm on the basis of the formation of nitrophenyl.

The measurement of GST activity was carried out according to the method of V. Habig et al. (1974) with some modifications. Aliquots of the samples (5 µl) were added to microplate wells containing 200 µl of 1 mM glutathione and 5 µl of 1 mM DNCB and incubated at 28 °C for 20 min. The activity of GST was determined spectrophotometrically on the basis of the formation of 5-(2,4-dinitrophenyl)-glutathione at a 340 nm wavelength.

Enzyme activity was measured in units of the transmission density ( $\Delta A$ ) of the incubation mixture during the reaction per 1 min and 1mg of protein. The protein concentration in the samples was determined by the method of M. Bradford (1976). To generate the calibration curve, bovine serum albumin was used.

### **Dopamine concentration measurements**

DA concentrations were measured at the 12 and 48 h pt in individual larval bodies (n = 10 per treatment). Mosquito larvae were homogenized in 30 µl of phosphate buffer and incubated in a Biosan TS 100 Thermoshaker for 10 minutes at 28 °C and 600 rpm, then incubated at room temperature for 20 minutes and centrifuged at 4 °C and 10.000 g for 10 minutes. The supernatants were transferred to clean tubes and centrifuged with the same settings for 5 minutes. Before transfer to the chromatograph, the samples were filtered.

DA concentrations were measured by an external standard method using an Agilent 1260 Infinity high-performance liquid chromatograph with an EsaCoulchem III electrochemical detector (cell model 5010A, potential 300 mV) according to the method of Gruntenko et al. (2005) with some modifications. Dopamine hydrochloride (Sigma-Aldrich) was used as a standard. Separation was performed in a ZorbaxSB-C18 column (4.6–250 mm, particles 5 µm) in isocratic mode. Mobile phase: 90% buffer (200 mg/l 1-OctaneSulfonicAcid (Sigma-Aldrich), 3.5 g/l KH<sub>2</sub>PO<sub>4</sub>) and 10%

acetonitrile. The flow rate was 1 ml/min. Chromatogram processing was performed using ChemStation software, and the amount of dopamine was determined by comparing the peak areas of the standard and the sample.

### Acid proteases

For the measurement of acid proteases, 50 µl of the homogenate supernatant was added to 250 µl of 0.1 M acetate buffer (pH 4.6) containing 0.3% hemoglobin (Sigma). The samples were incubated for 60 min at 27 °C, and the reaction was stopped by adding 500 µl of 5% TCA and cooling on ice for 10 min at 4 °C. The samples were centrifuged at 14000 g for 5 min at 4 °C, and enzyme activity was determined spectrophotometrically at a wavelength of 280 nm in a 96-well plate reader.

### Lysozyme-like activity

Lysozyme-like activity in the mosquito homogenate was determined through analysis of the lytic zone by diffusion into agar. Ten milliliters of Nutrient Agar (NA) (HiMedia, India) and *Micrococcus lysodeikticus* bacteria ( $1 \times 10^7$  cells/ml) were added to Petri dishes. The agar was perforated to create 2 mm-diameter wells, which were then filled with 3 µl of fresh hemolymph, followed by incubation at 37 °C for 24 h. A series of dilutions of chicken egg white lysozyme (EWL) (Sigma) (0.5 mg/ml, 0.2 mg/ml, 0.1 mg/ml, 0.005 mg/ml, 0.001 mg/ml) were added to each dish, allowing us to obtain a calibration curve based on these standards. Lytic activity was determined by measuring the diameter of the clear zone around each well and expressed as the equivalent of EWL (mg/ml) (Mohrig, Messner, 1968).

### CFU counts of *Metarhizium* and cultivated bacteria in infected larvae

Homogenates of the mosquitoes (3 larvae per sample) were suspended in sterile aqueous Tween-20 (0.03%) (1 ml), and the suspensions were then diluted 50-fold. Next, 100 µl aliquots were inoculated onto the surface of modified Sabouraud agar (10 g peptone, 40 g D-glucose anhydrous, 20 g agar, 1 g yeast extract) supplemented with an antibiotic cocktail (acetyltrimethyl ammonium bromide 0.35 g/L; cycloheximide 0.05 g/L; tetracycline 0.05 g/L; streptomycin 0.6 g/L; PanReacAppliChem, Germany) for the inhibition of bacteria and saprotrophic fungi. The Petri dishes were maintained at 28 °C in the dark. The colonies were then counted after 7 days. For the estimation of cultivated bacterial CFU counts, homogenates of larvae (3 larvae per sample) were suspended in 1 ml of 0.1M phosphate buffer. Then, the suspension was diluted to  $10^{-2}$ ,  $10^{-3}$  and  $10^{-4}$ . Aliquots of 100 µl of the larval dilutions were inoculated onto the surface of blood agar media (HiMedia, India). The Petri dishes were maintained at 28 °C. The colonies were counted after 48 h. Three samples of each treatment were used in the analysis.

### Statistical analysis

Data were analyzed using GraphPad Prism v.4.0 (GraphPad Software Inc., USA), Statistica 8 (StatSoft Inc., USA), PAST 3 (Hammer, Harper, Ryan, 2001) and AtteStat 12.5 (Gaidyshev, 2004). Differences between synergistic and additive effects were determined by comparing the expected and observed insect mortality using the  $\chi^2$  criterion (Tounou et al., 2008). An additive effect was recorded at  $\chi^2 < 3.84$  or a synergistic effect at  $\chi^2 > 3.84$  if the observed mortality was higher than expected, and an antagonistic was recorded at  $\chi^2 > 3.84$  if the observed mortality was lower than expected. The Kaplan-Meier test was used to calculate the median lethal time (presented as  $LT_{50} \pm SE$ ). A log-rank test was used to quantify differences in mortality

dynamics. As the distribution of the physiological parameters except for the dopamine concentration deviated from a normal distribution (Shapiro–Wilk test,  $P < 0.05$ ), we used the nonparametric equivalent of a two-way ANOVA: the Scheirer-Ray-Hare test (Scheirer *et al.*, 1976), followed by Dunn’s post hoc test. The data on dopamine concentrations passed the normality test (Shapiro–Wilk test,  $P > 0.05$ ) and were analyzed by two-way ANOVA followed by Tukey’s post hoc test. Differences between *Metarhizium* CFU counts were compared by t-tests.

## Results

### Synergy between avermectins and the fungus

Significant differences in the dynamics of larval mortality between the treatments were observed (log-rank test:  $\chi^2 = 397.3$  Df = 3,  $P < 0.0001$ ; Fig. 1). Treatment with avermectins or conidia of *M. robertsii* led to 57 and 55% mortality, respectively, whereas combined treatment led to 99% mortality at the 6th day postinfection. Mortality in the control treatment did not exceed 1%. The median lethal time (LT50) post-combined treatment ( $3 \pm 0.1$  d) occurred twice as fast as under treatment with avermectins ( $6 \pm 0.3$  d) or the fungus ( $6 \text{ d} \pm \text{inf.}$ ) alone ( $\chi^2 > 113.7$  Df = 1,  $P < 0.0001$ ).

From the 2nd to the 6th day pt, the avermectins and fungus interacted synergistically ( $\chi^2 > 18.5$ , df = 1,  $P < 0.001$ , ESM Table S1). These effects were consistently observed in four independent experiments.

### Colonization assay

At 48 h pt, we observed mass accumulation of *Metarhizium* conidia in the gut lumen (Fig. 2A). Germinated conidia were not detected in larvae at 48 h pt ( $n = 12$ ). However, in one sample (combined treatment), hyphal bodies were detected in the hemocoel (Fig. 2A). In the newly dead larvae after the fungal and combined treatments (4–6 days), we detected systemic colonization with hyphal bodies (Fig. 2B). Under the combined treatment, 83% hypha-positive larvae were found, while in the fungal treatment, 90% hypha-positive larvae were recorded. No significant differences between these treatments were observed ( $\chi^2 = 0.58$ , df = 1,  $P = 0.45$ ,  $n = 30$  larvae per treatment). No hyphal bodies were detected in the fungus-free treatments. A total of 70% and 60 % of larvae were overgrown with *Metarhizium* under incubation in moist chambers (Fig. 2C) after treatment with the fungus or the mixture (avermectins+fungus), respectively. Nongerminated conidia and hyphal bodies were detected in the water in which treated larvae were maintained (Fig. 2D).

### Phenoloxidase activity

At 12 h pt, we registered a significant decrease in PO activity under the influence of fungal infection (Scheirer-Ray-Haretest, effect of fungus:  $H_{1,52} = 12.6$ ,  $P = 0.00038$ ; Fig. 3). Avermectins did not significantly change PO activity ( $H_{1,52} = 0.3$ ,  $P = 0.54$ ). A stronger decrease in enzyme activity was observed after combined treatment, but a significant factor interaction was not revealed ( $H_{1,52} = 1.9$ ,  $P = 0.16$ ). At 48 h pt, we detected a significant increase in PO activity under the influence of avermectins ( $H_{1,32} = 5.33$ ,  $P = 0.02$ ). The effect of the fungus was not significant ( $H_{1,32} = 0.7$ ,  $P = 0.39$ ), but a tendency toward an interaction between the factors was revealed ( $H_{1,32} = 3.2$ ,  $P = 0.07$ ). This is explained by the inhibition of PO activity by the fungus alone (Dunn’s test,  $P = 0.01$ ,  $P = 0.04$ , compared to the control and avermectin



treatments, respectively) and by the tendency of increased enzyme activity after combined treatment.

### Dopamine concentration

The effects of the fungus or avermectins on the DA concentration at 12 h pt were not significant ( $H_{1,31} = 1.2$ ,  $P = 0.27$ ; Fig. 4), although a trend toward a factor interaction was revealed ( $H_{1,31} = 3.3$ ,  $P = 0.07$ ). This was due to a clear tendency to decrease the DA concentration after treatment with the fungus alone (Dunn's test,  $P = 0.07$  compared to fungus-free treatments) but not with the combination of the fungus and avermectins ( $P = 0.47$  compared to fungus-free treatments). At 48 h pt, we observed a significant decrease in the DA concentration under the influence of the fungus ( $H_{1,31} = 4.62$ ,  $P = 0.03$ ); however, there were no significant differences between the treatments (Dunn's test,  $p = 0.1$ ). No significant interaction effects between the factors on the DA concentration at 48 h pt were detected ( $H_{1,31} = 0.0$ ,  $P = 1.0$ ).

### Detoxifying enzymes

At 12 h pt, an interaction effect between the two factors (avermectins and the fungus) on GST activity was observed ( $H_{1,44} = 8.1$ ,  $P = 0.0043$ , Fig. 5A). Avermectins and the fungus alone significantly (1.5–2-fold) increased GST activity compared to untreated larvae (Dunn's test,  $P = 0.007$ ,  $P = 0.001$ , respectively), but after combined treatment, the enzyme activity did not significantly differ from that in the control. Similar patterns were registered for EST activity at 12 h pt (Fig. 5B). A trend toward an interaction between fungal infection and avermectins was revealed ( $H_{1,44} = 2.14$ ,  $P = 0.1$ ). In this case, EST was activated under the influence of avermectins alone (Dunn's test  $P = 0.002$ , compared to control), but fungal infection inhibited this activation. In particular, EST activity in the fungal and combined treatments did not differ from that in the control (Dunn's test  $P = 0.28$ ,  $P = 0.34$ , respectively).

At 48 h pt, we observed a significant increase in GST activity under the influence of avermectins ( $H_{1,32} = 4.2$ ,  $P = 0.03$ ). The effect of the fungus as well as the interaction between the factors on GST activity at this time point was not significant ( $H_{1,32} = 2.5$ ,  $P = 0.1$  and  $H_{1,32} = 0.61$ ,  $P = 0.43$ , respectively). EST activity nonsignificantly increased under the influence of avermectins ( $H_{1,32} = 1.85$ ,  $P = 0.17$ ). The effect of the fungus on enzyme activity was not significant ( $H_{1,32} = 0.001$ ,  $P = 0.96$ ), and no significant interactions between the factors were detected ( $H_{1,32} = 0.49$ ,  $P = 0.48$ ).

### Acid protease activity

At 12 h pt, a significant interaction between avermectins and the fungus on acid protease activity was observed ( $H_{1,48} = 14.8$ ;  $P = 0.00011$ ; Fig. 6). In particular, protease activity was increased after fungal treatment alone (3-fold compared to control,  $P = 0.0003$ ) but not after combined treatment. At 48 h pt, protease activity was strongly increased under the influence of avermectins ( $H_{1,43} = 27.4$ ,  $P = 0.000016$ ), and a trend toward an increase in enzyme activity was registered under the influence of the fungus ( $H_{1,43} = 3.18$ ,  $P = 0.07$ ). No significant interaction between the factors was revealed, although a trend toward the highest increase in protease activity was registered after the combined treatment.

### Lysozyme-like activity

At 12 h pt, we recorded a decrease in lysozyme-like activity under the influence of both fungal infection and avermectins (effect of fungus:  $H_{1,56} = 6.46$   $p = 0.011$ ; effect of avermectins:  $H_{1,56} =$

14.04  $p = 0.00017$ ; Fig. 7). The greatest decrease was observed after the combined treatment (Dunn's test:  $P < 0.001$  compared with the other treatments). At 48 h pt, a sharp (1.7–2.0-fold) increase in lysozyme-like activity was recorded under the influence of avermectins ( $H_{1,116} = 68.6$ ,  $P < 0.0000001$ ). The effect of the fungus on the level of the enzyme at 48 h pt was not significant. No significant interaction effect between the factors on the level of lysozyme was observed at 12 and 48 h pt ( $H = 0.23$ ,  $P = 0.63$ ).

### Fungal and bacterial CFUs

The plating of mosquito larval homogenates on modified Sabouraud agar showed significant differences in the *Metarhizium* CFUs between treatment with the fungus either alone or combined with avermectins (Fig. 8A). The *Metarhizium* CFU count in the fungal treatment was twice as high as that in the combined treatment ( $t = 6.4$ ,  $df = 18$ ,  $P = 0.001$ ). Homogenates of larvae from the fungus-free treatments (avermectin alone and control) did not form any fungal colonies.

The plating of larval homogenates on blood agar showed a significant (17–75-fold) increase in bacterial CFUs after treatment with the fungus and avermectins. A significant effect was registered for avermectins ( $H_{1,19} = 4.8$ ,  $P = 0.03$ ; Fig. 8B) but not for the fungus ( $H_{1,19} = 1.9$ ,  $P = 0.17$ ). However, a clear tendency toward an increase in CFUs was registered after treatment with the fungus alone (Dunn's test,  $P = 0.054$  compared to control). No significant interaction effect between the fungus and avermectins on bacterial CFUs was revealed ( $H_{1,19} = 1.9$ ,  $P = 0.17$ ).

### Discussion

We showed a synergistic effect between avermectins and *Metarhizium* fungi on aquatic invertebrates for the first time. A similar effect was shown previously only in terrestrial insects (Colorado potato beetle, cotton moth) (Anderson *et al.*, 1989; Asi *et al.*, 2010; Tomilova *et al.*, 2016), which are characterized by a completely different mode of fungal penetration (through the exo-skeleton). The accumulation of conidia of the fungus mainly in the gut lumen of mosquitoes coincides with studies of other researchers (Butt *et al.*, 2013). However, we report the first observation of systemic colonization of *Ae. aegypti* larvae after inoculation with *Metarhizium* conidia. It was previously suggested that only blastospores (and not conidia) are able to germinate from the gut lumen into the hemocoel of mosquito larvae (Alkhaibari *et al.*, 2016, 2018). Interestingly, the larvae treated with avermectins accumulated a lower amount of conidia, but this dose was sufficient for a synergistic effect on mortality. Apparently, the reduced accumulation of conidia was due to the specific binding of avermectins in the bodies of invertebrates with glutamate-dependent chlorine channels / receptors, which are located mostly in the swallowing and somatic muscles (Etter *et al.*, 1999; Forrester *et al.*, 1999). As a result, the larva's ability to swallow food, including the conidia of the fungus, is blocked.

We observed a decrease in PO activity and dopamine levels under the influence of the fungus, whereas in terrestrial arthropods, these enzymes are activated during mycoses (Ling & Yu, 2005; Yassine, Kamareddine & Osta, 2012; Yaroslavtseva *et al.*, 2017; Chertkova *et al.*, 2018). It has been suggested that DA release is associated with the general stress reactions related to the insect's responses to pathogens (Hirashima, Sukhanova & Rauschenbach, 2000; Chertkova *et al.*, 2018). In addition, DA is involved in the modulation of energetic metabolism and general defense mechanisms such as phagocytosis (Wu *et al.*, 2015). PO is involved in the inactivation of fungal propagules in the cuticle and hemocoel (Butt *et al.*, 2016). Dopamines are involved in the PO cascade (Andersen, 2010); however, synchronous and unidirectional changes in the levels of

PO and dopamines are not always observed during infections (Chertkova, 2016, unpublished data). Perhaps conidia entering the intestine of *Ae. aegypti* larvae are able to inhibit the initiation of the PO cascade. Interestingly, Alkhaibar et al. (2018) noted a short-term increase in PO activity in whole-body homogenates of *Culex quinquefasciatus* larvae after infection with conidia or blastospores of *M. brunneum* (4–6 h pt). It is possible that this reaction depends on the type and strain of the pathogen as well as the type and line of mosquitoes. We observed activation of detoxifying enzymes (GST, EST) in *Ae. aegypti* larvae at the early stages of toxicosis and infection (12 h pt) under mono-treatments with the fungus and avermectins. However, the combined treatment led to inhibition of the activation of GST and EST. A similar effect was observed at 12 h pt for antibacterial (lysozyme-like) and acid protease activities. Combined treatment leads to either inhibition or containment of the activation of these enzymes. GST and EST are used by insects to inactivate toxic products formed by insecticide-induced toxicosis (DeSilva et al., 1997; Boyer et al., 2006; Aponte et al., 2013) as well as under mycosis (Serebrov et al., 2006; Han et al., 2013) by reducing oxidative stress and speeding up the metabolism of toxins. Lysozyme inhibits the reproduction of Gram-positive bacteria (Abdou et al., 2007; Gandhe et al., 2007; Chapelle et al., 2009), which (e.g., Microbacteriaceae) are among the dominant bacteria in *Ae. aegypti* larvae (Kerri et al., 2014). It should also be noted that at 12 h pt of *Ae. aegypti* with blastospores of *M. brunneum*, a decrease in the expression levels of genes encoding defensins and cecropins (Alkhaibari et al., 2016), which inhibit the growth of both Gram-positive and Gram-negative bacteria and fungi, was observed (Jozefiak et al., 2017). The inhibition of acid protease activity under combined treatment may indicate disorders in food consumption and absorption. Disruption of food absorption and starvation can increase mortality from both fungi and insecticides (Furlong, Groden, 2003). Thus, we assume that the physiological causes of the observed synergism lie in the initial stages of the development of infection and toxicosis. In the later stages (48 h pt), we mainly observed activation of enzymes (PO, GST, EST, acid proteases and lysozyme-like activity), which apparently indicates destructive processes in tissues and organs under the action of both avermectins and fungi. The increase in PO activity on the second day after treatment with avermectins was probably due to the destruction of hemocytes and the release of intracellular proPO components. We have previously shown the cytotoxic effect of avermectins on hemocytes, leading to their death (Tomilova et al., 2016). Additionally, the cytostatic and cytotoxic effects of the avermectins complex on various cells of warm-blooded animals are well known (Sivkov et al., 1998; Kokoz et al., 1999; Korystov et al., 1999; Maioli et al., 2013). It should also be noted that an increase in PO in insects can be observed with the development of various bacterioses (e.g., Grizanova et al., 2014; Yaroslavtseva et al., 2017), which is consistent with the observed increase in CFUs under the influence of avermectins. An increase in GST under mycoses usually correlates with the severity of the infectious process (Dubovskiy et al., 2012; Tomilova et al., 2019) and confirms the results obtained by Butt et al. (2013) when studying the pathogenesis of *M. brunneum* in *Ae. aegypti* larvae. An increase in lysozyme-like activity under the action of avermectins could have occurred due to tissue destruction accompanied by the release of lysosome contents containing lysozyme (Zachary, Hoffmann, 1984). The activation of proteases at 48 h pt under the influence of avermectins is correlated with the increase in PO and lysozyme-like activity, which also indicates destructive changes in the tissues. We observed an increase in the number of cultivated bacteria in the larvae when treated with both avermectins and fungi. This effect may be associated with impaired intestinal peristalsis as

well as changes in the level of PO and antibacterial activity in the initial stages of toxicosis and fungal infection. Similar effects have been observed in terrestrial insects following topical infection by fungi (Wei et al., 2017; Ramirez et al., 2018; Polenogova et al., 2019) and are associated with the redistribution of immune responses between the cuticle and the intestines. In mosquito larvae, the fungus comes into direct contact with intestinal bacteria, which may exhibit fungistatic properties (Sivakumar et al., 2017; Zhang et al., 2017, etc.) or, alternatively, may act as synergists of fungi, as shown by Wei et al. (2017) in the imagoes of the mosquito *Anopheles stephensi*. It is possible that conflicting data on the systemic colonization of mosquito larvae by *Metarhizium* fungi are associated with differences in bacterial communities, which requires further research. In any case, fungi cannot successfully completely colonize an aquatic environment, and bacterial decomposition is observed in mosquito larvae, whereas fungus germination occurs in the air environment.

## Conclusions

In conclusion, this is the first study of the survival and physiological reactions of mosquito larvae under the combined action of avermectins and entomopathogenic fungi. The synergism observed under the combined action of these agents appears to be associated with physiological changes in the early stages of toxicosis and infection. In particular, inhibition of the activity of a number of enzymes is observed under the combined treatment associated with the detoxifying and immune systems. Systemic colonization of mosquito larvae by the fungus *Metarhizium* is shown for the first time in this study. Further investigations may be focused on studying the role of endosymbiotic mosquito bacteria in the development of toxicosis and mycoses as well as the development of preparative forms based on fungi and avermectins for regulating the number of mosquitoes in natural conditions.

## Acknowledgments

The authors are grateful to Dr. V.A. Shilo (Karasuk Station of the ISEA) for help in organizing the experiments.

## References

- Abdou, A.M., Higashiguchi, S., Aboueleinin, A.M., Kim, M., & Ibrahim, H.R. 2007. Antimicrobial peptides derived from hen egg lysozyme with inhibitory effect against *Bacillus* species. *Food Control*, 18(2):173-178. DOI: 10.1016/j.foodcont.2005.09.010.
- Alkhaibari, A.M., Carolino, A.T., Yavasoglu, S.I., Maffei, T., Mattoso, T.C., Bull, J.C., ... & Butt, T. M. 2016. *Metarhizium brunneum* blastospore pathogenesis in *Aedes aegypti* larvae: attack on several fronts accelerates mortality. *PLoS pathogens*, 12(7):e1005715. DOI: 10.1371/journal.ppat.1005715.
- Alkhaibari, A.M., Maffei, T., Bull, J.C., & Butt, T.M. 2018. Combined use of the entomopathogenic fungus, *Metarhizium brunneum*, and the mosquito predator, *Toxorhynchites brevipalpis*, for control of mosquito larvae: Is this a risky biocontrol strategy? *Journal of invertebrate pathology*, 153:38-50. DOI: 10.1016/j.jip.2018.02.003.
- Alout, H., Krajacich, B.J., Meyers, J.I., Grubaugh, N.D., Brackney, D.E., Kobylinski, K.C., ... & Dabiré, R.K. 2014. Evaluation of ivermectin mass drug administration for malaria transmission control across different West African environments. *Malaria journal*, 13(1):417. DOI: 10.1186/1475-2875-13-417.

Alves, S.N., Serrão, J.E., Mocelin, G., & Melo, A.L.D. 2004. Effect of ivermectin on the life cycle and larval fat body of *Culex quinquefasciatus*. *Brazilian archives of biology and technology*, 47(3):433-439. DOI: 10.1590/S1516-89132004000300014.

Andersen, S.O. 2010. Insect cuticular sclerotization: a review. *Insect biochemistry and molecular biology*, 40(3):166-178. DOI: 10.1016/j.ibmb.2009.10.007.

Anderson, T.E., Hajek, A.E., Roberts, D.W., Preisler, H.K., & Robertson, J.L. 1989. Colorado potato beetle (Coleoptera: Chrysomelidae): effects of combinations of *Beauveria bassiana* with insecticides. *Journal of Economic Entomology*, 82(1):83-89. DOI: 10.1093/jee/82.1.83.

Aponte, H.A., Penilla, R.P., Dzul-Manzanilla, F., Che-Mendoza, A., López, A.D., Solis, F., ... & Rodríguez, A.D. 2013. The pyrethroid resistance status and mechanisms in *Aedes aegypti* from the Guerrero state, Mexico. *Pesticide biochemistry and physiology*, 107(2):226-234. DOI: 10.1016/j.pestbp.2013.07.005.

Ashida, M., & Söderhäll, K. 1984. The prophenoloxidase activating system in crayfish. *Comparative Biochemistry and Physiology Part B: Comparative Biochemistry*, 77(1):21-26. DOI: 10.1016/0305-0491(84)90217-7.

Asi, M.R., Bashir, M.H., Afzal, M., Ashfaq, M., & Sahi, S. T. 2010. Compatibility of entomopathogenic fungi, *Metarhizium anisopliae* and *Paecilomyces fumosoroseus* with selective insecticides. *Pakistan Journal of Botany*, 42(6):4207-4214.

Badiane, T.S., Ndione, R.D., Toure, M., Seye, F., & Ndiaye, M. 2017. Larvicidal and Synergic Effects of two Biopesticides (*Azadirachta indica* and *Metarhizium anisopliae*) against Larvae of *Culex quinquefasciatus* (Diptera, Culicidae) (Say, 1823). *International Journal of Sciences*, 6:109-115. DOI: 10.18483/ijSci.1258.

Belevich, O.E., Yurchenko, Yu.A., Glupov, V.V., Kryukov, V.Yu. 2017. Effect of entomopathogenic fungus *Metarhizium robertsii* on non-target organisms, water bugs (Heteroptera: Corixidae, Naucoridae, Notonectidae). *Journal of Applied entomology and Zoology*, 52:439-445. DOI: 10.1007/s13355-017-0494-z.

Bhatt S., Gething P.W., Brady O.J., Messina J.P., Farlow A.W., Moyes C.L., Drake J.M., Brownstein J.S., Hoen A.G., Sankoh O., Myers M.F., George D.B., Jaenisch T., Wint G.R.W., Simmons C.P., Scott T.W., Farrar J.J., Hay S.I. 2013. The global distribution and burden of dengue. *Nature*, 496(7446):504-507. DOI: 10.1038/nature12060.

Boyer, S., David, J.P., Rey, D., Lemperiere, G., & Ravanel, P. 2006. Response of *Aedes aegypti* (Diptera: Culicidae) larvae to three xenobiotic exposures: larval tolerance and detoxifying enzyme activities. *Environmental Toxicology and Chemistry: An International Journal*, 25(2):470-476. DOI: 10.1897/05-267R2.1.

Boyer, S., Paris, M., Jegou, S., Lempérière, G., Ravanel, P. 2012. Influence of insecticide *Bacillus thuringiensis* subsp. *Israelensis* treatments on resistance and enzyme activities in *Aedes rusticus* larvae (Diptera: Culicidae). *Biol. Control*. 62:75–81. DOI: 10.1016/j.biocontrol.2012.02.001.

Bradford, M.M. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical biochemistry*, 72(1-2):248-254. DOI: 10.1016/0003-2697(76)90527-3.

Butt, T.M., Greenfield, B.P., Greig, C., Maffei, T.G., Taylor, J.W., Piasecka, J., ... & Quesada-Moraga, E. 2013. *Metarhizium anisopliae* pathogenesis of mosquito larvae: a verdict of accidental death. *PloS one*, 8(12):e81686. DOI: 10.1371/journal.pone.0081686.

Butt, T.M., Coates, C.J., Dubovskiy, I.M., & Ratcliffe, N.A. 2016. Entomopathogenic fungi: new insights into host–pathogen interactions. In Lovett B & Leger RJ St, eds. *Genetics and Molecular Biology of Entomopathogenic Fungi*. Academic Press, 307-364. DOI: 10.1016/j.jip.2018.02.020.

Campbell, W.C., Fisher, M.H., Stapley, E.O., Albers-Schonberg, G., & Jacob, T.A. 1983.

Ivermectin: a potent new antiparasitic agent. *Science*, 221(4613):823-828. DOI: 10.1126/science.6308762.

Chaccour, C.J., Hammann, F., Alustiza, M., Castejon, S., Tarimo, B.B., Abizanda, G., ... & Aldaz, A. 2017. Cytochrome P450/ABC transporter inhibition simultaneously enhances ivermectin pharmacokinetics in the mammal host and pharmacodynamics in *Anopheles gambiae*. *Scientific reports*, 7(1):8535. DOI: 10.1038/s41598-017-08906-x.

Chapelle, M., Girard, P.A., Cousserans, F., Volkoff, N.A., & Duvic, B. 2009. Lysozymes and lysozyme-like proteins from the fall armyworm, *Spodoptera frugiperda*. *Molecular immunology*, 47(2-3):261-269. DOI:10.1016/j.molimm.2009.09.028.

Chertkova, E.A., Dubovskiy, I.M., Yaroslavl'tseva, O.N., Grizanov, E.V., Kryukov, V.Y., Glupov, V.V. 2016. Changes in the dopamine levels of haemolymph of cabbage armyworm *Mamestra brassicae* L. (Lepidoptera: Noctuidae) and Colorado potato beetle *Leptinotarsa decemlineata* Say (Coleoptera: Chrysomelidae) at various pathogenesis. *Eurasian Entomological Journal*, 15(1):60-67. (In Russian).

Chertkova, E.A., Grizanov, E.V., & Dubovskiy, I.M. 2018. Bacterial and fungal infections induce bursts of dopamine in the haemolymph of the Colorado potato beetle *Leptinotarsa decemlineata* and greater wax moth *Galleria mellonella*. *Journal of invertebrate pathology*, 153:203-206.

Chouin-Carneiro, T., Vega-Rua, A., Vazeille, M., Yebakima, A., Girod, R., Goindin, D., Dupont-Rouzeyrol, M., Lourenco-de-Oliveira, R., & Failloux, A.B. 2016. Differential susceptibilities of *Aedes aegypti* and *Aedes albopictus* from the Americas to Zika virus. *PLoS neglected tropical diseases*, 10(3):e0004543. DOI: 10.1371/journal.pntd.0004543.

Crump, A., & Omura, S. 2011. Ivermectin, 'wonder drug' from Japan: the human use perspective. *Proceedings of the Japan Academy, Series B*, 87(2):13-28. DOI: 10.2183/pjab.87.13.

Delpuech, J.M., Frey, F., & Carton, Y. 1996. Action of insecticides on the cellular immune reaction of *Drosophila melanogaster* against the parasitoid *Leptopilina boulardi*. *Environmental Toxicology and Chemistry: An International Journal*, 15(12):2267-2271. DOI: 10.1002/etc.5620151221.

DeSilva, D., Hemingway, J., Ranson, H., & Vaughan, A. 1997. Resistance to Insecticides in Insect Vectors of Disease: *estA3*, a Novel Amplified Esterase Associated with Amplified *estA1* from Insecticide Resistant Strains of the Mosquito *Culex quinquefasciatus*. *Experimental parasitology*, 87(3):253-259. DOI: 10.1006/expr.1997.4245.

Dreyer, S.M., Morin, K.J., & Vaughan, J.A. 2018. Differential susceptibilities of *Anopheles albimanus* and *Anopheles stephensi* mosquitoes to ivermectin. *Malaria journal*, 17(1):148. DOI: 10.1186/s12936-018-2296-3.

Drinyaev, V.A., Kruglyak, E.B., Mosin, V.A., Ter-Simonyan, V.G., Novik, T.S., Rusakov, S.V., Berezina, N.S., Berezkina, N.E., Sterlina, T.S. 1999. Physicochemical and biological properties of aversectin C. *Appl. Biochem. Microbiol.*, 35(2):177-182.

Dubovskiy, I.M., Slyamova, N.D., Kryukov, V.Y., Yaroslavl'tseva, O.N., Levchenko, M.V., Belgibaeva, A.B., ... & Glupov, V.V. 2012. The activity of nonspecific esterases and glutathione-S-transferase in *Locusta migratoria* larvae infected with the fungus *Metarhizium anisopliae* (Ascomycota, Hypocreales). *Entomological review*, 92(1):27-31. DOI: 10.1134/S0013873812010022.

Dubovskiy, I.M., Whitten, M.M.A., Yaroslavl'tseva, O.N., Greig, C., Kryukov, V.Y., Grizanov, E.V., Mukherjee, K., Vilcinskis, A., Glupov, V.V., Butt, T.M. 2013. Can insects develop

resistance to insect pathogenic fungi? *PLoS One*, 8(4):e60248. DOI: 10.1371/journal.pone.0060248.

Etter, A., Cully, D.F., Liu, K.K., Reiss, B., Vassilatis, D.K., Schaeffer, J.M., & Arena, J.P. 1999. Picrotoxin blockade of invertebrate glutamate - gated chloride channels: subunit dependence and evidence for binding within the pore. *Journal of neurochemistry*, 72(1):318-326. DOI: 10.1111/jnc.1999.72.1.318.

Forrester, S.G., Hamdan, F.F., Prichard, R.K., & Beech, R.N. 1999. Cloning, sequencing, and developmental expression levels of a novel glutamate-gated chloride channel homologue in the parasitic nematode *Haemonchus contortus*. *Biochemical and biophysical research communications*, 254(3):529-534. DOI: 10.1006/bbrc.1998.0106.

Freitas, R.M.C.D., Faria, M.D.A., Alves, S.N., & Melo, A.L.D. 1996. Effects of ivermectin on *Culex quinquefasciatus* larvae. *Revista do Instituto de Medicina Tropical de São Paulo*, 38(4):293-298. DOI:10.1590/S0036-46651996000400010.

Furlong, M.J., & Groden, E. 2003. Starvation induced stress and the susceptibility of the Colorado potato beetle, *Leptinotarsa decemlineata*, to infection by *Beauveria bassiana*. *Journal of Invertebrate Pathology*, 83(2):127-138. DOI: 10.1016/S0022-2011(03)00066-1.

Gaidyshev, I.P. 2004. Solution of scientific and engineering tasks by means of Excel, VBA and C/C++. *St. Petersburg: BKhV-Petersburg*.

Gandhe, A.S., Janardhan, G., & Nagaraju, J. 2007. Immune upregulation of novel antibacterial proteins from silkmoths (Lepidoptera) that resemble lysozymes but lack muramidase activity. *Insect biochemistry and molecular biology*, 37(7):655-666. DOI: 10.1016/j.ibmb.2007.03.013.

Garrido-Jurado, I., Alkhaibari, A., Williams, S.R., Oatley-Radcliffe, D.L., Quesada-Moraga, E., & Butt, T.M. 2016. Toxicity testing of *Metarhizium* conidia and toxins against aquatic invertebrates. *Journal of pest science*, 89(2):557-564. DOI: 10.1007/s10340-015-0700-0.

Gorman, M.J., An, C., & Kanost, M.R. 2007. Characterization of tyrosine hydroxylase from *Manduca sexta*. *Insect biochemistry and molecular biology*, 37(12):1327-1337. DOI: 10.1016/j.ibmb.2007.08.006.

Greenfield, B.P., Peace, A., Evans, H., Dudley, E., Ansari, M.A., & Butt, T.M. 2015. Identification of *Metarhizium* strains highly efficacious against *Aedes*, *Anopheles* and *Culex* larvae. *Biocontrol science and technology*, 25(5):487-502. DOI: 10.1080/09583157.2014.989813.

Grizanov, E.V., Dubovskiy, I.M., Whitten, M.M.A., & Glupov, V.V. 2014. Contributions of cellular and humoral immunity of *Galleria mellonella* larvae in defence against oral infection by *Bacillus thuringiensis*. *Journal of invertebrate pathology*, 119:40-46. DOI: 10.1016/j.jip.2014.04.003.

Gruntenko, N.E., Karpova, E.K., Adonyeva, N.V., Chentsova, N.A., Faddeeva, N.V., Alekseev, A.A., & Rauschenbach, I.Y. 2005. Juvenile hormone, 20-hydroxyecdysone and dopamine interaction in *Drosophila virilis* reproduction under normal and nutritional stress conditions. *Journal of insect physiology*, 51(4):417-425.

Hammer, O., Harper, D.A., & Ryan, P.D. 2001. Palaeontological statistics software package for education and data analysis. *Palaeontologia Electronica*, 4(1):1-9.

Han, P., Jin, F., Dong, X., Fan, J., Qiu, B., & Ren, S. 2013. Transcript and protein profiling analysis of the destruxin A-induced response in larvae of *Plutella xylostella*. *PLoS One*, 8(4):e60771. DOI: 10.1371/journal.pone.0060771.

Habig, W.H., Pabst, M.J., & Jakoby, W.B. 1974. Glutathione S-transferases the first enzymatic step in mercapturic acid formation. *Journal of biological Chemistry*, 249(22), 7130-7139.

Hemingway, J., & Ranson, H. 2000. Insecticide resistance in insect vectors of human disease. *Annual review of entomology*, 45(1):371-391. DOI: 10.1146/annurev.ento.45.1.371.

Hirashima, A., Sukhanova, M. J., & Rauschenbach, I. Y. 2000. Biogenic amines in *Drosophila virilis* under stress conditions. *Bioscience, biotechnology, and biochemistry*, 64(12):2625-2630. DOI: 10.1271/bbb.64.2625.

Jozefiak, A., & Engberg, R.M. 2017. Insect proteins as a potential source of antimicrobial peptides in livestock production. A review. *Journal of Animal and Feed Sciences*, 26(2):87-99. DOI:10.22358/jafs/69998/2017

Coon, K.L., Vogel, K.J., Brown, M.R., & Strand, M.R. 2014. Mosquitoes rely on their gut microbiota for development. *Molecular ecology*, 23(11):2727-2739. DOI: 10.1111/mec.12771

Kokoz, Y.M., Tsyganova, V.G., Korystova, A.F., Grichenko, A.S., Zenchenko, K.I., Drinyaev, V.A., ... & Victorov, A.V. 1999. Selective cytostatic and neurotoxic effects of avermectins and activation of the GABA  $\alpha$  receptors. *Bioscience reports*, 19(6):535-546. DOI: 10.1023/A:1020262811459.

Korystov, Y.N., Mosin, V.A., Shaposhnikova, V.V., Levitman, M.K., Kudryavtsev, A.A., Kruglyak, E.B., ... & Drinyaev, V.A. 1999. A comparative study of the effects of aversectin C, abamectin and ivermectin on apoptosis of rat thymocytes induced by radiation and dexamethasone. *Acta Veterinaria Brno*, 68(1):23-29.

Ling, E., & Yu, X.Q. 2005. Prophenoloxidase binds to the surface of hemocytes and is involved in hemocyte melanization in *Manduca sexta*. *Insect biochemistry and molecular biology*, 35(12):1356-1366. DOI: 10.1016/j.ibmb.2005.08.007.

Maioli, M.A., de Medeiros, H.C., Guelfi, M., Trinca, V., Pereira, F.T., & Mingatto, F.E. 2013. The role of mitochondria and biotransformation in abamectin-induced cytotoxicity in isolated rat hepatocytes. *Toxicology in Vitro*, 27(2):570-579. DOI: 10.1016/j.tiv.2012.10.017.

Mohrig, W., & Messner, B. 1968. Lysozyme as antibacterial agent in honey and bees venom. *Acta biologica et medica Germanica*, 21(1):85.

Ortiz-Urquiza, A., Luo, Z., & Keyhani, N.O. 2015. Improving mycoinsecticides for insect biological control. *Applied microbiology and biotechnology*, 99(3):1057-1068. DOI: 10.1007/s00253-014-6270-x.

Paris, M., Tetreau, G., Laurent, F., Lelu, M., Despres, L., David, J.P. 2011. Persistence of *Bacillus thuringiensis israelensis* (Bti) in the environment induces resistance to multiple Bti toxins in mosquitoes. *Pest Management Science*. 67(1):122-8. DOI: 10.1002/ps.2046PMID:21162152.

Polenogova, O.V., Kabilov, M.R., Tyurin, M.V., Rotskaya, U.N., Krivopalov, A.V., Morozova, V.V., ... & Glupov, V.V. 2019. Parasitoid envenomation alters the *Galleria mellonella* midgut microbiota and immunity, thereby promoting fungal infection. *Scientific reports*, 9(1):4012. DOI: 10.1038/s41598-019-40301-6.

Prabhakaran, S.K., & Kamble, S.T. 1995. Purification and characterization of an esterase isozyme from insecticide resistant and susceptible strains of German cockroach, *Blattella germanica* (L.). *Insect biochemistry and molecular biology*, 25(4):519-524. DOI: 10.1016/0965-1748(94)00093-E.

Ramirez, J.L., Muturi, E.J., Dunlap, C., & Rooney, A.P. 2018. Strain-specific pathogenicity and subversion of phenoloxidase activity in the mosquito *Aedes aegypti* by members of the fungal entomopathogenic genus *Isaria*. *Scientific reports*, 8(1):9896. DOI: 10.1038/s41598-018-28210-6.



Robertson, J.L., & Preisler, H.K. Pesticide bioassays with arthropods. 1992. *CRC Boca Raton, FL*.

Scheirer, C.J., Ray, W.S., & Hare, N. 1976. The analysis of ranked data derived from completely randomized factorial designs. *Biometrics*, 32(2):429-434. DOI: 10.2307/2529511.

Scholte, E.J., Njiru, B.N., Smallegange, R.C., Takken, W., & Knols, B.G. 2003. Infection of malaria (*Anopheles gambiae* ss) and filariasis (*Culex quinquefasciatus*) vectors with the entomopathogenic fungus *Metarhizium anisopliae*. *Malaria journal*, 2(1):29. DOI: 10.1186/1475-2875-2-29.

Serebrov, V.V., Gerber, O.N., Malyarchuk, A.A., Martemyanov, V.V., Alekseev, A.A., Glupov, V.V. 2006. Effect of entomopathogenic fungi on detoxification enzyme activity in greater wax moth *Galleria mellonella* L. (Lepidoptera, Pyralidae) and role of detoxification enzymes in development of insect resistance to entomopathogenic fungi. *Biology Bulletin*, 33(6):712–718. (In Russian, with English summary). DOI: 10.1134/S1062359006060082.

Sivakumar, G., Rangeshwaran, R., Yandigeri, M.S., Mohan, M., Venkatesan, T., Ballal, C.R., ... & Verghese, A. 2017. Characterization and role of gut bacterium *Bacillus pumilus* on nutrition and defense of leafhopper (*Amrasca biguttula biguttula*) of cotton. *INDIAN J. Agr. Sci*, 87:534-539.

Sivkov G.S., Yakovlev V.V., Chashov I.A. 1998. Influence of the animal's immune response indicators. *Veterinarnaya Medicina*, 5:29-31. (in Russian).

Shoukat, R.F., Freed, S., Ahmad, K.W., & Rehman, A.U. 2018. Assessment of Binary Mixtures of Entomopathogenic Fungi and Chemical Insecticides on Biological Parameters of *Culex pipiens* (Diptera: Culicidae) under Laboratory and Field Conditions. *Pakistan Journal of Zoology*, 50(1):299-299. DOI: 10.17582/journal.pjz/2018.50.1.299.309.

Su, T., Thieme, J., Ocegueda, C., Ball, M., & Cheng, M.L. 2017. Resistance to *Lysinibacillus sphaericus* and other commonly used pesticides in *Culex pipiens* (Diptera: Culicidae) from Chico, California. *Journal of medical entomology*, 55(2):423-428. DOI: 10.1093/jme/tjx235.

Tilquin, M., Paris, M., Reynaud, S., Despres, L., Ravanel, P., Geremia, R.A., & Gury, J. 2008. Long lasting persistence of *Bacillus thuringiensis* subsp. *israelensis* (Bti) in mosquito natural habitats. *PLoS One*, 3(10):e3432. DOI: 10.1371/journal.pone.0003432.

Tolle, M.A. 2009. Mosquito-borne diseases. *Current problems in pediatric and adolescent health care*, 39(4):97-140. DOI: 10.1016/j.cppeds.2009.01.001.

Tomilova, O.G., Kryukov, V.Y., Duisembekov, B.A., Yaroslavl'tseva, O.N., Tyurin, M.V., Kryukova, N.A., ... & Glupov, V.V. 2016. Immune-physiological aspects of synergy between avermectins and the entomopathogenic fungus *Metarhizium robertsii* in Colorado potato beetle larvae. *Journal of invertebrate pathology*, 140:8-15. DOI: 10.1016/j.jip.2016.08.008.

Tomilova, O.G., Yaroslavl'tseva, O.N., Ganina, M.D., Tyurin, M.V., Chernyak, E.I., Senderskiy, I.V., ... & Glupov, V.V. 2019. Changes in antifungal defence systems during the intermoult period in the Colorado potato beetle. *Journal of insect physiology*, 116:106-117. DOI: 10.1016/j.jinsphys.2019.05.003.

Tounou, A.K., Kooyman, C., Douro-Kpindou, O.K., & Poehling, H.M. 2008. Interaction between *Paranosema locustae* and *Metarhizium anisopliae* var. *acridum*, two pathogens of the desert locust, *Schistocerca gregaria* under laboratory conditions. *Journal of invertebrate pathology*, 97(3):203-210. DOI: 10.1016/j.jip.2007.10.002.

Vivekanandhan, P., Karthi, S., Shivakumar, M.S., & Benelli, G. 2018. Synergistic effect of entomopathogenic fungus *Fusarium oxysporum* extract in combination with temephos against

three major mosquito vectors. *Pathogens and global health*, 112(1):37-46. DOI: 10.1080/20477724.2018.1438228.

Vontas, J., Ranson, H., & Alphey, L. 2010. Transcriptomics and disease vector control. *BMC biology*, 8(1):52. DOI: 10.1186/1741-7007-8-52.

Wei, G., Lai, Y., Wang, G., Chen, H., Li, F., & Wang, S. 2017. Insect pathogenic fungus interacts with the gut microbiota to accelerate mosquito mortality. *Proceedings of the National Academy of Sciences*, 114(23):5994-5999. DOI: 10.1073/pnas.1703546114.

Wu, S.F., Xu, G., Stanley, D., Huang, J., & Ye, G.Y. 2015. Dopamine modulates hemocyte phagocytosis via a D1-like receptor in the rice stem borer, *Chilo suppressalis*. *Scientific reports*, 5:12247. DOI: 10.1038/srep12247.

Yaroslavl'tseva, O.N., Dubovskiy, I.M., Khodyrev, V.P., Duisembekov, B.A., Kryukov, V.Y., & Glupov, V.V. 2017. Immunological mechanisms of synergy between fungus *Metarhizium robertsii* and bacteria *Bacillus thuringiensis* ssp. *morrisoni* on Colorado potato beetle larvae. *Journal of insect physiology*, 96:14-20. DOI: 10.1016/j.jinsphys.2016.10.004.

Yassine, H., Kamareddine, L., & Osta, M.A. 2012. The mosquito melanization response is implicated in defense against the entomopathogenic fungus *Beauveria bassiana*. *PLoS pathogens*, 8(11):e1003029. DOI: 10.1371/journal.ppat.1003029.


Zachary, D., & Hoffmann, D. 1984. Lysozyme is stored in the granules of certain haemocyte types in *Locusta*. *Journal of insect physiology*, 30(5):405-411. DOI: [10.1016/0022-1910\(84\)90098-2](https://doi.org/10.1016/0022-1910(84)90098-2).

Zhang, F., Sun, X.X., Zhang, X.C., Zhang, S., Lu, J., Xia, Y.M., ... & Wang, X.J. 2018. The interactions between gut microbiota and entomopathogenic fungi: a potential approach for biological control of *Blattella germanica* (L.). *Pest management science*, 74(2), 438-447. DOI: 10.1002/ps.47262018.

**Figure 1**(on next page)

Mortality dynamics of *Ae. aegypti* larvae after treatment with *M. robertsii* ( $1 \times 10^6$  conidia/ml), avermectins (0.00001%) and their combination.


The control was treated with distilled water. The asterisks (\*) indicate a synergistic effect ( $\chi^2 > 18.5$ ,  $df = 1$ ,  $P < 0.001$ , see Table S1).



# Figure 2

The colonization of *Ae. aegypti* by *Metarhizium*


A - accumulation of conidia in the intestine and colonization of the hemocoel by hyphal bodies; B - colonization of the fat body; C - mosquito larvae with germinated *Metarhizium* conidia in a dampening chamber; D - nongerminated conidia in a sample of water in which mosquito larvae were maintained. Scale bar: 20  $\mu$ m.



# Figure 3(on next page)

Activity of PO in the whole-body homogenates of *Ae. aegypti* larvae after treatment with *M. robertsii*, avermectins and their combination.

In the control treatment, equal amounts of water were added. Boxes denote interquartile ranges (25–75%) bisected horizontally by median values; whiskers indicate min and max values. Significant differences are indicated with different letters within a time point.




# Figure 4(on next page)

Dopamine concentration in whole-body homogenates of *Ae. aegypti* larvae after treatment with *M. robertsii*, avermectins and their combination. In the control treatment, equal amounts of water were added.

Boxes denote interquartile ranges (25–75%) bisected horizontally by median values; whiskers show min and max values. Significant differences are indicated by different letters within a time point.







# Figure 5(on next page)

GST (A) and EST (B) activity in whole-body homogenates of *Ae. aegypti* larvae after treatment with *M. robertsii*, avermectins and their combination. In the control treatment, equal amounts of water were added.


Boxes denote interquartile ranges (25–75%) bisected horizontally by median values; whiskers show min and max values. Significant differences are indicated by different letters within a time point.



# Figure 6(on next page)

Acid protease activity in whole-body homogenates of *Ae. aegypti* larvae after treatment with *M. robertsii*, avermectins and their combination. In the control treatment, equal amounts of water were added.


Boxes denote interquartile ranges (25–75%) bisected horizontally by median values; whiskers show min and max values. Significant differences are indicated by different letters within a time point.



# Figure 7 (on next page)

Lysozyme-like activity in whole-body homogenates of *Ae. aegypti* larvae after treatment with *M. robertsii*, avermectins and their combination. In the control treatment, equal amounts of water were added.

Boxes denote interquartile ranges (25–75%) bisected horizontally by median values; whiskers show min and max values. Significant differences are indicated by different letters within a time point.



# Figure 8(on next page)


Colony forming units of *M. robertsii* (A) and cultivable bacteria (B) in whole-body homogenates of *Ae. aegypti* larvae after treatment with *M. robertsii*, avermectins and their combination.

Boxes denote interquartile ranges (25 – 75 %) bisected horizontally by median values.

Whiskers show min and max values. Significant differences are indicated by different letters.



**A**



**B**

