

# Comparative proteomic analysis of pepper (*Capsicum annuum* L.) seedlings under selenium stress

Chenghao Zhang<sup>1,2</sup>, Baoyu Xu<sup>1</sup>, Wei Geng<sup>3</sup>, Yunde Shen<sup>4</sup>, Dongji Xuan<sup>4</sup>, Qixian Lai<sup>5</sup>, Chenjia Shen<sup>6</sup>, Chengwu Jin<sup>7</sup> and Chenliang Yu<sup>1</sup>

<sup>1</sup> Institute of Agricultural Equipment, Zhejiang Academy of Agricultural Sciences, Hangzhou, Zhejiang, China

<sup>2</sup> Key Laboratory of Creative Agriculture, Ministry of Agriculture, Zhejiang Academy of Agricultural Science, Hangzhou, Zhejiang, China

<sup>3</sup> Vegetable Research Institute, Zhejiang Academy of Agricultural Sciences, Hangzhou, Zhejiang, China

<sup>4</sup> College of Mechanical and Electrical Engineering, Wenzhou University, Wenzhou, Zhejiang, China

<sup>5</sup> Key Laboratory of Creative Agriculture, Ministry of Agriculture, Zhejiang Academy of Agricultural Sciences, Hangzhou, Zhejiang, China

<sup>6</sup> College of Life and Environmental Science, Hangzhou Normal University, Hangzhou, Zhejiang, China

<sup>7</sup> School of Food Engineering, Ludong University, Yantai, Shandong, China

## ABSTRACT

Selenium (Se) is an essential trace element for human and animal health. Se fertilizer has been used to increase the Se content in crops to meet the Se requirements in humans and animals. To address the challenge of Se poisoning in plants, the mechanisms underlying Se-induced stress in plants must be understood. Here, to elucidate the effects of Se stress on the protein levels in pepper, we used an integrated approach involving tandem mass tag labeling, high performance liquid chromatography fractionation, and mass spectrometry-based analysis. A total of 4,693 proteins were identified, 3,938 of which yielded quantitative information. Among them, the expression of 172 proteins was up-regulated, and the expression of 28 proteins was down-regulated in the Se/mock treatment comparison. According to the above data, we performed a systematic bioinformatics analysis of all identified proteins and differentially expressed proteins (DEPs). The DEPs were most strongly associated with the terms “metabolic process,” “posttranslational modification, protein turnover, chaperones,” and “protein processing in endoplasmic reticulum” according to Gene Ontology, eukaryotic orthologous groups classification, and Kyoto Encyclopedia of Genes and Genomes enrichment analysis, respectively. Furthermore, several heat shock proteins were identified as DEPs. These results provide insights that may facilitate further studies on the pepper proteome expressed downstream of the Se stress response. Our data revealed that the responses of pepper to Se stress involve various pathways.

Submitted 23 April 2019  
Accepted 9 October 2019  
Published 27 November 2019

Corresponding authors  
Chengwu Jin, [jinchwu@ldu.edu.cn](mailto:jinchwu@ldu.edu.cn)  
Chenliang Yu, [21007030@zju.edu.cn](mailto:21007030@zju.edu.cn)

Academic editor  
Gilda Eslami

Additional Information and  
Declarations can be found on  
page 15

DOI [10.7717/peerj.8020](https://doi.org/10.7717/peerj.8020)

© Copyright  
2019 Zhang et al.

Distributed under  
Creative Commons CC-BY 4.0

## OPEN ACCESS

**Subjects** Agricultural Science, Bioinformatics, Molecular Biology, Environmental Contamination and Remediation, Environmental Impacts

**Keywords** Pepper, Heat shock proteins, Selenium stress, Metabolic process, Differential expression protein

## INTRODUCTION

Selenium (Se) is a trace element that is essential for human and animal health, and it is an active component of numerous enzymes in human metabolism (Sager, 2006; Semnani et al., 2010). Because of its important protective effects in animals and plants, many studies on Se have been conducted in a broad range of fields including medicine, agriculture, and nutrition (Chen et al., 2002; Thavarajah, Ruszkowski & Vandenberg, 2008). Although there is no direct evidence that Se is necessary for plant growth, Se plays a key role in plant growth and development. Se can improve antioxidant enzyme activity and enhance the tolerance of *Rumex patientia* × *R. tianshanicus* seedlings to salt stress (Kong, Wang & Bi, 2005). In addition, Se enhances plant resistance to abiotic stresses, including heavy metals (Kumar et al., 2012), waterlogging (Wang, 2011), chilling (Chu, Yao & Zhang, 2010), high temperature (Djanaguiraman, Prasad & Seppanen, 2010) and drying (Pukacka, Ratajczak & Kalemba, 2011). Se also plays a critical role in plant resistance to biological stress. Plants with high Se content in grasslands can resist invasion by herbivores (Quinn et al., 2008).

The properties of Se facilitate the formation of stable compound structures with multiple oxidation states (+2, +4, and +6), covalent bonding to non-metals (such as carbon), and strong coordination with metals such as cadmium (Fernandes et al., 2018). The main forms of Se taken up by plants are selenate (VI) and selenite (IV), whereas the Se forms in soil are influenced by the soil pH and oxidation potential (Elrashidi et al., 1987). The uptake and transport mechanisms of the two major valences of Se ( $\text{SeO}_4^{2-}$  and  $\text{SeO}_3^{2-}$ ) in soil differ (White, 2016). The chemical properties of selenate and sulfate are similar (Shibagaki et al., 2002). These compounds are antagonistic during plant uptake, and the sulfate transporter regulates the uptake of selenite (Shibagaki et al., 2002; El Kassis et al., 2007; White, 2016). Currently, the mechanism of selenite absorption by plants is unclear. Most studies suggest that the mechanism of absorption of selenite is similar to that of phosphate, but selenite absorption is negatively correlated with phosphate absorption (Zhang et al., 2014; Song et al., 2017).

At present, Se poisoning incidents in plants have rarely been reported. Preliminary studies have shown that the toxic effect of Se on plants is similar to that of heavy metals to some extent and can hinder plant growth and metabolism. In agricultural environments, excessive Se has been found to decrease radish seeds by 14% and radish yield by 8–9% (Hladun et al., 2013). Se stress in barley hinders plant growth and significantly decreases fresh weight, water content, and photosynthetic capacity (Molnárová & Fargašová, 2009). Paciolla, De Leonardis & Dipierro (2011) have observed that Se treatment (8–16 mg/L) significantly inhibits barley germination (Paciolla, De Leonardis & Dipierro, 2011). In addition, Se (four to six mg/L) significantly inhibits root and bud growth in soybean seedlings, whereas root growth in lettuce and ryegrass is completely inhibited even at a concentration of one mg/L (Hartikainen et al., 1997; Aggarwal et al., 2011). The above

results demonstrate that Se causes toxicity in plants. However, current understating of the molecular mechanism of Se toxicity remains limited ([El-Ramady et al., 2015](#); [Galinha et al., 2015](#); [Jia et al., 2019](#)). Elemental Se and Se compounds are increasingly accumulating in surface soil and water. Excessive amounts of Se pose a potential risk in agricultural production ([Kuppusamy et al., 2018](#); [Jia et al., 2019](#)).

Pepper (*Capsicum annuum* L.), an economically important vegetable in the *Solanaceae* family, has been used as a spice in China and Korea for decades ([Choi et al., 2005](#)). Recently, a novel tandem mass spectrometry (MS/MS)-based tandem mass tag (TMT) labeling strategy was developed for large-scale protein quantification ([Hao et al., 2017](#); [Xu et al., 2017](#)). Relatively limited proteomic data on pepper under Se stress have been reported. In the present study, we used a TMT labeling-based quantitative proteomics approach to identify differentially expressed proteins (DEPs) under Se treatment. Our data enabled the identification and exploration of the roles of candidate proteins associated with Se stress resistance.

## MATERIALS AND METHODS

### Plant materials and Se treatments

Pepper seeds (*C. annuum* 8 #, a cultivar provided by pepper breeding group in Fujian Agriculture and Forestry University) were sterilized with 1% sodium hypochlorite for 30 min and grown in steam-sterilized soil. The seedlings were grown in a greenhouse under the following conditions: 12 h light ( $150 \mu\text{m}^2 \text{s}^{-1}$ ) at 26 °C, 12 h dark at 23 °C, and relative humidity of 60%. Seedlings were irrigated with half-strength Hoagland solution (pH 5.6). Pepper plants at the four true leaf stages were used for Se treatment. Seedlings were sprayed with half-strength Hoagland solution containing 0 or 100 ppm  $\text{Na}_2\text{SeO}_4$ . After 24 h, the shoots were collected for protein extraction.

### Protein extraction and trypsin digestion

Samples were removed from storage at  $-80^\circ\text{C}$ , and fixed amounts of tissue samples were ground to powder while liquid nitrogen was added. The samples in each group were treated with four volumes of phenol extraction buffer (containing 10 mM dithiothreitol, 1% protease inhibitor, and two mM EDTA), then sonicated three times on ice with a high intensity ultrasonic processor (Scientz, Ningbo, China). The supernatant was centrifuged for 10 min at 4 °C and  $5,500\times g$  with an equal volume of Tris equilibrium phenol. The supernatant was collected and precipitated overnight with five volumes of 0.1M ammonium acetate/methanol. The protein precipitate was washed with methanol and acetone successively. Finally, the precipitate was re-dissolved in 8M urea, and the protein concentration was determined with a BCA kit (code P0010; Beyotime, Beijing, China) according to the manufacturer's instructions.

For digestion, the final concentration of dithiothreitol in protein solution was five mM, and reduction was performed by incubation at 56 °C for 30 min. The mixture was then alkylated with 11 mM iodoacetamide at room temperature for 15 min. Finally, the urea concentration of the sample was diluted to less than 2M by addition of 100 mM triethyl ammonium bicarbonate. Trypsin was added at a mass ratio of 1:50 (trypsin:protein), and

enzymatic hydrolysis was carried out overnight at 37 °C. Trypsin was then added at a mass ratio of 1:100 (trypsin:protein), and enzymatic hydrolysis continued for 4 h.

### TMT labeling and HPLC fractionation

The trypsinized peptide segments were desalted with a Strata X C18 column (Phenomenex, Torrance, CA, USA) and then freeze-dried under vacuum. The peptides were dissolved in 0.5M triethyl ammonium bicarbonate and labeled with a TMT kit (ThermoFisher, Shanghai, China) according to the manufacturer's instructions. The procedure was as follows: the labeled reagent was dissolved in acetonitrile after thawing, incubated at room temperature for 2 h after mixing with the peptide segments, then desalinated after mixing with the labeled peptide segment and freeze-dried in a vacuum.

Peptide segments were classified with high pH reverse-phase high performance liquid chromatography (HPLC) with an Agilent 300 Extend C18 column (five  $\mu\text{m}$  diameter, 4.6 mm inner diameter, 250 mm length) (Agilent, Shanghai, China). The gradient of peptide segments was 8–32% acetonitrile (pH 9.0), and more than 60 min was required to separate the peptide segments into 60 components. The peptide segments were then merged into 18 components, which were then vacuum freeze-dried.

### LC-MS/MS analysis

The tryptic peptides were dissolved in 0.1% formic acid (solvent A) and directly loaded onto an Acclaim PepMap 100 reverse-phase pre-column (ThermoFisher, Shanghai, China). The gradient comprised an increase from 6% to 23% solvent B (0.1% formic acid in 98% acetonitrile) over 26 min, from 23% to 35% in 8 min, and to 80% in 3 min; then the concentration was held at 80% for the last 3 min. All steps were performed at a constant flow rate of 500 nL/min on an EASY-nLC 1000 ultra-HPLC system.

After equilibration, the peptides were ionized with an NSI ion source and analyzed by MS/MS (Q Exactive<sup>TM</sup> mass spectrometer; ThermoFisher, Shanghai, China) coupled online to ultra-HPLC. The ion source voltage was set to 2.0 kV, and the parent ions of peptide segments and their secondary fragments were detected and analyzed with a high resolution Orbitrap. The scanning range of the primary MS was set to 350–1,800 m/z, and the scanning resolution was set to 70,000. The scanning range of the secondary MS was set to 100 m/z, and the scanning resolution of the secondary MS was set to 17,500. The data acquisition mode used a data-dependent scanning program; that is, the first 20 peptide ions with the highest signal intensity were selected to enter the higher collision dissociation collision pool in turn after the first scan, and the fragmentation energy was 28% for fragmentation. Similarly, secondary MS analysis was carried out sequentially. To improve the efficiency of MS, AGC was set to 5E4, the signal threshold was set to 2E4, the maximum injection time was set to 100 ms, and the dynamic elimination time of MS/MS scanning was set to 30 s to avoid repeated scanning of parent ions.

### Database search

The resulting MS/MS data were processed and searched against a public proteome *C. annuum* database (<https://www.uniprot.org/uniprot/?query=proteome:UP000222542>)

with the Maxquant search engine (v.1.5.2.8, <https://maxquant.org/>) concatenated with a reverse decoy database. The retrieval parameter settings were as follows: trypsin/P was used for digestion; the number of missing digestion sites was set to two; and the minimum length of peptide segments was set to seven amino acids. The maximum number of modifications of the peptide segment was set to five. The mass error tolerance of the primary parent ions of the first search and main search was set to 20 and 5 ppm, respectively. The mass error tolerance was 0.02 Da. Cysteine alkylation was set as a fixed modification. Oxidation on Met was specified as a variable modification. The quantitative method was set to TMT-6plex, and the false discovery rate for protein identification and PSM identification was set to 1%.

### Protein annotation


The Gene Ontology (GO) annotation proteome was derived from the UniProt-GOA database (<http://www.ebi.ac.uk/GOA/>). Subsequently, proteins were classified according to GO annotation on the basis of three categories: biological process, cellular component, and molecular function. The Kyoto Encyclopedia of Genes and Genomes (KEGG) database was used to annotate protein pathways. KEGG online service tools KAAS ([http://www.genome.jp/kaas-bin/kaas\\_main](http://www.genome.jp/kaas-bin/kaas_main)) were used to annotate the KEGG database descriptions of proteins, and then the annotation results were mapped on the KEGG pathway database by using the KEGG online service tool KEGG mapper (<http://www.kegg.jp/kegg/mapper.html>). For domain annotation, the InterProScan database (<http://www.ebi.ac.uk/interpro/>) was used to annotate the domain functional descriptions of identified proteins. For subcellular localization, WoLFSPORT (a subcellular localization predication software ([http://www.genscript.com/psort/wolf\\_psort.html](http://www.genscript.com/psort/wolf_psort.html))) was used. WoLFSPORT is an updated version of PSORT/PSORT II for the prediction of eukaryotic sequences.

### Statistical analysis

For GO enrichment and pathway analysis, a two-tailed Fisher's exact test was used to test the enrichment of the DEPs against all identified proteins. A GO term or KEGG pathway with a *P*-value < 0.05 was considered significant. The proteins with TMT intensity values were considered quantified, and the minimal PIF was set as 0.75. Statistical analyses were carried out in SPSS ver. 19.0 (SPSS Inc., Chicago, IL, USA). All reported values represent the averages of three replicates with the standard deviation (mean ± SD).

### Quantitative real-time PCR validation

Total RNA was extracted using a Ultrapure RNA Kit according to the manufacturer's protocol (Code: CW0597, CWBIO, Beijing, China). First-strand cDNA synthesis was carried out using a SuperScript™ IV First-Strand Synthesis System according to the manufacturer's protocol (Code:18091050; ThermoFisher, Waltham, MA, USA). QRT-PCR was performed on ABI 7500 Real-Time PCR System (Roche, Basel, Switzerland) using UltraSYBR Mixture(High ROX)Kit (Code: CW2602, CWBIO, Beijing, China) with the primers listed in [Table S1](#). The CaACTIN (Capana12g001934) was used as an internal



**Figure 1** Quality control (QC) validation of Mass spectrometer (MS) data. (A) Heatmap of Pearson correlation coefficients from all quantified proteins between each pair of samples. Protein were extracted in three biological replicates for each sample group. All protein samples were trypsin digested and analyzed by HPLC-MS/MS. (B) Two-dimensional scatter plot of PCA (principal component analysis) distribution of all samples using quantified proteins. (C) Relationship between molecular weight and coverage of proteins identified by mass spectrometry. (D) Mass error distribution of all identified peptides. (E) Basic statistical data of MS results. (F) Length distribution of all identified phosphorylated peptides. [Full-size](#) DOI: 10.7717/peerj.8020/fig-1

standard to calculate relative fold-differences based on comparative cycle threshold ( $2^{-\Delta\Delta C_t}$ ) values.

## RESULTS

### Primary MS data and quantitative proteome analysis

A total of 252,213 secondary spectra were obtained by MS analysis of the mock treated and Se treated pepper seedlings. A Pearson correlation coefficient analysis indicated high replicability of the experiment (Fig. 1A). A search of the proteome *C. annuum* database indicated 47,316 spectra were available, and the utilization rate of the spectra was 18.8%. A total of 24,048 peptide fragments were identified by spectral analysis, including 21,704 unique peptide fragments (Fig. 1B). Most of the peptides contained 7–20 amino acids, results consistent with the trypsin enzymatic hydrolysis and high-energy collision dissociation fragmentation (Fig. 1C). The molecular weights of the proteins were


negatively correlated with their coverage (Fig. 1D). The first-order mass error of most spectra was less than 10 ppm, in agreement with the high accuracy of orbital well MS. The results indicated high mass accuracy of the MS data (Fig. 1E). Principal component analysis of the quantitative protein data for all samples is presented in Fig. 1F. Detailed information on the identified peptides, including amino acid sequences, protein descriptions, carried charge of peptide, The maximal posterior error probability for peptides (PEP) is listed in Table S2.

A total of 4,693 proteins were identified, among which 3,938 were quantifiable. To understand the functions and characteristics of the proteins identified and quantified in the data, we performed detailed annotation analysis on the basis of GO, protein domain, KEGG pathway, KOG functional classification (eukaryotic orthologous groups), and subcellular localization (Table S3).

### Effects of Se treatment on the global proteome of pepper seedlings

A total of 200 DEPs were identified with a fold-difference expression threshold of 1.5 (Se/Mock ratio  $\geq 1.5$  or  $\leq 0.667$ ) and a *t*-test *P*-value  $< 0.05$  (Table S4). All identified proteins and DEPs under Se treatment were grouped into three GO categories (biological process, cellular component, and molecular function) (Fig. 2A). In the biological process category, 1,594 identified proteins and 67 DEPs were involved in “metabolic processes,” 1,423 identified proteins and 46 DEPs were involved in “cellular processes,” and 1,107 identified proteins and 46 DEPs were involved in “single-organism processes.” In the molecular function category, 1,959 identified proteins and 74 DEPs had “catalytic activities,” 1,899 identified proteins and 74 DEPs had “binding activities,” and 178 identified proteins and one DEP had “structural activities.” In the cellular components category, 737 identified proteins and 10 DEPs were “cell”-related proteins; 418 identified proteins and three DEPs were “macromolecular complex”-related proteins; and 402 identified proteins and four DEPs were “organelle”-related proteins. The distribution of the GO annotations of the up-regulated and down-regulated DEPs is shown in Fig. S1. We also used WoLFPSORT (<http://wolffpsort.org/>) software to determine the subcellular location prediction and classification statistics for all identified proteins and DEPs (Fig. 2B), which were grouped according to their subcellular localizations. All identified proteins were classified into 15 subcellular components, including 1,728 chloroplast-localized proteins, 1,283 cytoplasm-localized proteins, and 784 nuclear-localized proteins. For the DEPs, 13 subcellular components were identified, including 72 cytoplasm-localized DEPs, 44 chloroplast-localized DEPs, and 42 nuclear-localized DEPs.

The expression profiles of the DEPs in six samples are presented in a heat map (Fig. 3A). To reveal the changing trends among the six samples, we assigned all DEPs to one of three clusters (I–III) by using MeV (<https://sourceforge.net/projects/mev-tm4/>) with the K-means method (Fig. 3B). The proteins in clusters I and II showed up-regulation, whereas the proteins in cluster III were down-regulated in the Se stress treatment replicates. Among the DEPs, 172 were up-regulated and 28 were down-regulated (Figs. 3C and 3D). The 16.6 kDa heat shock protein (HSP) (A0A1U8FR15) and two 18.5 kDa class I HSPs (A0A1U8DWR1, A0A1U8E6M6) were up-regulated over ninefold by Se treatment




**Figure 2** Classification of all identified proteins and DEPs. (A) GO analysis of all identified proteins and DEPs. All proteins were classified by GO terms based on three categories: molecular function, biological process and cellular component. (B) Subcellular classify of all identified proteins and DEPs. [Full-size !\[\]\(5fd6ef84f97f42d7f8b34275f1b65312\_img.jpg\) DOI: 10.7717/peerj.8020/fig-2](https://doi.org/10.7717/peerj.8020/fig-2)

compared with the mock treatment. In addition, a glycine-rich protein (A0A2G2ZTC5) and histone H1 protein were down-regulated more than twofold by Se treatment compared with the mock treatment. A total of 136 DEPs were classified into 20 KOG terms. “Post-translational modification, protein turnover, chaperones” contained the largest DEPs (Fig. 4).

### Enrichment analysis of DEPs under Se treatment

To determine whether the DEPs were significantly enriched in some functional types, we performed an enrichment analysis of DEPs by using GO classification, KEGG pathways, and protein domains. Among the DEPs, most up-regulated proteins were enriched in “sequence-specific DNA binding,” “iron ion binding,” “prephenate dehydrogenase (NAD<sup>+</sup>/NADP<sup>+</sup>) activity,” “nucleic acid binding transcription factor activity,” and “apoplast” (Fig. 5A). For the down-regulated proteins, the top five enriched GO terms were






**Figure 3** Impacts Se stress treatment on proteome levels in pepper. (A) Expression profiles of the DEPs response to Se stress. (B–D) All DEPs were analyzed and clustered into three major Clusters by K-means method. (E) Volcano plot of DEPs. (F) The numbers of up- and down-regulated proteins in the Se treatment seedlings compared to the mock seedlings. [Full-size !\[\]\(fcc3264021d438d9732560e78099f674\_img.jpg\) DOI: 10.7717/peerj.8020/fig-3](https://doi.org/10.7717/peerj.8020/fig-3)

“thiamine-containing compound metabolic process,” “chlorophyll metabolic process,” “porphyrin-containing compound biosynthetic process,” “water-soluble vitamin biosynthetic process,” and “vitamin biosynthetic process” (Fig. 5B).

Under Se treatment, 131 DEPs were grouped into different KEGG pathways, seven of which were enriched ( $P < 0.05$ ). For the up-regulated proteins, the DEPs were associated with “protein processing in endoplasmic reticulum,” “endocytosis,” “sesquiterpenoid and triterpenoid biosynthesis,” “SNARE interactions in vesicular transport,” and “plant-pathogen interaction” (Fig. 6A). For the down-regulated proteins, the DEPs were associated with “thiamine metabolism” and “porphyrin and chlorophyll metabolism” (Fig. 6B). In the present study, 32 DEPs in the Se treated pepper were identified to be involved in nine metabolic pathways, most of which were significantly up-regulated by Se treatment. However, in the “thiamine metabolism” pathway, four proteins (A0A2G3ALC4, A0A1U8FC73, A0A2G3A131, and A0A2G2Z8I0) were significantly down-regulated by Se treatment (Table 1).




**Figure 4** KOG functional classification chart of differentially expressed proteins.

Full-size DOI: [10.7717/peerj.8020/fig-4](https://doi.org/10.7717/peerj.8020/fig-4)


Protein domain enrichment analysis revealed that 25 protein domains were enriched in the DEPs (Fig. 7A). The five most enriched protein domains were “alpha crystallin/Hsp20 domain” (13 proteins), “HSP20-like chaperone” (14 proteins), “target SNARE coiled-coil homology domain” (four proteins), “HSP 70 kDa” (five proteins), and “DnaJ domain” (six proteins). Many HSPs were identified according to the results of the protein domain enrichment analysis. In total, 23 HSPs were significantly up-regulated (Fig. 7B). The expression levels of some HSP genes were basically consistent with the proteomic analyses (Fig. S2).

## DISCUSSION

With the increased accumulation of Se in the soil as a result of anthropogenic activity, Se, which has toxic effects on plants at certain concentrations, has gradually become a potential environmental risk factor. In studies of Se, quantitative proteomics analyzed through MS have been applied primarily in analysis of bio-transformation of Se-containing compounds in experimental models such as animals, yeast, and cancer cells (Zhang *et al.*, 2010; Sinha *et al.*, 2016). In general, proteomic analysis technologies have rarely been applied in the study of Se metabolism in plants. In the present study,




**Figure 5** GO enrichment analysis of DEPs. Distribution of the up-regulated (A) and down-regulated (B) proteins with GO enrichment analysis. Full-size [DOI: 10.7717/peerj.8020/fig-5](https://doi.org/10.7717/peerj.8020/fig-5)



**Figure 6** KEGG enrichment analysis of the DEPs in pepper after Se stress treatment. (A) Significantly enriched KEGG terms of the up-regulated proteins. (B) Significantly enriched KEGG terms of the down-regulated proteins. Full-size [DOI: 10.7717/peerj.8020/fig-6](https://doi.org/10.7717/peerj.8020/fig-6)

**Table 1** Identification of the DEPs involved in metabolic pathways.

Protein accession	Protein description	Ratio	P-value	MW (kDa)
Butanoate metabolism				
A0A1U8E166	“Hydroxymethylglutaryl-CoA lyase, mitochondrial	1.788	0.00039971	45.332
A0A2G2YVX0	Glutamate decarboxylase	4.962	1.6822E-07	59.504
Cysteine and methionine metabolism				
A0A2G3AEL6	L-lactate dehydrogenase	4.042	0.00132081	37.54
A0A2G3ADN2	1-aminocyclopropane-1-carboxylate synthase	3.506	0.00199508	54.867
A0A1U8FJ05	1-aminocyclopropane-1-carboxylate oxidase 4	1.601	1.67031E-05	36.33
A0A1U8FEU9	1-aminocyclopropane-1-carboxylate oxidase 1	3.69	1.6569E-06	36.059
A0A1U8EYM1	Tyrosine aminotransferase	2.399	0.00038372	47.177
A0A1U8E953	Arginine decarboxylase	3.279	4.8968E-06	78.211
A0A1U8FBD0	Proline dehydrogenase	2.603	0.00046451	55.32
A0A2G2ZVP6	Lipoxygenase	2.359	2.6782E-07	102.6
A0A2G2YXE3	Glyoxysomal fatty acid beta-oxidation multifunctional protein MFP-a	4.247	0.00027878	101.81
Glycerophospholipid metabolism				
A0A2G2YY43	Glycerol-3-phosphate 2-O-acyltransferase 4	0.625	0.00053958	51.031
A0A1U8G1E3	Glycerophosphodiester phosphodiesterase GDPD2	2.767	2.4846E-07	42.766
A0A1U8H0F1	Glycerol-3-phosphate acyltransferase 5	2.083	0.00026256	55.181
A0A2G2ZQ18	Putative choline kinase 1	2.506	0.000083157	40.256
Linoleic acid metabolism				
A0A2G2ZVP6	Lipoxygenase	2.359	2.6782E-07	102.6
A0A2G2ZBY6	Lipoxygenase	3.646	2.0736E-08	97.904
Phenylpropanoid biosynthesis				
A0A2G2YQ27	Phenylalanine ammonia-lyase	4.291	3.8524E-08	78.308
A0A1U8DW23	Peroxidase	1.569	0.000022029	36.129
A0A2G2YUF1	Retinal dehydrogenase 1	2.003	3.8333E-06	54.692
A0A2G3A835	Caffeoyl-CoA O-methyltransferase 1	1.769	0.000020361	27.232
Phenylalanine, tyrosine and tryptophan biosynthesis				
A0A1U8FBU3	Arogenate dehydrogenase 1, chloroplastic	1.76	0.0026381	42.449
A0A1U8EYM1	Tyrosine aminotransferase	2.399	0.00038372	47.177
A0A2G2YRI9	Arogenate dehydrogenase 1	2.856	1.4056E-06	45.635
Ubiquinone and other terpenoid-quinone biosynthesis				
A0A1U8FWD5	Putative NAD(P)H dehydrogenase (Quinone) FQR1-like 1	1.719	0.0124045	21.674
A0A1U8EYM1	Tyrosine aminotransferase	2.399	0.00038372	47.177
Sesquiterpenoid and triterpenoid biosynthesis				
A0A1U8HFR8	Vetispiradiene synthase 1	6.712	1.67142E-05	64.165
A0A1U8EWI6	Uncharacterized protein	3.689	1.67363E-05	56.854
Thiamine metabolism				
A0A2G3ALC4	1-deoxy-D-xylulose-5-phosphate synthase, chloroplastic	0.607	0.000102066	76.896
A0A1U8FC73	Thiamine thiazole synthase, chloroplastic	0.548	0.000136529	38.071
A0A2G3A131	Adenylate kinase 4	0.604	4.9351E-06	26.487
A0A2G2Z8I0	Phosphomethylpyrimidine synthase	0.612	0.00125775	70.093



**Figure 7** Domain enrichment analysis of the DEPs in pepper after Se stress treatment. (A) Protein domain enrichment bubble plot of DEPs. (B) The accumulation of HSP proteins after Se stress treatment. Full-size [DOI: 10.7717/peerj.8020/fig-7](https://doi.org/10.7717/peerj.8020/fig-7)

we conducted a TMT-based quantitative proteomic analysis of the responses of pepper shoots to Se stress. Many DEPs were identified, and a group of proteins potentially involved in Se stress responses were identified. The Se-responsive DEPs and the associated metabolic pathways may play critical roles in Se stress signaling and responses in pepper.

Se is chemically similar to sulfur and is absorbed by plants through similar metabolic pathways (Van Hoewyk et al., 2008; Cakir, Turgut-Kara & Ari, 2016). Most plants non-specifically absorb Se from the environment through sulfate transporters and assimilate Se into organic forms of Se through the S metabolic pathway (Cappa et al., 2014). To date, studies on Se-tolerance mechanisms have focused on some Se-hyperaccumulating plants (Freeman et al., 2010; Sabbagh & Van Hoewyk, 2012; Cappa et al., 2015). In *Cardamine hupingshanensis* seedlings, the expression of the *sulfite oxidase* (SOX) gene in the roots is up-regulated after the addition of selenite, thus indicating that selenite may first be converted to selenate, and then the selenate is metabolized (Zhou et al., 2018).

Plants have evolved several efficient and complex strategies for dealing with different abiotic stresses, including Se stress ([Chen et al., 2002](#); [Lyons, Stangoulis & Graham, 2004](#); [Pérez-Clemente et al., 2013](#); [Schiavon et al., 2013](#)). GO analysis revealed that 14 proteins were associated with “response to stimulus,” including 11 up- and three down-regulated proteins significantly altered by Se stress in pepper. Among the proteins, a peroxidase (A0A1U8DW23) and a glutathione peroxidase (A0A2G3AE16) were up-regulated over 1.5-fold by Se treatment, thus suggesting that reactive oxygen species accumulate after Se treatment. In the present study, the expression of two pathogenesis-related proteins (STH-21, A0A1U8GEH7; PR4, I6VW44) significantly decreased. Pathogenesis-related proteins are a class of stress-tolerant proteins that are promising tools for plant genetic engineering ([Ali et al., 2018](#)). KOG analysis revealed that 36 proteins were related to the term “posttranslational modification, protein turnover, chaperones.” The identified DEPs may be involved in plant responses to Se stress. In pepper, nine “amino acid transport and metabolism”-related proteins and 10 “secondary metabolites biosynthesis, transport and catabolism”-associated proteins were identified. The changes in the DEPs suggested that Se stress may influence metabolite content in pepper.

Plants are thought to produce specific metabolites in response to Se stress ([Fernandes et al., 2018](#); [Pilon-Smits et al., 2009](#)). Transcriptomic analysis of Se-treated *Arabidopsis thaliana* has revealed that sulfur content decreases while the expression of sulfur absorption and metabolism genes increases, and the signaling pathways for ethylene and jasmonic acid respond to Se stress ([Van Hoewyk et al., 2008](#)). Small RNA and degradome sequencing analysis of *Astragalus chyrsochlorus* callus has revealed that miR167a, miR319, miR1507a, miR4346, miR7767-3p, miR7800, miR9748, and miR-n93 target transcription factors, disease resistance proteins, cysteine synthase, plant hormone signal transduction, plant-pathogen interaction, and sulfur metabolism pathways in response to Se stimuli ([Cakir, Candar-Cakir & Zhang, 2016](#)). In the present study, KEGG enrichment analysis revealed that seven pathways, including the sesquiterpenoid and triterpenoid biosynthesis pathway, plant-pathogen interaction, and thiamine metabolism pathway, were enriched after Se treatment. In addition, 32 DEPs involved in nine metabolic pathways were identified. Nine up-regulated DEPs were associated with cysteine and methionine metabolism. The main cause of plant Se poisoning is thought to be the incorrect synthesis of selenomethionine and selenocysteine (SeCys) into proteins, thereby causing changes or instability in the structures of proteins ([Sabbagh & Van Hoewyk, 2012](#); [Van Hoewyk, 2013](#)). High concentrations of SeCys in cells would lead to Se poisoning, and SeCys transformation is a direct means of Se detoxification ([Pilon et al., 2003](#); [Tamaoki, Freeman & Pilon-Smits, 2008](#); [Sabbagh & Van Hoewyk, 2012](#)). Terpenoids, the most diverse class of chemicals produced by plants, are involved in protection against various abiotic factors ([Lange, 2015](#); [Tholl, 2015](#)). For example, terpenoids are considered to be important defensive metabolites in *Eucalyptus froggattii* seedlings ([Goodger, Heskes & Woodrow, 2013](#)). Four upregulated DEPs were associated with “ubiquinone and other terpenoid-quinone biosynthesis” and “sesquiterpenoid and triterpenoid biosynthesis,” thus indicating that Se stress influenced the accumulation of some terpenoids.

Heat shock proteins were initially defined as proteins rapidly up-regulated by heat stress (Hartl & Hayer-Hartl, 2002). Studies increasingly show that HSP concentrations in plants increase rapidly as environmental conditions deteriorate (Murakami et al., 2004; Hu, Hu & Han, 2009; Lee, Yun & Kwon, 2012). HSPs are a class of evolutionarily conserved proteins that can be divided into five families according to molecular weight and sequence homology: small HSPs (molecular weight from 15 to 42 kDa), HSP60, HSP70, HSP90, and HSP100 (Boston, Viitanen & Vierling, 1996; Wang et al., 2004; Waters, 2013). HSP90 and HSP70 are essential for plant resistance to pathogen infections (Kanzaki et al., 2003; Noël et al., 2007; Chen et al., 2010). AtHSP17.6A is induced by osmotic stress, and PtHSP17.8 is involved in tolerance to heat and salt stress (Sun et al., 2001; Li et al., 2016). Among the Se-stress-induced DEPs in pepper, several HSPs were identified. A total of 14 HSP20, four HSP40, and five HSP70 proteins were significantly upregulated by salt stress, thus suggesting molecular cross-talk between heat shock responses and Se stress.

## CONCLUSIONS

A TMT-based proteomic method was used to investigate changes in protein levels between control and Se treated pepper seedlings. In total, 4,693 proteins and 200 DEPs were identified. A number of DEPs were found to be mainly involved in responses to stress and metabolic processes. Our results provide basic tools for identifying candidate proteins and the molecular mechanisms of the Se stress response in pepper plants.

## ACKNOWLEDGEMENTS

We are grateful to the PTM Biolabs company for technical support. We thank International Science Editing for editing this manuscript.

## ADDITIONAL INFORMATION AND DECLARATIONS

### Funding

This work was funded by the Natural Science Foundation of Zhejiang Province, China, Grant No. LQ17C150003. The National Natural Science Foundation of China, Grant No. 31701967, the key research and development program of Zhejiang Province, China, Grant No. 2017C02018 and the key research and development program of Hangzhou, China, Grant No. 20180416A07. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

### Grant Disclosures

The following grant information was disclosed by the authors:

Natural Science Foundation of Zhejiang Province: LQ17C150003.

National Natural Science Foundation of China: 31701967.

Key research and development program of Zhejiang Province, China: 2017C02018.

Key research and development program of Hangzhou, China: 20180416A07.

### Competing Interests

The authors declare that they have no competing interests.

## Author Contributions

- Chenghao Zhang conceived and designed the experiments, analyzed the data, contributed reagents/materials/analysis tools, prepared figures and/or tables, approved the final draft.
- Baoyu Xu performed the experiments, contributed reagents/materials/analysis tools, prepared figures and/or tables, approved the final draft.
- Wei Geng performed the experiments, contributed reagents/materials/analysis tools, prepared figures and/or tables, approved the final draft.
- Yunde Shen performed the experiments, contributed reagents/materials/analysis tools, prepared figures and/or tables, approved the final draft.
- Dongji Xuan analyzed the data, authored or reviewed drafts of the paper, approved the final draft.
- Qixian Lai analyzed the data, authored or reviewed drafts of the paper, approved the final draft.
- Chenjia Shen analyzed the data, authored or reviewed drafts of the paper, approved the final draft.
- Chengwu Jin conceived and designed the experiments, analyzed the data, contributed reagents/materials/analysis tools, prepared figures and/or tables, approved the final draft.
- Chenliang Yu conceived and designed the experiments, contributed reagents/materials/analysis tools, prepared figures and/or tables, approved the final draft.

## Data Availability

The following information was supplied regarding data availability:

The raw data is available at ProteomeXchange: [PXD013257](https://proteomecentral.proteomexchange.org/protein/PXD013257).

## Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.8020#supplemental-information>.

## REFERENCES

- Aggarwal M, Sharma S, Kaur N, Pathania D, Bhandhari K, Kaushal N, Kaur R, Singh K, Srivastava A, Nayyar H. 2011. Exogenous proline application reduces phytotoxic effects of selenium by minimising oxidative stress and improves growth in bean (*Phaseolus vulgaris* L.) seedlings. *Biological Trace Element Research* **140**(3):354–367 DOI [10.1007/s12011-010-8699-9](https://doi.org/10.1007/s12011-010-8699-9).
- Ali S, Ganai BA, Kamili AN, Bhat AA, Mir ZA, Bhat JA, Tyagi A, Islam ST, Mushtaq M, Yadav P, Rawat S, Grover A. 2018. Pathogenesis-related proteins and peptides as promising tools for engineering plants with multiple stress tolerance. *Microbiological Research* **212–213**:29–37 DOI [10.1016/j.micres.2018.04.008](https://doi.org/10.1016/j.micres.2018.04.008).
- Boston RS, Viitanen PV, Vierling E. 1996. Molecular chaperones and protein folding in plants. *Plant Molecular Biology* **32**(1–2):191–222 DOI [10.1007/BF00039383](https://doi.org/10.1007/BF00039383).
- Cakir O, Candar-Cakir B, Zhang B. 2016. Small RNA and degradome sequencing reveals important microRNA function in *Astragalus chrysochlorus* response to selenium stimuli. *Plant Biotechnology Journal* **14**(2):543–556 DOI [10.1111/pbi.12397](https://doi.org/10.1111/pbi.12397).



- Cakir O, Turgut-Kara N, Ari S. 2016.** Selenium induced selenocysteine methyltransferase gene expression and antioxidant enzyme activities in *Astragalus chrysochlorus*. *Acta Botanica Croatica* 75(1):11–16 DOI 10.1515/botcro-2016-0015.
- Cappa JJ, Cappa PJ, El Mehdawi AF, McAleer JM, Simmons MP, Pilon-Smits EAH. 2014.** Characterization of selenium and sulfur accumulation across the genus *Stanleya* (Brassicaceae): a field survey and common-garden experiment. *American Journal of Botany* 101(5):830–839 DOI 10.3732/ajb.1400041.
- Cappa JJ, Yetter C, Fakra S, Cappa PJ, DeTar R, Landes C, Pilon-Smits EAH, Simmons MP. 2015.** Evolution of selenium hyperaccumulation in *Stanleya* (Brassicaceae) as inferred from phylogeny, physiology and X-ray microprobe analysis. *New Phytologist* 205(2):583–595 DOI 10.1111/nph.13071.
- Chen L, Hamada S, Fujiwara M, Zhu T, Thao NP, Wong HL, Krishna P, Ueda T, Kaku H, Shibuya N, Kawasaki T, Shimamoto K. 2010.** The Hop/Sti1-Hsp90 chaperone complex facilitates the maturation and transport of a PAMP receptor in rice innate immunity. *Cell Host & Microbe* 7(3):185–196 DOI 10.1016/j.chom.2010.02.008.
- Chen L, Yang F, Xu J, Hu Y, Hu Q, Zhang Y, Pan G. 2002.** Determination of selenium concentration of rice in China and effect of fertilization of selenite and selenate on selenium content of rice. *Journal of Agricultural and Food Chemistry* 50(18):5128–5130 DOI 10.1021/jf0201374.
- Choi GS, Kim JH, Lee DH, Kim JS, Ryu KH. 2005.** Occurrence and distribution of viruses infecting pepper in Korea. *Plant Pathology Journal* 21(3):258–261 DOI 10.5423/PPJ.2005.21.3.258.
- Chu J, Yao X, Zhang Z. 2010.** Responses of wheat seedlings to exogenous selenium supply under cold stress. *Biological Trace Element Research* 136(3):355–363 DOI 10.1007/s12011-009-8542-3.
- Djanaguiraman M, Prasad PVV, Seppanen M. 2010.** Selenium protects sorghum leaves from oxidative damage under high temperature stress by enhancing antioxidant defense system. *Plant Physiology and Biochemistry* 48(12):999–1007 DOI 10.1016/j.plaphy.2010.09.009.
- El Kassis E, Cathala N, Rouached H, Fourcroy P, Berthomieu P, Terry N, Davidian JC. 2007.** Characterization of a selenate-resistant *Arabidopsis* mutant. Root growth as a potential target for selenate toxicity. *Plant Physiology* 143(3):1231–1241 DOI 10.1104/pp.106.091462.
- El-Ramady HR, Domokos-Szabolcsy É, Shalaby TA, Prokisch J, Fári M. 2015.** Selenium in agriculture: water, air, soil, plants, food, animals and nanoselenium. In: Lichtfouse E, ed. *CO2 Sequestration, Biofuels and Depollution*. Berlin: Springer International Publishing, 153–232.
- Elrashidi MA, Adriano DC, Workman SM, Lindsay WL. 1987.** Chemical-equilibria of selenium in soils - a theoretical development. *Soil Science* 144(2):141–152 DOI 10.1097/00010694-198708000-00008.
- Fernandes J, Hu X, Smith MR, Go YM, Jones DP. 2018.** Selenium at the redox interface of the genome, metabolome and exposome. *Free Radical Biology and Medicine* 127:215–227 DOI 10.1016/j.freeradbiomed.2018.06.002.
- Freeman JL, Tamaoki M, Stushnoff C, Quinn CF, Cappa JJ, Devonshire J, Fakra SC, Marcus MA, McGrath SP, Van Hoewyk D, Pilon-Smits EAH. 2010.** Molecular mechanisms of selenium tolerance and hyperaccumulation in *Stanleya pinnata*. *Plant Physiology* 153(4):1630–1652 DOI 10.1104/pp.110.156570.
- Galinha C, Sánchez-Martínez MJ, Pacheco AMG, Freitas MC, Coutinho Jé, Maças B, Almeida AS, Pérez-Corona MT, Madrid Y, Wolterbeek HT. 2015.** Characterization of selenium-enriched wheat by agronomic biofortification. *Journal of Food Science and Technology* 52(7):4236–4245 DOI 10.1007/s13197-014-1503-7.

- Goodger JQD, Heskes AM, Woodrow IE. 2013.** Contrasting ontogenetic trajectories for phenolic and terpenoid defences in *Eucalyptus froggattii*. *Annals of Botany* **112**(4):651–659 DOI [10.1093/aob/mct010](https://doi.org/10.1093/aob/mct010).
- Hao J, Guo H, Shi X, Wang Y, Wan QH, Song YB, Zhang L, Dong M, Shen CJ. 2017.** Comparative proteomic analyses of two *Taxus* species (*Taxus X media* and *Taxus mairei*) reveals variations in the metabolisms associated with paclitaxel and other metabolites. *Plant and Cell Physiology* **58**(11):1878–1890 DOI [10.1093/pcp/pcx128](https://doi.org/10.1093/pcp/pcx128).
- Hartikainen H, Ekholm P, Piironen V, Xue T, Koivu T, Yli-Halla M. 1997.** Quality of the ryegrass and lettuce yields as affected by selenium fertilization. *Agricultural and Food Science* **6**(5–6):381–387 DOI [10.23986/afsci.72801](https://doi.org/10.23986/afsci.72801).
- Hartl FU, Hayer-Hartl M. 2002.** Molecular chaperones in the cytosol: from nascent chain to folded protein. *Science* **295**(5561):1852–1858 DOI [10.1126/science.1068408](https://doi.org/10.1126/science.1068408).
- Hladun KR, Parker DR, Tran KD, Trumble JT. 2013.** Effects of selenium accumulation on phytotoxicity, herbivory, and pollination ecology in radish (*Raphanus sativus* L.). *Environmental Pollution* **172**:70–75 DOI [10.1016/j.envpol.2012.08.009](https://doi.org/10.1016/j.envpol.2012.08.009).
- Hu W, Hu G, Han B. 2009.** Genome-wide survey and expression profiling of heat shock proteins and heat shock factors revealed overlapped and stress specific response under abiotic stresses in rice. *Plant Science* **176**(4):583–590 DOI [10.1016/j.plantsci.2009.01.016](https://doi.org/10.1016/j.plantsci.2009.01.016).
- Jia M, Zhang Y, Huang B, Zhang H. 2019.** Source apportionment of selenium and influence factors on its bioavailability in intensively managed greenhouse soil: a case study in the east bank of the Dianchi Lake, China. *Ecotoxicology and Environmental Safety* **170**:238–245 DOI [10.1016/j.ecoenv.2018.11.133](https://doi.org/10.1016/j.ecoenv.2018.11.133).
- Kanzaki H, Saitoh H, Ito A, Fujisawa S, Kamoun S, Katou S, Yoshioka H, Terauchi R. 2003.** Cytosolic HSP90 and HSP70 are essential components of INF1-mediated hypersensitive response and non-host resistance to *Pseudomonas cichorii* in *Nicotiana benthamiana*. *Molecular Plant Pathology* **4**(5):383–391 DOI [10.1046/j.1364-3703.2003.00186.x](https://doi.org/10.1046/j.1364-3703.2003.00186.x).
- Kong L, Wang M, Bi D. 2005.** Selenium modulates the activities of antioxidant enzymes, osmotic homeostasis and promotes the growth of sorrel seedlings under salt stress. *Plant Growth Regulation* **45**(2):155–163 DOI [10.1007/s10725-005-1893-7](https://doi.org/10.1007/s10725-005-1893-7).
- Kumar M, Bijo AJ, Baghel RS, Reddy CRK, Jha B. 2012.** Selenium and spermine alleviate cadmium induced toxicity in the red seaweed *Gracilaria dura* by regulating antioxidants and DNA methylation. *Plant Physiology and Biochemistry* **51**:129–138 DOI [10.1016/j.plaphy.2011.10.016](https://doi.org/10.1016/j.plaphy.2011.10.016).
- Kuppusamy S, Yoon YE, Song YK, Kim JH, Kim HT, Yong BL. 2018.** Does long-term application of fertilizers enhance the micronutrient density in soil and crop?—evidence from a field trial conducted on a 47-year-old rice paddy. *Journal of Soils and Sediments* **18**(1):49–62 DOI [10.1007/s11368-017-1743-z](https://doi.org/10.1007/s11368-017-1743-z).
- Lange BM. 2015.** The evolution of plant secretory structures and emergence of terpenoid chemical diversity. *Annual Review of Plant Biology* **66**(1):139–159 DOI [10.1146/annurev-arplant-043014-114639](https://doi.org/10.1146/annurev-arplant-043014-114639).
- Lee J-H, Yun HS, Kwon C. 2012.** Molecular communications between plant heat shock responses and disease resistance. *Molecules and Cells* **34**(2):109–116 DOI [10.1007/s10059-012-0121-3](https://doi.org/10.1007/s10059-012-0121-3).
- Li J, Zhang J, Jia H, Li Y, Xu X, Wang L, Lu M. 2016.** The *Populus trichocarpa* PtHSP17.8 involved in heat and salt stress tolerances. *Plant Cell Reports* **35**(8):1587–1599 DOI [10.1007/s00299-016-1973-3](https://doi.org/10.1007/s00299-016-1973-3).
- Lyons GH, Stangoulis JCR, Graham RD. 2004.** Exploiting micronutrient interaction to optimize biofortification programs: the case for inclusion of selenium and iodine in the HarvestPlus program. *Nutrition Reviews* **62**(6):247–252 DOI [10.1111/j.1753-4887.2004.tb00047.x](https://doi.org/10.1111/j.1753-4887.2004.tb00047.x).

- Molnárová M, Fargašová A. 2009. Se (IV) phytotoxicity for monocotyledonae cereals (*Hordeum vulgare* L., *Triticum aestivum* L.) and dicotyledonae crops (*Sinapis alba* L., *Brassica napus* L.). *Journal of Hazardous Materials* 172(2–3):854–861 DOI 10.1016/j.jhazmat.2009.07.096.
- Murakami T, Matsuba S, Funatsuki H, Kawaguchi K, Saruyama H, Tanida M, Sato Y. 2004. Over-expression of a small heat shock protein, sHSP17.7, confers both heat tolerance and UV-B resistance to rice plants. *Molecular Breeding* 13(2):165–175 DOI 10.1023/b:molb.0000018764.30795.c1.
- Noël LD, Cagna G, Stuttmann J, Wirthmüller L, Betsuyaku S, Witte C-P, Bhat R, Pochon N, Colby T, Parker JE. 2007. Interaction between SGT1 and cytosolic/nuclear HSC70 chaperones regulates *Arabidopsis* immune responses. *Plant Cell* 19(12):4061–4076 DOI 10.1105/tpc.107.051896.
- Paciolla C, De Leonardis S, Dipierro S. 2011. Effects of selenite and selenate on the antioxidant systems in *Senecio scandens* L. *Plant Biosystems - An International Journal Dealing with All Aspects of Plant Biology* 145(1):253–259 DOI 10.1080/11263504.2010.509942.
- Pilon M, Owen JD, Garifullina GF, Kurihara T, Mihara H, Esaki N, Pilon-Smits EAH. 2003. Enhanced selenium tolerance and accumulation in transgenic *Arabidopsis* expressing a mouse selenocysteine lyase. *Plant Physiology* 131(3):1250–1257 DOI 10.1104/pp.102.014639.
- Pilon-Smits EAH, Quinn CF, Tapken W, Malagoli M, Schiavon M. 2009. Physiological functions of beneficial elements. *Current Opinion in Plant Biology* 12(3):267–274 DOI 10.1016/j.pbi.2009.04.009.
- Pukacka S, Ratajczak E, Kalembe E. 2011. The protective role of selenium in recalcitrant *Acer saccharium* L. seeds subjected to desiccation. *Journal of Plant Physiology* 168(3):220–225 DOI 10.1016/j.jplph.2010.07.021.
- Pérez-Clemente RM, Vives V, Zandalinas SI, López-Climent MF, Muñoz V, Gómez-Cadenas A. 2013. Biotechnological approaches to study plant responses to stress. *BioMed Research International* 2013(12):1–10 DOI 10.1155/2013/654120.
- Quinn CF, Freeman JL, Galeas ML, Klamper EM, Pilon-Smits EAH. 2008. The role of selenium in protecting plants against prairie dog herbivory: implications for the evolution of selenium hyperaccumulation. *Oecologia* 155(2):267–275 DOI 10.1007/s00442-007-0907-8.
- Sabbagh M, Van Hoewyk D. 2012. Malformed selenoproteins are removed by the ubiquitin–proteasome pathway in *Stanleya pinnata*. *Plant and Cell Physiology* 53(3):555–564 DOI 10.1093/pcp/pcs015.
- Sager M. 2006. Selenium in agriculture, food, and nutrition. *Pure and Applied Chemistry* 78(1):111–133 DOI 10.1351/pac200678010111.
- Schiavon M, Dall’Acqua S, Mietto A, Pilon-Smits EAH, Sambo P, Masi A, Malagoli M. 2013. Selenium fertilization alters the chemical composition and antioxidant constituents of tomato (*Solanum lycopersicon* L.). *Journal of Agricultural and Food Chemistry* 61(44):10542–10554 DOI 10.1021/jf4031822.
- Semnani S, Roshandel G, Zendeabad A, Keshtkar A, Rahimzadeh H, Abdolahi N, Besharat S, Moradi A, Mirkarimi H, Hasheminasab S. 2010. Soils selenium level and esophageal cancer: an ecological study in a high risk area for esophageal cancer. *Journal of Trace Elements in Medicine and Biology* 24(3):174–177 DOI 10.1016/j.jtemb.2010.03.002.
- Shibagaki N, Rose A, McDermott JP, Fujiwara T, Hayashi H, Yoneyama T, Davies JP. 2002. Selenate-resistant mutants of *Arabidopsis thaliana* identify *Sultr1;2*, a sulfate transporter required for efficient transport of sulfate into roots. *Plant Journal* 29(4):475–486 DOI 10.1046/j.0960-7412.2001.01232.x.
- Sinha I, Karagoz K, Fogle RL, Hollenbeak CS, Zea AH, Arga KY, Stanley AE, Hawkes WC, Sinha R. 2016. “Omics” of selenium biology: a prospective study of plasma proteome network

- before and after selenized-yeast supplementation in healthy men. *OMICS: A Journal of Integrative Biology* **20**(4):202–213 DOI [10.1089/omi.2015.0187](https://doi.org/10.1089/omi.2015.0187).
- Song Z, Shao H, Huang H, Shen Y, Wang L, Wu F, Han D, Song J, Jia H. 2017.** Overexpression of the phosphate transporter gene OsPT8 improves the Pi and selenium contents in *Nicotiana tabacum*. *Environmental and Experimental Botany* **137**:158–165 DOI [10.1016/j.envexpbot.2017.02.011](https://doi.org/10.1016/j.envexpbot.2017.02.011).
- Sun W, Bernard C, Van De Cotte B, Van Montagu M, Verbruggen N. 2001.** At-HSP17.6A, encoding a small heat-shock protein in *Arabidopsis*, can enhance osmotolerance upon overexpression. *Plant Journal* **27**(5):407–415 DOI [10.1046/j.1365-3113X.2001.01107.x](https://doi.org/10.1046/j.1365-3113X.2001.01107.x).
- Tamaoki M, Freeman JL, Pilon-Smits EAH. 2008.** Cooperative ethylene and jasmonic acid signaling regulates selenite resistance in *Arabidopsis*. *Plant Physiology* **146**(3):1219–1230 DOI [10.1104/pp.107.110742](https://doi.org/10.1104/pp.107.110742).
- Thavarajah D, Ruszkowski J, Vandenberg A. 2008.** High potential for selenium biofortification of lentils (*Lens culinaris* L.). *Journal of Agricultural and Food Chemistry* **56**(22):10747–10753 DOI [10.1021/jf802307h](https://doi.org/10.1021/jf802307h).
- Tholl D. 2015.** Biosynthesis and biological functions of terpenoids in plants. *Advances in Biochemical Engineering/Biotechnology* **148**:63–106 DOI [10.1007/10\\_2014\\_295](https://doi.org/10.1007/10_2014_295).
- Van Hoewyk D. 2013.** A tale of two toxicities: malformed selenoproteins and oxidative stress both contribute to selenium stress in plants. *Annals of Botany* **112**(6):965–972 DOI [10.1093/aob/mct163](https://doi.org/10.1093/aob/mct163).
- Van Hoewyk D, Takahashi H, Inoue E, Hess A, Tamaoki M, Pilon-Smits EA. 2008.** Transcriptome analyses give insights into selenium-stress responses and selenium tolerance mechanisms in *Arabidopsis*. *Physiologia Plantarum* **132**(2):236–253 DOI [10.1111/j.1399-3054.2007.01002.x](https://doi.org/10.1111/j.1399-3054.2007.01002.x).
- Wang C-Q. 2011.** Water-stress mitigation by selenium in *Trifolium repens* L. *Journal of Plant Nutrition and Soil Science* **174**(2):276–282 DOI [10.1002/jpln.200900011](https://doi.org/10.1002/jpln.200900011).
- Wang W, Vinocur B, Shoseyov O, Altman A. 2004.** Role of plant heat-shock proteins and molecular chaperones in the abiotic stress response. *Trends in Plant Science* **9**(5):244–252 DOI [10.1016/j.tplants.2004.03.006](https://doi.org/10.1016/j.tplants.2004.03.006).
- Waters ER. 2013.** The evolution, function, structure, and expression of the plant sHSPs. *Journal of Experimental Botany* **64**(2):391–403 DOI [10.1093/jxb/ers355](https://doi.org/10.1093/jxb/ers355).
- White PJ. 2016.** Selenium accumulation by plants. *Annals of Botany* **117**:217–235.
- Xu DB, Yuan HW, Tong YF, Zhao L, Qiu LL, Guo WB, Shen CJ, Liu HJ, Yan DL, Zheng BS. 2017.** Comparative proteomic analysis of the graft unions in Hickory (*Carya cathayensis*) provides insights into response mechanisms to grafting process. *Frontiers in Plant Science* **8**:676 DOI [10.3389/fpls.2017.00676](https://doi.org/10.3389/fpls.2017.00676).
- Zhang L, Hu B, Li W, Che R, Deng K, Li H, Yu F, Ling H, Li Y, Chu C. 2014.** OsPT2, a phosphate transporter, is involved in the active uptake of selenite in rice. *New Phytologist* **201**(4):1183–1191 DOI [10.1111/nph.12596](https://doi.org/10.1111/nph.12596).
- Zhang J, Wang L, Anderson LB, Witthuhn B, Xu Y, Lu J. 2010.** Proteomic profiling of potential molecular targets of methyl-selenium compounds in the transgenic adenocarcinoma of mouse prostate model. *Cancer Prevention Research* **3**(8):994–1006 DOI [10.1158/1940-6207.CAPR-09-0261](https://doi.org/10.1158/1940-6207.CAPR-09-0261).
- Zhou Y, Tang Q, Wu M, Mou D, Liu H, Wang S, Zhang C, Ding L, Luo J. 2018.** Comparative transcriptomics provides novel insights into the mechanisms of selenium tolerance in the hyperaccumulator plant *Cardamine hupingshanensis*. *Scientific Reports* **8**(1):2789 DOI [10.1038/s41598-018-21268-2](https://doi.org/10.1038/s41598-018-21268-2).