

Wild plant species growing closely connected in a subalpine meadow host distinct root-associated bacterial communities.

Kristin Aleklett, Jonathan W Leff, Noah Fierer, Miranda Hart

Plant roots are known to harbor large and diverse communities of bacteria. It has been suggested that plant identity can structure these root-associated communities, but few studies have specifically assessed how the composition of root microbiota varies within and between plant species growing under natural conditions. We assessed the community composition of endophytic and epiphytic bacteria through high throughput sequencing using 16S rDNA derived from root tissues collected from a population of a wild, clonal plant (Orange hawkweed – *Pilosella aurantiaca*) as well as two neighboring plant species (Oxeye daisy – *Leucanthemum vulgare* and Alsike clover – *Trifolium hybridum*).

Our first goal was to determine if plant species growing in close proximity, under similar environmental conditions, still hosted unique root microbiota. Our results showed that plants of different species host distinct bacterial communities in their roots. In terms of community composition, *Betaproteobacteria* (especially the family *Oxalobacteraceae*) were found to dominate in the root microbiota of *L. vulgare* and *T. hybridum* samples, whereas the root microbiota of *P. aurantiaca* had a more heterogeneous distribution of bacterial abundances where *Gammaproteobacteria* and *Acidobacteria* occupied a larger portion of the community.

We also explored the extent of individual variance within each plant species investigated, and found that in the plant species thought to have the least genetic variance among individuals (*P.aurantiaca*) still hosted just as diverse microbial communities. Whether all plant species host their own distinct root microbiota and plants more closely related to each other share more similar bacterial communities still remains to be fully explored, but among the plants examined in this experiment there was no trend that the two species belonging to the same family shared more similarities in terms of bacterial community composition.

1 Kristin Aleklett¹, Jonathan W. Leff², Noah Fierer², Miranda Hart^{1*}

2 ¹*Department of Biology, University of British Columbia – Okanagan CANADA, ²Cooperative*

3 *Institute for Research in Environmental Sciences and the Department of Ecology and*

4 *Evolutionary Biology, University of Colorado, Boulder, Colorado, USA*

5 **Corresponding author: Miranda Hart, miranda.hart@ubc.ca, +1-250-807-9398*

6 Introduction

Plant roots function as distinct habitats within the soil and bacterial communities in root systems have repeatedly been shown to differ from those of the surrounding bulk soil (Smalla et al. 2001; Haichar et al. 2008; Gottel et al. 2011; Lundberg et al. 2012). Even though root associated bacterial communities (both *rhizospheric* – in the soil surrounding the roots, *epiphytic* – living at the surface of roots and *endophytic* – living inside root tissues) have been under investigation for many years, there is still little consensus in how these communities are formed and what determines their composition (Berg and Smalla 2009; Aleklett and Hart 2013; Bulgarelli et al. 2013).

Traditionally, the composition of bacterial communities living in association with plants has been attributed to environmental factors. For example, soil type has been suggested as the strongest determinant of community structure in root associated microbial communities (De Ridder-Duine et al. 2005; Singh et al. 2007; Lundberg et al. 2012; Bulgarelli et al. 2013). At the same time, it has also been argued that the host plant may play an equally large role in determining the composition of its root microbiota (Marschner et al. 2005; Costa et al. 2006; Hartmann et al. 2008; Doornbos et al. 2011), especially endophytic bacterial communities (Haichar et al. 2008).

Recent work has demonstrated that hosts can alter their root microbiota by regulating soil conditions in the vicinity of the root system through root exudation of sugars, phenolics and amino acids that could also function as signaling molecules with the microbes in the surrounding soil (Chaparro et al. 2013). Since root exudation patterns and composition can be associated with plant gene expression, variation in host genetics has the potential to create large differences in the chemical profile of plants and consequently the composition of microbes able to inhabit the root system. Several studies have found that different plant species or genotypes of the same species host distinct microbial communities (Bailey et al. 2005; Marschner et al. 2005; van Overbeek and van Elsas 2008; Schweitzer et al. 2008; Micallef et al. 2009a; Manter et al. 2010; Becklin et al.

2012; Peiffer et al. 2013). Even studies where soil type was considered to have the strongest effect on structuring the root microbiota, differences in bacterial community composition between genotypes was still detected (Bulgarelli et al. 2012; Lundberg et al. 2012).

The root environment varies greatly among plant species (Bardgett et al. 2014); these differences may lead to the selection of distinct bacterial communities. Plants can differ in terms of both root lifespan (Roumet et al. 2006), root architecture (Hodge et al. 2009), root surface structure and components and patterns of root exudation (Bais et al. 2006). Root exudates are known to provide a food source for the microbes (Farrar et al. 2003), instigators of symbiotic associations (such as mycorrhizal infection or nodule formation) (Bais et al. 2004), and defend the plant against pathogens (Doornbos et al. 2011). All these plant characteristics could contribute to shaping root systems of different plants into local habitats and potentially distinct niches for microbial colonizers.

The role of intra-species variation among root associated microbial communities has been overlooked, but might represent a significant proportion of variation in natural systems (Bell et al. 2013). Since we know that natural populations exhibit variation in root exudation patterns and root morphology, one would expect there to be variation among individual plants in their root microbiota as well (Micallef et al. 2009). But variation among plants might also be driven by environmental heterogeneity because we know that small-scale environmental heterogeneity exists in soil systems. Is this variation static across plant taxa, or do different taxa exhibit more variation than others? If plant genetics are determining bacterial community composition, then certainly, populations with low genetic diversity (i.e. asexually reproducing, or metapopulations), would be expected to have less variation than sexually reproducing populations with high levels of gene flow. Because we sampled a plant species known to reproduce clonally through stolons and apomixis (*P. aurantiaca*), we also examined whether individuals within that species had less

dispersion in their microbial community composition than individuals within the other two plant species.

The majority of studies characterizing bacterial communities in the root microbiota have been conducted with model plants in artificial greenhouse settings or agricultural contexts (Marschner and Yang 2001; Garbeva et al. 2004; Micallef et al. 2009; Manter et al. 2010; Doornbos et al. 2011; Lundberg et al. 2012) where the study of genetically modified plants have been especially informing when it comes to understanding slight differences between plant genotypes (van Overbeek and van Elsas 2008; Weinert et al. 2009; Inceoğlu et al. 2010). While these studies are crucial for understanding the mechanistic basis of plant:microbe interactions, they do not reflect how natural environmental conditions contribute to variation in bacterial community composition across individual plants, particularly in complex environments where a wide diversity of plants and biota are interacting.

In this study, we explored variation in bacterial community composition between individual root systems of neighboring plants in a common field in order to determine how much variation exists within and between plant taxa. We sampled the root microbiota of three plant species growing within 10 meters from each other in a field and asked – are bacterial root communities distinct among plant species growing in a common location, close enough to be exposed to the same environmental conditions? And – do certain plant species contain more intra-species variance in bacterial communities than others?

Methods

Field site and target plant

Field site description

Samples were collected in August, 2011, from a subalpine meadow near Chute Lake, British Columbia, Canada (49.698859N,-119.533133W). The sampling area has not been used for

agriculture or forestry but is in proximity to a forestry road as well as a camp site. Since it contains a high number of invasive plant species, it could also be considered a disturbed site. The soil at the site was determined to be a sandy loam, and the site is classified under the biogeoclimatic zone Interior Douglas Fir, dry warm (IDFdw) (Biogeoclimatic Ecosystem Classification (BEC) and Ecology Research program of the British Columbia). This zone is characterized by cold winters and dry, warm summers with low amounts of precipitation due to the rain shadow created by the Coast, Cascade and Columbia mountains. Because the climate is very dry in the summers, forest fires are a common feature in the landscape, and the site is located close to the area that was badly burnt during a large fire in Okanagan Mountain Park in 2003. The plant community sampled for this experiment was growing up on a rock plateau, away from the tree coverage, completely exposed to the elements (see Fig.1 in additional material). The dominant vegetation at the field site consisted of Orange hawkweed (*Pilosella aurantiaca* (L.) F.W. Schultz and Schultz-Bip), Hairy vetch (*Vicia villosa*, Roth), Oxeye daisy (*Leucanthemum vulgare*, Lam.), Wild strawberry (*Fragaria virginiana*, Duchesne) Timothy (*Phleum pratense*, L.) and Alsike clover (*Trifolium hybridum*, L.) growing homogenously, but at different abundances across the field.

Target plant

Our target plant was *P. aurantiaca* (formerly known as *Hieracium aurantiacum*), which is native to Europe and invasive in North America. Because of *P.aurantiaca*'s stoloniferous reproduction, the species is able to form dense mats in the landscape, making it hard for other plant species to compete for space through seed germination in the field (Giroday and Baker 2006). This is one of the features that have made the species a successful invasive plant within British Columbia (Giroday and Baker 2006). Genetic diversity within *P. aurantiaca* has previously been examined across 48 locations in North America, and results showed that there were only three genotypes, of

which two were found only in isolated locations (one in Alaska and one in Oregon) (Loomis and Fishman 2009). By choosing to work with a plant expressing this low diversity in wild populations, we hoped to minimize genetic variance within the population that we sampled.

To clarify the role of host identity and intra-species variance in bacterial root microbiota, we additionally sampled two of the co-occurring plant species, *L. vulgare* and *T. hybridum* that were in the same developmental stage (flowering) as *P. aurantiaca*.

Experimental design

Root systems of *P. aurantiaca* were collected one meter apart along two 10 m transects (n=20) in order to explore within species variance of a clonal plant. To be able to compare bacterial community composition between species, additional samples of *T. hybridum* (n=10) and *L. vulgare* (n=10) were also collected where present along the transects, several of which were growing within centimeters of *P. aurantiaca* samples. Because *T. hybridum* and *L. vulgare* were not as abundant within the field, their original sample size was lower. For statistical comparisons between species however, an equal sample size of 8 plants/species was used throughout analyses.

Each root system was rinsed from surrounding rhizospheric soil in de-ionized water in order to separate it from roots of neighboring plants. Root systems were then cut up in pieces and a subsample of root tissue, representative of the whole root system, including young fresh roots as well as older root tissues (with no exclusion of nodules in *T. hybridum*), was collected and further used for classification of bacterial community composition. Since no further treatment was performed in order to remove rhizoplane microbes, we assume that the communities extracted could be of either endophytic or epiphytic origin.

Bacterial community analysis

126 *Amplification and sequencing of target gene*

127 DNA from all collected plant tissues (0.25g/sample) was extracted using a PowerSoil DNA
 128 Isolation Kit (MoBio Laboratories Inc., USA) according to the manufacturer's protocol.
 129 Microbial diversity and the relative abundances of individual taxa were assessed by barcoded
 130 pyrosequencing of a portion of the 16S rRNA gene. Each DNA sample was amplified in triplicate
 131 through PCR reactions using the protocol described in Fierer et al. (2008) except with a different
 132 primer pair. The forward primer contained the 454 Life Sciences primer B sequence, the bacterial
 133 primer 799f (Chelius and Triplett 2001) and a two-base linker sequence ('AG'). The reverse
 134 primer contained the 454 Life Sciences primer A sequence, a unique 12 bp error-correcting Golay
 135 barcode (Fierer et al. 2008), a 'GT' linker sequence, and the 'universal' bacterial primer 1115r
 136 (Reysenbach and Pace 1995). The targeted gene region has shown to be appropriate for accurate
 137 taxonomic classification of bacterial sequences and the primers are designed to exclude
 138 chloroplasts from plant tissues in the samples (Redford et al. 2010). Amplicons were visualized
 139 via gel electrophoresis purified and quantified. Amplicons from all samples were then combined
 140 in equimolar ratios into a single tube. Samples were sequenced at Engencore (University of South
 141 Carolina) on a Roche GS-FLX sequencer running the Titanium chemistry.

142 *Processing raw sequence data*

143 All sequences were de-multiplexed and further analyzed using the Quantitative Insights Into
 144 Microbial Ecology (QIIME) toolkit (Caporaso et al. 2010). Operational Taxonomic Units (OTUs)
 145 were defined at the level of $\geq 97\%$ similarity and the taxonomy assigned by comparing a
 146 representative sequence from each OTU to the Greengenes database (February 14th 2012 version)
 147 (DeSantis et al. 2006) using the Basic Local Alignment Search Tool (BLAST) classifier (Altschul
 148 et al. 1990). In order to correct for differences in the number of sequences analyzed per sample, a
 149 randomly selected subset of 400 sequences per root sample was used to compare relative

differences in taxonomic diversity. Only samples from which we obtained a minimum of 400 bacterial sequences per sample or more were considered in the study, eliminating 3 samples from the study (one *P. aurantiaca* and two *L. vulgare*). Though 400 sequences cannot fully capture the rare biosphere, it allowed us to compare samples while still maintaining as many samples as possible. It has previously been shown that studies of bacterial communities show similar results even at a lower rarefaction (Hamady and Knight 2009; Kuczynski et al. 2010). In fact, re-analyzing our data set with a higher rarefaction limit showed the same general trends but drastically lowered our number of samples available to analyze.

Statistics

Differences in community composition between samples were calculated using phylogenetic metric (UniFrac) where weighted UniFrac shows an emphasis on the more abundant taxa in samples and un-weighted UniFrac treats all taxa the same (Lozupone et al. 2007; Hamady et al. 2010). As a comparison, we also included a taxonomic metric (Bray-Curtis distance) to explore whether dissimilarity patterns were the same in terms of presence/absence of taxa. Before calculating Bray-Curtis distances, all relative abundances were log-transformed. 2-D scatterplots of Principal Coordinates Analysis (PCoA) generated in PRIMER-E (Clarke and Gorley, 2006) were used to visualize the greatest amount of variability in the pair-wise distances between samples.

We tested for variance among host plants in their root microbiota using a 2-way PerMANOVA (Anderson 2005) with host species and transect as factors and weighted and un-weighted UniFrac as well as Bray Curtis as our dissimilarity metrics. All analyses were permuted 9999 times. Since it has been shown that PerMANOVA is not robust when sampling efforts are un-equal (Anderson and Walsh 2013), we subsampled 8 samples from each species which were used for both PerMANOVA and PermDISP analyses.

Variability in community composition within each of the three species was analyzed through PermDISP (Anderson 2004) (9999 permutations), creating a centroid for each species and measuring the average spread of samples belonging to that species from the centroid. A large spread (high average) would indicate a high variability in community composition among individuals within the species (Anderson 2004).

Results

Variation between host species

When comparing the phylogenetic overlap between bacterial root microbiota (UniFrac) across three different species of plant hosts growing in a common field, bacterial communities from samples of the same plant species were significantly more similar to each other than to bacterial communities sampled from plants of the two other species (Table 1). This was true for both weighted (Pseudo-F=8.54 $p=0.0001$) and un-weighted UniFrac (Pseudo-F=1.66 $p=0.0001$) as well as Bray Curtis dissimilarities (Pseudo-F=2.27 $p=0.0001$) (Table 1). These patterns were also evident from the principal coordinates analyses which showed little overlap between samples of different plant species (Fig.1).

Variation within host species

Plant species differed in how much variance there was among bacterial communities of individual root samples (Fig.1). There was a significant difference between plant species in the amount of compositional dissimilarity of bacterial taxa between individual plants (Bray Curtis: $F=9.56$ $p=0.02$). That is, the amount of dispersion of individual plants from the centroid differed among plant species. In this case, *P. aurantiaca* exhibited the most variance among plant individuals, while *T. hybridum* showed the least (Table 2). This was not the case when the same data was analyzed using phylogenetic measures, since we could not detect any significant

198 difference in dispersion among plant taxa (UniFrac: weighted $F=2.57$ $p=0.15$; un-weighted
199 $F=2.56$ $p=0.63$) (Table 2).

200 *Relative abundance of taxa across hosts*

201 A total number of 4384 unique OTUs were analyzed within the rarefied data set. A taxonomic
202 summary, showing the average abundance of bacterial phyla in *P. aurantiaca*, *T. hybridum* and *L.*
203 *vulgare* samples, illustrates the compositional differences between root systems of different plant
204 species (Fig.2). In *P. aurantiaca*, the most abundant phylum was *Betaproteobacteria* which made
205 up, on average, 29% out of all sequences found in *P. aurantiaca* samples, followed by
206 *Bacteroidetes* (19%), *Alphaproteobacteria* (16%) and *Actinobacteria* (12%). In *T. hybridum*,
207 *Betaproteobacteria* made up, on average, 51% of the all bacterial sequences found in the species,
208 followed by *Alphaproteobacteria* (21%) and *Bacteroidetes* (16%). Bacterial communities in *L.*
209 *vulgare* samples were, similarly to *T. hybridum*, dominated by *Betaproteobacteria* (50%),
210 followed by *Bacteroidetes* (18%) and *Alphaproteobacteria* (12%). A table showing the relative
211 distribution of the 14 most abundant OTUs across all samples as well as their relative abundance
212 within samples of the different species is given in Table 3.

213 A closer examination of the relative abundances of all *Betaproteobacteria* found in samples
214 showed that while *Burkholderiales* was the predominant order across all three species, *T.*
215 *hybridum* and *L. vulgare* samples were heavily dominated by bacteria of the family
216 *Oxalobacteriaceae* – especially bacteria of the genus *Herbaspirillum* (11% of the total bacterial
217 community in *T. hybridum* and 18% in *L. vulgare*) (Fig.3).

218 **Discussion**

219 *Host specificity*

Our study shows that root bacterial communities vary significantly between plants belonging to three different species, growing in close proximity to each other in natural plant communities. Because the plant roots collected in our study were growing in close proximity to each other it is unlikely that variation in soil conditions (usually thought to be one of the main drivers of microbial community structure) is a significant source of variation among our samples. While these results need to be further confirmed across multiple field sites in order to draw general conclusions about whether this is just a local trend or a general pattern across plant species and soil types, our results support previous work showing bacterial host plant specificity in roots of agricultural crops (Marschner and Yang 2001; Wieland et al. 2001; Haichar et al. 2008) and wild grass species (Kuske and Ticknor 2002; Osanai et al. 2012).

Although all plant species investigated in this study (*P. aurantiaca*, *T. hybridum* and *L. vulgare*), are perennial, there are significant differences in root morphology between the species. For example, *P. aurantiaca* and *L. vulgare* (both belonging to the family *Asteraceae*) have creeping root stocks and produce fibrous root systems whereas *T. hybridum* (family *Fabaceae*) grows a branching tap root system that is known to form nodules with nitrogen fixing bacteria. This variation in root morphology could contribute to the differences in abundance and composition of bacteria in our results (Bardgett et al. 2014). For example, roots that penetrate deeper soil may encounter different microbes than those in shallow layers (Fierer et al. 2003). Similarly, the thickness and/or texture of the root surface (i.e. woody, fibrous) may be more or less penetrable to colonizing bacteria.

Part of the variance seen in bacterial community composition between the three plant species could also be caused by species-specific root exudation patterns. For example, several members of the *Asteraceae* family are known to produce allelochemicals that could affect the bacterial community as well as surrounding plants (Alford et al. 2009). However, these differences are difficult to assess in wild plant communities, especially when roots of different

plant species grow in close proximity to each other with entangled root systems. In our study, the roots grew so intimately that exudation from one plant species could have influenced root systems of neighboring plants.

Individual variation in root microbiota

We know that genetic differences between plants, even at the genotype level, can affect the composition of the root microbiota (Bailey et al. 2005; Schweitzer et al. 2008; Peiffer et al. 2013). Thus, we would expect variation in bacterial community composition among individuals within a population of plants, even when they are growing in a common environment, due to genetic variation in the population in terms of root traits and exudation chemistry, among other factors. Though there is a potential for clonality in *T. hybridum*, we still predicted less individual variance in the root microbiota among individuals from *P. aurantiaca* (thought to consist of mainly one genotype across all of North America (Loomis and Fishman 2009)) than within the two co-occurring out-crossing plant species with presumed higher genetic diversity (*L. vulgare* and *T. hybridum*).

Though our data show a significant difference in compositional turnover within different plant species, it rejects the hypothesis that *P. aurantiaca* had the most similar root communities across individuals. Comparing the average dispersion of bacterial community composition for the three plant species, there was no indication that *P. aurantiaca* had a smaller dispersion than the two other plant species (Table 2). Instead, it shows that *P. aurantiaca* had the highest variation within a species comparing dispersion based on taxonomic differences (Table 2). The fact that we could not detect any differences in dispersion when using phylogenetic metrics suggests that individual root systems differ more in terms of which taxa are present or absent than how related they are, or that there is little phylogenetic conservatism at the individual level.

Overall, this study shows that the extent of individual variation seen in root microbiota varies between species, but that a plant species thought to be more genetically homogenous does not necessarily host more homogeneous root communities. It also indicates that individual variation in bacterial community composition in root systems is determined, not only by plant genetics, but also by small scale variance in the surrounding environment (soil chemistry, humidity, biotic interactions etc.) and potentially, events throughout the plant's life that could affect root colonization (Alekklett andand Hart 2013).

Bacterial community composition

Similar patterns of bacterial community composition to what we found in our plants, growing in a subalpine meadow in Canada, have been reported in rhizosphere samples of other studies. For example, roots tissues of the plant species that we sampled were mainly dominated by *Betaproteobacteria*, (Fig. 2), especially members of the order *Burkholderiales* and the family *Oxalobacteriaceae*, which represented as much as 32% of the total bacterial community in *L. vulgare* (Fig. 3). Seed- and root-colonizing populations of *Oxalobacteriaceae* have previously shown to be responsive to plant species (Green et al. 2007), supporting our data of plant species hosting distinct bacterial communities. Dominance by these taxa in root systems has also been reported in other studies. For example, roots of *Arabidopsis thaliana*, examined at the same taxonomic resolution by Lundberg et al. (2012), were dominated by *Betaproteobacteria* and *Oxalobacteriaceae* both in samples of rhizosphere soil as well as in the endophytic root compartment. In sphagnum mosses, *Burkholderiales* has also been documented as one of the dominant bacterial groups, thought to be behind the production of anti-fungal compounds and anti-microbial properties of the sphagnum mosses (Opelt et al. 2007).

Other studies have found *Actinobacteria* to dominate in root tissues of plants (e.g. Ottesen et al. 2013), especially in communities of the endophytic compartment (Bodenhause et al.

2013). In our study, *Actinobacteria* represented at most 12% out of the total bacterial community in the plant species that we sampled (Fig. 2) and was mainly found in samples of *P. aurantiaca* that, in general, were less dominated by beta-proteobacteria.

The dominance of sequences belonging to the genus *Herbaspirillum* was further emphasized when we examined the fourteen most abundant OTUs across all samples (Table 3). *Herbaspirillum* spp. are known to colonize apoplastic or intracellular spaces of plant tissues and several species have shown the ability to fix nitrogen (Schmid et al. 2006). While it is believed that this nitrogen fixing ability could be beneficial to their plant host, it has also been documented that certain *Herbaspirillum* strains are mild pathogens and a causative agent of “mottled stripe disease” in crops such as sugar-cane (Schmid et al. 2006). Besides *Herbaspirillum*, we also saw high abundances of sequences belonging to *Limnohabitans* and *Cytophaga* (Table 3), two genera more commonly associated with bacterial communities in fresh water (Kirchman 2002; Simek et al. 2010) as well as the species *Methylibium petroleiphilum*, a recognized methylotroph (Kane et al. 2007) and *Janthinobacterium lividum*, known to thrive in soils (Shivaji et al. 1991) and produce antibiotics (Johnson et al. 1990). The high presence of these groups in our samples could be due to the inclusion of epiphytic members of the root microbiota, where bacteria associated with water films and soil particles of the root surface would be expected.

In comparison, Bodenhausen and colleagues (2013) found that a *Flavobacterium* of the phylum *Bacteroidetes* stood out as the single most abundant OTU in endophytic root samples, making up 10.15% of the total community. Though bacteria of the phylum *Bacteroidetes* represented a significant part of the community in root samples of the three plant species sampled in our study (Fig.2), they were by no means the most dominant taxonomic group in any of the species (Table 3).

As the genus *Trifolium* are known to be hosts of nitrogen fixing bacteria that form nodules in their roots, *T.hybridum* samples were expected to host larger populations of

317 *Alphaproteobacteria*, specifically belonging to the order *Rhizobiales* which is a common
 318 symbiont of legumes (Masson-Boivin et al. 2009). This expected pattern was not evident in our
 319 results though. The only OTU belonging to the order *Rhizobiales* detected in notable abundances
 320 in our study was a *Bradyrhizobium* taxa which made up 1% of the collective community of *T.*
 321 *hybridum* samples and 1.5% in *P. aurantiaca* samples (Table 3). Instead, it was evident that the *T.*
 322 *hybridum* community was dominated by the family *Oxalobacteraceae* (23.48%) (Fig.3) and
 323 specifically one OTU of the genus *Herbaspirillum* (10.75%) (Table 3), which is mainly known to
 324 colonize roots of non-leguminous plants, and have nitrogen fixing properties (Baldani et al. 1997).
 325 What stands out though is that this group of bacteria was even more predominant in *L. vulgare*
 326 samples, where *Oxalobacteraceae* made up 32.47% of the community and the same
 327 *Herbaspirillum* OTU represented 18.26% of the total community.

328 *Variance in relative abundances of bacterial taxa across plant species*

329 We observed differences in the evenness of bacterial taxa across host plants. While the roots of *L.*
 330 *vulgare* and *T. hybridum* seemed dominated by a few select groups of microbes, samples of *P.*
 331 *aurantiaca* supported communities with abundances more evenly distributed among bacterial
 332 taxa (Fig. 2; Table 3). Though few studies have looked specifically at variance in bacterial
 333 evenness between plant species, it could be an important source of variation. For example,
 334 dominance of single taxon may indicate specialized plant/bacterial associations whereas high
 335 evenness in community composition could reflect generalist associations among plants and
 336 bacteria. Alternatively, differences in evenness may result from microbial interactions within the
 337 plant, not driven by the plant but microbial competition for plant resources.

338 *Conclusions*

In this study, we showed that plant identity plays a major role in explaining the variation seen in root microbiota both between and within plant species growing under natural conditions. Further studies across a larger set of wild plant species and natural sites as well as more detailed investigations of the effect of plant genetics versus plant phenotypic traits on bacterial community assembly could help resolve the relative contribution of host identity at an individual level in shaping the root microbiota. It would also allow us to draw further conclusions as to whether plants more related to each other actually host more similar bacterial communities across plant species and families.

The results of our study speak of how intimately related bacterial communities are with their host plants. Root systems of wild plants are never alone; they are constantly surrounded by the roots of other plants, entangled in the soil, competing for resources and space. Yet, our results show that bacterial communities associated with roots of plants growing in a common field are distinct between plant species.

Ultimately, we are not able to tell exactly why these three plant species have such distinct bacterial root communities, but further studies linking metabolomics of wild plants with bacterial community composition would be useful for better understanding how plants affect bacterial community assembly.

Acknowledgements

We would like to thank Monika Gorzelak for helping out with sampling efforts in the field.

References

Aleklett K, Hart M. 2013. The root microbiota—a fingerprint in the soil? *Plant and Soil* 370: 671–686. doi:10.1007/s11104-013-1647-7.

- 361 Alford ÉR, Vivanco JM, Paschke MW. 2009. The effects of flavonoid allelochemicals from
362 knapweeds on legume-rhizobia candidates for restoration. *Restoration Ecology* 17: 506–514.
363 doi:10.1111/j.1526-100X.2008.00405.x.
- 364 Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool.
365 *Journal of Molecular Biology* 215: 403–10. doi:10.1016/S0022-2836(05)80360-2.
- 366 Anderson MJ. 2004. PERMDISP: a FORTRAN computer program for permutational analysis of
367 multivariate dispersions (for any two-factor ANOVA design) using permutation tests.
368 Department of Statistics, University of Auckland, New Zealand
- 369 Anderson MJ. 2005. PERMANOVA: a FORTRAN computer program for permutational
370 multivariate analysis of variance. Department of Statistics, University of Auckland, New
371 Zealand.
- 372 Anderson MJ, Walsh DCI. 2013. PERMANOVA, ANOSIM, and the Mantel test in the face of
373 heterogeneous dispersions: What null hypothesis are you testing? *Ecological Monographs*
374 83: 557–574.
- 375 Bailey JK, Deckert R, Schweitzer JA, Rehill BJ, Lindroth RL, Gehring C, Whitham TG. 2005.
376 Host plant genetics affect hidden ecological players: links among *Populus*, condensed
377 tannins, and fungal endophyte infection. *Canadian Journal of Botany* 83: 356–361.
378 doi:10.1139/b05-008. Bais HP, Park S-W, Weir TL, Callaway RM, Vivanco JM. 2004. How
379 plants communicate using the underground information superhighway. *Trends in Plant*
380 *Science* 9: 26–32. doi:10.1016/j.tplants.2003.11.008.
- 381 Bais HP, Weir TL, Perry LG, Gilroy S, Vivanco JM. 2006. The role of root exudates in
382 rhizosphere interactions with plants and other organisms. *Annual Review of Plant Biology*
383 57: 233–66. doi:10.1146/annurev.arplant.57.032905.105159.
- 384 Baldani J, Caruso L, Baldani V, Goi S, Döbereiner J. 1997. Recent advances in BNF with non-
385 legume plants. *Soil Biology and Biochemistry* 29: 911–922.
- 386 Bardgett RD, Mommer L, De Vries FT. 2014. Going underground: root traits as drivers of
387 ecosystem processes. *Trends in Ecology and Evolution* 29: 692–699.
- 388 Becklin KM, Hertweck KL, Jumpponen A. 2012. Host identity impacts rhizosphere fungal
389 communities associated with three alpine plant species. *Microbial Ecology* 63: 682–93.
390 doi:10.1007/s00248-011-9968-7.
- 391 Bell C, Carrillo Y, Boot CM, Rocca JD, Pendall E, Wallenstein MD. 2013. Rhizosphere
392 stoichiometry: are C : N : P ratios of plants, soils, and enzymes conserved at the plant
393 species-level? *New Phytologist*: 1–13. doi:10.1111/nph.12531.
- 394 Berg G, Smalla K. 2009. Plant species and soil type cooperatively shape the structure and
395 function of microbial communities in the rhizosphere. *FEMS Microbiology Ecology* 68: 1–
396 13. doi:10.1111/j.1574-6941.2009.00654.x.

- 397 Bodenhausen N, Horton MW, Bergelson J. 2013. Bacterial communities associated with the
398 leaves and the roots of *Arabidopsis thaliana*. *PloS one* 8: e56329.
399 doi:10.1371/journal.pone.0056329.
- 400 Bulgarelli D, Rott M, Schlaeppi K, Ver Loren van Themaat E, Ahmadinejad N, Assenza F, Rauf
401 P, Huettel B, Reinhardt R, Schmelzer E, Peplies J, Gloeckner FO, Amann R, Eickhorst T,
402 Schulze-Lefert P. 2012. Revealing structure and assembly cues for *Arabidopsis* root-
403 inhabiting bacterial microbiota. *Nature* 488: 91–5. doi:10.1038/nature11336.
- 404 Bulgarelli D, Schlaeppi K, Spaepen S, Ver Loren van Themaat E, Schulze-Lefert P. 2013.
405 Structure and functions of the bacterial microbiota of plants. *Annual Review of Plant*
406 *Biology* 64: 807–38. doi:10.1146/annurev-arplant-050312-120106.
- 407 Caporaso J, Kuczynski GJ, Stombaugh J, Bittinger K, Bushman FD, Costello EK, Fierer N, Peña
408 AG, Goodrich JK, Gordon JI, Huttley GA, Kelley ST, Knights D, Koenig JE, Ley RE,
409 Lozupone CA, McDonald D, Muegge BD, Pirrung M, Reeder J, Sevinsky JR, Turnbaugh PJ,
410 Walters WA, Widmann J, Yatsunenko T, Zaneveld J, Knight R. 2010. QIIME allows analysis
411 of high-throughput community sequencing data. *Nature Methods* 7: 335–6.
412 doi:10.1038/nmeth.f.303.
- 413 Chaparro JM, Badri DV, Bakker MG, Sugiyama A, Manter DK, Vivanco JM. 2013. Root
414 exudation of phytochemicals in *Arabidopsis* follows specific patterns that are
415 developmentally programmed and correlate with soil microbial functions. *PloS one* 8:
416 e55731. doi:10.1371/journal.pone.0055731.
- 417 Chelius MK, Triplett EW. 2001. The diversity of archaea and bacteria in association with the
418 roots of *Zea mays* L. *Microbial Ecology* 41: 252–263. doi:10.1007/s002480000087.
- 419 Clarke KR, Gorley RN. 2006. PRIMER v6: User Manual/Tutorial. Plymouth Marine Laboratory,
420 Plymouth, UK.
- 421 Costa R, Götz M, Mrotzek N, Lottmann J, Berg G and Smalla K. 2006. Effects of site and plant
422 species on rhizosphere community structure as revealed by molecular analysis of microbial
423 guilds. *FEMS Microbiology Ecology* 56: 236–49. doi:10.1111/j.1574-6941.2005.00026.x.
- 424 DeSantis TZ, Hugenholtz P, Larsen N, Rojas M, Brodie EL, Keller K, Huber T, Dalevi D, Hu P,
425 Andersen GL. 2006. Greengenes, a chimera-checked 16S rRNA gene database and
426 workbench compatible with ARB. *Applied and Environmental Microbiology* 72: 5069–72.
427 doi:10.1128/AEM.03006-05.
- 428 Doornbos RF, van Loon LC, Bakker PAHM. 2011. Impact of root exudates and plant defense
429 signaling on bacterial communities in the rhizosphere. A review. *Agronomy for Sustainable*
430 *Development* 32: 227–243. doi:10.1007/s13593-011-0028-y.
- 431 Farrar J, Hawes M, Jones D, Lindow S. 2003. How roots control the flux of carbon to the
432 rhizosphere. *Ecology* 84: 827–837.

- 433 Fierer N, Hamady M, Lauber CL, Knight R. 2008. The influence of sex, handedness, and
434 washing on the diversity of hand surface bacteria. *Proceedings of the Natural Academy of*
435 *Science USA* 105: 17994–9. doi:10.1073/pnas.0807920105.
- 436 Fierer N, Schimel JP, Holden PA. 2003. Variations in microbial community composition through
437 two soil depth profiles. *Soil Biology and Biochemistry* 35: 167–176. doi:10.1016/S0038-
438 0717(02)00251-1.
- 439 Garbeva P, van Veen JA and van Elsas JD. 2004. Microbial diversity in soil: selection microbial
440 populations by plant and soil type and implications for disease suppressiveness. *Annual*
441 *review of phytopathology* 42: 243–70. doi:10.1146/annurev.phyto.42.012604.135455.
- 442 Giroday H, Baker V. 2006. Invasive hawkweeds (*Hieracium* spp.) in Northeastern British
443 Columbia - Invasive plant risk assessment and literature search. *British Columbia Ministry*
444 *of Forests and Range*, Prince George, Canada.
- 445 Gottel, NR, Castro HF, Kerley M, Yang Z, Pelletier DA, Podar M, Karpinets T, Uberbacher E,
446 Tuskan GA, Vilgalys R, Doktycz MJ, Schadt CW. 2011. Distinct microbial communities
447 within the endosphere and rhizosphere of *Populus deltoides* roots across contrasting soil
448 types. *Applied and Environmental Microbiology* 77: 5934–44. doi:10.1128/AEM.05255-11.
- 449 Green SJ, Michel FC, Hadar Y, Minz D. 2007. Contrasting patterns of seed and root colonization
450 by bacteria from the genus *Chryseobacterium* and from the family *Oxalobacteraceae*. *The*
451 *ISME Journal* 1: 291–9. doi:10.1038/ismej.2007.33.
- 452 Haichar FEZ, Marol C, Berge O, Rangel-Castro JI, Prosser JI, Balesdent J, Heulin T, Achouak W.
453 2008. Plant host habitat and root exudates shape soil bacterial community structure. *The*
454 *ISME journal* 2: 1221–30. doi:10.1038/ismej.2008.80.
- 455 Hamady M, Knight R. 2009. Microbial community profiling for human microbiome projects:
456 Tools, techniques, and challenges. *Genome research*, 19, 1141–52.
- 457 Hamady M, Lozupone C, R. Knight. 2010. Fast UniFrac: facilitating high-throughput
458 phylogenetic analyses of microbial communities including analysis of pyrosequencing and
459 PhyloChip data. *The ISME Journal* 4: 17–27. doi:10.1038/ismej.2009.97.
- 460 Hartmann A, Schmid M, van Tuinen D, Berg G. 2008. Plant-driven selection of microbes. *Plant*
461 *and Soil* 321: 235–257. doi:10.1007/s11104-008-9814-y.
- 462 Hodge A, Berta G, Doussan C, Merchan F, Crespi M. 2009. Plant root growth, architecture and
463 function. *Plant and Soil* 321: 153–187. doi:10.1007/s11104-009-9929-9.
- 464 Inceoglu O, Salles JF, van Overbeek L, van Elsas JD. 2010. Effects of plant genotype and growth
465 stage on the betaproteobacterial communities associated with different potato cultivars in
466 two fields. *Applied and Environmental Microbiology*, 76, 3675–84.
- 467 Kane SR, Chakicherla AY, Chain PSG, Schmidt R, Shin MW, Legler TC, Scow KM, Larimer
468 FW, Lucas SM, Richardson PM, Hristova KR. 2007. Whole-genome analysis of the methyl

- 469 tert-butyl ether-degrading beta-proteobacterium *Methylibium petroleiphilum* PM1. *Journal*
470 *of Bacteriology* 189: 1931–45. doi:10.1128/JB.01259-06.
- 471 Kirchman DL. 2002. The ecology of Cytophaga-Flavobacteria in aquatic environments. *FEMS*
472 *Microbiology Ecology* 39: 91–100. doi:10.1111/j.1574-6941.2002.tb00910.x.
- 473 Kuczynski J, Liu Z, Lozupone C, McDonald D, Fierer N, Knight R. 2010. Microbial community
474 resemblance methods differ in their ability to detect biologically relevant patterns. *Nature*
475 *methods* 7: 813–9.
- 476 Kuske C, Ticknor L. 2002. Comparison of soil bacterial communities in rhizospheres of three
477 plant species and the interspaces in an arid grassland. *Applied and Environmental*
478 *Microbiology* 68: 1854–1863. doi:10.1128/AEM.68.4.1854–1863.2002.
- 479 Loomis ES, Fishman L. 2009. A continent wide clone: population genetic variation of the
480 invasive plant *Hieracium aurantiacum* (Orange Hawkweed; Asteraceae) in North America.
481 *International Journal of Plant Science* 170: 759–765. doi:10.1086/599241.
- 482 Lozupone CA, Hamady M, Kelley ST, Knight R. 2007. Quantitative and qualitative beta diversity
483 measures lead to different insights into factors that structure microbial communities. *Applied*
484 *and Environmental Microbiology* 73: 1576–85. doi:10.1128/AEM.01996-06.
- 485 Lundberg DS, Lebeis SL, Paredes SH, Yourstone S, Gehring J, Malfatti S, Tremblay J,
486 Engelbrektson A, Kunin V, del Rio TG, Edgar RC, Eickhorst T, Ley RE, Hugenholtz P,
487 Tringe SG, Dangl JL. 2012. Defining the core *Arabidopsis thaliana* root microbiome. *Nature*
488 488: 86–90. doi:10.1038/nature11237.
- 489 Manter DK, Delgado JA, Holm DG, Stong RA. 2010. Pyrosequencing reveals a highly diverse
490 and cultivar-specific bacterial endophyte community in potato roots. *Microbial Ecology* 60:
491 157–66. doi:10.1007/s00248-010-9658-x.
- 492 Marschner P, Grierson PF, Rengel Z. 2005. Microbial community composition and functioning in
493 the rhizosphere of three *Banksia* species in native woodland in Western Australia. *Applied*
494 *Soil Ecology* 28: 191–201. doi:10.1016/j.apsoil.2004.09.001.
- 495 Marschner P, Yang C. 2001. Soil and plant specific effects on bacterial community composition
496 in the rhizosphere. *Soil Biology and Biochemistry* 33: 1437–1445.
- 497 Masson-Boivin C, Giraud E, Perret X, Batut J. 2009. Establishing nitrogen-fixing symbiosis with
498 legumes: how many rhizobium recipes? *Trends in Microbiology* 17: 458–66.
- 499 Micallef SA, Channer S, Shiaris MP, Colón-Carmona A. 2009. Plant age and genotype impact the
500 progression of bacterial community succession in the *Arabidopsis* rhizosphere. *Plant*
501 *signaling and behavior* 4: 777–80. doi:10.1093/jxb/erp053.
- 502 Osanai Y, Bougoure DS, Hayden HL, Hovenden MJ. 2012. Co-occurring grass species differ in
503 their associated microbial community composition in a temperate native grassland. *Plant*
504 *and Soil* 368: 419–431. doi:10.1007/s11104-012-1529-4.

- 505 Ottesen AR, González Peña A, White JR, Pettengill JB, Li C, Allard S, Rideout S, Allard M, Hill
506 T, Evans P, Strain E, Musser S, Knight R, Brown E. 2013. Baseline survey of the anatomical
507 microbial ecology of an important food plant: *Solanum lycopersicum* (tomato). *BMC*
508 *Microbiology* 13: 114. doi:10.1186/1471-2180-13-114.
- 509 Opelt K, Chobot V, Hadacek F, Schönmann S, Eberl L and Berg G. 2007. Investigations of the
510 structure and function of bacterial communities associated with *Sphagnum* mosses.
511 *Environmental microbiology* 9: 2795–809. doi:10.1111/j.1462-2920.2007.01391.x.
- 512 van Overbeek L, van Elsas JD. 2008. Effects of plant genotype and growth stage on the structure
513 of bacterial communities associated with potato (*Solanum tuberosum* L.). *FEMS*
514 *Microbiology Ecology* 64: 283–96. doi:10.1111/j.1574-6941.2008.00469.x.
- 515 Peiffer JA, Spor A, Koren O, Jin Z, Tringe SG, Dangl JL, Buckler ES, Ley RE. 2013. Diversity
516 and heritability of the maize rhizosphere microbiome under field conditions. *Proceedings of*
517 *the Natural Academy of Science USA* 110: 6548–53. doi:10.1073/pnas.1302837110.
- 518 Redford, A. J., Bowers RM, Knight R, Y. Linhart and N. Fierer. 2010. The ecology of the
519 phyllosphere: geographic and phylogenetic variability in the distribution of bacteria on tree
520 leaves. *Environmental microbiology* 12: 2885–93. doi:10.1111/j.1462-2920.2010.02258.x.
- 521 de Ridder-Duine AS, Kowalchuk GA, Klein Gunnewiek PJA, Smant W, van Veen JA, de Boer W.
522 2005. Rhizosphere bacterial community composition in natural stands of *Carex arenaria*
523 (sand sedge) is determined by bulk soil community composition. *Soil Biology and*
524 *Biochemistry* 37: 349–357. doi:10.1016/j.soilbio.2004.08.005.
- 525 Roumet C, Urcelay C., Díaz S. 2006. Suites of root traits differ between annual and perennial
526 species growing in the field. *The New Phytologist* 170: 357–68. doi:10.1111/j.1469-
527 8137.2006.01667.x.
- 528 Schmid M, Baldani JIVO, Hartmann A. 2006. The Prokaryotes. Pages 141–150 (M. Dworkin, S.
529 Falkow, E. Rosenberg, K.-H. Schleifer, and E. Stackebrandt, Eds.). 141–150, New York, NY.
- 530 Schweitzer JA, Bailey JK, Fischer DG, LeRoy CJ, Lonsdorf EV, Whitham TG, Hart SC. 2008.
531 Plant–soil–microorganism interactions: heritable relationship between plant genotype and
532 associated soil microorganisms. *Ecology* 89: 773–781. doi:10.1890/07-0337.1.
- 533 Shivaji S, Ray M, Kumar G, Reddy G, Saisree L, Wynn-Williams D. 1991. Identification of
534 *Janthinobacterium lividum* from the soils of the islands of Scotia Ridge and from Antarctic
535 peninsula. *Polar biology* 11: 267–271.
- 536 Simek K, Kasalicky V, Jezbera J, Jezberová J, Hejzlar J, Hahn MW. 2010. Broad habitat range of
537 the phylogenetically narrow R-BT065 cluster, representing a core group of the
538 Betaproteobacterial genus *Limnochabitans*. *Applied and Environmental Microbiology* 76:
539 631–9. doi:10.1128/AEM.02203-09.
- 540 Singh, B. K., S. Munro, J. M. Potts and P. Millard. 2007. Influence of grass species and soil type
541 on rhizosphere microbial community structure in grassland soils. *Applied Soil Ecology* 36:
542 147–155. doi:10.1016/j.apsoil.2007.01.004.

- 543 Smalla K, Wieland G, Buchner A, Zock A, Parzy J, Kaiser S, Roskot N. 2001. Bulk and
 544 rhizosphere soil bacterial communities studied by denaturing gradient gel electrophoresis :
 545 plant-dependent enrichment and seasonal shifts revealed. *Applied and Environmental*
 546 *Microbiology* 67: 4742–4751. doi:10.1128/AEM.67.10.4742.
- 547 Weinert N, Meincke R, Gottwald C, Heuer H, Gomes NCM, Schlöter M, Berg G, Smalla K.
 548 2009. Rhizosphere communities of genetically modified zeaxanthin-accumulating potato
 549 plants and their parent cultivar differ less than those of different potato cultivars. *Applied*
 550 *and Environmental Microbiology* 75: 3859-3865.
- 551 Wieland G, Neumann R, Backhaus H. 2001. Variation of microbial communities in soil,
 552 rhizosphere, and rhizoplane in response to crop species, soil type, and crop development.
 553 *Applied and Environmental Microbiology* 67: 5849–5854. doi:10.1128/AEM.67.12.5849–
 554 5854.2001.

Table 1(on next page)

PerMANOVA results, comparing bacterial community resemblance between plant species and transects using different diversity metrics.

Factor	Diversity metric	Pseudo-F	P (perm)
<i>Species</i>	Weighted UniFrac	6.15	0.0001
	Un-weighted UniFrac	1.43	0.0001
	Bray Curtis	1.88	0.0001
<i>Transect</i>	Weighted UniFrac	1.22	0.23
	Un-weighted UniFrac	1.06	0.22
	Bray Curtis	1.07	0.24
<i>SpXTr</i>	Weighted UniFrac	1.54	0.09
	Un-weighted UniFrac	1.03	0.28
	Bray Curtis	1.06	0.20

1

Principal coordinates analysis plot illustrating the phylogenetic overlap in root prokaryotic community composition among samples from three different plant species.

Phylogenetic overlap between communities was assessed using weighted UniFrac. Community composition was significantly different among plant species ($P < 0.001$; PerMANOVA).




Table 2 (on next page)

PermDISP results showing the average spread from centroid and standard error (SE) for samples of each species.

The PERMANOVA (P(permanova)) values are assessing whether there is a significant difference between species in sample dispersion, using different diversity metrics.

Diversity metric	F	P (perm)	Species	Average	SE
Weighted UniFrac	3.10	0.11	<i>P. aurantiaca</i>	9.96 E ⁻²	4.28 E ⁻³
			<i>T. hybridum</i>	8.57 E ⁻²	5.70 E ⁻³
			<i>L. vulgare</i>	0.12	1.29 E ⁻²
Un-weighted UniFrac	1.04	0.37	<i>P. aurantiaca</i>	0.47	3.39 E ⁻³
			<i>T. hybridum</i>	0.46	5.42 E ⁻³
			<i>L. vulgare</i>	0.47	7.25 E ⁻³
Bray Curtis	9.30	0.005	<i>P. aurantiaca</i>	51.43	0.73
			<i>T. hybridum</i>	47.13	0.44
			<i>L. vulgare</i>	51.05	1.05

2

Comparison of the average bacterial community composition and relative abundances, at the phylum level (*Proteobacteria* divided into class) in root samples from three different plant species.

Results show a strong dominance of sequences belonging to *Betaproteobacteria* in all three plant species, but especially in *T. hybridum* (51%) and *L. vulgare* (50%). The phyla representing less than 1% out of the total community were grouped as “Other” and consisted of: NKB19, *Nitrospirae*, PAUC34f, *Cyanobacteria*, *Elusimicrobia*, *Fibrobacteres*, *Chlamydiae*, SC4, *Spirochaetes* and *Thermi*. Sequences not matching the database were recorded as “No blast hit”.




Table 3(on next page)

The core root microbiota represented by the fourteen OTUs with the highest abundances across all samples

Values are calculated as the average percent out of the 400 sequences recorded for each sample, across all species (Total %). As a comparison, data is also included for what percentage (on average) the fourteen OTUs make up within the bacterial communities of the three plant species examined (*P. aurantiaca*, *T. hybridum* and *L. vulgare*).

# OTU ID	Total %	<i>P. aurantiaca</i> %	<i>T. hybridum</i> %	<i>L. vulgare</i> %	Phylum	Class	Order	Family	Genus	Species
1537	7.6	2.8	9.4	16.8	Proteobacteria	Beta-proteobacteria	Burkholderiales	Oxalobacteraceae	Herbaspirillum	
19032	3.1	3.2	3.9	2.2	Proteobacteria	Beta-proteobacteria	Burkholderiales			
22328	2.7	1.0	5.1	3.8	Proteobacteria	Beta-proteobacteria	Burkholderiales	Comamonadaceae	Limnohabitans	
30435	1.5	1.7	1.3	1.2	Proteobacteria	Beta-proteobacteria	Burkholderiales		Methylibium	Methylibium petroleiphilum
4453	1.3	0.8	1.7	2.1	Proteobacteria	Beta-proteobacteria	Burkholderiales	Oxalobacteraceae	Janthinobacterium	Janthinobacterium lividum
1231	1.2	1.9	0.5	0.4	Bacteroidetes	Sphingobacteria	Sphingobacteriales	Flexibacteraceae	Cytophaga	
22285	1.0	1.5	1.0	0.4	Proteobacteria	Alpha-proteobacteria	Rhizobiales	Bradyrhizobiaceae	Bradyrhizobium	
20009	1.0	1.5	0.2	0.6	Actinobacteria	Actinobacteria	Actinomycetales	Thermomonosporaceae	Actinocorallia	Actinocorallia longicatena
25072	0.9	1.0	1.0	0.7	Proteobacteria	Gamma-proteobacteria	Xanthomonadales	Xanthomonadaceae	Rhodanobacter	Rhodanobacter lindaniclasticus
1340	0.9	1.0	1.0	0.6	Bacteroidetes	Sphingobacteria	Sphingobacteriales	Sphingobacteriaceae	Sphingobacterium	Sphingobacterium faecium
5213	0.8	0.6	0.9	1.3	Bacteroidetes	Sphingobacteria	Sphingobacteriales	Sphingobacteriaceae		
29492	0.8	1.3	0.1	0.6	Proteobacteria	Gamma-proteobacteria	Chromatiales	Sinobacteraceae		
26917	0.8	1.1	0.3	0.7	Chloroflexi	Chloroflexi	Roseiflexales	Kouleothrixaceae	Kouleothrix	
1184	0.8	0.4	1.3	1.1	Bacteroidetes	Flavobacteria	Flavobacteriales	Flavobacteriaceae	Chryseobacterium	

3

Average relative abundances of *Betaproteobacteria* families and *Oxalobacteraceae* genera found in root samples of the three plant species.

Values are given as the percentage of sequences belonging to a certain taxa out of the total average bacterial community for each of the three plant species (rarefied at 400 sequences/sample). The heat map is colour coded from blue (low abundance) to red (high abundance).

Taxon	P.aurantiaca	T.hybridum	L.vulgare
Unclassified Betaproteobacteria	2.33	1.43	1.50
Unclassified Burkholderiales	9.92	13.65	7.19
Burkholderiales_Alcaligenaceae	0.32	0.53	0.31
Burkholderiales_Burkholderiaceae	0.57	0.63	0.31
Burkholderiales_Comamonadaceae	2.67	9.78	5.88
Burkholderiales_Oxalobacteraceae	9.28	23.48	32.47
Gallionellales_Gallionellaceae	0.08	0.00	0.09
Hydrogenophilaes_Hydrogenophilaceae	0.01	0.05	0.00
Unclassified Methylophilales	0.05	0.03	0.03
Methylophilales_Methylophilaceae	1.37	0.70	0.88
Unclassified Rhodocyclales	0.76	0.08	0.44
Rhodocyclales_Rhodocyclaceae	1.87	0.35	1.22
Total percentage Betaproteobacteria	29%	51%	50%

	P.aurantiaca	T.hybridum	L.vulgare
Unclassified Oxalobacteraceae	3.39	6.93	8.94
Oxalobacteraceae_Collimonas	0.38	0.88	0.72
Oxalobacteraceae_Herbaspisillum	3.21	10.75	18.28
Oxalobacteraceae_Anthinobacterium	1.17	2.13	2.66
Oxalobacteraceae_Massilia	0.61	1.50	1.13
Oxalobacteraceae_Oxalobacter	0.50	1.30	0.75