

Rehabilitation time has greater influences on soil mechanical composition and erodibility than does rehabilitation land type in the hilly-gully region of the Loess Plateau, China

Leilei Qiao^{1▲}, Wenjing Chen^{1,2▲}, Yang Wu², Hongfei Liu², Jiaoyang Zhang^{3,4}, Guobin Liu^{1, 3}, Sha Xue^{1, 3*}

¹ State Key Laboratory of Soil Erosion and Dryland Farming on the Loess Plateau, Institute of Soil and Water Conservation, Northwest A&F University, Yangling 712100, P. R. China.

² Collage of Forestry, Northwest A&F University, Yangling 712100, P. R. China.

³ Institute of Soil and Water Conservation, Chinese Academy of Sciences and Ministry of Water Resources, Yangling 712100, P. R. China.

⁴ University of Chinese Academy of Sciences, Beijing, China.

Address correspond Institute of Soil and Water Conservation, Northwest A&F University, Yangling 712100, P. R. China. E-mail: xuesha100@163.com

▲Joint first authors

Rehabilitation time has greater influences on soil mechanical composition and erodibility than does rehabilitation land type in the hilly-gully region of the Loess Plateau, China

Background. The major landscape in the hilly-gully region of the Loess Plateau is greatly affected by vegetation rehabilitation on abandoned cropland. Although many studies have shown that the rehabilitation have greatly improved soil conditions and protected them from erosion, these effectiveness were not always in consensus possibly due to the land type of vegetation or to the rehabilitation time. To close this gap, we conducted a long term experiment as follows.

Methods. In this study, we analysed four land types of vegetation rehabilitation (shrub land, woodland, naturally revegetated grassland, and orchard land) with different rehabilitation times and investigated the mechanical composition and erodibility of the soil. Areas of slope cropland and natural forest were selected as controls.

Results. The results showed that soil depth, rehabilitation time and rehabilitation land type had strong impacts on soil mechanical composition, micro-aggregation and erodibility. Following rehabilitation, naturally revegetated grassland and shrub land had lower fractal dimensions of particle size distribution (fractal dimensions of PSD), fractal dimensions of micro-aggregation, and erodibility (K factor) than did cropland. Compared to the positive effects of rehabilitation mainly happened in the topsoil layer at other rehabilitation land type, that of woodland happened in the deeper soil layer.

Besides, the indispensable rehabilitation time for the significant improvement of soil condition was shorter at naturally revegetated grassland than that at shrub land and woodland.

Discussion. Although rehabilitation time was more influential than was rehabilitation land type or soil depth, the differences among the rehabilitation land types showed that naturally revegetated grassland with native plants is the most time-saving rehabilitation vegetation for the Loess Plateau in the conversion from slope cropland. The success of rehabilitation in this forestry practice was mainly contributed by the suited species of rehabilitation land type to the local climate and soil. Based on the differences of rehabilitation effectiveness resulting from land type, we should be cautious to choose land types for the rehabilitation of soil conditions in the Loess Plateau.

Key word: erosion; vegetation rehabilitation; fractal dimension; Loess Plateau

Introduction

Soil erosion initiated by either natural or human factors is a serious environmental problem in many regions of the world. It not only caused the degradation of soil quality but also the destruction of ecosystem function and safety (Chen & Peng 2000; Lian et al., 2013; Borrelli et al. 2017). Severe soil erosion is a serious challenge in the Loess Plateau of China (Sun et al., 2016b). To alleviate soil erosion and restore the local ecological environment, the Chinese government implemented the “Grain for Green” programme in 1999 to convert degraded cropland to forest and grassland (Uchida et al. 2005; Chen et al., 2007a; Uchida et al. 2009;

Eliminado: parts

Zhang et al., 2011a; Song et al., 2015). This programme has greatly decreased soil loss (*Chen et al., 2007b*).

Several studies had examined the effects of plant species changes, land preparation, rainfall intensity, anthropogenic disturbance, afforestation, and land abandonment on the mechanical composition and erodibility of the soil (*Koulouri & Giourga 2007; Keesstra et al., 2009; Xia et al., 2009; Zhang et al., 2011b; Moora et al. 2014; Yu et al., 2017*). Soil mechanical composition and micro-aggregate stability were shaped by complicated geophysical and environmental processes and responded to land-use changes, thereby affecting soil hydrological and mechanical functioning and soil erosion (*Wang et al., 2005; Alagöz & Yilmaz 2009; Xiao et al., 2014; Wang et al., 2016*). Many studies have reported positive impacts of vegetation rehabilitation of sloped cropland on soil conditions and soil resistance to erosion (*Xu et al., 2013; Ziadat & Taimeh 2013; Xiao et al., 2014; Fu et al., 2015; Sun et al., 2016b*). Different plant species, with differences in morphology, architecture and other biological characteristics, show variation in their effectiveness for vegetation rehabilitation (*Bochet & García-Fayos 2004; Ghestem et al. 2014; Fu et al., 2015*). However, local precipitation, parent material, disturbance and their interaction and sampling time can influence vegetation rehabilitation and make interpretation of results challenging. Thus, long-term research on the dynamics of soil erosion is necessary to understand the effects of vegetation rehabilitation on soil physical condition while accounting for confounding factors. However, several studies have focused on the effects of different rehabilitation patterns or the dynamic changes following rehabilitation in a certain

land type but have not clearly identified the impacts of the various rehabilitation land type on the soil mechanical composition and erodibility during a long-time scale. Soil erosion, solution transformation and soil-moisture are influenced by soil particle size distribution (PSD) (Mazaheri & Mahmoodabadi 2012; Yu et al., 2015). Land use could influence soil structure and physical and biochemical activity through PSD affected by water erosion (Basic et al., 2004; Su et al., 2004). Therefore, its variation remains to be characterized for understanding and evaluating soil structure and dynamics and the effects of land use on soil structure. Fractal theory, an effective and reliable tool, can be used to characterize it (Chen and Zhou, 2013).

For the past fifty years, people pressurized by an increasing population into converting native grasslands into farmlands in the most parts of Loess Plateau of China, which caused the loss of most of the topsoil in many locations (Wei et al. 2006; Zhou et al. 2006). The "Grain for Green" Programs (GGP) launched by Chinese government aimed at reducing soil erosion through replacing degraded cropland with forest and grassland. Since then, a sloped cropland was abandoned and restored naturally and artificially. As we know, herbs, rather than trees or shrubs, were dominant on the Loess Plateau due to its special geological characteristics in a long historical period. A proper choice of rehabilitation land type for the success of afforestation is the key thing (Lu et al., 2003; Jiang et al., 2013). Thus, the difference of this rehabilitation effectiveness between native vegetation (naturally revegetated native grass) and common forestry afforestation (artificial ecological forest, artificial economic forest, and artificial shrub) deserves our attention. In this study, we

Con formato: Color de fuente: Texto 1

Eliminado: n

Eliminado: artificial

collected comprehensive and long-term data on historic vegetation (e.g., forest, shrubland and grassland) with different rehabilitation times (1) to elucidate the effects of rehabilitation land type, time and soil depth on soil mechanical condition and erodibility; (2) to clearly identify the key influencing factors.

Materials and methods

Experimental area

This study was conducted in Ansai County, Shannxi Province, China (36°31'-37°20'N, 108°52'-109°26'E; 1,012-1,731 m a.s.l.), which lies in the middle part of the Loess Plateau. This region has a typical semiarid continental climate with a mean annual temperature of 8.8°C, meaning that monthly temperature ranges from 22.5°C in July to 7°C in January and an annual precipitation of 549.1 mm, which mainly occurs between July and September. The landform is characterized by a deeply incised hilly-gully Loess landscape. The soil in this area is mainly Huangmian soil, a Calcic Cambisol classified in the WRB reference system (FAO/UNESCO/ISRIC, 1988), originating from wind-blown deposits and characterised by yellow color, absence of bedding, silty texture, looseness, macroporosity, and wetness-induced collapsibility (Xiao *et al.*, 2014). This type of soil is characterized by weak cohesion (Sun *et al.*, 2016a), which makes it highly susceptible to severe soil erosion.

So far, native vegetation on the Loess Plateau remains controversial, because the vegetation cover of the Loess Plateau has been changing greatly during the historical period. The research about paleo-pedology, phytolith, organic carbon stable isotope

Comentario [GP1]: Could you please insert a reference here? It is important to see how much updated these data are. Thanks.

and pollen records showed that herbs, rather than trees or shrubs, were dominant on the Loess Plateau in both the cold-dry period and the warm-humid period, owing to specific lithological property with thick loess which can not support an extensive forest development, even during the climatic optimum in this area (Lu et al., 2003; Jiang et al., 2013). Overgrazing, deforestation and other land-use patterns led to severe damage to the ecological environment and severe soil erosion by the middle of the last century. Since the late 1950s, the land use type in this region changed remarkably by the GGP, sloping cropland has been replanted with woodland (*Robinia pseudoacacia*), shrubland (*Caragana korshinskii*, *Hippophae rhamnoides*), artificial grassland (*Medicago sativa*) and naturally revegetated grassland to control soil erosion (Sun et al., 2016b). Much of barren lands and degraded croplands with slopes over 15° were rebuilt, accounting for about 14% of hilly Loess Plateau (Xu et al., 2018). Artificially and naturally rebuilt grassland, shrub land, and planted woodland consist of the main land use types in the region.

In this region, before restoration the soil is weakly cohesive and thus prone to erosion, with erosion modulus of 10,000-12,000 mg km⁻² yr⁻¹ (Liu, 1999). Recently, some soil and water conservation measure such as revegetation have effectively reduced soil erosion and have successfully restored some degraded ecosystems in this area, where the representative vegetation includes woody plants such as *Robinia pseudoacacia*, shrubs as *Hippophae rhamnoides* and *Caragana microphylla* and herbaceous plants such as *Artemisia sacrorum* and *Stipa bungeana* (Sun et al., 2016a). The wood land area has increased from <5% to >40% since 1980 (Xu et al., 2018)

(Figure 1).

Sampling and data collection

Based on the investigation of the history of land use, we selected four types of vegetation rehabilitation of cropland abandoned between July 10 and September 10 in 2011 and 2012, the season in which plant community biomass peaks. The four types were naturally revegetated grassland (with rehabilitation times of 2yr, 5yr, 8yr, 11yr, 15yr, 18yr, 26yr and 30yr), planted shrubland (with rehabilitation times of 5yr, 10yr, 20yr, 30yr, 36yr, 47yr), planted woodland (with rehabilitation times of 5yr, 10yr, 20yr, 37yr and 56yr), and orchard land (with rehabilitation times of 5yr, 10yr and 20yr). All these sites suffering similar farming practices before conversion, and the farmlands had more than 200 years of cultivation history (*Deng et al., 2016*). Thus, sloping cropland sites were selected as representative of the original condition for the rehabilitation chronosequences of the naturally revegetated grassland, revegetated shrub land, woodland and orchard land. In this area, the climax vegetation is the *Quercus liaotungensis* Koidz (*Zhang et al., 2011b*), which ~~was~~ naturally regenerated on abandoned land from grassland to shrub land and climax forest (*Q. liaotungensis*) over about 150 years, based on previous research of secondary forests in this area. So, we considered it as representative of the soil-dominated climax community in vegetation rehabilitation to assess the effectiveness of vegetation rehabilitation. These selected sites offered representativeness, typicality and consistency and had similar slope gradients, slope aspects, and topography. The properties of the experimental sites are shown in Table 1.

Eliminado: were

184 The size of the plots were varied with the plant communities to match their
185 spatial distribution: the replicated plots of 20 ×20m were established in each site of
186 planted woodland (*Robinia pseudoacacia*), while the replicated plots of 10 ×10m
187 were established in each shrub land site (*Caragana microphylla*, *Hippophae*
188 *rhamnoides*) and in each orchard land site. The smaller replicate plots (2 ×2m) were
189 randomly established in each naturally revegetated grassland site (including *Artemisia*
190 *sacrorum*, *A. capillaries*, *A. giraldii*, *Aneurolepidium dasystachys*, *Bothriochloa*
191 *ischaemum*, *Heteropappus altaicus*, *Lespedeza bicolor*, *Stipa bungeana*, *Setaria*
192 *viridis*, and other grasses). The plots were separated by at least 50m.

193 We choose four random sampling to avoid the sampling error. At each sampling
194 plot, after removing ground litter, five soil layers (0-10, 10-20, 20-30, 30-50, and
195 50-100cm) were separately collected with a soil drilling sampler (4cm diameter). The
196 soil samples from the same layer of the same plot were mixed to form one sample.
197 The samples were divided into two parts and were passed through 2mm screens for
198 removing roots, gravel, and coarse fragments. Then each sample was brought to
199 laboratory. One part was naturally air-dried to measure the organic carbon and analyse
200 soil organic carbon (SOC), total nitrogen (TN), and total phosphorus (TP) contents,
201 particle size distributions and micro-aggregates. The other part was stored in a
202 refrigerator at 4°C to analyse water-soluble amounts (carbon, nitrogen), microbial
203 biomass (carbon, nitrogen), enzyme activity as well as other variables not reported in
204 this paper.

Physical and chemical analyses

The soil bulk density (BD) of each soil layer was measured with the cutting ring method (Ding et al., 2019). SOC was determined using the dichromate oxidation method (Nelson & Sommers, 1982), and TN was determined using the Kjeldahl method (Bremner, 1982). For soil PSD (particle-size distribution) and micro-aggregate analysis, soil samples were analysed by a laser diffraction technique using a Longbench Mastersizer 2000 (Malvern Instruments, Malvern, England) (Xiao et al., 2014). There are some differences between the pretreatment methods for determining soil PSD and micro-aggregate. For soil PSD, soil samples were pretreated with 6% H₂O₂ and 10% HCL to remove organic matter and carbonates and oxides and were soaked in distilled water for 24 h, then mechanically dispersed with 0.4% Calgon by an ultrasonic bath for 5 min. For micro-aggregate determination, the soil samples were soaked in distilled water for 24 h and mechanically dispersed in ultrasonication for 5 min (Xiao et al., 2014). Soil PSD was described in terms of the percentage of sand (0.05 – 2 mm), fine silt (0.002 – 0.020 mm), coarse silt (0.02 – 0.05 mm) and clay (<0.002 mm). The size grades of the micro-aggregates were classified to be the same as that of the PSD.

Fractal features

The fractal dimension of the PSD and micro-aggregation were calculated by the following formula (Tyler & Wheatcraft 1992):

$$V(r < R_i)/V_T = (R_i/R_{max})^{3-D}$$

where r is the particle diameter, R_i is the particle size of subinterval i in the particle size grading, $V(r < R_i)$ is the total volume of soil particles with diameter less than R_i , V_T is the sum volume of soil particles, and R_{max} is the maximum diameter of soil particles.

Erodibility (K)

Soil erodibility was measured by the K factor in the EPIC model using SOC content and soil PSD (Williams *et al.*, 1984) and was calculated as follows:

$$K = \{0.2 + 0.3 \exp[-0.0256 \text{SAN}(1 - 0.01 \text{SIL})]\} \times \left(\frac{\text{SIL}}{\text{CLA} + \text{SIL}} \right)^{0.3} \times \left(1.0 - \frac{0.25 C}{C + \exp(3.72 - 2.95 C)} \right) \times \left(1.0 - \frac{0.25 C}{\text{SNI} + \exp(-5.51 + 22.9 \text{SNI})} \right)$$

where SAN, SIL, and CLA are the sand (%), silt (%), and clay (%) fractions, respectively; C is the soil organic carbon content (%); and SNI = 1-SAN/100.

Statistical analysis

Three-way ANOVA was performed to test the effects of rehabilitation land type (naturally revegetated grassland, woodland, shrub land, orchard land), rehabilitation time (years since sloping cropland abandonment) and soil depth (0-10cm, 10-20cm, 20-30cm, 30-50cm, and 50-100cm) on soil mechanical composition and erodibility. Significance was evaluated at the 0.05 level ($P < 0.05$). Duncan's (D) post hoc test was used to perform multiple comparisons when significance of the ANOVA was observed. The differences between the natural forest and various types of vegetation rehabilitation at each last restoration year were examined by student's t test. Pearson

correlation analysis was used to analyse the correlations among particle fractal dimension, micro-aggregate fractal dimension, erodibility and soil nutrients, soil texture, and recovery time. In addition, linear regression analysis was used to determine the relationships between each of particle fractal dimension, micro-aggregate fractal dimension, and erodibility and recovery time in each soil layer between 0 and 100 cm. All of the above statistical analyses were conducted using SPSS 17.0 (SPSS Inc., Chicago, IL, USA) and R version 3.4.4.

Results

Rehabilitation time, and rehabilitation land type had significant effects on the soil PSD fractal dimension and K factor, only the rehabilitation land type had significant effects on the soil micro-aggregate fractal dimension (Table 2). PSD fractal dimension, micro-aggregate fractal dimension and erodibility showed trends of decline since cropland in all land types. However, PSD fractal dimension, micro-aggregate fractal dimension and erodibility varied among the land types (Figure 2-4).

The rehabilitation pattern in naturally revegetated grassland

Fractal dimensions of PSD and K factor began to show greater decreases in naturally revegetated grassland than in cropland at 5th, 11th years, respectively, since cropland abandonment. And these trends mainly occurred in 0-20cm, gradually weaken with depth. Overall, the minimum time before significant decreases appeared in the particle fractal dimension, soil micro-aggregate fractal dimension and

erodibility varied among the different rehabilitation land types; in general, the times were shorter for naturally revegetated grassland than for the other land types (Figure 2-4) (Table 1-3). Naturally revegetated grassland did not differ from natural forest in fractal dimensions of micro-aggregation or K factor (20-100cm) over rehabilitation time (Figure 5-7) (Table 1-3). Linear regression revealed that fractal dimensions of PSD decreased with the number of years since farmland conversion in the 0-50cm (except in 20-30cm)(Figure 2-4) (Table 1-3).

The rehabilitation pattern in Wood land

Fractal dimensions of PSD, fractal dimensions of micro-aggregation and K factor began to show greater decrease in woodland than in cropland in the 10th, 10th and 37th respectively since cropland abandonment (Figure 2-4) (Table 1-3). Over rehabilitation time, woodland did not differ from natural forest in fractal dimensions of PSD (30-100cm) and fractal dimensions of micro-aggregation (0-100cm), K factor in woodland did not differ from natural forest (0-30cm) and even was lower than that in natural forest (30-100cm) (Figure 5-7) (Table 1-3). Compared to fractal dimensions of micro-aggregation and K factor, fractal dimensions of PSD in this site showed a clear decreasing trend at all soil layers (Figure 2-4) (Table 1-3).

The rehabilitation pattern in Shrub land

Fractal dimensions of PSD, fractal dimensions of micro-aggregation and K factor began to show greater decreases in shrub land than in cropland in the 20th, 36th and

30th years, respectively, since cropland abandonment, and tended to have significant differences with cropland since then (Figure 2-4) (Table 1-3). The decreasing trend of fractal dimensions of PSD and fractal dimensions of micro-aggregation occurred in 0-100cm, but it gradually weakened with increasing depth. After rehabilitation, naturally revegetated grassland were also higher than natural forest in fractal dimensions of PSD (0-50cm), fractal dimensions of micro-aggregation (0-30cm) and K factor (0-10cm) (Figure 5-7) (Table 1-3), and had no difference with that of natural forest at the deep layer. Linear regression indicated that fractal dimensions of PSD decreased with the number of years since farmland conversion in the 0-100 cm (except 30-50cm) (Figure 2-4) (Table 1-3).

The rehabilitation pattern in Orchard land

Fractal dimensions of PSD and fractal dimensions of micro-aggregation showed a trend of lower levels in orchard land than in cropland, but there are no significant differences between them (Figure 2-4) (Table 1-3). However, K factor in this site didn't decrease after a long-term rehabilitation. Following rehabilitation, fractal dimensions of PSD, fractal dimensions of micro-aggregation, and K factor were significantly higher in orchard land than in natural forest at shallow soil layer (Figure 5-7) (Table 1-3). Linear regression revealed that fractal dimensions of PSD decreased with the number of years since farmland conversion in the 0-10 cm soil layers (Figure 2-4) (Table 1-3).

Discussion

Effects of rehabilitation time on soil mechanical composition and erodibility

In our study, rehabilitation time was a key factor in driving changes in soil mechanical condition, erodibility and properties (Table 2) (Figure 8). The fractal dimensions of PSD, K factor of the soil in the various rehabilitation land types showed decreasing trends following rehabilitation (Figure 2-4) (Table 1-3). These changes were mainly due to the large amounts of soil nutrients released by residues and decomposing dead roots, and they promote plant growth and rehabilitation succession (Guo *et al.*, 2013). Fractal dimensions of PSD, K factor positively correlated with SOC, and the improvement of mechanical conditions were mainly explained by the soil nutrient levels (Table 3). Soil organic matter, as a binding agent, favoured soil structure stabilization and infiltration and protected it from erosion (García-Orenes *et al.* 2012). In addition, well-developed root systems played a vital role in soil mechanical functioning and actively exude substrates, such as polysaccharides, phenolic compounds, and polygalacturonic acid that affect soil particle cohesion and aggregation (Hodge *et al.* 2009; Sun *et al.*, 2016b). Furthermore, by directly binding soil particles in situ, plant roots prevented soil from being blown or washed away (Reubens *et al.*, 2007). Dense canopies and ground litter following rehabilitation protected soil aggregates from breakdown and prevent particles from being washed away by raindrop energy and runoff (Zuazo & Pleguezuelo 2009; Wang *et al.*, 2008).

Effects of rehabilitation land type on soil mechanical composition and erodibility

Vegetation rehabilitation type was the most influential factor in driving soil mechanical composition and erodibility (Table 2) (Figure 8). Variation in vegetation recovery patterns and plant traits led to variation in the production and release of soil organic matter, which affect soil crusting, splash, aggregate size and stability. For example, high levels of soil organic matter reduced soil vulnerability to detachment by surface flow, rain splash and other erosion-inducing phenomena (Xiao *et al.*, 2014; Fu *et al.*, 2015). Litters with different chemical composition among different plant species would impact decomposition rates and the release of soil organic matter (Ayres *et al.*, 2006). Thus naturally revegetated grassland site dominated with high quality litter showed faster circulating rates than shrub and tree sites dominated with relative low quality litter. This phenomenon may explain why the recovery time needed to reach significant improvements of soil mechanical conditions and soil erosion was shorter for naturally revegetated grassland than for woodland and shrub land. Our results are in accordance with the study of Yu *et al.* (2015), which showed that high concentrations of soil organic matter greatly affected the fractal dimensions of PSD and generally facilitated the improvement of soil structure. The species of naturally revegetated grassland on the whole corresponds with the native vegetation species on Loess Plateau during a long historical period (Lu *et al.*, 2003; Jiang *et al.*, 2013). They belong to Poaceae and Asteraceae families which are tolerant to drought, cold and grazing due to the characteristics of low water requirements, fibrous root

Con formato: Color de fuente: Texto 1

Eliminado: on

system, are fully suited to the local arid or semi-arid climates and soil (Lu et al., 2003; Jiang et al., 2013). Thick loess were mainly caused by the loosely cemented silt (Liu 1985; Yang & Ding 2008) which allows rainwater to infiltrate quickly (Yang et al. 2012). Thus naturally revegetated grass as native Loess Plateau vegetation were the best selected species for rehabilitation of soil conditions. Cespedes et al.,(2012) even found that afforestation with fast growing exotic species showing its negative effects on soil, compared to native grassland.

Eliminado:

In addition, vegetation alleviates erosion of soil by its canopy effectively reducing water-induced soil erosion (Mohammad & Adam 2010; Wei et al., 2010). However, that effectiveness was different from various land types. In naturally revegetated grassland site, the lower vegetation layer was more effective in reducing the kinetic energy of rainfall striking the soil surface than the tall vegetation in shrub land and woodland. Owing to the lack of roots at deep soil layer, the naturally revegetated grassland only showed positive effect at the shallow soil layer (0-10cm). However, due to the stronger stretching ability of the trees roots, woodland site also showed the potential of alleviate soil erosion at deep layer. The soil loss in orchard land was continued over a long time owing to human disturbances (such as production management and tillage practices) and the absence of surface cover protection. This leads to the breakdown of shallow soil aggregates and the washing away of soil particles by raindrop energy and runoff (Wang et al., 2008).

Effects of soil depth on soil fractal dimension and erodibility

In the analysed four types of vegetation rehabilitation, soil depth had large influences on soil mechanical composition and erodibility (Table 2) (Figure. 8) being consistent with previous studies (*Xiao et al., 2014*). In our study, the positive effects of vegetation recovery mainly occurred in the topsoil with the higher reduction rates of fractal dimensions of PSD, fractal dimensions of micro-aggregation and K factor in the topsoil than in the subsoil (Figures 2-4) (Table 1-3). This pattern was resulting from the variation in plant root distribution density decreasing along soil depth (*Reubens et al., 2007*); thus, the deeper soil layers were, the weaker the improvements of soil conditions were (*Sun et al., 2014*). In addition, soil nutrients accumulated near the soil surface due to the decomposition of vegetation litter and by influence of the biogeochemical cycling (*Wang et al., 2014*).

Conclusion

Our study suggested that vegetation rehabilitation time, type and soil depth significantly affects soil mechanical composition and erosion. Following the conversion of sloping cropland to naturally revegetated grassland, shrub land or woodland, the soil structure gradually recovered, and the resistance of the soil against erosive forces gradually increased, primarily within the topsoil. For the conversion of sloping cropland, the natural restoration process of grass represents a more efficient rehabilitation practice than does the planting of other vegetation types. The key point of success rehabilitation project is whether the selection of species fit current climatic

and geological conditions, such as the naturally revegetated grass in our study. Based on the differences of rehabilitation effectiveness among the rehabilitation land type, it is important to carefully select land types for the rehabilitation of soil conditions in the Loess Plateau. Our study, conducted at the regional scale, revealed the effects of vegetation rehabilitation on soil erosion in the Loess Plateau, China, but it strongly contributes to our understanding of the mechanisms through which rehabilitation improves soil quality and provides a suggestion for ecosystem management in arid and semi-arid regions.

Acknowledgements

We thank the anonymous referees and editors of the journal who provided valuable comments and suggestions on our manuscript. This research was funded by the National Key Research and Development Program of China (2016YFC0501707).

References

- Alagöz Z, Yilmaz E. 2009.** Effects of different sources of organic matter on soil aggregate formation and stability: A laboratory study on a Lithic Rhodoxeralf from Turkey. *Soil and Tillage Research* **103**: 419-424.
- Ayres E, Dromph KM, Bardgett RD. 2006.** Do plant species encourage soil biota that specialise in the rapid decomposition of their litter? *Soil Biology & Biochemistry* **38**: 183-186.
- Basic F, Kisic I, Mesic M, Nestroy O, Butorac A. 2004.** Tillage and crop management effects on soil erosion in central Croatia. *Soil & Tillage Research* **78**: 197-206.

416 **Blanco-Canqui H, Lal R. 2008.** Principles of soil conservation and management.

417 *Springer Science & Business Media.*

418 **Bochet E, García-Fayos P. 2004.** Factors Controlling Vegetation Establishment and

419 Water Erosion on Motorway Slopes in Valencia, Spain. *Restoration Ecology* **12**:

420 166-174.

421 **Borrelli P, Robinson DA, Fleischer LR, Lugato E, Ballabio C, Alewell C,**

422 **Meusburger K, Modugno S, Schütt B, Ferro V, Bagarello V, Oost KV,**

423 **Montanarella L, Panagos P. 2017.** An assessment of the global impact of 21st

424 century land use change on soil erosion. *Nature Communications* **8**:2013.

425 **Bremner JM, Mulvaney CS. 1982.** Nitrogen-total. Agronomy monography 9. In:

426 Page A L, Miller R H, Keeney D R. *Methods of Soil Analysis, Part 2, Chemical*

427 *and Microbial Properties, Madison: Agronomy Society of America, 595–624.*

428 **Céspedes-Payret C, Pinheiro G, Gutierrez O, Panario D. 2012.** Land use change in

429 a temperate grassland soil: Afforestation effects on chemical properties and their

430 ecological and mineralogical implications. *Science of the Total Environment* **438**:

431 549-557.

432 **Chen F, Peng BZ. 2000.** The effect of land use changes on soil conditions in arid

433 region. *Chinese Geographical Science* **10**: 226-230.

434 **Chen L, Fu GB, Huang Z, Huang Y, Gui L. 2007.** Effect of land use conversion on

435 soil organic carbon sequestration in the loess hilly area, loess plateau of China.

436 *Ecological Research* **22**: 641-648 DOI 10.1007/s11284-006-0065-1.

437 **Chen L, Wei W, Fu B, Lu Y. 2007.** Soil and Water Conservation on the Loess

Eliminado: n

439 Plateau in China: Review and Perspective. *Progress in Physical Geography* **31**:
440 389-403 DOI 10.1177/0309133307081290.

441 **Chen, X, Zhou, J. 2013.** Volume-based soil particle fractal relation with soil
442 erodibility in a small watershed of purple soil. *Environmental Earth Sciences* **70**:
443 1735-1746.

444 **Deng, L, Wang, K, Tang, Z, Shanguan Z. 2016.** Soil organic carbon dynamics
445 following natural vegetation restoration: Evidence from stable carbon
446 isotopes ($\delta^{13}\text{C}$). *Agriculture Ecosystems & Environment* **221**: 235-244.

447 **Ding L, Wang P, Zhang W, Zhang Y, Li S, Wei X, Yang F. 2019.** Shrub
448 Encroachment Shapes Soil Nutrient Concentration, Stoichiometry and Carbon
449 Storage in an Abandoned Subalpine Grassland. *Sustainability* **11**: 1732.

450 **Fu BJ, Wang YF, Lu YH, He CS, Chen LD, Song CJ.c 2015.** The effects of
451 land-use combinations on soil erosion: a case study in the Loess Plateau of China.
452 *Progress in Physical Geography* **33**: 793-804.

453 **FAO/UNESCO/ISRIC. 1988.** Soil map of the world; revised legend. *World Soil*
454 *ResourceReport*, vol. 60. Rome, Food and Agriculture Organization of the
455 United Nations.

456 **García-Orenes F, Roldán A, Mataix-Solera J, Cerdà A, Campoy M, Arcenegui V,**
457 **Caravaca F. 2012.** Soil structural stability and erosion rates influenced by
458 agricultural management practices in a semi-arid Mediterranean agro-ecosystem.

459 *Soil Use and Management* **28**: 571-579.

460 **Gao L, Bowker MA, Xu M, Sun H, Tuo D, Zhao Y. 2017.** Biological soil crusts
461 decrease erodibility by modifying inherent soil properties on the Loess Plateau,
462 China. *Soil Biology & Biochemistry* **105**: 49-58.

463 **Ghestem M, Veylon G, Bernard A, Vanel Q, Stokes A. 2014.** Influence of plant root
464 system morphology and architectural traits on soil shear resistance. *Plant and*
465 *Soil* **377**: 43-61.

466 **Guo C, Dannenmann M, Gasche R, Zeller B, Papen H, Polle A, Rennenberg H,**
467 **Simon J. 2013.** Preferential use of root litter compared to leaf litter by beech
468 seedlings and soil microorganisms. *Plant & Soil* **368**: 519-534.

469 **Hodge A, Berta G, Doussan C, Merchan F, Crespi M. 2009.** Plant root growth,
470 architecture and function. *Plant and Soil* **321**: 153-187.

471 **Hu F, Liu J, Xu C, Wang Z, Liu G, Li H, Zhao S. 2018.** Soil internal forces initiate
472 aggregate breakdown and splash erosion. *Geoderma* **320**: 43-51.

473 **Hu J, Lu Y, Fu B, Comber AJ, Harris P. 2017.** Quantifying the effect of ecological
474 restoration on runoff and sediment yields: A meta-analysis for the Loess Plateau
475 of China. *Progress in Physical Geography-Earth* **41**: 753-774.

476 **Jiang W, Cheng Y, Yang X, Yang S. 2013.** Chinese Loess Plateau vegetation since
477 the Last Glacial Maximum and its implications for vegetation restoration.
478 *Journal of Applied Ecology* **50**: 440-448.

479 **Keesstra SD, Dam OV, Verstraeten G, Huissteden JV. 2009.** Changing sediment
480 dynamics due to natural reforestation in the Dragonja catchment, SW Slovenia.
481 *Catena* **78(1)**: 60-71.

482 **Koulouri M, Giourga C. 2007.** Land abandonment and slope gradient as key factors
483 of soil erosion in Mediterranean terraced lands. *Catena* **69**: 274-281.

484 **Lian J, Zhao XY, Zuo XA, Wang SK, Wang XY, Luo YQ. 2013.** Land cover
485 changes and the effects of cultivation on soil properties in Shelihu wetland,
486 Horqin Sandy Land, Northern China. *Journal of Arid Land* **5**: 71-79.

487 **Liu, GB. 1999.** Soil conservation and sustainable agriculture on the Loess Plateau:
488 Challenges and prospects. *Ambio*: 663-668.

489 **Liu SL, Guo XD, Fu BJ, Lian G, Wang J. 2007.** The effect of environmental
490 variables on soil characteristics at different scales in the transition zone of the
491 Loess Plateau in China. *Soil Use and Management* **23**: 92-99.

492 **Liu D S. 1985.** Loess and the Environment. *China Ocean Press*, Beijing.

493 **Chen H, Shao M, Li Y. 2008.** Soil desiccation in the Loess Plateau of China.
494 *Geoderma* **143**: 91-100.

495 **Lü H, Liu D, Guo Z. 2003.** Natural vegetation of geological and historical periods in
496 Loess Plateau. *Chinese Science Bulletin* **48**: 411-416.

497 **Mazaheri MR, Mahmoodabadi M. 2012.** Study on infiltration rate based on primary
498 particle size distribution data in arid and semiarid region soils. *Arabian Journal*

499 *of Geosciences* **5**: 1039-1046.

500 **Mohammad AG, Adam MA. 2010.** The impact of vegetative cover type on runoff
501 and soil erosion under different land uses. *Catena* **81**: 97-103.

502 **Moora M, Davison J, Õpik M, Metsis M, Saks Ü, Jairus T, Vasar M, Zobel M.**
503 **2014.** Anthropogenic land use shapes the composition and phylogenetic structure
504 of soil arbuscular mycorrhizal fungal communities. *FEMS microbiology ecology*
505 **90**: 609-621.

506 **Nelson DW, Sommers LE. 1982.** Total carbon, organic carbon, and organic matter.
507 Agronomy nonograph 9. In: Page A L, Miller RH, Keeney DR. Methods of Soil
508 Analysis, Part 2, Chemical and Microbial Properties. *Madison: Agronomy*
509 *Society of America*, 539–552.

510 **Parysow P, Wang G, Gertner G, Anderson AB. 2003.** Spatial uncertainty analysis
511 for mapping soil erodibility based on joint sequential simulation. *Catena* **53**:
512 65-78.

513 **Reubens B, Poesen J, Danjon F, Geudens G, Muys B. 2007.** The role of fine and
514 coarse roots in shallow slope stability and soil erosion control with a focus on
515 root system architecture: a review. *Trees* **21**: 385-402.

516 **Song Z, Zhang C, Liu G, Dong Q, Xue S. 2015.** Fractal Feature of Particle-Size
517 Distribution in the Rhizospheres and Bulk Soils during Natural Recovery on the
518 Loess Plateau, China. *Plos One* **10**: e0138057.

519 **Su YZ, Zhao HL, Zhao WZ, Zhang TH. 2004.** Fractal features of soil particle size
520 distribution and the implication for indicating desertification. *Geoderma* **122:**
521 43-49.

522 **Sun C, Liu G, Xue S. 2016.** Land-Use Conversion Changes the Multifractal Features
523 of Particle-Size Distribution on the Loess Plateau of China. *International*
524 *Journal of Environmental Research & Public Health* **13:** 785.

525 **Sun C, Liu G, Xue S. 2016.** Natural succession of grassland on the Loess Plateau of
526 China affects multifractal characteristics of soil particle-size distribution and soil
527 nutrients. *Ecological Research* **31:** 891-902.

528 **Sun W, Shao Q, Liu J, Zhai J. 2014.** Assessing the effects of land use and
529 topography on soil erosion on the Loess Plateau in China. *Catena* **121:** 151-163

530 **Tyler SW, Wheatcraft SW. 1992.** Fractal Scaling of Soil Particle-Size Distributions:
531 Analysis and Limitations. *Soil Science Society of America Journal* **56:** 362-369 .

532 **Uchida E, Rozelle S, Xu J. 2009.** Conservation payments, liquidity constraints, and
533 off-farm labor: impact of the Grain-for-Green Program on rural households in
534 China. *American Journal of Agricultural Economics* **91:** 70-86.

535 **Uchida E, Xu J, Rozelle S. 2005.** Grain for Green: Cost-Effectiveness and
536 Sustainability of China's Conservation Set-Aside Program. *Land Economics* **81:**
537 247-264.

538 **Wang B, Liu GB, Xue S, Zhu B. 2011.** Changes in soil physico-chemical and

539 microbiological properties during natural succession on abandoned farmland in
540 the Loess Plateau. *Environmental Earth Sciences* **62**: 915-925.

541 **Wang D, Fu B, Zhao W, Hu H, Wang Y. 2008.** Multifractal characteristics of soil
542 particle size distribution under different land-use types on the Loess Plateau,
543 China. *Catena* **72**: 29-36.

544 **Wang G, Gertner G, Liu X, Anderson A. 2001.** Uncertainty assessment of soil
545 erodibility factor for revised universal soil loss equation. *Catena* **46**: 1-14.

546 **Wang HM, Wang WJ, Chen H, Zhang Z, Mao Z, Zu YG. 2014.** Temporal changes
547 of soil physic-chemical properties at different soil depths during larch
548 afforestation by multivariate analysis of covariance. *Ecology & Evolution* **4**:
549 1039-1048.

550 **Wang JG, Yang W, Yu B, Li ZX, Cai CF, Ma RM. 2016.** Estimating the influence
551 of related soil properties on macro- and micro-aggregate stability in ultisols of
552 south-central China. *Catena* **137**: 545-553.

553 **Wang XD, Liu SZ, Liu GC. 2005.** Fractal Characteristics of Soil under Different
554 Land-Use Patterns in the Arid and Semi-Arid Region of the Western Tibet
555 Plateau, China. *Wuhan University Journal of Natural Sciences* **10**: 785-790.

556 **Wei J, Zhou J, Tian JL, He XB, Tang KL. 2006** Decoupling soil erosion and human
557 activities on the Chinese Loess Plateau in the 20th century. *Catena* **68**:10-15.

558 **Wei W, Chen L, Fu B, Chen J. 2010.** Water erosion response to rainfall and land use

559 in different drought-level years in a loess hilly area of China. *Catena* **81**: 24-31.

560 **Williams JR, Jones CA, Dyke PT, 1984.** A modeling approach to determining the
561 relationship between erosion and soil productivity. *Transactions of the ASAE* **27**:
562 129-144.

563 **Xia L, Zhang GC, Heathman GC, Wang YQ, Huang CH. 2009.** Fractal features of
564 soil particle-size distribution as affected by plant communities in the forested
565 region of Mountain Yimeng, China. *Geoderma* **154**: 123-130.

566 **Xiao L, Xue S, Liu GB, Zhang C. 2014.** Fractal features of soil profiles under
567 different land use patterns on the Loess Plateau, China. *Journal of Arid Land* **6**:
568 550-560.

569 **Xu M, Wang Z, Zhao Y. 2018.** Stratification ratio of soil organic carbon as an
570 indicator of carbon sequestration and soil quality in ecological restoration.
571 *Restoration Ecology* **26**: 555-562.

572 **Xu G, Li Z, Li P. 2013.** Fractal features of soil particle-size distribution and total soil
573 nitrogen distribution in a typical watershed in the source area of the middle Dan
574 River, China. *Catena* **101**: 17-23.

575 **Yang SL, Ding ZL, Wang X, Tang ZH, Gu ZY. 2012.** Negative $\delta^{18}\text{O}$ - $\delta^{13}\text{C}$
576 relationship of pedogenic carbonate from northern China indicates a strong
577 response of C_3/C_4 biomass to the seasonality of Asian monsoon precipitation.
578 *Palaeogeography, Palaeoclimatology, Palaeoecology* **317–318**: 32–40.

579 **Yang SL, Ding ZL. 2008.** Advance-retreat history of the East-Asian summer
580 monsoon rainfall belt over Northern China during the last two glacial-interglacial
581 cycles. *Earth and Planetary Science Letters* **274**: 499–510.

582 **Yu J, Lv X, Bin M, Wu H, Du S, Zhou M, Yang Y, Han G. 2015.** Fractal features of
583 soil particle size distribution in newly formed wetlands in the Yellow River Delta.
584 *Scientific Reports* **5**: 10540.

585 **Zhang C, Xue S, Liu GB, Song, ZL. 2011.** A comparison of soil qualities of
586 different revegetation types in the Loess Plateau, China. *Plant & Soil* **347**:
587 163-178.

588 **Zhang GH, Liu GB, Wang GL, Wang, YX. 2011.** Effects of Vegetation Cover and
589 Rainfall Intensity on Sediment-Bound Nutrient Loss, Size Composition and
590 Volume Fractal Dimension of Sediment Particles. *Pedosphere* **21**: 676-684.

591 **Su ZA, Zhang JH, Nie XJ. 2010.** Effect of Soil Erosion on Soil Properties and Crop
592 Yields on Slopes in the Sichuan Basin, China. *Pedosphere* **20**: 736-746.

593 **Zheng FL. 2006.** Effect of Vegetation Changes on Soil Erosion on the Loess Plateau.
594 *Pedosphere* **16**: 420-427.

595 **Ziadat FM, Taimeh AY. 2013.** Effect of rainfall intensity, slope, land use and
596 antecedent soil moisture on soil erosion in an arid environment. *Land*
597 *Degradation & Development* **24**: 582-590.

598 **Zhu B, Li Z, Li P, Liu G, Xue S. 2010.** Soil erodibility, microbial biomass, and

599 physical–chemical property changes during long-term natural vegetation
600 rehabilitation: a case study in the Loess Plateau, China. *Ecological Research* **25**:
601 531-541.

602 **Zhuang J, Mccarthy JF, Perfect E, Mayer LM, Jastrow JD. 2008.** Soil Water
603 Hysteresis in Water-Stable Microaggregates as Affected by Organic Matter. *Soil*
604 *Science Society of America Journal* **72**: 212-220.

605 **Zhang YW, Deng L, Yan WM, Shangguan ZP. 2016.** Interaction of soil water
606 storage dynamics and long-term natural vegetation succession on the Loess
607 Plateau, China. *Catena* **137**: 52-60.

608 **Zuazo VHD, Pleguezuelo CRR. 2009.** Soil-erosion and runoff prevention by plant
609 covers: a review. *Sustainable agriculture: Springer*, 785-811.

610 **Zhou ZC, Shangguan ZP, Zhao D. 2006.** Modeling vegetation coverage and soil
611 erosion in the Loess Plateau Area of China. *Ecol Model* **198**:263-268.