

Effectiveness of water-saving technologies for the restoration of endemic *Opuntia* cacti in the Galápagos Islands, Ecuador (#38957)

1

First submission

Guidance from your Editor

Please submit by **8 Aug 2019** for the benefit of the authors (and your \$200 publishing discount).



Structure and Criteria

Please read the 'Structure and Criteria' page for general guidance.



Custom checks

Make sure you include the custom checks shown below, in your review.



Raw data check

Review the raw data. Download from the [materials page](#).



Image check

Check that figures and images have not been inappropriately manipulated.

Privacy reminder: If uploading an annotated PDF, remove identifiable information to remain anonymous.

Files

Download and review all files from the [materials page](#).

5 Figure file(s)
2 Table file(s)
1 Raw data file(s)
1 Other file(s)

! Custom checks

Field study

- ! Have you checked the authors [field study permits](#)?
- ! Are the field study permits appropriate?



Structure and Criteria

Structure your review

The review form is divided into 5 sections. Please consider these when composing your review:

1. BASIC REPORTING
2. EXPERIMENTAL DESIGN
3. VALIDITY OF THE FINDINGS
4. General comments
5. Confidential notes to the editor

 You can also annotate this PDF and upload it as part of your review

When ready [submit online](#).

Editorial Criteria

Use these criteria points to structure your review. The full detailed editorial criteria is on your [guidance page](#).

BASIC REPORTING

-  Clear, unambiguous, professional English language used throughout.
-  Intro & background to show context. Literature well referenced & relevant.
-  Structure conforms to [PeerJ standards](#), discipline norm, or improved for clarity.
-  Figures are relevant, high quality, well labelled & described.
-  Raw data supplied (see [PeerJ policy](#)).

EXPERIMENTAL DESIGN

-  Original primary research within [Scope of the journal](#).
-  Research question well defined, relevant & meaningful. It is stated how the research fills an identified knowledge gap.
-  Rigorous investigation performed to a high technical & ethical standard.
-  Methods described with sufficient detail & information to replicate.

VALIDITY OF THE FINDINGS

-  Impact and novelty not assessed. Negative/inconclusive results accepted. *Meaningful* replication encouraged where rationale & benefit to literature is clearly stated.
-  All underlying data have been provided; they are robust, statistically sound, & controlled.
-  Speculation is welcome, but should be identified as such.
-  Conclusions are well stated, linked to original research question & limited to supporting results.

Standout reviewing tips

3



The best reviewers use these techniques

Tip

Support criticisms with evidence from the text or from other sources

Example

Smith et al (J of Methodology, 2005, V3, pp 123) have shown that the analysis you use in Lines 241-250 is not the most appropriate for this situation. Please explain why you used this method.

Give specific suggestions on how to improve the manuscript

Your introduction needs more detail. I suggest that you improve the description at lines 57- 86 to provide more justification for your study (specifically, you should expand upon the knowledge gap being filled).

Comment on language and grammar issues

The English language should be improved to ensure that an international audience can clearly understand your text. Some examples where the language could be improved include lines 23, 77, 121, 128 – the current phrasing makes comprehension difficult.

Organize by importance of the issues, and number your points

1. Your most important issue
2. The next most important item
3. ...
4. The least important points

Please provide constructive criticism, and avoid personal opinions

I thank you for providing the raw data, however your supplemental files need more descriptive metadata identifiers to be useful to future readers. Although your results are compelling, the data analysis should be improved in the following ways: AA, BB, CC

Comment on strengths (as well as weaknesses) of the manuscript

I commend the authors for their extensive data set, compiled over many years of detailed fieldwork. In addition, the manuscript is clearly written in professional, unambiguous language. If there is a weakness, it is in the statistical analysis (as I have noted above) which should be improved upon before Acceptance.

Effectiveness of water-saving technologies for the restoration of endemic *Opuntia* cacti in the Galápagos Islands, Ecuador

Patricia I Tapia¹, Luka Negoita², James P Gibbs³, Patricia Jaramillo⁴

¹ School of Natural and Environmental Sciences, University of Newcastle-upon-Tyne, Tyne and Wear, United Kingdom

² Charles Darwin Research Station, Galápagos Verde 2050, Charles Darwin Foundation, Santa Cruz, Galápagos, Ecuador

³ Department of Environmental and Forest Biology, State University of New York College of Environmental Science and Forestry, Syracuse, New York, United States

⁴ Facultad de Ciencias, Universidad de Málaga, Málaga, Spain

Corresponding Author: Patricia Jaramillo
Email address: patricia.jaramillo@fcdarwin.org.ec

Restoration of keystone species is a primary strategy used to combat biodiversity loss and recover ecological services. This is particularly true for oceanic islands, which despite their small land mass, host a large fraction of the planet's imperiled species. The endemic *Opuntia* spp. cacti are one example and a major focus for restoration in the Galápagos archipelago, Ecuador. These cacti are keystone species that support much of the unique vertebrate animal community in arid zones, yet human activities have substantially reduced *Opuntia* populations. Extreme aridity poses a major obstacle for restoring *Opuntia* populations yet water-saving technologies may aid restoration efforts. The aim of this study was to evaluate current restoration efforts and the utility of two water-saving technologies as tools for restoring *Opuntia* populations in the Galápagos archipelago. We planted 1425 seedlings between 2013 and 2018, of which 66% had survived by the end of 2018. Compared with no-technology controls, seedlings planted with Groasis Waterboxx® water-saving technology (polypropylene trays with water reservoir and protective refuge for germinants) had increased survival on one island (Plaza Sur) and growth rate on four islands whereas the "Cocoon" water-saving technology (similar technology but made of biodegradable fiber) did not affect growth and actually reduced seedling survival. Survival and growth rate were also influenced by vegetation zone, altitude, and precipitation in ways largely contingent on island. Overall, our findings suggest that water-saving technologies are not always universally applicable but can substantially increase the survival and growth rate of seedlings in certain conditions, providing in some circumstances a useful tool for improving restoration outcomes for rare plants of arid ecosystems.

Effectiveness of water-saving technologies for the restoration of endemic *Opuntia* cacti in the Galápagos Islands, Ecuador

Patricia Isabela Tapia¹, Luka Negoita², James P Gibbs³, and Patricia Jaramillo^{2,4}

¹ School of Natural and Environmental Sciences, Newcastle University, Newcastle Upon Tyne, Tyne and Wear, United Kingdom

² Galápagos Verde 2050, Charles Darwin Research Station, Charles Darwin Foundation, Santa Cruz, Galápagos, Ecuador

³ Department of Environmental and Forest Biology, State University of New York, Syracuse, NY, USA

⁴ Facultad de Ciencias, Universidad de Málaga, Málaga, España

Corresponding Author:

Patricia Jaramillo^{2,4}

Charles Darwin Research Station, Charles Darwin Foundation, Santa Cruz, Galápagos, Ecuador

Email address: patricia.jaramillo@fcdarwin.org.ec

Abstract

Restoration of keystone species is a primary strategy used to combat biodiversity loss and recover ecological services. This is particularly true for oceanic islands, which despite their small land mass, host a large fraction of the planet's imperiled species. The endemic *Opuntia* spp. cacti are one example and a major focus for restoration in the Galápagos archipelago, Ecuador. These cacti are keystone species that support much of the unique vertebrate animal community in arid zones, yet human activities have substantially reduced *Opuntia* populations. Extreme aridity poses a major obstacle for restoring *Opuntia* populations yet water-saving technologies may aid restoration efforts. The aim of this study was to evaluate current restoration efforts and the utility of two water-saving technologies as tools for restoring *Opuntia* populations in the Galápagos

archipelago. We planted 1425 seedlings between 2013 and 2018, of which 66% had survived by the end of 2018. Compared with no-technology controls, seedlings planted with Groasis Waterboxx® water-saving technology (polypropylene trays with water reservoir and protective refuge for germinants) had increased survival on one island (Plaza Sur) and growth rate on four islands whereas the “Cocoon” water-saving technology (similar technology but made of biodegradable fiber) did not affect growth and actually reduced seedling survival. Survival and growth rate were also influenced by vegetation zone, altitude, and precipitation in ways largely contingent on island. Overall, our findings suggest that water-saving technologies are not always universally applicable but can substantially increase the survival and growth rate of seedlings in certain conditions, providing in some circumstances a useful tool for improving restoration outcomes for rare plants of arid ecosystems.

Introduction

The restoration of previously abundant keystone species is one way to combat loss of biodiversity and ecological services (Grime, 1998). This is particularly true on oceanic islands, which comprise little of the planet’s land mass yet host a disproportionate amount of its imperiled species (Myers et al., 2000; Campbell & Donlan, 2005). The Galápagos archipelago is a case in point: its land area is minimal (8006 km²) yet it hosts a remarkable array of endemic life forms with as many as 60% of its 168 endemic plant species now threatened with extinction (Tye, 2007; Black, 1973). Active restoration programs are underway throughout the archipelago. For example, Project Isabela (1997-2006), was the world’s largest restoration effort at the time and dedicated to eradicating introduced mammal herbivores on multiple islands in the archipelago (Carrion et al., 2011).

The *Opuntia* spp. cacti (prickly pear cactus) are a major focus for restoration in the Galápagos archipelago, Ecuador, which hosts six endemic species, with 14 total taxa when including varieties. Human impact in the Galápagos archipelago has steadily increased over the last 200 years (Jaramillo, 1998), resulting in declines of *Opuntia* populations on these islands (Snell, Snell & Stone, 1994). Several factors have been attributed as the primary threats to *Opuntias* including herbivory by introduced mammals (Snell, Snell & Stone, 1994), extinction of keystone predators that once regulated numbers of cactivores (Sulloway & Noonan, 2015), and

the increased intensity of El Niño events (Snell, Snell & Stone, 1994; Hicks & Mauchamp, 1996). *Opuntia* cacti provide many ecosystem services for other native and endemic species (Grant & Grant, 1981; Hicks & Mauchamp, 1995, 1996; Gibbs, Marquez & Sterling, 2008). Examples include Galápagos giant tortoises and land iguanas that depend on *Opuntia* cacti as a food source while also contributing to *Opuntia* regeneration through seed dispersal (Hamann, 1993; Snell, Snell & Stone, 1994; Gibbs, Marquez & Sterling, 2008; Gibbs, Sterling & Zabala, 2010). Efforts are being made to protect and restore populations of these imperiled cacti (Hicks & Mauchamp, 1996) but it is not clear which factors most control *Opuntia* populations (Sullo way & Noonan, 2015). *Opuntia* declines on Plaza Sur Island, for example, are especially pronounced and regeneration remains low despite goat eradication in the 1960s (Snell, Snell & Stone, 1994). Nonetheless, active planting of these species is critical for preventing extinction until their threats are better understood and eliminated.

Severe aridity poses a major obstacle for restoring plant communities over much of Galápagos, including the restoration of xerophytes such as cacti. The lowland zones of the archipelago, where *Opuntias* are most common and historically abundant (Browne et al., 2003), can receive less than 10 cm rainfall annually (Trueman & d'Ozouville, 2010). “Water-saving” technologies are tools that may help increase survival and growth of planted cactus seedlings while reducing the need for manual watering in these arid environments of the Galápagos (Jaramillo, 2015; Jaramillo, Cueva, Jiménez, & Ortiz, 2014; Jaramillo et al., 2015; Hoff, 2014; Jaramillo, Tapia, & Gibbs, 2018; Peyrusson, 2018, Faruqi et al., 2018; Kulkarni, 2011). Although these technologies show much promise, there remains a dearth in formal scientific studies evaluating their efficacy (but see Liu, Li, & Ren, 2014). Therefore, the objective of the current study was to determine the success of current restoration efforts and evaluate the utility of two water-saving technologies as tools for restoring *Opuntia* populations in the Galápagos archipelago. Through this objective we hope to better understand the utility of water-saving technologies for restoring these and other keystone plant species in arid island ecosystems throughout the world.

Materials & Methods

Study Area, Focal Species, and Water-saving Technologies

The Galápagos archipelago is located in the Pacific Ocean, about 1000 km west of the coast of mainland Ecuador (1°39' N, 92°0' W to 1°26' S, 89°14' W, WGS 84, Fig. 1) (DPNG, 2014). Our study focused on measuring the utility of water-saving technologies for enhancing cactus growth and survival of four endemic *Opuntia* taxa within the archipelago: *Opuntia echios* var. *echios* Howell, *Opuntia echios* var. *gigantea* Howell, *Opuntia megasperma* var. *megasperma* Howell, and *Opuntia megasperma* var. *orientalis* Howell (Hicks & Mauchamp, 1996). The water-saving technologies used in this study function by sheltering seedlings and ground around them from the heat of the sun while storing and providing water. We evaluated two technologies: Groasis Waterboxx® (Groasis), a protective polypropylene box that collects rainwater that it provides to the plant (Hoff, 2014); and the Cocoon system, a 99% biodegradable box that contains and provides water to the plant similar to Groasis, but Cocoon is only filled with water at the time of planting (Land Life Company, 2015; Faruqi et al., 2018). These water-saving technologies have been proposed as a tool to assist plant restoration of *Opuntia* taxa via “Galápagos Verde 2050” (GV2050), a project started by the Charles Darwin Foundation in 2013 with the mission of restoring degraded ecosystems and aiding with sustainable agriculture in the Galápagos archipelago (Jaramillo et al., 2014, 2015, 2017). GV2050 seeks to restore ecosystems by using a data-informed experimental approach for understanding the best conditions, mechanisms, and tools for successful plantings of native and endemic species (Jaramillo et al., 2015).

Planting and Data Collection

A total of 1425 total cacti (1137 *Opuntia echios* var. *echios*, 68 *Opuntia echios* var. *gigantea*, 24 *Opuntia megasperma* var. *megasperma*, and 196 *Opuntia megasperma* var. *orientalis*) were planted on six islands (Baltra, Española, Floreana, Plaza Sur, San Cristóbal, and Santa Cruz) between 2013 and 2018 (Table 1). Permission to plant Opuntias within protected sites on these islands was granted by the Dirección del Parque Nacional Galápagos (DPNG) through permit number PC-11-19 (Table 2). To evaluate the factors most important for successful *Opuntia* restoration data were used only from *Opuntias* that were grown from seed and planted using either Groasis, Cocoon, or control (no technology) treatments on Floreana, Santa Cruz, Baltra, and Plaza Sur islands yielding a sample of 1029 *Opuntia* individuals of three taxa (Table 1).

Planting sites on each island were selected based on locations where historic *Opuntia* populations were known to have thrived but are now in decline (Sulloway & Noonan 2015; Sulloway et al., 2013; Table 2). Seedlings were sown from seeds collected in each respective planting location using standardized seed collection and stratification techniques and grown for one year at the Charles Darwin Research Station, Santa Cruz Island, before transferring to each island (Jaramillo, 2019; Jaramillo, Tapia & Gibbs, 2017). Each seedling was randomly assigned a treatment of either control (no technology), Groasis, or Cocoon, ensuring a representative sample of replicates within each treatment and site. The number of controls was maintained at one control for every five Groasis or Cocoon technology treatment replicates. Plantings were conducted according to established methods for installing Groasis, Cocoon, and controls (Jaramillo et al., 2017). Wire fences were secured and maintained around each individual planting on Plaza Sur and Baltra islands to prevent land iguana herbivory. Planting site covariates were recorded at time of planting: altitude (elevation), soil type (rocky sand, rocky clay, rich clay, and rich sandy clay), vegetation zone (arid, littoral, and transitional; Johnson & Raven, 1973), and treatment (control, Groasis, and Cocoon). Growth (vegetative height) and qualitative plant state (“good,” “regular,” “poor,” and dead) were noted during each repeated visit approximately every six weeks following planting.

Two measures were used to evaluate restoration success (Menendez & Jaramillo, 2015). Two-year survival was quantified as whether or not a seedling survived for at least two years after planting—the period of greatest mortality risk. Only seedlings planted before 2017 (at least two years since planting) were included in that analysis. Relative growth rate was also calculated based on the vegetative height of each seedling over time. Whereas survival is the primary metric for establishing success of population restoration, growth rate can indicate the speed of ecosystem recovery due to the rate of increase in the biomass of a keystone species (Grime, 1998). An additional environmental covariate of total precipitation across the six months following planting was compiled based on available climate data from 2013 to 2018 (Trueman & D’Ozouville, 2010; CDF, 2018).

Data analysis

All statistical analyses were conducted using the R statistical software package v3.3.3 (R Core Team, 2017). To test the overall effect of water-saving technologies on the restoration of

Opuntia cacti, a model comparison approach was implemented using fixed- and mixed-effects regression models of the form:

2-year survival logistic fixed-effect model:

$$\begin{aligned} 2YearSurvival \\ = \alpha + \beta_1 \times treatment + \beta_2 \times 6MonthPrecip + \beta_3 \times Zone + \beta_4 \times Altitude \\ + \beta_5 \times island \end{aligned}$$

Relative growth rate linear mixed-effect model:

$$\begin{aligned} \log(RGR) \\ = \alpha + \beta_1 \times treatment + \beta_2 \times 6MonthPrecip + \beta_3 \times SoilType + \beta_4 \times Zone + \\ \beta_5 \times Altitude + \beta_6 \times PlantAge + \beta_6 \times island + N(0, \sigma^2_{PlantID}) \end{aligned}$$

The growth rate model is a general linear mixed-effects regression fit using the ‘lme4’ package (Bates et al., 2015). Relative growth rate (RGR) was calculated as the relative rate of increase in height over time and was log-transformed after adding one to meet assumptions of normality. Plant age was included in the model to account for the fact that RGR changes as seedlings get older. Plant ID is included as a random effect. Random effects account for within-group correlation that results from non-independent data points (Pinheiro & Bates, 2000). For example, our growth data are based on repeated measures of each individual plant, which means that growth measurements are not independent within an individual plant. The random effect for Plant ID allows us to include all observations in our analysis by accounting for this non-independence. The two-year survival model tested the overall survival of each seedlings two years after planting and was fit using a generalized linear model function with a binomial family logit function in the ‘base’ package (R Core Team, 2017). Because only one data point was available for each plant, the lower sample size required a simpler model in which soil type was removed in order to allow the model to converge successfully and no random effects were necessary. These models were then compared to null models using the likelihood-ratio to test for the effect of treatment on growth rate and survival. Null models were the same as the models listed above except for the exclusion of technology treatment. A significant difference between the two models indicates that the variable that was excluded (i.e., treatment) is a significantly important predictor.

We examined the relative effect of each variable within the growth rate and survival models to assess the relative importance of technologies as well as other environmental factors such as soil type and altitude. All continuous variables in our models were standardized by subtracting the mean and dividing by two times the standard deviation in order to relativize the effect of each variable coefficient on growth rate and two-year survival (Gelman, 2008). Confidence intervals (95%) for each coefficient in each full model were then bootstrapped using the 'boot' package in R (Canty, 2017) and plotted for visual comparison. P-values were generated using the Satterthwaite method in the 'lmerTest' package in R (Kuznetsova et al., 2017). P-values generated from mixed-effect models are not always accurate, but we include these values for the sake of highlighting the degree to which variables differ in their relative importance. Furthermore, all significance values generated in this way were consistent with bootstrapped confidence interval results. Coefficients for logistic models were back-transformed to odds ratio by exponentiating and subtracting one. In this way the coefficient values can be interpreted as the proportional effect of each variable on increasing (or decreasing if negative) the probability of two-year survival. Each model was fit using data from all four islands included in the analysis (Baltra, Floreana, Santa Cruz, and Plaza Sur), but due to high control treatment mortality on Plaza Sur, the models were also tested using data that *excluded* Plaza Sur as well as using data *exclusively from* Plaza Sur. When testing with data exclusively from Plaza Sur, "island" was removed from the models and treatment type consisted of only Groasis and controls because no Cocoons were used on Plaza Sur. Finally, the current state of all planted individuals included in the analysis (up through 2018) was plotted as stacked bar plots to visualize rates of survival between islands and treatments.

Results

General outcomes

Of the 1425 *Opuntia* spp. individuals planted between 2013 and 2018, (most plantings were made in 2015 and 2016, Fig. 2), 943 *Opuntias* remained alive by the end of 2018 (66% overall survival, Fig. 2). On Plaza Sur, 737 *Opuntia* individuals were planted between 2015 and 2018 with 452 survivors by the end of 2018 (an increase of 135% from the last recorded population estimates of 334 in 2014; Fig. 3). Survival of seedling plantings on Plaza Sur was 26.8% ($n = 82$)

for controls and 62.2% ($n = 519$) for Groasis (Fig. 4a). Survival of seedling plantings on Floreana was 66.7% ($n = 3$) for controls and 31.2% ($n = 16$) for Groasis (Fig. 4b). Survival of seedling plantings on Baltra was 79.7% ($n = 74$) for controls, 45% ($n = 20$) for Cocoon, and 65.5% ($n = 255$) for Groasis (Fig. 4c). Survival of seedlings planted on Santa Cruz was 77.8% ($n = 9$) for controls, 27.8% ($n = 18$) for Cocoon, and 72.7% ($n = 33$) for Groasis (Fig. 4d).

Outcomes across all islands

Model comparisons: Treatment type (Groasis, Cocoon, or Control) was associated with growth rate of *Opuntia* species ($\chi^2 (2) = 60.77$, $P < 0.001$) and two-year survival rate of *Opuntia* seedlings ($\chi^2 (2) = 154.73$, $P < 0.001$). In the two-year survival logistic regression, altitude (1.14, $P < 0.001$), littoral zone (14.91, $P < 0.001$), transitional zone (13.17, $P = 0.035$), and six-month precipitation (-0.38, $P = 0.004$) had odds ratios with confidence intervals that did not overlap zero (Fig. 5a). Groasis technology had a positive odds ratio of 0.73 ($P < 0.001$), while Cocoon had a negative odds ratio of -0.95 ($P < 0.001$) (Fig. 5a). In the growth rate regression, littoral zone (0.48, $P < 0.001$), plant age (-0.53, $P < 0.001$), and six-month precipitation (0.25, $P = 0.031$) all had effect sizes with confidence intervals that did not overlap zero (Fig. 5b). Groasis technology had a positive effect size with a coefficient of 0.54 ($P < 0.001$), while Cocoon had an insignificant coefficient ($P = 0.160$) (Fig. 5b).

Outcomes on Plaza Sur Island only

Model comparisons: On Plaza Sur Island, treatment type (Groasis or Control) was associated with growth rate of *Opuntia* species ($\chi^2 (1) = 18.92$, $P = 0.001$) and two-year survival rate of *Opuntia* seedlings ($\chi^2 (1) = 57.93$, $P < 0.001$). In the two-year survival logistic regression, littoral zone (310.5, $P < 0.001$), altitude (1.32, $P < 0.001$), and six-month precipitation (-0.62, $P < 0.001$) had odds ratios with confidence intervals that did not overlap zero (Fig. 5c). Groasis technology had a positive odds ratio of 3.19 ($P < 0.001$) (Fig. 5c). In the growth rate regression, littoral zone (0.49, $P < 0.001$), plant age (-0.3, $P < 0.001$), six-month precipitation (-0.22, $P = 0.001$), and altitude (0.18, $P = 0.012$) all had effect sizes with confidence intervals that did not overlap zero (Fig. 5d). Groasis technology had a positive effect size with a coefficient of 0.46 ($P < 0.001$) (Fig. 5d).

Outcomes on all islands excluding Plaza Sur

Model comparisons: Treatment type (Groasis, Cocoon, or Control) was associated with growth rate of *Opuntia* species ($\chi^2 (2) = 23.62$, $P < 0.001$), but not with two-year survival rate of *Opuntia* seedlings ($\chi^2 (2) = 43.31$, $P > 0.001$). In the two-year survival logistic regression, transition zone (-0.99 , $P < 0.001$) had a negative odds ratio with confidence intervals that did not overlap zero (Fig. 5e). Both Groasis and Cocoon technologies had significant negative odds ratios of -0.31 ($P = 0.034$) and -0.9 ($P > 0.001$) respectively (Fig. 5e). In the growth rate regression, plant age (-0.78 , $P < 0.001$) and six-month precipitation (0.51 , $P < 0.001$) had effect sizes with confidence intervals that did not overlap zero (Fig. 5f). Groasis technology had a positive effect size with a coefficient of 0.45 ($P < 0.001$), while cocoon had an insignificant coefficient ($P > 0.05$) (Fig. 5f).

Discussion

Water-saving technologies enhanced survival and growth of *Opuntia* plantings, but benefits of these technologies were highly contingent upon planting environment. For example, Groasis technology was effective at increasing growth rate across islands overall, but was only effective at aiding survival on Plaza Sur Island where Groasis increased the probability of two-year survival of seedlings more than three-fold (319%) (Fig. 5). Cocoon technology, however, provided no improvement in growth rate and actually reduced probability of two-year survival of seedlings by 95% overall (Fig. 5). Altogether, our *Opuntia* restoration efforts have increased the population of *Opuntia* spp. in the Galápagos archipelago by 943 individuals (66% survival of 1425 plantings), more than doubling the population of *Opuntia* cacti on Plaza Sur Island, from 334 to 786 in just four years (Fig. 3).

These results emphasize the species- and site-specific contingencies of applying water-saving technologies for plant restorations. For example, Cocoon technology did not provide any advantage when planting *Opuntias* in the Galápagos archipelago. This is despite the fact that in other systems and with other species Cocoon has been shown to increase survival rates in planted trees from 0-20% to 75-95% (Faruqi et al., 2018). One possible explanation is that *Opuntia* cacti have a short initial rooting depth compared to other species (Snyman, 2005), and this may reduce access to the water available from the Cocoon (Land Life Company, 2015). *Acacia macracantha*,

for example, has much deeper roots and has had much greater success when planted with Cocoon technology in the Galápagos (GV2050, *unpublished data*).

Although Groasis technology helped increase growth rate of *Opuntias* overall, it had a clear, positive effect on the survival of *Opuntias* only on Plaza Sur Island. A likely factor contributing to this is that compared to other islands, the majority of *Opuntias* were planted on Plaza Sur preceding the greatest period of drought in the Galápagos over the last five years (Appendix 1; CDF, 2018). Despite fairly regular seasonal patterns of water availability in the Galápagos (Snell & Rea, 1999; Restrepo et al., 2012), there remains much variability, especially ~~that~~ caused by El Niño events (Trueman & d'Ozouville, 2010). In this way Groasis may have the greatest advantage when ensuring water availability for *Opuntias* during periods of especially severe drought. *Opuntia* cacti are typically more resistant to desiccation and water stress compared to other species that do not have physiological adaptations for surviving low-water desert conditions (Racine & Downhower, 1974; Dubrovsky, 1998), and this may explain why Groasis was only effective for *Opuntia* cacti under extreme drought. These findings support the idea that water availability for *Opuntias* plays less of a role in survival than previously assumed (Coronel, 2002; Jaramillo, Tapia & Gibbs, 2018; Racine & Downhower, 1974). This does not negate the value of the Cocoon or Groasis technology for restoration overall, but rather presents the important observation that water-saving technologies such as Cocoon and Groasis should be considered on a case-by-case basis and tested with each species and in different environmental conditions before making expansive planting efforts. Groasis technology may provide a form of insurance for the unpredictability of extreme drought events and the benefits of using Groasis technology may in some cases outweigh the costs in the long run.

Site co-variables also affected *Opuntia* survival and growth. In particular, vegetation zone, altitude, and precipitation were important predictors of *Opuntia* survival and growth but as with water-saving technologies, these effects were highly contingent on island. *Opuntias* had a greater survival and growth rate in the littoral vegetation zone on Plaza Sur but had greater survival in the arid vegetation zone on other islands. This effect may be due to an interaction between environmental and biotic factors unique to Plaza Sur or other islands. For example, Plaza Sur has especially high land iguana densities speculated to be due to the loss of its main predator from the island, the Galápagos hawk (Sulloway & Noonan, 2015). This high herbivore density may

help keep invasive plant species in check on Plaza Sur—species that may otherwise shade out *Opuntia* seedlings on other islands (Schofield, 1973, Hicks & Mauchamp, 1996, 2000).

Surprisingly, six-month precipitation did not have a positive effect on seedling survival in any of our analyses, and actually decreased survival of seedlings planted on Plaza Sur. This finding contradicts conclusions from previous work by Coronel (2002) who found that precipitation during the six months following planting was an important factor for *Opuntia* survival. Coronel (2002), however, found that the negative effect of desiccation was mostly evident in *Opuntias* grown from cladodes rather than seeds as in the current analysis. Furthermore, most seedlings were planted on Plaza Sur at the start of a long period of drought so there was not as much variation in precipitation on Plaza Sur seedlings to fully test its effects. Altitude was only a significant predictor of survival and growth rate on Plaza Sur (Fig. 5). This may be in part because altitude is closely associated with vegetation zone, and this can account for some of the altitude effect. That said, it is not clear what is driving the positive effect of altitude on Plaza Sur. Although littoral zone has a positive impact on survival and growth, seedlings that are too low in elevation are more exposed to ocean salt spray which can increase seedling mortality (Boyce, 1954). Soil type had no significant effects on growth rate (Fig. 5), suggesting that, at least for *Opuntias*, substrate is of less importance for growth rate than factors such as vegetation zone or altitude. The effect of soil type on survival could not be tested with the current data due to limitations in sample size.

The observational aspects of our study have some inherent limits. Although it seems likely that extreme drought was the primary driver of control treatment seedling mortality on Plaza Sur, other effects cannot be ruled out. Plaza Sur is a small island (the smallest island by far of the four in this analysis: only 13 ha, with the next larger being Baltra at 2100 ha), which could increase the exposure of seedlings to salt spray, exposure to sea lion activity, as well as a suite of other effects associated with small islands (Lomolino & Weiser, 2001). It may also be that the high concentration of land iguanas and sea lions (Jaramillo pers. obs.) has impacted the edaphic environment of the island through their excrement as can be common on seabird islands (Rajakaruna et al., 2009). Thus, the small area and low variation in altitude, precipitation, and vegetation zones associated with Plaza Sur plantings suggests that any significant effect of these factors within Plaza Sur be taken cautiously when generalizing to *Opuntia* restoration beyond this island. The experimental treatments of the study involving water-saving technologies,

however, do suggest that extreme drought is the most probable hypothesis for the high control mortality on Plaza Sur. Another important caveat is that taxon effects are confounded with island effects. With one exception, each island had a particular species or variety of *Opuntia* (Table 1). It is possible that some of the island-based differences are actually due to slightly different environmental requirements of the *Opuntia* taxa used in this study.

In conclusion, this study underlines the importance of considering the specific circumstances and methodologies that affect successful restoration. Water-saving technologies such as the Groasis Waterboxx® and Cocoon are promising systems for restoring species in arid environments but should not be assumed to function equally well in all environments and with all species. Even within one system, as in the current study, the benefits of Groasis vary tremendously and likely depend on the precipitation available following plantings. It is possible that species already adapted for low water conditions, such as cacti, have a much lower threshold at which Groasis or other water-saving technologies provide a benefit. Future evaluations of these technologies should monitor precipitation to test whether there is a threshold level of drought where these technologies become more effective. In some cases and for some species there may be no threshold for effective use as with the Cocoon technology for *Opuntias*. Preliminary plantings coupled with extensive environmental and experimental data collection is essential before large-scale planting efforts are initiated with water-saving technologies. The Galápagos Verde 2050 project of the Charles Darwin Foundation presents a model for data-informed adaptive management and conservation. We hope this model may inspire other restoration efforts to adopt similar data-informed approaches. Continued monitoring and accounting for context-specific contingencies in restoration work is essential (Cabin, 2007) and future restoration efforts should continually adapt management protocols based on current results (Parma et al., 1998).

Acknowledgments

This is the work of the entire Galápagos Verde 2050 project team of the Charles Darwin Foundation, particularly María Guerrero, Esme Plunkett, and Paúl Mayorga. Assistance and advice was also provided by: Jorge Carrión, Danny Rueda, Jeffreys Málaga, Milton Chugcho, Rafael Chango, Jibson Valle, Francisco Calva, Edie Rosero and Francisco Moreno from DPNG. Novarino Castillo provided valuable field assistance. Institutional support was provided by

DPNG (Dirección del Parque Nacional Galápagos), ECOGAL (Aeropuerto Ecológico de Baltra), FAE (Fuerza Aérea Ecuatoriana), ABG (Agencia de Regulación y Control de la Bioseguridad y Cuarentena para Galápagos), GAD (Gobiernos Autónomos Descentralizados from Floreana and Santa Cruz). Washington Tapia and Felipe Cruz † provided critical advice and encouragement. Steve Rushton and Theodora Commandeur provided valuable comments on an earlier version of the manuscript.

Funding

Funding was provided by the ComON Foundation, The Leona M and Harry B. Helmsley Charitable Trust, and the BESS Forest Club.

References

- Aldridge, C.L., Boyce, M.S. & Baydack, R.K. (2004). Adaptive management of prairie grouse: how do we get there? *Wildlife Society Bulletin* 32: 92-103.
- Bates, D., Mächler, M., Bolker, B. & Walker, S. (2015). Fitting linear mixed-effects models using {lme4}. *Journal of Statistical Software* 67: 1-48.
- Bensted-Smith, R. (2002). A biodiversity vision for the Galapagos Islands. Charles Darwin Foundation and World Wildlife Fund, Puerto Ayora, Galápagos.
- Bisconti, M., Landini, W., Bianucci, G., Cantalamessa, G., Carnevale, G., Ragaini, L., & Valleri, G. (2001). Biogeographic relationships of the Galápagos terrestrial biota: Parsimony analyses of endemism based on reptiles, land birds and Scalesia land plants. *Journal of Biogeography* 28: 495-510.
- Black, J. (1973). Galápagos archipiélago del Ecuador. Quito, Ecuador: Imprenta Europa.
- Boyce, S.G. (1954). The salt spray community. *Ecological Society of America* 24: 29–67.
- Browne, B.R.A., Anderson, D.J., White, M.D., Johnson, M.A. (2003). Evidence for low genetic divergence among Galápagos *Opuntia* cactus species. *Noticias de Galápagos* 62: 11–15.
- Cabin, R. J. (2007). Science-driven restoration: a square grid on a round earth? *Restoration Ecology* 15: 1–7.
- Campbell, K. & Donlan, C.J. (2005). Feral goat eradication on islands. *Conservation Biology* 19: 1362–1374.
- Campbell, K., Baxter, G.S., Murray, P.J., Coblenz, B.E. & Donlan, C.J. (2007) Development of

397 a prolonged estrus effect for use in Judas goats. *Applied Animal Behaviour Science* 102:12-
398 23.

399 Canty, A. & Ripley, B. (2017). Boot: bootstrap R (S-Plus) functions. R package version 1.3-19.

400 Cardinale, B.J., Duffy, J.E., Gonzalez, A., Hooper, D.U., Perrings, C., Venail, P., Narwani, A.,
401 Mace, G.M., Tilman, D., Wardle, D.A., Kinzig, A.P., Daily, G.C., Loreau, M., Grace, J.B.,
402 Larigauderie, A., Srivastava, D.S. & Naeem, S. (2012). Biodiversity loss and its impact on
403 humanity. *Nature* 486: 59–67.

404 Carrion V., Donlan C.J., Campbell K.J., Lavoie C., Cruz F. 2011. Archipelago-Wide Island
405 Restoration in the Gala Islands : Reducing Costs of Invasive Mammal Eradication Programs
406 and Reinvasion Risk. *PLOS One* 6: 1–7.

407 Charles Darwin Foundation (2018). Charles Darwin Foundation meteorological database – base
408 de datos meteorológicos de la FCD. Online data portal - portal de datos en línea:
409 <http://www.darwinfoundation.org/datazone/climate/> Last updated December 31, 2018.

410 Chapin, F.S., Zavaleta, E.S., Eviner, V.T., Naylor, R.L., Vitousek, P.M., Reynolds, H.L.,
411 Hooper, D.U., Lavorel, S., Sala, O.E., Hobbie, S.E., Mack, M.C. & Díaz, S. (2000).
412 Consequences of changing biodiversity. *Nature* 405: 234-42.

413 Coronel, V. (2002). Distribución y re-establecimiento de *Opuntia megasperma* var. *Orientalis*
414 Howell. (Cactaceae) en punta Cevallos, Isla Española-Galápagos. Facultad de Ciencia y
415 Tecnología, Escuela de Biología del Medio Ambiente. Universidad del Azuay, Cuenca,
416 Azuay, Ecuador

417 Cruz, F., Donlan, C., Campbell, K., & Carrión, V. (2005). Conservation action in the Galápagos:
418 feral pig (*Sus scrofa*) eradication from Santiago Island. *Biological Conservation* 121:
419 473-478.

420 Cruz, F., Carrion, V., Campbell, K.J., Lavoie, C. & Donlan, C.J. (2009) Bio-economics of large-
421 scale eradication of feral goats from Santiago Island, Galápagos. *Journal of Wildlife*
422 *Management* 73: 191-200.

423 Defaa, C., Achour, A., Mousadik, A. & Msanda, F. (2015) Effets de l’hydrogel sur la survie et la
424 croissance des plantules d’arganier sur une parcelle de régénération en climat aride.
425 *Journal of Applied Biosciences* 92: 8586 – 8594.

426 Dirección del Parque Nacional Galápagos (2014). Plan de manejo de las áreas protegidas de
427 Galápagos para el buen vivir. Dirección del Parque Nacional Galápagos, Puerto Ayora,
428 Galápagos, Ecuador.

- 429 Dubrovsky, J.G., North, G.B. & Nobel, P.S. (1998). Root growth, developmental changes in the
430 apex, and hydraulic conductivity for *Opuntia ficus-indica* during drought. *New*
431 *Phytologist* 138: 75-82.
- 432 Eckhard, R.C. (1972). Introduced plants and animals in the Galápagos Islands. *BioScience* 22:
433 585-590.
- 434 Faruqi, S., Wu, A., Brolis, E., Anchondo, A., & Batista, A. (2018). The business of planting
435 trees: a growing investment opportunity. World Resources Institute and The Nature
436 Conservancy. Retrieved from: <https://www.wri.org/publication/business-of-planting-trees>
- 437 Ferwerda, W.H. (2015). 4 returns, 3 zones, 20 years: a holistic framework for ecological
438 restoration by people and business for next generations. RSM/IUCN CEM.
- 439 Ferwerda, W. & Moolenaar, S. (2016). Four returns: a long-term holistic framework for integrated
440 landscape management and restoration involving business. *Solutions* 7: 36-41.
- 441 Gelman, A. (2008). Scaling regression inputs by dividing by two standard deviations. *Statistics*
442 *in Medicine* 27: 2865-2873.
- 443 Gibbs, J. P., Marquez, C., & Sterling, E. J. (2008). The role of endangered species reintroduction
444 in ecosystem restoration: tortoise–cactus interactions on Española Island, Galápagos.
445 *Restoration Ecology* 16: 88-93.
- 446 Gibbs, J.P., Sterling, E.J., Zabala, F.J. (2010). Giant tortoises as ecological engineers: A long-
447 term quasi-experiment in the Galápagos Islands. *Biotropica* 42: 208-214.
- 448 González-Pérez, F. & Cubrero-Pardo, P. (2010). Efecto de las actividades turísticas sobre el
449 comportamiento de fauna representativa de las Islas Galápagos, Ecuador. *Latin American*
450 *Journal of Aquatic Research* 38: 493-500.
- 451 Grant, B.R. & Grant, P.R. (1981). Exploitation of *Opuntia* cactus by birds on the Galápagos.
452 *Oecologia* 49: 179–187.
- 453 Grime, J. (1998). Benefits of plant diversity to ecosystems: immediate, filter and founder effects.
454 *Journal of Ecology* 86: 902-910.
- 455 Hamann, O. (1993). On vegetation recovery, goats and giant tortoises on Pinta Island,
456 Galápagos, Ecuador. *Biodiversity and Conservation* 2: 138-151.

- Hanson, C., Campbell, K. (2013). Floreana Island ecological restoration: rodent and cat eradication feasibility analysis version 6.0. Island Conservation, Santa Cruz, California, 85 pp.
- Helsen, P., Verdyck, P., Tye, A., Desender, K., Van Houtten, N., & Van Dongens, S. (2007). Isolation and characterization of polymorphic microsatellite markers in Galápagos prickly pear (*Opuntia*) cactus species. *Molecular Ecology Notes* 7: 454-456.
- Hicks, D.J. & Mauchamp, A. (1995). Size-dependent predation by feral mammals on Galápagos *Opuntia*. *Noticias de Galápagos* 53: 26-28.
- Hicks, D.J. & Mauchamp, A. (1996). Evolution and conservation biology of the Galápagos opuntias (Cactaceae). *Haseltonia* 4: 89-102.
- Hicks, D.J. & Mauchamp, A. (2000). Population structure and growth patterns of *Opuntia echios* var. *gigantea* along an elevational gradient in the Galápagos Islands. *Biotropica* 32: 235-243.
- Hoff, P. (2014). Groasis technology: manual de instrucciones de plantación, pp. 1-27.
- Jaramillo, P. (1998). Impact of human activities on the native plant life in Galápagos National Park. In Ospina, P. & Muñoz, E. (ed.) *Galápagos Report*. Fundación Natura and World Wildlife Fund, Quito, pp. 50–5.
- Jaramillo, P. (2015). Water-saving technology: the key to sustainable agriculture and horticulture in Galápagos to BESS Forest Club. Galápagos Verde 2050, Charles Darwin Foundation, pp. 1-12.
- Jaramillo, P., Cueva, P., Jiménez, E., & Ortiz, J. (2014). Galápagos Verde 2050. Charles Darwin Foundation, Puerto Ayora, Galápagos, Ecuador.
- Jaramillo, P., Lorenz, S., Ortiz, G., Ortiz, J., Rueda, D., Gibbs, J.P., & Tapia, W. (2015). Galápagos Verde 2050: An opportunity to restore degraded ecosystems and promote sustainable agriculture in the Archipelago. Biodiversity and Ecosystem Restoration: GNPD, GCREG, CDF, and GC. In Cayot, L., Cruz, D. & Knab, R. (ed.). *Galápagos Report 2013 – 2014*, pp. 133-143.
- Jaramillo, P., Tapia, W., & Gibbs, J.P. (2017). Action plan for the ecological restoration of Baltra and Plaza Sur Islands. *Charles Darwin Foundation*, pp. 1-56.
- Jaramillo, P., Tapia, W., & Gibbs, J.P. (2018). Galápagos Verde 2050: restauración ecológica de ecosistemas degradados y agricultura sostenible utilizando tecnologías ahorradoras de agua. *In prep*. Fundación Charles Darwin, Puerto Ayora, Isla Santa Cruz.

Jaramillo, P., Tapia, W., & Tye, A. (2018). *Opuntia megasperma* var. *orientalis* Howell. In FCD & WWF-Ecuador (ed.) *Atlas de Galápagos, Ecuador*. Quito, Ecuador: FCD & WWF-Ecuador, pp. 58-59.

Jaramillo, P. (2019). Manual de procesos y protocolos del proyecto Galápagos Verde 2050. *In prep.* Fundación Charles Darwin, Puerto Ayora, Galápagos, Ecuador.

Johnson, M.P., Raven, P.H. (1973). Species number and endemism: The Galápagos archipelago revisited. *Science* 179: 893-895.

Kulkarni, S. (2011). Innovative technologies for water saving in irrigated agriculture. *International Journal of Water Resources and Arid Environments* 1: 226-231.

Kuznetsova, A., Brockhoff, P.B. & Christensen, R.H.B. (2017). lmerTest Package: tests in linear mixed effects models. *Journal of Statistical Software* 82: 1-26.

Land Life Company (2015). Benefits of the COCOON technology. Retrieved from <https://landlifecompany.com>

Lavoie C, Donlan C, Campbell K, Cruz F, Carrión V (2007). Geographic tools for eradication programs of insular non-native mammals. *Biological Invasions* 9:139-148.

Larson, E.R., Howell, S., Kareiva, P., Armsworth, P.R. (2016). Constraints of philanthropy on determining the distribution of biodiversity conservation funding. *Conservation Biology* 30: 206-215.

Liu M, Zhiyang L, Ren W (2014) Research on the effect of waterboxx technology on  **Haloxylon** **ammodendron** afforestation in arid and semiarid areas. *Advanced Science, Engineering and Medicine* 6: 236-239.

Lomolino, M.V. & Weiser, M.D. (2001). Towards a more general species-area relationship diversity on all islands, great and small. *Journal of Biogeography* 28: 431-445.

McDonald-Madden, E., Baxter, P.W.J. & Possingham, H.P. (2008). Making robust decisions for conservation with restricted money and knowledge. *Journal of Applied Ecology* 45: 1630-1638.

Menendez, Y., & Jaramillo, P. (2015). Aplicación Android y plataforma virtual del proyecto Galápagos Verde 2050.

Myers, N., Mittermeier, R.A., Mittermeier, C.G., da Fonseca, G.A.B., Kent, J. (2000). Biodiversity hotspots for conservation priorities. *Nature* 403:853-858.

Parma, A.M. & NCEAS Working Group on Population Management (1998). What can adaptive

- management do for our fish, forests, food, and biodiversity? *Integrative Biology: Issues, News, and Reviews* 1: 16-26.
- Peyrusson, F. (2018). Effect of hydrogel on the plants growth. *Université Catholique Louvain, Belgium*.
- Pinheiro, J.C. & Bates, D.M. (2000). Mixed-effects models in S and S-PLUS. Springer. New York.
- Poulakakis, N., Rusello, M., Geists, D., & Caccone, A. (2012). Unravelling the peculiarities of island life: vicariance, dispersal and the diversification of the extinct and extant giant Galápagos tortoises. *Molecular Ecology* 21: 160-173.
- R Core Team (2017). R development Core Team. R: a language for statistical computing 55: 275-286.
- Racine, C.H., & Downhower, J. F. (1974). Vegetative and reproductive strategies of *Opuntia* (Cactaceae) in the Galápagos Islands. *Biotropica* 6: 175-186.
- Rajakaruna, N., Pope, N., Perez-Orozco, J., & Harris, T.B. (2009). Ornithocoprophilous plants of Mount Desert Rock, a remote bird-nesting island in the Gulf of Maine, USA. *Rhodora* 111: 417-448.
- Restrepo, A., Colinvaux, P., Bush, M., Correa-Metrio, A., Conroy, J., Gardener, M.R., Jaramillo, P., Steinitz-Kannan, M., Overpeck, J. (2012). Impacts of climate variability and human colonization on the vegetation of the Galápagos Islands. *Ecology* 93:1853-1866.
- Schofield, E.K. (1973). Galápagos flora: the threat of introduced plants. *Biological Conservation* 5: 48-51.
- Snell, H.L., Snell, H.M., & Stone, P. (1994). Accelerated mortality of *Opuntia* on Isla Plaza Sur: another threat from an introduced vertebrate? *Noticias de Galápagos* 53, 19-20.
- Snell, H.M., Stone, P.A., & Snell, H.L. (1996). A summary of geographical characteristics of the Galápagos Islands. *Journal of Biogeography* 23: 619-624.
- Snell, H. & Rea, S. (1999). The 1997–98 El Niño in Galápagos: can 34 years of data estimate 120 years of pattern? *Noticias de Galápagos*: 11–20.
- Snyman, H.A. (2005). A case study on in situ rooting profiles and water-use efficiency of cactus pears, *Opuntia ficus-indica* and *O. robusta*. *Journal of the Professional Association for Cactus Development*: 1-21.
- Sulloway, F.J. & Noonan, K. (2015). *Opuntia* cactus loss in the Galápagos Islands, 1957-2014

- 550 (Pérdida de cactus *Opuntia* en las Islas Galápagos, 1957-2014). Technical Report, 1–30.
- 551 Sulloway, F.J., Noonan, K.M., Noonan, D.A. & Olila, K.J. (2013). Documenting ecological
552 changes in the Galápagos since Darwin’s visit. Project Proposal, 1-32.
- 553 Trueman, M. & D’Ozouville, N. (2010). Characterizing the Galápagos terrestrial climate in the
554 face of global climate change. *Galápagos Research* 67: 26–37.
- 555 Tye, A. (2007). La flora endémica de Galápagos: aumentan las especies amenazadas. In: FCD,
556 PNG, INGALA, editors. *Informe Galápagos 2006-2007*. Puerto Ayora, Galápagos,
557 Ecuador. p 101-107.

Table 1(on next page)

Total number of *Opuntia* spp. individuals planted by island by Galápagos Verde 2050 (2013-2018).

Numbers in parentheses ‘()’ are the number of individuals used in the current study analysis (Figures 4 & 5).

1 **Table 1. Total number of *Opuntia* spp. individuals planted by island by Galápagos Verde**
 2 **2050 (2013-2018).** Numbers in parentheses ‘()’ are the number of individuals used in the current
 3 study analysis (Figures 4 & 5).

Species	Baltra	Española	Floreana	Plaza Sur	San Cristóbal	Santa Cruz
<i>Opuntia echios</i> var. <i>echios</i>	400 (349)	—	—	737 (601)	—	—
<i>Opuntia echios</i> var. <i>gigantea</i>	—	—	—	—	—	68 (60)
<i>Opuntia megasperma</i> var. <i>megasperma</i>	—	—	20 (19)	—	4 (0)	—
<i>Opuntia megasperma</i> var. <i>orientalis</i>	—	196 (0)	—	—	—	—

4

Table 2 (on next page)

List of all sites of Galápagos Verde 2050 *Opuntia* spp. restoration and number of *Opuntia* spp. individuals planted (2013-2018).

Numbers in parentheses ‘()’ represent the percent of individuals that have survived through 2018.

Table 2. List of all sites of Galápagos Verde 2050 *Opuntia* spp. restoration and number of *Opuntia* spp. individuals planted (2013-2018). Numbers in parentheses ‘()’ represent the percent of individuals that have survived through 2018.

Island	Site Name	# Planted	UTM East ¹	UTM North ¹
Baltra (70%)	Antiguo basurero	158 (69%)	804668	9950436
	Casa de piedra	125 (74%)	802460	9948203
	Jardín ecológico Aeropuerto	1 (100%)	804100	9950795
	Parque Eólico	116 (68%)	803992	9950909
Española (79%)	Las Tunas	196 (79%)	199759*	9849118*
Floreana (40%)	Botadero de basura	3 (33%)	781054	9858587
	Cementerio	7 (29%)	780322	9858645
	Escuela Amazonas	5 (40%)	779594	9858865
	Gobierno Parroquial Floreana	1 (0%)	779530	9859029
	Oficina Técnica Parque	4 (75%)	779531	9859244
	Nacional Galápagos			
Plaza Sur (61%)	Centro	254 (62%)	815800	9935365
	Los Lobos Este	253 (47%)	815936	815936
	Oeste Cerro Colorado	230 (76%)	815304	9935602
San Cristóbal (100%)	CA Jacinto Gordillo	4 (100%)	209711*	9900150*
Santa Cruz (65%)	Colegio Nacional Galápagos	2 (50%)	798782	9918296
	Espacio Verde ABG	8 (88%)	797864	9918887
	Fundación Charles Darwin	51 (67%)	800106	9917856
	Oficina Técnica Parque	7 (29%)	799811	9917994
	Nacional Galápagos			

¹ UTM Zone = 15M, datum = WGS84

* UTM Zone = 16M

Figure 1(on next page)

Map of the Galápagos Islands, Ecuador.

Islands included in the current study are darkened and labeled in bold.




Figure 2 (on next page)

Total *Opuntia* spp. restoration from 2013 to 2018 across Baltra, Española, Floreana, Plaza Sur, San Cristóbal, and Santa Cruz islands.

Values above bars indicate total surviving individuals by the end of each year (y-axis values).

Values at the bottom indicate the total number of individuals planted each year.




Figure 3(on next page)

Approximate *Opuntia echios* var. *echios* population on Plaza Sur island from 1980 to 2018.

Redrawn from Snell et al. (1994) with 2014 addition from Sulloway and Noonan (2015), and 2015-2018 values based on estimate from 2014 (334) plus surviving individuals from Galápagos Verde 2050 (GV2050, green shading) replanting efforts.





Figure 4(on next page)

State of each planted *Opuntia* individual by the end of 2018 within each island.


A. Plaza Sur; B. Floreana; C. Baltra; D. Santa Cruz. “N” indicates the total number of individuals within each treatment on each island. Figure based on only those data used in the current analysis.

A




B

Manuscript to be reviewed



C



D




Figure 5(on next page)

Plots of the relative effect of variable parameters on two-year survival and growth rate of planted *Opuntia* individuals.

A. all islands two-year survival; B. all islands growth rate; C. Plaza Sur island two-year survival; D. Plaza Sur island growth rate; E. all islands excluding Plaza Sur two-year survival; and F. all islands excluding Plaza Sur growth rate. Each point represents coefficient estimate +/- bootstrapped 95% confidence intervals. P-values are generated based on the Satterthwaite method (* $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$). Values for two-year survival models are converted to odds ratio by exponentiating coefficients and subtracting one. Analyses are based on data from Baltra, Floreana, Plaza Sur, and Santa Cruz islands. Littoral zone values in A. and C. fall outside the scale of those boxes, so confidence intervals are presented as text.

