

# Polyphenol-rich *Trapa quadrispinosa* pericarp extract ameliorates high-fat diet induced non-alcoholic fatty liver disease by regulating lipid metabolism and insulin resistance in mice

Tunyu Jian<sup>1</sup>, Han Lü<sup>Corresp., 1</sup>, Xiaoqin Ding<sup>1</sup>, Yuexian Wu<sup>1</sup>, Yuanyuan Zuo<sup>1</sup>, Jiawei Li<sup>1</sup>, Jian Chen<sup>Corresp., 1, 2</sup>, Hong Gu<sup>3</sup>

<sup>1</sup> Institute of Botany, Jiangsu Province and Chinese Academy of Sciences, Nanjing, China

<sup>2</sup> Department of Food Science and Technology, College of Light Industry and Food Engineering, Nanjing Forestry University, Nanjing, China

<sup>3</sup> Department of Colorectal Surgery, Jiangyin Hospital of Traditional Chinese Medicine, Jiangyin, China

Corresponding Authors: Han Lü, Jian Chen

Email address: xiaohan1814@163.com, chenjian80@aliyun.com

In China, *Trapa quadrispinosa* (also called water caltrop) has long been used as a function food and folk medicine to treat diabetes mellitus for years. In the present study, the extract of *T. quadrispinosa* pericarp (TQPE) which mainly contains hydrolysable tannins was prepared to investigate the potential therapeutic action in non-alcoholic fatty liver disease (NAFLD) mice induced by high fat-diet (HFD). After the administration of TQPE (15, 30 mg/kg/day) for 8 weeks, the increased weight of body and liver were significantly suppressed. TQPE also ameliorated liver lipid deposition and reduced lipids parameters of blood in mice. Moreover, TQPE attenuated oxidative stress and showed a hepatoprotective effect in mice. TQPE was also found to decrease the value of homeostatic model assessment for insulin resistance. In addition, TQPE administration increased the phosphorylation of AMP-activated protein kinase (AMPK) and Acetyl-CoA carboxylase (ACC) and inhibited sterol regulatory element-binding protein (SREBP) in the liver tissue. Meanwhile, TQPE elevated insulin receptor substrate-1 (IRs-1) and protein kinase B (Akt) phosphorylation. These results reflected that, as a nature product, TQPE is a potential agent for suppressing the process of NAFLD via regulation of the AMPK/SREBP/ACC and IRs-1/Akt pathways.

# Polyphenol-rich *Trapa quadrispinosa* pericarp extract ameliorates high-fat diet induced non-alcoholic fatty liver disease by regulating lipid metabolism and insulin resistance in mice

Tunyu Jian<sup>1</sup>, Han Lü<sup>1\*</sup>, Xiaoqin Ding<sup>1</sup>, Yuexian Wu<sup>1</sup>, Yuanyuan Zuo<sup>1</sup>, Jiawei Li<sup>1</sup>, Jian Chen<sup>1,2\*</sup>, Hong Gu<sup>3</sup>

<sup>1</sup> Institute of Botany, Jiangsu Province and Chinese Academy of Sciences, Nanjing, China

<sup>2</sup> Department of Food Science and Technology, College of Light Industry and Food Engineering, Nanjing Forestry University, Nanjing, China

<sup>3</sup> Department of Colorectal Surgery, Jiangyin Hospital of Traditional Chinese Medicine, Jiangyin, China

Corresponding Author:

Han Lü

No.1 Qianhuhoucun, Nanjing, Jiangsu, 210014, China

Email address: xiaohan1814@163.com

Jian Chen

No.1 Qianhuhoucun, Nanjing, Jiangsu, 210014, China

Email address: chenjian80@aliyun.com;

## Abstract

In China, *Trapa quadrispinosa* (also called water caltrop) has long been used as a function food and folk medicine to treat diabetes mellitus for years. In the present study, the extract of *T. quadrispinosa* pericarp (TQPE) which mainly contains hydrolysable tannins was prepared to investigate the potential therapeutic action in non-alcoholic fatty liver disease (NAFLD) mice induced by high fat-diet (HFD). After the administration of TQPE (15, 30 mg/kg/day) for 8 weeks, the increased weight of body and liver were significantly suppressed. TQPE also ameliorated liver lipid deposition and reduced lipids parameters of blood in mice. Moreover, TQPE attenuated oxidative stress and showed a hepatoprotective effect in mice. TQPE was also found to decrease the value of homeostatic model assessment for insulin resistance. In addition, TQPE administration increased the phosphorylation of AMP-activated protein kinase (AMPK) and Acetyl-CoA carboxylase (ACC) and inhibited sterol regulatory element-binding protein (SREBP) in the liver tissue. Meanwhile, TQPE elevated insulin receptor substrate-1 (IRs-1) and protein kinase B (Akt) phosphorylation. These results reflected that, as a nature product, TQPE is

a potential agent for suppressing the process of NAFLD via regulation of the AMPK/SREBP/ACC and IRs-1/Akt pathways.

## Introduction

As a common chronic liver disease, Non-alcoholic fatty liver disease (NAFLD) is defined by pathological accumulation of lipid in the liver without excess alcohol consumption (Golabi *et al.* 2017). Being a hepatic manifestation of metabolic syndrome, it is similar to those chronic metabolic disorders, such as obesity, insulin resistance, type 2 diabetes mellitus (T2DM), inflammation and cardiovascular disease (Bagherniya *et al.* 2018). NAFLD increases the risk of progressive liver injury, which appears as a continuum disease progression, from simple steatosis to liver failure and hepatocellular carcinoma (Suolang *et al.* 2019). NAFLD has emerged as a worldwide serious public health burden, epidemiology of NAFLD have highlighted surprisingly high prevalence in many countries (the estimated prevalence is 25-30% in adults) (Moore 2019; Ratziu 2018). Therefore, there is a great demand for exploring effective therapeutic agents to treat and prevent NAFLD.

The recent evidence indicated that fat accumulation and insulin resistance (IR) are intensely associated with the development and progression of NAFLD (Araujo *et al.* 2018; Fan *et al.* 2018; Jian *et al.* 2018). As a highly evolutionarily conserved sensor of cellular energy status, AMP-activated protein kinase (AMPK) plays a critical role in regulating hepatic lipid metabolism including lipolysis, glucose transport and gluconeogenesis (Brown & Goldstein 1997). Sterol regulatory element-binding protein (SREBP), a key transcription factor in regulating liver lipid synthesis, is the downstream of AMPK (Li *et al.* 2011). Acetyl-CoA carboxylase (ACC), a member of lipogenic factor, is the downstream target of SREBP. AMPK activation phosphorylates and inhibits ACC in adipose and hepatic tissues thus downregulate fatty acid synthesis (Bijland *et al.* 2013; Zhang *et al.* 2018). In NAFLD model of many studies, it was observed that the inhibition of phosphorylation of AMPK led to lipid accumulation by increasing SREBP and inhibiting ACC phosphorylation (Chen *et al.* 2019; Li *et al.* 2018b; Park *et al.* 2019; Zhou *et al.* 2017). Besides, IR is also strongly associated with hepatic lipid accumulation in NAFLD. Insulin signaling transduction is dependent on insulin receptor substrate-1 (IRs-1), and phosphorylation of IRs-1 give rise to insulin pathway activation (Fu *et al.* 2018; Saez-Lara *et al.* 2016). Moreover, for insulin signaling cascade conduction, Protein kinase B (Akt) is another essential factor. Impairment of Akt activity has been demonstrated under NAFLD condition, thus activated Akt (increased phosphorylation) could ameliorate hepatic steatosis and improve IR in NAFLD model (Fan *et al.* 2018; Jung *et al.* 2018). Therefore, targeting regulation of AMPK and insulin signaling pathway might be a new and useful therapeutic approach to drop lipid accumulation and insulin resistance in NAFLD.

Nowadays, pharmacological studies have significantly expanded to screen natural products for exploration of novel pharmaceutical agents. Many studies revealed that medicinal plant extracts, herb formulas have remarkable therapeutic effect on NAFLD (Bagherniya *et al.* 2018; Chen *et al.* 2017; Li *et al.* 2018a; Suolang *et al.* 2019). *Trapa quadrispinosa*, also called water

chestnut or water caltrop, is a floating-leaf aquatic plant, which is commonly cultivated in the south of China, as well as in India and south east Asia. The fruit of water caltrop, is a function food and folk medicine, which could be used to treat metabolic syndrome such as diabetes mellitus (DM). But the pericarps of *T. quadrispinosa* were usually discarded in large quantities after the seeds had been harvested. Interestingly, recent researches have demonstrated that the pericarps of water caltrop also displayed multiple biological activities, including hypoglycemic (Huang *et al.* 2016), anti-tumor (Lin *et al.* 2013), anti-inflammatory (Kim *et al.* 2015), anti-oxidant effects and hepatprotective activity (Kim *et al.* 2014). To our knowledge, the therapeutic effect of *Trapa quadrispinosa* pericarps extract (TQPE) in high-fat diet (HFD) induced NAFLD, remains unknown.

The purpose of the present study was designed to determine whether *Trapa quadrispinosa* pericarps extract (TQPE) could attenuate NAFLD induced by HFD in mice, also explore a possible mechanism of this action.

## Materials & Methods

### Chemicals and Reagents

HPLC grade methanol used for the mobile phase in HPLC-DAD/QTOF analysis was obtained from Tedia Co. Inc. (Fairfield, OH, USA). Formic acid in HPLC grade was taken from Acros Organics (Geel, Belgium). Pure water was produced from a Milli-Q system (Millipore, Bedford, MA, USA). Five standard compounds isolated from *T. quadrispinosa* were prepared and identified according to our previous study (Lv *et al.* 2019). Gallic acid was obtained from National Institutes for Food and Drug Control of China (Beijing, China). The other reagents were obtained from Sinopharm Chemical Reagent Co. Ltd. (Shanghai, China). A kit from KeyGen Biotechnology (Nanjing, China) was used for the protein extraction and BCA protein assay. Besides, the kits for detecting blood glucose, high-density lipoprotein cholesterol (HDL-c), low-density lipoprotein cholesterol (LDL-c), total cholesterol (TC), triglycerides (TG), aspartate aminotransferase (AST), alanine aminotransferase (ALT), superoxide dismutase (SOD) and malonaldehyde (MDA) were all obtained from Nanjing Jiancheng Bioengineering Institute (Nanjing, China). Determination of insulin by enzyme-linked immunosorbent assay (ELISA) commercially kits were purchased from Beyotime Institute of Biotechnology (Haimen, China).

The antibodies of AMPK $\alpha$ , p-AMPK $\alpha$  (Thr172), ACC, p-ACC, IRs-, p-IRs-1 (Tyr895), Akt, p-Akt (Ser473) for western blotting (WB) were purchased from Cell Signaling Technology (Danvers, MA, USA). SREBP was obtained from Santa Cruz Biotechnology (Santa Cruz, USA). Anti-rabbit/anti-mouse IgG and HRP-linked antibody were purchased from Cell Signaling Technology (Danvers, USA).

### Preparation and characterization of *Trapa quadrispinosa* pericarp extract (TQPE)

The air-dried *T. quadrispinosa* (collected in Shandong Zaozhuang of China) pericarp powder (10.0 kg) was extracted twice with fifty litres 80% (v/v) ethanol by soaking at room temperature for 14 days. After concentrating in vacuum at 50 °C, the extract was suspended in distilled water and then partitioned by petroleum ether, ethyl acetate and normal-butanol successively. The ethyl

acetate extract was first concentrated and then dissolved in water, and later applied to a column packed with macroreticular resin XAD16 (The Dow Chemical Company). The column was orderly eluted with a gradient of H<sub>2</sub>O, 10%, 40%, 60% and 95% ethanol solution. The 40% ethanol elution was collected, concentrated and dried and finally the TQPE (175g) was obtained.

According to previous reported method, the total phenolic content of TQPE was measured by the Folin-Ciocalteu phenol reagent (Singleton & Rossi 1965). Briefly, 1.0 mg of TQPE extract was solved in 1.0 ml of methanol was used as test solution. After incubation with Folin-Ciocalteu phenol reagent, the test solution was added 20% sodium carbonate to develop a color and its absorbance was measured at 755 nm. The final value was expressed as gallic acid (standard) equivalent.

5.0 mg/ml of TQPE (dissolved in methanol) was injected in an Agilent 6530 accurate-mass quadrupole time-of-flight system (Agilent Technologies, CA, USA) for the HPLC-QTOF/MS analysis. Separation was conducted by an Agilent ZORBAX SB-C18 column (1.8  $\mu$ m, 4.6  $\times$  100 mm; Waldbronn, Germany). The mobile phase was composed of Methanol (A) and 0.1% formic acid (B) under gradient conditions (0-50 min, 10-50%A; 50-65 min, 50-100% A) at 1 ml/min. The QTOF-MS equipped with an electrospray ion source operated in the negative ion mode at 4 kV capillary voltage, 10 L/min drying gas and 350  $^{\circ}$ C gas temperature. Mass range of scanning was 100-1000 m/z. MassHunter Qualitative Analysis software (Agilent Technologies, USA) was used to perform measurement and analysis.

Furthermore, HPLC-DAD was performed by Dionex Ultimate 3000 HPLC system (Thermo Fisher Scientific, USA) equipped with a diode array-UV detector at 210 nm. Nucleosil C18 column (4.6 mm  $\times$  150 mm, 3  $\mu$ m, ThermoFisher Scientific, USA) was used for separation. The mobile phase was composed of Acetonitrile (A) and 0.05% trifluoroacetic acid (B) under gradient conditions as followed: 0-5 min, 5%A; 5-15 min, 5-10% A; 15-55min, 10-25% A; 55-65min, 25-100% A, the flow rate of HPLC was 1 ml/min.

# **Animals**

The male ICR mice were obtained from Shanghai Sino-British SIPPR/BK Lab Animal Co., Ltd. (China) and they were maintained in controlled condition as our previous study described (Jian *et al.* 2018).

# **Experimental protocol**

At first stage, mice were provided either a normal chow diet (10% of calories) or a high fat diet (60% of calories) for 4 weeks, which were obtained from Shanghai Lab-oratory Animal Co. Ltd. (Shanghai, China). Then, the high fat diet-fed mice were randomly divided into 3 groups: one control group and two test groups, the test groups were administered with low dose of TQPE (15 mg/kg) and high dose of TQPE (30 mg/kg) per day. Normal control groups (CON) and high fat diet-fed mice control (HFD) group were treated with solution of 0.5% CMC-Na. All mice in each group were administrated relevant reagents for another 8 weeks. In the end of experimental period, all the mice were fasted for 12 h, then anesthetized and euthanized by isoflurane gas. Thereafter, the serum was separated from blood and the liver was removed and weighted. All the samples were stored at -80 $^{\circ}$ C condition before use.

## Biochemical parameters analysis

According to the commercial kits instructions, serum TC, HDL, LDL, TG, SOD, MDA, ALT and AST concentrations were determined in a microplate reader (Molecular Device, CA, USA). Serum insulin level was determined by using ELISA commercially kits. The HOMA-IR (Homeostatic model assessment for insulin resistance) value which was calculated as fasting blood glucose (mmol/L)  $\times$  fasting serum insulin (mIU/L)/22.5, was represented insulin sensitivity.

## Liver pathological examination

Liver tissues were fixed overnight at 4 °C in 4% paraformaldehyde and embedded in paraffin. Sections (4  $\mu$ m) were prepared for hematoxylin & eosin (H&E) stain. The histological images were observed and photographed under an optical microscope (Carl Zeiss, Germany).

## Western blot analysis of liver tissue

The total protein from the liver tissues was extracted using a commercial kit. Proteins from the liver supernatant were quantified by a BCA kit. 25  $\mu$ g of total proteins was separated on 8-10% SDS-PAGE gel then transferred onto a PVDF membrane from Millipore (MA, USA). The membrane was blocked for 2 h in 5% nonfat milk powder solution at room temperature. Then the membranes were incubated with primary antibody at 4°C overnight. At the second day, the membranes were incubated with secondary antibody at room temperature for 1-1.5 h. The Membrane-bound antibodies complexes were noticed by ECL detection (Santa Cruz Biotechnology, CA, USA). Tanon 5200 image system (Tanon, Shanghai, China) was used to examine the signal.  $\beta$ -actin was employed to demonstrate the standard proteins quantity. ImageJ software (NIH) was applied to quantify the density of western blotting bands

## Statistical analysis

All results were expressed as mean  $\pm$  SE from 10 mice. All statistical analyses were done through the GraphPad Prism Software (Version 7.02, CA, USA). One-way ANOVA with a post-hoc test was used for the statistical significance evaluation.  $P < 0.05$  was indicated statistical significance.

# Results

## Analysis of chemical constituents and content of TQPE by HPLC-QTOF/MS and HPLC-DAD

According to Folin-Ciocalteu phenol test, the most abundant compound in the TQPE was polyphenolics (hydrolysable tannins), the content of which was accounting for  $91.7 \pm 2.1\%$  (gallic acid equivalents). To furtherly understand the potential bioactive components in TQPE, a HPLC-QTOF/MS assay was carried out. The base peak chromatogram of TQPE was illustrated in Figure 1A. Eight phenolic compounds ( $R_t = 9.45, 23.07, 24.15, 24.76, 30.72, 35.58, 36.52$  and  $47.65$  min) were tentatively identified according to previous studies (Hatano *et al.* 1990; Huang *et al.* 2016). Their structures were shown in Figure 1B. The predominated polyphenols in TPQE was gallic acid and its derivatives, which included hydrolysable gallotannins and gallates. We also used HPLC-DAD method to calculate the content of five polyphenols in TPQE, which was

showed in Table 1. Gallic acid (Compound 1), 1,2,3-tri-*O*-galloyl- $\beta$ -D-glucopyranose (Compound 2), 1,2,3-tetra-*O*-galloyl-4,6-*O*-hexahydroxydiphenoyl- $\beta$ -D-glucopyranoside (Compound 6), 1,2,3,6-tetra-*O*-galloyl- $\beta$ -D-glucopyranose (Compound 7), 1,2,3,4,6-penta-*O*-hexahydroxydiphenoyl- $\beta$ -D-glucopyranoside (Compound 8) were accounted for  $696.56 \pm 4.61$  mg/g.

### **Effect of TQPE on the body and liver weight of HFD mice**

As showed in Figure 2A and Figure 2B, compared with the normal group, body weight in HFD group and their liver to body weight ratio rose obviously ( $P < 0.001$ ). Administration of TQPE (15 and 30 mg/kg) significantly inhibited HFD induced growth in body weight and liver to body weight ratios ( $P < 0.01$ ). Treatment with TQPE reversed the continuing weight gain in both body and liver.

### **Histopathological examination**

Normal control group livers showed glossy and resilient appearance. In contrast, as showed in Figure 2C, the NAFLD mice livers were enlarged with yellow necrosis foci appearing. After the livers were treated with TQPE, the liver appearance improved in a dose-dependent manner (Figure 2C).

Figure 2D displayed liver histology photo sections. In the control group, liver tissues showed normal liver lobular structure without fatty accumulation or inflammatory. In HFD treated mice, liver sections revealed significantly higher damage of hepatic lobule structures compared with normal mice, while lipid droplets were also spotted in liver cells. As displayed in Figure 2D, after treating with TQPE in 15 and 30 mg/kg, hepatocyte swelling as well as quantities and volumes of lipid droplet were all alleviated. Morphology of liver lobular structure almost regained normal status, especially in the mice treated with 30 mg/kg TQPE.

### **TQPE ameliorates Lipid Parameters in NAFLD mice**

Relevant lipids parameters were detected to evaluate the effects of TQPE on lipid metabolism in NAFLD mice. TC, TG and LDL content were markedly elevated ( $P < 0.001$ ; Figure 3A, B and D) along with a decrease in HDL level (Figure 3C;  $P < 0.05$ ) in NAFLD mice compared to the normal group. In Figure 3A-C, the data showed that administration of TQPE (15 mg/kg) significantly reduced TC, TG and elevated HDL compared with the HFD group ( $P < 0.01$ ). Furthermore, after treatment with a high dose of TQPE at 30 mg/kg, besides above effects, it dramatically decreased LDL but increased HDL level (Figure 3A-D;  $P < 0.001$ ).

### **TQPE changes the oxidative stress balance and reduces liver injury in NAFLD mice**

As demonstrated in Figure 4A and B, HFD group mice showed less adequate SOD and surplus level of MDA in the serum compared to the control group. This phenomenon implying an increased oxidative stress and a decrease of antioxidant capacity in HFD-induced NAFLD mice. Treatment with TQPE ameliorated this symptom by significantly raising SOD and reducing MDA level ( $P < 0.001$ , Figure 4A and B). In Figure 4C and D, compared with normal mice, the contents of ALT and AST were seen elevating in the NAFLD mice. Administration of TQPE (15 and 30 mg/kg) significantly reduced ALT and AST levels indicating the recovery from liver damage induced by HFD-induced NAFLD mice ( $P < 0.001$ ).

## Effects of TQPE on serum insulin and Homeostatic Model Assessment of Insulin Resistance (HOMA-IR)

Insulin concentration in serum and HOMA-IR value were calculated to evaluate the effect of TQPE on insulin blood level and action.

In HFD mice, both the serum insulin concentration and HOMA-IR value were increased significantly, which indicating that there was an insulin resistance in NAFLD mice ( $P < 0.001$ , Figure 5). Both the value of serum insulin concentration and HOMA-IR were significantly decreased after the treatment of TQPE (15 and 30 mg/kg) in comparison with the HFD mice ( $P < 0.001$ , Figure 5A and 5B). The data indicated that TQPE suppressed HFD-induced IR in NAFLD mice.

## TQPE changes pathways participated in lipid metabolism and insulin resistance

For further investigate the mechanism of TQPE preventing NAFLD, we detected the change of proteins expression including AMPK, SREBP, ACC, IRS-1 and Akt in the liver tissue, which were involved in lipid metabolism and insulin resistance. In Figure 6A-C, the phosphorylation of AMPK and ACC were decreased along with an increase in SREBP in HFD group in comparison with control group ( $P < 0.001$ ). Meanwhile, compared with control group, in HFD mice liver, it showed a drop of IRS-1 and Akt phosphorylation ( $P < 0.001$ ; Figure 6D and E). Treatment with TQPE at a high dose of 30 mg/kg markedly recovered the phosphorylation level of AMPK, ACC and inhibited the expression of SREBP in HFD-induced NAFLD mice ( $P < 0.001$ ; Figure 6A-C). Additionally, TQPE (15 and 30 mg/kg) administration kept the similar level of IRS-1 and Akt phosphorylation than the detected in the control mice ( $P < 0.01$ ; Figure 6D and E).

## Discussion

The current study investigated the therapeutic effect of *Trapa quadrispinosa* pericarp extract (TQPE), on lipid accumulation and insulin resistance in HFD-induced NAFLD, and worked out the possible mechanism. We concluded that TQPE attenuated HFD-induced lipid accumulation in the liver, probably was mediated by the AMPK activation and SREBP-mediated lipogenesis inhibition. TQPE also improved IR in NAFLD possibly through upregulating the levels of p-IRS-1 and p-Akt protein (Figure 7).

It is widely accepted and confirmed that hepatic steatosis is the earliest stage of NAFLD, which is manifested as accumulation of lipid in the liver. In nutrient oversupply states, ectopic lipid accumulation is closely related to hyperglycemia and hyperlipidemia. AMPK is a crucial key and important metabolic sensor and regulator of lipid and glucose metabolism in diverse tissues and cells (Cheng *et al.* 2016; Day *et al.* 2017). Activated AMPK promotes energy production, meanwhile, and represses ATP-consuming processes, to maintain energy status balance (Gonzales *et al.* 1992; Ruderman & Prentki 2004). Both SREBP and ACC are two downstream effectors of AMPK. For instance, AMPK activation inhibits fatty acid synthesis via a reduction in the transcriptional activation of SREBP, which is a key transcription factor during *de novo* lipogenesis in the regulation of lipogenic genes including *acc1* in the liver (Ahmed & Byrne 2007; Zhou *et al.* 2017). AMPK activation also could switch off fatty acid synthesis through ACC phosphorylation in adipose and hepatic tissues (Bijland *et al.* 2013; Zhang *et al.*



2018). As described, AMPK activation might reduce hepatic lipid accumulation via lipogenesis inhibition, indicating that AMPK is a potential therapeutic target for the treatment of hepatic steatosis in NAFLD (Zhou *et al.* 2017). Recent studies showed that many compounds are able to activate AMPK in animal and cellular models to improve lipid metabolism in NAFLD, by inhibiting SREBP and increasing phosphorylates of ACC (Chen *et al.* 2019; Kang *et al.* 2019; Park *et al.* 2019). In our study, we explored the lipid contents comprising TC, TG and LDL, the concentration of which were significantly lowered after TQPE treatment, while the HDL level was enhanced in HFD-induced NAFLD ( $P < 0.01$ ; Figure 3). We also found that TQPE at dose of 30 mg/kg/day could obviously activate the phosphorylation of AMPK and ACC, while inhibit SREBP expression ( $P < 0.001$ ; Figure 7A-C). These results demonstrated that AMPK axis is associated with the protective effects of TQPE in lipid metabolism under NAFLD status.

Impaired responsiveness to insulin evokes insulin resistance in NAFLD, hence, the improvement of insulin sensitivity is important for the treatment of NAFLD. Insulin receptor substrate-1 (IRS-1) is a core factor in insulin signaling transduction. Research showed that phosphorylation of IRS-1 at Ser accounts for IR, while phosphorylation of IRS-1 at Tyr is required for responses of insulin stimulates (Bhattacharyya *et al.* 2015; Xiao *et al.* 2018). In metabolic disorders, such as NAFLD and T2DM, impaired IRS-1 level was observed, for instance, lipid accumulation in the liver was closely associated with increased serine phosphorylation of IRS-1 (Araujo *et al.* 2018; Dallak 2018; Dong *et al.* 2019). Restoration of IRS-1 might be a beneficial treatment for curing NAFLD (Yang *et al.* 2018; Zhou *et al.* 2018). Protein kinase B (Akt) as downstream of IRS-1, plays an essential role in insulin signaling cascade. In response to upstream signal of IRS-1 on Tyr, activated Akt increases hepatic glucose uptake, glycogen synthesis and decreases lipogenesis (Cignarelli *et al.* 2019; Geidl-Flueck & Gerber 2017; Guo 2014). Moreover, hepatic Akt activation enhances the insulin sensitivity (Cignarelli *et al.* 2019; Ke *et al.* 2015). Therefore, IRS-1/Akt is important for hepatic insulin signaling, regulates this pathway might be beneficial for treating NAFLD. In the current work, the level of insulin and HOMA-IR value increased significantly in HFD-induced NAFLD ( $P < 0.001$ ; Figure 5). This mechanism supported by decreased hepatic phosphorylation of IRS-1 on Tyr ( $P < 0.001$ ; Figure 6D). In parallel, there was also a significant decline in phosphorylation of Akt ( $P < 0.001$ ; Figure 6E) in NAFLD mouse model. TQPE administration with different doses (15 and 30 mg/kg/day) reversed the high level of insulin and HOMA-IR value compared with NAFLD mice ( $P < 0.001$ ; Figure 5). This effect might be associated with restore phosphorylation of IRS-1 and Akt ( $P < 0.05$ ; Figure 6D, E) by TQPE to improve IR in NAFLD model.

In addition, ALT and AST are two hepatic enzymes, which are well-known key biochemical markers for detecting liver damage in NAFLD mode (Katsagoni *et al.* 2017). In our study, HFD treatment induced increasing of ALT and AST, while the administration of TQPE (15 and 30 mg/kg/day) could revise these parameters ( $P < 0.001$ ; Figure 4C, D). In comparison with a dose of 2000 mg/kg of *T. natans* pericarp extract displaying antihyperglycemic effect in rats (Yasuda *et al.* 2014), and based on exploratory studies of 50-100 mg/kg doses, we found that a lower dose of 30 mg/kg of TQPE, also exerted a therapeutic action on HFD-induced NAFLD

in mice, while no significant abnormal changes were found in the present study, so we stepped down doses for this study to provide a good safety margin. According to USA Food and Drug Administration's instruction (Food and Drug Administration. 2005), the dose of 30 mg/kg TQPE used in human is approximately 200 mg/day, thus this dose is acceptable for patients compared to other clinical trials used (250-960 mg/day) (Chen *et al.* 2016; Qiu *et al.* 2013; Ried *et al.* 2013).

## Conclusions

The pericarps of *T. quadrispinosa* is a kind of agricultural waste, that are usually discarded in large quantities after the seeds had been harvested. From economic aspect, the present study helps to discover polyphenol-rich extraction from this waste with medicinal value. The present study demonstrated that *Trapa quadrispinosa* pericarp extract (TQPE) administration significantly improve the metabolic parameters including the insulin resistance in high-fat diet-induced NAFLD mice. The protective mechanisms of TQPE may be attributed to regulate AMPK/SREBP/ACC and IRs-1/Akt signal pathways, which are involved in lipid metabolism and IR, respectively. TQPE treatment provides a novel therapeutic strategy to prevent HFD-induced NAFLD, however, there were still some limitations in this study i) the amount of food eaten was not directly measured, however, observation by caretakers showed that the animals were feeding and the loss of weight was not simply related to decreased eating; ii) the active entity in the extract which is absorbed and found in the plasma of the mice was not determined. In further studies, we warranted to determine the bioavailability of the active entity, and evaluate the mechanisms by which polyphenolic mediates AMPK/SREBP/ACC and IRs-1/Akt in HFD-induced NAFLD.

## Acknowledgements

The authors would like to thank Mr. Shirong Yang (Centre for instrumental analysis, College of Forestry, Nanjing Forestry University, Nanjing, China) for his useful help in the compound structure analysis.

## ADDITIONAL INFORMATION AND DECLARATIONS

### Funding

This research was funded by the National Natural Science Foundation of China (No. 81703224; No. 81773885; No. 31770366), Jiangsu Province Science and Technology Modern Agricultural Plan (No. BE2016383). Jiangsu Province Independent Innovation in Agricultural science and technology fund (CX(18)3042). And Natural Science Foundation of the Jiangsu Province (BK20160603). The authors are also indebted to the supports from Jiangsu Scientific and Technological Innovations Platform (Jiangsu Provincial Service Center for Antidiabetic Drug Screening) and Jiangsu Key Laboratory for the Research and Utilization of Plant Resources (No. JSPKLB201825; No. JSPKLB201833).

## Competing Interests

The authors declare there are no competing interests.

## Author Contributions

Formal analysis, T.Y.J. and X.Q.D.; Investigation, T.Y.J., H.L., X.Q.D and Y.X.W.; Methodology, Y.Y.Z. and J.W.L.; Project administration, H.L. and J.C.; Resources, H.G.; Writing - original draft, T.Y.J. and H.L.; Writing – review & editing, J.C..

## Animal Ethics

The following information was supplied relating to ethical approvals (i.e., approving body and any reference numbers):

All animal experiments for the study followed the Guide for the Care and Use of Laboratory Animals was approved by the Animal Ethics Committee of China Pharmaceutical University (certificate number: SYXK2016-0011, approval date: 27 January 2016 to 26 January 2021).

## References

- Ahmed MH, and Byrne CD. 2007. Modulation of sterol regulatory element binding proteins (SREBPs) as potential treatments for non-alcoholic fatty liver disease (NAFLD). *Drug Discovery Today* 12:740-747.10.1016/j.drudis.2007.07.009
- Araujo LCC, Feitosa KB, Murata GM, Furigo IC, Teixeira SA, Lucena CF, Ribeiro LM, Muscara MN, Costa SKP, Donato J, Jr., Bordin S, Curi R, and Carvalho CRO. 2018. Uncaria tomentosa improves insulin sensitivity and inflammation in experimental NAFLD. *Scientific Reports* 8:11013.10.1038/s41598-018-29044-y
- Bagherniya M, Nobili V, Blesso CN, and Sahebkar A. 2018. Medicinal plants and bioactive natural compounds in the treatment of non-alcoholic fatty liver disease: A clinical review. *Pharmacological Research* 130:213-240.10.1016/j.phrs.2017.12.020
- Bhattacharyya S, Feferman L, and Tobacman JK. 2015. Carrageenan Inhibits Insulin Signaling through GRB10-mediated Decrease in Tyr(P)-IRS1 and through Inflammation-induced Increase in Ser(P)307-IRS1. *Journal of Biological Chemistry* 290:10764-10774.10.1074/jbc.M114.630053
- Bijland S, Mancini SJ, and Salt IP. 2013. Role of AMP-activated protein kinase in adipose tissue metabolism and inflammation. *Clinical Science (London, England: 1979)* 124:491-507.10.1042/CS20120536
- Brown MS, and Goldstein JL. 1997. The SREBP pathway: regulation of cholesterol metabolism by proteolysis of a membrane-bound transcription factor. *Cell* 89:331-340
- Chen IJ, Liu CY, Chiu JP, and Hsu CH. 2016. Therapeutic effect of high-dose green tea extract on weight reduction: A randomized, double-blind, placebo-controlled clinical trial. *Clinical Nutrition* 35:592-599.10.1016/j.clnu.2015.05.003
- Chen K, Chen X, Xue H, Zhang P, Fang W, Chen X, and Ling W. 2019. Coenzyme Q10 attenuates high-fat diet-induced non-alcoholic fatty liver disease through activation of the AMPK pathway. *Food Funct* 10:814-823.10.1039/c8fo01236a
- Chen Q, Wang T, Li J, Wang S, Qiu F, Yu H, Zhang Y, and Wang T. 2017. Effects of Natural Products on Fructose-Induced Nonalcoholic Fatty Liver Disease (NAFLD). *Nutrients* 9.10.3390/nu9020096

- Cheng S, So WY, Zhang D, Cheng Q, Boucher BJ, and Leung PS. 2016. Calcitriol Reduces Hepatic Triglyceride Accumulation and Glucose Output Through Ca<sup>2+</sup>/CaMKK $\beta$ /AMPK Activation Under Insulin-Resistant Conditions in Type 2 Diabetes Mellitus. *Current Molecular Medicine* 16:747-758
- Cignarelli A, Genchi VA, Perrini S, Natalicchio A, Laviola L, and Giorgino F. 2019. Insulin and Insulin Receptors in Adipose Tissue Development. *Int J Mol Sci* 20.10.3390/ijms20030759
- Dallak MA. 2018. Acylated ghrelin induces but deacylated ghrelin prevents hepatic steatosis and insulin resistance in lean rats: Effects on DAG/ PKC/JNK pathway. *Biomedicine and Pharmacotherapy* 105:299-311.10.1016/j.biopha.2018.05.098
- Day EA, Ford RJ, and Steinberg GR. 2017. AMPK as a Therapeutic Target for Treating Metabolic Diseases. *Trends in Endocrinology and Metabolism* 28:545-560.10.1016/j.tem.2017.05.004
- Dong R, Yang X, Wang C, Liu K, Liu Z, Ma X, Sun H, Huo X, Fu T, and Meng Q. 2019. Yangonin protects against non-alcoholic fatty liver disease through farnesoid X receptor. *Phytomedicine* 53:134-142.10.1016/j.phymed.2018.09.006
- Fan Y, He Z, Wang W, Li J, Hu A, Li L, Yan L, Li Z, and Yin Q. 2018. Tangganjian decoction ameliorates type 2 diabetes mellitus and nonalcoholic fatty liver disease in rats by activating the IRS/PI3K/AKT signaling pathway. *Biomedicine and Pharmacotherapy* 106:733-737.10.1016/j.biopha.2018.06.089
- Food and Drug Administration. 2005. Guidance for industry: estimating the maximum safe starting dose in initial clinical trials for therapeutics in adult healthy volunteers. *Center for Drug Evaluation and Research (CDER)*.7
- Fu D, Cui H, and Zhang Y. 2018. Lack of CIC-2 Alleviates High Fat Diet-Induced Insulin Resistance and Non-Alcoholic Fatty Liver Disease. *Cellular Physiology and Biochemistry* 45:2187-2198.10.1159/000488164
- Geidl-Flueck B, and Gerber PA. 2017. Insights into the Hexose Liver Metabolism-Glucose versus Fructose. *Nutrients* 9.10.3390/nu9091026
- Golabi P, Bush H, and Younossi ZM. 2017. Treatment Strategies for Nonalcoholic Fatty Liver Disease and Nonalcoholic Steatohepatitis. *Clinics in Liver Disease* 21:739-753.10.1016/j.cld.2017.06.010
- Gonzales GF, Garcia-Hjarles M, and Velasquez G. 1992. Hyperprolactinaemia and hyperserotoninaemia: their relationship to seminal quality. *Andrologia* 24:95-100
- Guo S. 2014. Insulin signaling, resistance, and the metabolic syndrome: insights from mouse models into disease mechanisms. *Journal of Endocrinology* 220:T1-T23.10.1530/JOE-13-0327
- Hatano T, Okonogi A, Yazaki K, and Okuda T. 1990. Trapanins a and b, oligomeric hydrolyzable tannins from trapa japonica flerov. *Chem Pharm Bull* 38:2707-2711.10.1016/j.chemosphere.2016.04.049
- Huang HC, Chao CL, Liaw CC, Hwang SY, Kuo YH, Chang TC, Chao CH, Chen CJ, and Kuo YH. 2016. Hypoglycemic Constituents Isolated from Trapa natans L. Pericarps. *Journal of Agricultural and Food Chemistry* 64:3794-3803.10.1021/acs.jafc.6b01208
- Jian T, Ding X, Wu Y, Ren B, Li W, Lv H, and Chen J. 2018. Hepatoprotective Effect of Loquat Leaf Flavonoids in PM2.5-Induced Non-Alcoholic Fatty Liver Disease via Regulation of IRs-1/Akt and CYP2E1/JNK Pathways. *Int J Mol Sci* 19.10.3390/ijms19103005
- Jung TW, Park HS, Choi GH, Kim D, and Lee T. 2018. beta-aminoisobutyric acid attenuates LPS-induced inflammation and insulin resistance in adipocytes through AMPK-mediated pathway. *Journal of Biomedical Science* 25:27.10.1186/s12929-018-0431-7
- Kang MC, Ding Y, Kim HS, Jeon YJ, and Lee SH. 2019. Inhibition of Adipogenesis by Diploretohydroxycarmalol (DPHC) through AMPK Activation in Adipocytes. *Marine Drugs* 17.10.3390/md17010044

- 449 Katsagoni CN, Georgoulis M, Papatheodoridis GV, Panagiotakos DB, and Kontogianni MD.  
450 2017. Effects of lifestyle interventions on clinical characteristics of patients with non-  
451 alcoholic fatty liver disease: A meta-analysis. *Metabolism* 68:119-  
452 132.10.1016/j.metabol.2016.12.006
- 453 Ke B, Zhao Z, Ye X, Gao Z, Manganiello V, Wu B, and Ye J. 2015. Inactivation of NF-kappaB  
454 p65 (RelA) in Liver Improves Insulin Sensitivity and Inhibits cAMP/PKA Pathway.  
455 *Diabetes* 64:3355-3362.10.2337/db15-0242
- 456 Kim B, Kim JE, Choi BK, and Kim HS. 2015. Anti-Inflammatory Effects of Water Chestnut  
457 Extract on Cytokine Responses via Nuclear Factor-kappaB-signaling Pathway.  
458 *Biomolecules & Therapeutics* 23:90-97.10.4062/biomolther.2014.080
- 459 Kim YS, Hwang JW, Han YK, Kwon HJ, Hong H, Kim EH, Moon SH, Jeon BT, and Park PJ.  
460 2014. Antioxidant activity and protective effects of *Trapa japonica* pericarp extracts  
461 against tert-butylhydroperoxide-induced oxidative damage in Chang cells. *Food and*  
462 *Chemical Toxicology* 64:49-56.10.1016/j.fct.2013.11.018
- 463 Li Q, Li HJ, Xu T, Du H, Huan Gang CL, Fan G, and Zhang Y. 2018a. Natural Medicines Used  
464 in the Traditional Tibetan Medical System for the Treatment of Liver Diseases. *Frontiers*  
465 *in Pharmacology* 9:29.10.3389/fphar.2018.00029
- 466 Li Y, Xu S, Mihaylova MM, Zheng B, Hou X, Jiang B, Park O, Luo Z, Lefai E, Shyy JY, Gao B,  
467 Wierzbicki M, Verbeuren TJ, Shaw RJ, Cohen RA, and Zang M. 2011. AMPK  
468 phosphorylates and inhibits SREBP activity to attenuate hepatic steatosis and  
469 atherosclerosis in diet-induced insulin-resistant mice. *Cell Metab* 13:376-  
470 388.10.1016/j.cmet.2011.03.009
- 471 Li YC, Qiao JY, Wang BY, Bai M, Shen JD, and Cheng YX. 2018b. Paeoniflorin Ameliorates  
472 Fructose-Induced Insulin Resistance and Hepatic Steatosis by Activating LKB1/AMPK  
473 and AKT Pathways. *Nutrients* 10.10.3390/nu10081024
- 474 Lin Q, Shen J, Ning Y, Shen S, and Das UN. 2013. Inhibitory effects of water caltrop pericarps  
475 on the growth of human gastric cancer cells in vitro. *Current Pharmaceutical Design*  
476 19:7473-7478
- 477 Lv H, Jian T, and Ding X. 2019. *Trapa natans* pericarp extract ameliorates hyperglycemia and  
478 hyperlipidemia in type 2 diabetic mice. *Revista Brasileira de*  
479 *Farmacognosia*.10.1016/j.bjp.2019.04.011
- 480 Moore JB. 2019. From sugar to liver fat and public health: systems biology driven studies in  
481 understanding non-alcoholic fatty liver disease pathogenesis. *Proceedings of the*  
482 *Nutrition Society*:1-15.10.1017/S0029665119000570
- 483 Park M, Yoo JH, Lee YS, and Lee HJ. 2019. *Lonicera caerulea* Extract Attenuates Non-  
484 Alcoholic Fatty Liver Disease in Free Fatty Acid-Induced HepG2 Hepatocytes and in  
485 High Fat Diet-Fed Mice. *Nutrients* 11.10.3390/nu11030494
- 486 Qiu F, Jiang J, Ma Y, Wang G, Gao C, Zhang X, Zhang L, Liu S, He M, Zhu L, Ye Y, Li Q, and  
487 Miao P. 2013. Opposite Effects of Single-Dose and Multidose Administration of the  
488 Ethanol Extract of Danshen on CYP3A in Healthy Volunteers. *Evidence-Based*  
489 *Complementary and Alternative Medicine* 2013:730734.10.1155/2013/730734
- 490 Ratzliff V. 2018. The painful reality of end-stage liver disease in NASH. *Lancet Gastroenterol*  
491 *Hepatol* 3:8-10.10.1016/S2468-1253(17)30365-5
- 492 Ried K, Frank OR, and Stocks NP. 2013. Aged garlic extract reduces blood pressure in  
493 hypertensives: a dose-response trial. *European Journal of Clinical Nutrition* 67:64-  
494 70.10.1038/ejcn.2012.178
- 495 Ruderman N, and Prentki M. 2004. AMP kinase and malonyl-CoA: targets for therapy of the  
496 metabolic syndrome. *Nature Reviews: Drug Discovery* 3:340-351.10.1038/nrd1344
- 497 Saez-Lara MJ, Robles-Sanchez C, Ruiz-Ojeda FJ, Plaza-Diaz J, and Gil A. 2016. Effects of  
498 Probiotics and Synbiotics on Obesity, Insulin Resistance Syndrome, Type 2 Diabetes

- and Non-Alcoholic Fatty Liver Disease: A Review of Human Clinical Trials. *Int J Mol Sci* 17.10.3390/ijms17060928
- Singleton V, and Rossi JN, M. H. 1965. Colorimetry of Total Phenolics with Phosphomolybdic-Phosphotungstic Acid Reagents. *Am J Enol Viticult* 16:144-158.10.1186/s13063-017-1796-8
- Suolang PC, Liu BQ, Chen J, De J, Nima ZB, and Dunzhu CR. 2019. Protective effect and mechanism of Qiwei Tiexie capsule on 3T3-L1 adipocytes cells and rats with nonalcoholic fatty liver disease by regulating LXRalpha, PPARgamma, and NF-kappaB-iNOS-NO signaling pathways. *Journal of Ethnopharmacology* 236:316-325.10.1016/j.jep.2019.03.006
- Xiao XH, Wang YD, Qi XY, Wang YY, Li JY, Li H, Zhang PY, Liao HL, Li MH, Liao ZZ, Yang J, Xu CX, Wen GB, and Liu JH. 2018. Zinc alpha2 glycoprotein protects against obesity-induced hepatic steatosis. *International Journal of Obesity (2005)* 42:1418-1430.10.1038/s41366-018-0151-9
- Yang Y, Wang J, Zhang Y, Li J, and Sun W. 2018. Black Sesame Seeds Ethanol Extract Ameliorates Hepatic Lipid Accumulation, Oxidative Stress, and Insulin Resistance in Fructose-Induced Nonalcoholic Fatty Liver Disease. *Journal of Agricultural and Food Chemistry* 66:10458-10469.10.1021/acs.jafc.8b04210
- Yasuda M, Yasutake K, Hino M, Ohwatari H, Ohmagari N, Takedomi K, Tanaka T, and Nonaka G. 2014. Inhibitory effects of polyphenols from water chestnut (*Trapa japonica*) husk on glycolytic enzymes and postprandial blood glucose elevation in mice. *Food Chemistry* 165:42-49.10.1016/j.foodchem.2014.05.083
- Zhang D, Xie T, and Leung PS. 2018. Irisin Ameliorates Glucolipotoxicity-Associated beta-Cell Dysfunction and Apoptosis via AMPK Signaling and Anti-Inflammatory Actions. *Cellular Physiology and Biochemistry* 51:924-937.10.1159/000495395
- Zhou B, Zhou DL, Wei XH, Zhong RY, Xu J, and Sun L. 2017. Astragaloside IV attenuates free fatty acid-induced ER stress and lipid accumulation in hepatocytes via AMPK activation. *Acta Pharmacologica Sinica* 38:998-1008.10.1038/aps.2016.175
- Zhou Y, Ding YL, Zhang JL, Zhang P, Wang JQ, and Li ZH. 2018. Alpinetin improved high fat diet-induced non-alcoholic fatty liver disease (NAFLD) through improving oxidative stress, inflammatory response and lipid metabolism. *Biomedicine and Pharmacotherapy* 97:1397-1408.10.1016/j.biopha.2017.10.035

**Table 1** (on next page)

Concentration of five polyphenols in TPQE by HPLC-DAD analysis (n=3)

**Table 1.** Concentration of five polyphenols in TPQE by HPLC-DAD analysis (n=3)






Peak number	Compound name	Concentration (mg/g)
1	Gallic acid	27.66 ± 0.99
2	1,2,3-tri- <i>O</i> -galloyl- $\beta$ -D-glucopyranose	19.89 ± 0.07
6	1,2,3-tetra- <i>O</i> -galloyl-4,6- <i>O</i> -hexahydroxydiphenoyl- $\beta$ -D-glucopyranoside	182.12 ± 2.38
7	1,2,3,6-tetra- <i>O</i> -galloyl- $\beta$ -D-glucopyranose	461.91±7.28
8	1,2,3,4,6-penta- <i>O</i> -hexahydroxydiphenoyl- $\beta$ -D-glucopyranoside	4.04±0.09



# Figure 1

Figures 1


The chromatogram of TQPE analyzed by HPLC-QTOF **(A)** and the structures of eight dominated polyphenols in TQPE **(B-E)**



# Figure 2

Figure 2


Body weight **(A)**, ratios of liver to body weight (%) **(B)**, appearance of the liver **(C-F)**, histological analysis of liver tissues (H&E, 200× magnification) **(G-J)**. All the results were presented as mean  $\pm$  SE (n=10).  $^{###} P < 0.001$  vs. the CON group;  $^{**} P < 0.01$ ,  $^{***} P < 0.001$  vs. the HFD group.



# Figure 3

Figure 3


Effects of TQPE in HFD-induced NAFLD on plasma total cholesterol (TC) **(A)**, triglycerides (TG) **(B)**, high-density lipoprotein cholesterol (HDL) **(C)**, and low-density lipoprotein cholesterol (LDL) **(D)** were measured. All the results were presented as mean  $\pm$  SE (n=10). #  $P < 0.05$ , ###  $P < 0.001$  vs. the CON group; \*\*  $P < 0.01$ , \*\*\*  $P < 0.001$  vs. the HFD group.



# Figure 4

Figure 4


TQPE administration inhibits oxidative stress and liver injury in NAFLD. The levels of SOD **(A)**, MDA **(B)**, ALT **(C)** and AST **(D)** in the serum were measured. All the results were presented as mean  $\pm$  SE (n=10).  $^{###} P < 0.001$  vs. the CON group;  $^{***} P < 0.001$  vs. the HFD group.



# Figure 5

Figure 5

Effects of TQPE on insulin and HOMA-IR in HFD-induced NAFLD. The levels of insulin **(A)** in the serum and HOMA-IR value **(B)** were measured. All the results were presented as mean  $\pm$  SE (n=10).  $^{###} P < 0.001$  vs. the CON group;  $^{***} P < 0.001$  vs. the HFD group.




# Figure 6

Figure 6

Effects of TQPE on AMPK, SREBP, ACC, IRS-1 and Akt in the liver tissue of HFD-induced NAFLD mice. p-AMPK, AMPK **(A-B)**, SREBP **(C-D)**, p-ACC, ACC **(E-F)**, p-IRS-1, IRS-1 **(G-H)**, p-Akt and Akt **(I-J)** expression were probed by western blotting. Results were presented as mean  $\pm$  SE ( $n = 3$ ).  $^{###} P < 0.001$  vs. the CON group;  $^{*} P < 0.05$ ,  $^{**} P < 0.01$ ,  $^{***} P < 0.001$  vs. the HFD group.





# Figure 7

Figure 7

A schematic illustration of how TQPE interfered with lipid accumulation and IR in HFD-induced NAFLD mouse.

