

Prognostic analysis of very early onset pancreatic cancer: a population based analysis

Dongjun Dai¹, Yanmei Wang¹, Xinyang Hu¹, Hongchuan Jin², Xian Wang^{Corresp. 1}

¹ Department of Medical Oncology, Sir Run Run Shaw Hospital, Medical School of Zhejiang University, Zhejiang University, Hangzhou, Zhejiang, China

² Laboratory of Cancer Biology, Key Lab of Biotherapy, Sir Run Run Shaw Hospital, Medical School of Zhejiang University, Zhejiang University, Hangzhou, Zhejiang, China

Corresponding Author: Xian Wang

Email address: wangx118@zju.edu.cn

Background : We aimed to use competing risk model to assess whether very early onset pancreatic cancer (VEOPC) (< 45 years) had a worse prognosis than older pancreatic cancer (PC) patients, and to build a competing risk nomogram for predicting the risk of death of VEOPC.

Methods: We selected pancreatic adenocarcinoma (PDAC) patients as our cohort from Surveillance, Epidemiology, and End Results (SEER) database. The impact of cancer specific death was estimated by competing risk analysis. Multivariate Fine-Gray regression for proportional hazards modeling of the subdistribution hazard (SH) model based nomogram was constructed, which was internally validated by discrimination and calibration with 1,000 bootstraps.

Results: Our cohort included 1,386 VEOPC patients and 53,940 older patients. We observed that in unresectablePDAC patients, VEOPC had better cancer specific survival (CSS) than each older group (45-59 years, 60-69 years, 70-79 years and > 79 years). There was no significant prognostic difference between VEOPC and each older group in resectablePDAC. Our competing nomogram showed well discrimination and calibration by internal validation.

Conclusion: For unresectable PDAC patients, VEOPC had better CSS than older patients. Our competing risk nomogram might be an easy-to-use tool for the specific death prediction of VEOPC patients with PDAC.

Prognostic analysis of very early onset pancreatic cancer: a population based analysis

Dongjun Dai¹, Yanmei Wang¹, Xinyang Hu¹, Hongchuan Jin², Xian Wang^{1,*}

¹ Department of Medical Oncology, Sir Run Run Shaw Hospital, Medical School of Zhejiang University, Zhejiang University, Hangzhou, China;

² Laboratory of Cancer Biology, Key Lab of Biotherapy, Sir Run Run Shaw Hospital, Medical School of Zhejiang University, Zhejiang University, Hangzhou, China

Corresponding Author:

Xian Wang ¹

Department of Medical Oncology, Sir Run Run Shaw Hospital, Medical School of Zhejiang University, Zhejiang University, 3 East Qingchun Road, Hangzhou, China

Email address: wangx118@zju.edu.cn

Abstract

Background: We aimed to use competing risk model to assess whether very early onset pancreatic cancer (VEOPC) (< 45 years) had a worse prognosis than older pancreatic cancer (PC) patients, and to build a competing risk nomogram for predicting the risk of death of VEOPC.

Methods: We selected pancreatic adenocarcinoma (PDAC) patients as our cohort from Surveillance, Epidemiology, and End Results (SEER) database. The impact of cancer specific death was estimated by competing risk analysis. Multivariate Fine-Gray regression for proportional hazards modeling of the subdistribution hazard (SH) model based nomogram was constructed, which was internally validated by discrimination and calibration with 1,000 bootstraps.

Results: Our cohort included 1,386 VEOPC patients and 53,940 older patients. We observed that in unresectable PDAC patients, VEOPC had better cancer specific survival (CSS) than each older group (45-59 years, 60-69 years, 70-79 years and > 79 years). There was no significant prognostic difference between VEOPC and each older group in resectable PDAC. Our competing nomogram showed well discrimination and calibration by internal validation.

Conclusion: For unresectable PDAC patients, VEOPC had better CSS than older patients. Our competing risk nomogram might be an easy-to-use tool for the specific death prediction of VEOPC patients with PDAC.

List of abbreviations: PC, pancreatic cancer; VEOPC, very early onset pancreatic cancer; BMI, body mass index; OS, overall survival; PDAC; pancreatic adenocarcinoma; SEER, Surveillance, Epidemiology, and End Results; NOS, not otherwise specified; LNR, lymph nodes ratio; AJCC, American Joint Committee on Cancer; CID, cumulative incidences of death; CIF, cumulative incidences function; CSS, cancer specific survival; HR, hazard ratio; 95%CI, 95%confidence index; AUC, the area under the curve of receiver operating characteristic curve.

Introduction

Pancreatic cancer (PC) patients have a 5 years survival lower than 10%(Siegel et al. 2018). According to American cancer society, the average age of PC patients at diagnosis is 70 years. Less than 3% of PC patients are under the age of 45 years at diagnosis (Surveillance 2019a), which was defined as very early onset PC (VEOPC)(McWilliams et al. 2016). Duo to its rareness, few data were reported regarding the clinicopathological features and treatment response of VEOPC patients(Grosfeld et al. 1990; Jaksic et al. 1992; Lack et al. 1983; McWilliams et al. 2016; Shorter et al. 2002).

The identification of prognostic factors of PC might improve the prediction of the survival and the selection of therapy for PC patients. The well-known PC prognostic factors comprised resection margins, lymph nodes status and lymph node ratio, perineural and blood vessel invasion, tumor localization and body mass index (BMI) and treatment such as chemotherapy(Bilici 2014; Lin et al. 2016). However, whether age is a prognostic factor of PC is currently controversial. Some studies found “younger” PC patients (with different definition of age < 45 years, age < 50 years, age < 60 years and age < 70 years) had better overall survival (OS) than the older PC patients(Baxter et al. 2007; He et al. 2013; Saad et al. 2018; Sugiura et al. 2017). While the other studies found younger age (with different definition of age < 65 years and age < 70 years)(Barbas et al. 2012; Miyazaki et al. 2016; van der Geest et al. 2016) was not associated with the survival of PC patients. To be noted, these studies had limited number of VEOPC patients or did not treat the VEOPC as a group.

Nomogram is a valuable tool to use different clinical variables to determine a statistical prognostic model that generates a probability of clinical outcomes for a single patient (Balachandran et al. 2015). Nomogram have been applied in various types of cancers (Fakhry et al. 2017; Huang et al. 2016; Liang et al. 2015; Rose et al. 2015). Nowadays, nomograms are commonly used to estimate prognosis in oncology and medicine, which fit the trend of personalized medicine (Balachandran et al. 2015). It is worthy to construct a nomogram for predicting the prognosis of VEOPC patients.

Pancreatic adenocarcinoma (PDAC) is the main type of PC, which has a worse prognosis than other histology types. The current study selected PDAC patients as our cohort. To be noted, the competing event of death is widely existing in the prognostic analysis of cancer. The failure to account for such competing events, such as Kaplan–Meier method and standard Cox proportional hazards regression analysis, might result in an overestimate of the cumulative incidences (Satagopan et al. 2004). The competing risk model considers both the disease-specific death and other causes of death, which was fit to the analysis in the presence of competing risk events (Kim 2007). The competing risk analysis was often missed in previous studies when included with very aged PC patients. The aim of our study was to use competing risk model to estimate the prognostic value of age in PDAC patients and construct competing nomogram for cancer-specific death prediction of VEOPC patients by using the data from Surveillance, Epidemiology, and End Results (SEER) database of the National Cancer Institute of United States, which currently collects and publishes cancer incidence and survival data from population-based cancer registries covering approximately 34.6 percent of the United States population (Surveillance 2019b).

Methods

Cohort selection

The cohort was selected by SEER*Stat 8.3.5 software (SEER ID: daid) with the following criteria: (1) it should be a primary PDAC patients (ICD-O-3 histology codes 8140 and 8500) registered equal or after year of 2004; (2) it should be a patient who has a survival time over than 0 month with records of survival status; (3) it should have at least one year follow up time to capture enough events to ensure a meaningful analysis. We also excluded few patients with unclear information of race, surgery, number of regional lymph nodes examined, radiotherapy and marital status. Finally, we excluded 1 patient with cancer in the islets of langerhans. The detailed data selection process was shown in Supplemental Figure 1.

Study variables and endpoints

We included 14 variables into our analysis, which were listed as the follows: age at diagnosis, sex (Females and males), race (Caucasian, African American, American Indian/Alaska Native and Asian or Pacific Islander), tumor location (Head of pancreas, body of pancreas, tail of pancreas, pancreatic duct, other specified parts of pancreas, overlapping lesion of pancreas and Pancreas, NOS (Not otherwise specified)), surgery experience (No surgery, local or partial pancreatectomy, local or partial pancreatectomy and duodenectomy, total pancreatectomy with or without gastrectomy or duodenectomy, pancreatectomy NOS or surgery NOS), tumor size, LNR (Lymph nodes ratio, which is defined as the ratio of the number of positive lymph nodes to the total number of lymph nodes examined), 6th American Joint Committee on Cancer (AJCC) tumor TNM stage, grade, radiotherapy and chemotherapy experience and marital status. The value of age at diagnosis, tumor size and LNR were classified into small categorical variables to fit the linear assumption. The widowed or single (Never married or having a domestic partner) or divorced or separated patients was defined as "Single". The median observed survival time was referred to median follow-up time.

Statistical analyses

The difference of each variable between VEOPC and other PC patients was evaluated by Chi-Squared tests. Cumulative incidences of death (CID) was estimated for deaths caused by cancer or other reasons by using the cumulative incidences function (CIF) analysis. Multivariate SH model was used to assess the cancer specific survival (CSS). All the variables were included in the multivariate analysis. Hazard ratio (HR) and 95%confidence index (95%CI) were calculated.

Multivariate SH model-based nomogram was constructed by multivariate logistic regression model to predict the 6 months, 12 months, 18 months and 24 months cancer specific death of VEOPC patients. The internal validation of nomogram was performed by discrimination and calibration with 1,000 time bootstraps(Balachandran et al. 2015). The discrimination was estimated by the area under the curve of receiver operating characteristic curve (AUC, which was also referred as C-statistics). The AUC ranges from 0.5-1.0, with 0.5 indicates the outcomes is totally random and 1.0 indicates the perfect discrimination. The calibration was assessed by calibration curves, which shows how close the nomogram estimated risk was to the observed risk. All the statistical analyses were performed by "R" version 3.6.0. The CIF test and multivariate SH analysis were performed by "R" package "cmprsk". The competing risk nomogram was constructed by

“R” package “mstate” and plotted by “R” package “regplot”. The calibration curve was drawn by “R” package “riskRegression”. The AUC was calculated by “R” package “riskRegression” and plotted by “R” package “ggplot2”. A two tailed p-value less than 0.05 was considered statistically significant.

Results

Cohort selection

We included the patients diagnosed equal to or before 2015 to ensure an enough follow-up time. There were 1,386 VEOPC patients and 53,940 older PDAC patients (≥ 45 years). The median survival time of VEOPC and older patients were 9 and 7 months, respectively. Significant differences were found between VEOPC and other PC patients. The VEOPC patients had more males, fewer Caucasians, more surgery rates, fewer LNR, higher TNM stage, higher grade, more experience of chemotherapy and radiotherapy and fewer of married status ($p < 0.001$, Table 1) than the older patients. We also provided a table with characteristics for all patients (Supplemental Table 1). We divided the older patients into 4 groups with different age ranges (45-59 years, 60-69 years, 70-79 years and >79 years) for further analysis.

Prognostic analysis of age in PDAC patients

The CIF plot showed VEOPC patients had a decreased risk of cancer death than older patients ($p < 0.001$, Figure 1a). Further subgroup analyses by surgery showed that VEOPC patients had better prognosis than the older patients whether they performed surgery or not ($p < 0.001$, Figure 1b-c). Moreover, multivariate SH model with all the included variables showed that in unresectable PDAC patients, the VEOPC had better CSS than older patients. While in the resected PDAC patients, no significant difference was found between VEOPC and older patients (Table 2).

Independent prognostic factors of CSS of VEOPC

As shown in Table 3, multivariate SH model was used to find independent prognostic factors for the CSS of VEOPC patients. We found the female, race of Asian or Pacific Islander, PC located in pancreatic duct, no surgery, high LNR, high M stage, high grade, lack of chemotherapy, and single status were risk factors of the VEOPC patients ($P < 0.05$).

Nomogram development and validation

The nomogram was constructed by multivariate SH model. A weighted total score calculated from each variable was used to estimate the 6 months, 12 months, 18 months and 24 months cancer specific death of VEOPC patients (Figure 2). The sex, race, location, surgery strategy, tumor size, LNR, TNM stage, grade, chemotherapy and marital status were found to be significantly important for the prognosis of VEOPC patients (Variables with asterisk in Figure 2). The calibration plots showed well correlation between observed CSS and nomogram predicted CSS (Figure 3a-d). Time dependent AUC plot suggested well discrimination of our nomogram ($AUC \geq 0.736$, Figure 3e). Furthermore, we equally stratified the whole VEOPC into three groups based on their level of competing nomogram scores and performed a CIF analysis. The CIF plot (Figure 4) showed VEOPC with higher scores of our competing nomogram was at high risk of cancer related death. It indicated that our competing nomogram was useful for risk stratification of the prognosis of VEOPC patients. The detailed competing nomogram scores for all variables were listed in Supplemental Table 2.

Discussion

No previous competing analysis was performed to analyze the prognostic difference between VEOPC and older PC patients. The current study used competing risk model to estimate the prognostic difference between VEOPC and older patients by using the data from SEER. We found that among the unresectable PDAC patients, the VEOPC had better CSS than older patients. And among resected PDAC patients, there was no significant prognostic difference between VEOPC and older patients. We also built the first competing nomogram for primary VEOPC patients with histology type of PDAC.

Age was observed as an independent prognostic factor for many cancers. Previous studies identified that younger patients had worse prognosis than older patients in colorectal cancer (van Eeghen et al. 2015) and breast cancer (Chen et al. 2016) while younger patients had better prognosis than older patients in lung cancer (Arnold et al. 2016), liver cancer (Zhang & Sun 2015) and prostate cancer (Pettersson et al. 2018). A study found that the <45 years PDAC patients had better survival than > 70 years PDAC patients (He et al. 2013). Another study found <50 years PDAC patients had no significant difference on prognosis when compared with their whole cohort (Tingstedt et al. 2011). To be noted, these studies only included with limited samples. A SEER based study found the risk of cancer-specific death decreases with age in resectable PDAC patients (He et al. 2018). However, we found the age was not an independent prognostic factor for resectable PDAC patients. This difference might be caused by the difference of inclusion

criteria that our study included more variables in the multivariate SH model. We provided a detailed inclusion workflow in the Supplemental Figure 1. We found the VEOPC patients, despite with a higher TNM stage, had a better CSS than the older patients in unresectable patients (Supplemental Table 3). It might be explained by the well accepted fact that younger patients had more tolerance to chemotherapy. Indeed, in the unresectable group, the VEOPC patients had a 17.5% higher rate to have chemotherapy than the older patients.

Males had higher rate of PC incidence in United States(Yadav & Lowenfels 2013). We observed the males had 15% higher rate than the females in VEOPC. Previous study found males had worse prognosis than females in patients under 60 years but not in VEOPC patients(McWilliams et al. 2016). It might be caused by the sample size. Our study had much higher amount of VEOPC patients than this study and we found the males had worse CSS than females.

Marriage status was linked to the improvements in cardiovascular, endocrine, immune function, and cancer prognosis(Aizer et al. 2013; Gallo et al. 2003; Herberman & Ortaldo 1981). Previous study observed that unmarried patients were at significantly higher risk of presentation with metastatic cancer, undertreatment, and death resulting from their cancer(Aizer et al. 2013). We found the VEOPC patients with married status had better CSS than patients with single status, which provided new hint to the association between support care and cancer prognosis.

Our SH model based nomogram was well validated by discrimination and calibration. The variables included in our nomogram were easy to be obtained, allowing a feasible translation into the future clinical use. To be noted, we included both the significant variables and the insignificant variables into our nomogram, since if a nomogram only includes statistically significant variables, it might tend to exert an inappropriately large influence, leading to falsely narrowed confidence intervals, which would make this nomogram appear more accurate than it is(Nicholas J Vogelzang 2005). Patients with higher competing nomogram scores of our study were found to have higher CID, indicating that our nomogram was useful for risk stratification of the prognosis of VEOPC patients.

Our study had some limitations. First, the detailed strategy of chemotherapy or radiotherapy was missed in current study as well as other therapies such as immunotherapy. Second, although the current nomogram was well validated by internal validation, there was a lack of an external validation by cohort other than SEER. Third, it should be noted that only factors available in the SEER registry were examined. The results from this study might only fit the population from United States. Fourth, our study

has no exploration of molecular mechanism, which was of interest but cannot be conducted at this time with data currently available.

Conclusions

The current study showed that for unresectable PDAC, VEOPC had better CSS than older patients. For the VEOPC patients, we built a SH model based nomogram to estimate their CSS. This nomogram was shown to have good discrimination and calibration.

References

- Aizer AA, Chen MH, McCarthy EP, Mendu ML, Koo S, Wilhite TJ, Graham PL, Choueiri TK, Hoffman KE, Martin NE, Hu JC, and Nguyen PL. 2013. Marital status and survival in patients with cancer. *J Clin Oncol* 31:3869-3876. 10.1200/JCO.2013.49.6489
- Arnold BN, Thomas DC, Rosen JE, Salazar MC, Blasberg JD, Boffa DJ, Detterbeck FC, and Kim AW. 2016. Lung Cancer in the Very Young: Treatment and Survival in the National Cancer Data Base. *J Thorac Oncol* 11:1121-1131. 10.1016/j.jtho.2016.03.023
- Balachandran VP, Gonen M, Smith JJ, and DeMatteo RP. 2015. Nomograms in oncology: more than meets the eye. *Lancet Oncol* 16:e173-180. 10.1016/S1470-2045(14)71116-7
- Barbas AS, Turley RS, Ceppa EP, Reddy SK, Blazer DG, 3rd, Clary BM, Pappas TN, Tyler DS, White RR, and Lagoo SA. 2012. Comparison of outcomes and the use of multimodality therapy in young and elderly people undergoing surgical resection of pancreatic cancer. *J Am Geriatr Soc* 60:344-350. 10.1111/j.1532-5415.2011.03785.x
- Baxter NN, Whitson BA, and Tuttle TM. 2007. Trends in the treatment and outcome of pancreatic cancer in the United States. *Ann Surg Oncol* 14:1320-1326. 10.1245/s10434-006-9249-8
- Bilici A. 2014. Prognostic factors related with survival in patients with pancreatic adenocarcinoma. *World J Gastroenterol* 20:10802-10812. 10.3748/wjg.v20.i31.10802
- Chen HL, Zhou MQ, Tian W, Meng KX, and He HF. 2016. Effect of Age on Breast Cancer Patient Prognoses: A Population-Based Study Using the SEER 18 Database. *PLoS One* 11:e0165409. 10.1371/journal.pone.0165409
- Fakhry C, Zhang Q, Nguyen-Tan PF, Rosenthal DI, Weber RS, Lambert L, Trotti AM, 3rd, Barrett WL, Thorstad WL, Jones CU, Yom SS, Wong SJ, Ridge JA, Rao SSD, Bonner JA, Vigneault E, Raben D, Kudrimoti MR, Harris J, Le QT, and Gillison ML. 2017. Development and Validation of Nomograms Predictive of Overall and Progression-Free Survival in Patients With Oropharyngeal Cancer. *J Clin Oncol* 35:4057-4065. 10.1200/JCO.2016.72.0748
- Gallo LC, Troxel WM, Matthews KA, and Kuller LH. 2003. Marital status and quality in middle-aged women: Associations with levels and trajectories of cardiovascular risk factors. *Health Psychol* 22:453-463. 10.1037/0278-6133.22.5.453
- Grosfeld JL, Vane DW, Rescorla FJ, McGuire W, and West KW. 1990. Pancreatic tumors in childhood: analysis of 13 cases. *J Pediatr Surg* 25:1057-1062.

He C, Zhang Y, Cai Z, Lin X, and Li S. 2018. Overall survival and cancer-specific survival in patients with surgically resected pancreatic head adenocarcinoma: A competing risk nomogram analysis. *J Cancer* 9:3156-3167. 10.7150/jca.25494

He J, Edil BH, Cameron JL, Schulick RD, Hruban RH, Herman JM, Zheng L, Iacobuzio-Donahue C, Ahuja N, Pawlik TM, and Wolfgang CL. 2013. Young patients undergoing resection of pancreatic cancer fare better than their older counterparts. *J Gastrointest Surg* 17:339-344. 10.1007/s11605-012-2066-4

Herberman RB, and Ortaldo JR. 1981. Natural killer cells: their roles in defenses against disease. *Science* 214:24-30. 10.1126/science.7025208

Huang YQ, Liang CH, He L, Tian J, Liang CS, Chen X, Ma ZL, and Liu ZY. 2016. Development and Validation of a Radiomics Nomogram for Preoperative Prediction of Lymph Node Metastasis in Colorectal Cancer. *J Clin Oncol* 34:2157-2164. 10.1200/JCO.2015.65.9128

Jaksic T, Yaman M, Thorner P, Wesson DK, Filler RM, and Shandling B. 1992. A 20-year review of pediatric pancreatic tumors. *J Pediatr Surg* 27:1315-1317.

Kim HT. 2007. Cumulative incidence in competing risks data and competing risks regression analysis. *Clin Cancer Res* 13:559-565. 10.1158/1078-0432.CCR-06-1210

Lack EE, Cassady JR, Levey R, and Vawter GF. 1983. Tumors of the exocrine pancreas in children and adolescents. A clinical and pathologic study of eight cases. *Am J Surg Pathol* 7:319-327.

Liang W, Zhang L, Jiang G, Wang Q, Liu L, Liu D, Wang Z, Zhu Z, Deng Q, Xiong X, Shao W, Shi X, and He J. 2015. Development and validation of a nomogram for predicting survival in patients with resected non-small-cell lung cancer. *J Clin Oncol* 33:861-869. 10.1200/JCO.2014.56.6661

Lin JY, Zhang XM, Kou JT, Fa H, Zhang XX, Dai Y, and He Q. 2016. Analysis of prognostic factors for pancreatic head cancer according to para-aortic lymph node. *Cancer Med* 5:2701-2707. 10.1002/cam4.853

McWilliams RR, Maisonneuve P, Bamlet WR, Petersen GM, Li D, Risch HA, Yu H, Fontham ET, Lockett B, Bosetti C, Negri E, La Vecchia C, Talamini R, Bueno de Mesquita HB, Bracci P, Gallinger S, Neale RE, and Lowenfels AB. 2016. Risk Factors for Early-Onset and Very-Early-Onset Pancreatic Adenocarcinoma: A Pancreatic Cancer Case-Control Consortium (PanC4) Analysis. *Pancreas* 45:311-316. 10.1097/MPA.0000000000000392

Miyazaki Y, Kokudo T, Amikura K, Kageyama Y, Takahashi A, Ohkohchi N, and Sakamoto H. 2016. Age does not affect complications and overall survival rate after pancreaticoduodenectomy: Single-center experience and systematic review of literature. *Biosci Trends* 10:300-306. 10.5582/bst.2016.01093

Nicholas J Vogelzang PTS, William U Shipley, Frans MJ Debruyne, W. Marston Linehan. 2005. *Comprehensive Textbook of Genitourinary Oncology 3rd Edition*: LWW; Third edition (September 15, 2005).

Pettersson A, Robinson D, Garmo H, Holmberg L, and Stattin P. 2018. Age at diagnosis and prostate cancer treatment and prognosis: a population-based cohort study. *Ann Oncol* 29:377-385. 10.1093/annonc/mdx742

Rose PG, Java J, Whitney CW, Stehman FB, Lanciano R, Thomas GM, and DiSilvestro PA. 2015. Nomograms Predicting Progression-Free Survival, Overall Survival, and Pelvic Recurrence in Locally Advanced Cervical Cancer Developed From an Analysis of Identifiable Prognostic Factors in Patients From NRG Oncology/Gynecologic Oncology Group Randomized Trials of Chemoradiotherapy. *J Clin Oncol* 33:2136-2142. 10.1200/JCO.2014.57.7122

Saad AM, Turk T, Al-Husseini MJ, and Abdel-Rahman O. 2018. Trends in pancreatic adenocarcinoma incidence and mortality in the United States in the last four decades; a SEER-based study. *BMC Cancer* 18:688. 10.1186/s12885-018-4610-4

- Satagopan JM, Ben-Porat L, Berwick M, Robson M, Kutler D, and Auerbach AD. 2004. A note on competing risks in survival data analysis. *Br J Cancer* 91:1229-1235. 10.1038/sj.bjc.6602102
- Shorter NA, Glick RD, Klimstra DS, Brennan MF, and Laquaglia MP. 2002. Malignant pancreatic tumors in childhood and adolescence: The Memorial Sloan-Kettering experience, 1967 to present. *J Pediatr Surg* 37:887-892.
- Siegel RL, Miller KD, and Jemal A. 2018. Cancer statistics, 2018. *CA Cancer J Clin* 68:7-30. 10.3322/caac.21442
- Sugiura T, Okamura Y, Ito T, Yamamoto Y, Ashida R, and Uesaka K. 2017. Impact of Patient Age on the Postoperative Survival in Pancreatic Head Cancer. *Ann Surg Oncol* 24:3220-3228. 10.1245/s10434-017-5994-0
- Surveillance E, and End Results Program. 2019a. Cancer Stat Facts: Pancreatic Cancer. Available at <https://seer.cancer.gov/statfacts/html/pancreas.html>.
- Surveillance E, and End Results Program. 2019b. Overview of the SEER Program. Available at <https://seer.cancer.gov/about/overview.html> (accessed November 28th 2019).
- Tingstedt B, Weitkamper C, and Andersson R. 2011. Early onset pancreatic cancer: a controlled trial. *Ann Gastroenterol* 24:206-212.
- van der Geest LG, Besselink MG, van Gestel YR, Busch OR, de Hingh IH, de Jong KP, Molenaar IQ, and Lemmens VE. 2016. Pancreatic cancer surgery in elderly patients: Balancing between short-term harm and long-term benefit. A population-based study in the Netherlands. *Acta Oncol* 55:278-285. 10.3109/0284186X.2015.1105381
- van Eeghen EE, Bakker SD, van Bochove A, and Loffeld RJ. 2015. Impact of age and comorbidity on survival in colorectal cancer. *J Gastrointest Oncol* 6:605-612. 10.3978/j.issn.2078-6891.2015.070
- Yadav D, and Lowenfels AB. 2013. The epidemiology of pancreatitis and pancreatic cancer. *Gastroenterology* 144:1252-1261. 10.1053/j.gastro.2013.01.068
- Zhang W, and Sun B. 2015. Impact of age on the survival of patients with liver cancer: an analysis of 27,255 patients in the SEER database. *Oncotarget* 6:633-641. 10.18632/oncotarget.2719

Figure legends

Figure 1. CIF plots of the association between age and the prognosis of PDAC patients
Legends: a-c, The CIF plots of the association between age and the prognosis of all cohort (a), patients with surgery (b) or without surgery (c).

Figure 2. Nomogram for predicting 6 months, 12 months, 18 months and 24 months cancer specific death of VEOPC patients

Legends: The nomogram is used by summing the points identified on the top scale for each independent variable and drawing a vertical line from the total points scale to the 6 months, 12 months, 18 months and 24 months cancer specific death to obtain the probability of survival. The total points projected to the bottom scale indicate the probability of the 6 months, 12 months, 18 months and 24 months cancer specific death. Sex: 1 = Male, 2 = Female; Race: 1 = Caucasian, 2 = African American, N = Other race;

Location: 250 = Head of pancreas, 251 = Body of pancreas, 252 = Tail of pancreas, 253 = Pancreatic duct, 257 = Other specified parts of pancreas, 258 = Overlapping lesion of pancreas, 259 = Pancreas, NOS; Surgery, 0 = No surgery, 1 = Local or partial pancreatectomy, 2 = Local or partial pancreatectomy and duodenectomy, 3 = Total pancreatectomy with or without gastrectomy or duodenectomy, 4 = Pancreatectomy NOS or surgery NOS; Tumor size: 2 = ≤ 2 cm, 4 = 2cm to 4cm, 6 = 4cm to 6cm, 8 = > 6 cm, 9 = Unknown; LNR: 0 = 0, 1 = ≤ 0.2 , 2 = 0.2-0.4, 3 = > 0.4 , 4 = No nodes were examined; T stage: 1 = T1-T2, 2 = T3-T4, 9 = Unknown; N stage: 0 = NO, 1 = N1, 9 = Unknown; M stage: 0 = M0, 1 = M1, 9 = Unknown; Grade: 1 = I-II, 2 = III-IV, 9 = Unknown; Chemotherapy: 0 = none/unknown and 1 = yes; Radiotherapy: 0 = none/unknown or refused, 1 = beam radiation or combination of beam with implants or isotopes or radiation with method or source not specified or radioactive implants or radioisotopes and N = Recommended, unknown if administered; Marital status: 0 = married; 1 = widowed or single (never married or having a domestic partner) or divorced or separated. The asterisk means significant.

Figure 3. Calibration curves and time dependent AUC plot for internal validation of SH based nomograms

Legends: a-d, the calibration plots for 6 months (a), 12 months (b), 18 months (c) and 24 months (d) cancer specific death prediction of VEOPC patients; The x-axis shows the nomogram predicted probability while the y-axis is the actual survival estimated by the SH method, the black thick line overlaps the grey line indicating near perfect calibration; e, time dependent AUC plots of SH-based nomograms.

Figure 4. CIF plot based on the competing nomogram scores for the VEOPC patients

Legends: the whole cohort of VEOPC patients were divided into three equally distributed groups. The groups were named “High”, “Intermediate” and “Low” according to the levels of patients’ competing nomogram scores.

Table 1(on next page)

The demographic and treatment data of VEOPC and older PC (≥ 45 years) patients

All variables showed significant differences between VEOPC and older PC patients ($p < 0.001$).

Table 1. The demographic and treatment data of VEOPC and older PC (>=45 years) patients

Characteristics	VEOPC (N = 1386)		Older PC (N = 53932)	
	No. of patients	%	No. of patients	%
Sex				
Male	796	57.4%	27337	50.7%
Female	590	42.6%	26595	49.3%
Race				
Caucasian	1026	74.0%	43089	79.9%
African American	207	14.9%	6522	12.1%
American Indian/Alaska Native	13	0.9%	287	0.5%
Asian or Pacific Islander	140	10.1%	4034	7.5%
Location				
Head of pancreas	738	53.2%	29410	54.5%
Body of pancreas	154	11.1%	6921	12.8%
Tail of pancreas	182	13.1%	6458	12.0%
Pancreatic duct	12	0.9%	319	0.6%
Other specified parts of pancreas	18	1.3%	805	1.5%
Overlapping lesion of pancreas	113	8.2%	4049	7.5%
Pancreas, NOS	169	12.2%	5970	11.1%
Surgery				
No surgery	1040	75.0%	41989	77.9%
Local or partial pancreatectomy	43	3.1%	1517	2.8%
Local or partial pancreatectomy and duodenectomy	242	17.5%	8175	15.2%
Total pancreatectomy with or without gastrectomy or duodenectomy	50	3.6%	1978	3.7%
Pancreatectomy NOS or surgery NOS	11	0.8%	273	0.5%
Tumor size (cm)				
<=2	144	10.4%	4883	9.1%
2 to 4	524	37.8%	23278	43.2%
4 to 6	333	24.0%	12388	23.0%
>6	104	7.5%	3840	7.1%
Unknown	281	20.3%	9543	17.7%

LNR				
0	137	9.9%	4542	8.4%
<=0.2	126	9.1%	3983	7.4%
0.2-0.4	57	4.1%	2180	4.0%
0.4-1	55	4.0%	2116	3.9%
No nodes were examined	1011	72.9%	41111	76.2%
T stage				
T1-T2	275	19.8%	11774	21.8%
T3-T4	865	62.4%	33301	61.7%
Unknown	246	17.7%	8857	16.4%
N stage				
NO	603	43.5%	26845	49.8%
N1	584	42.1%	19325	35.8%
Unknown	199	14.4%	7762	14.4%
M stage				
M0	582	42.0%	26570	49.3%
M1	786	56.7%	25678	47.6%
Unknown	18	1.3%	1684	3.1%
Grade				
I-II	326	23.5%	11939	22.1%
III-IV	256	18.5%	8755	16.2%
Unknown	804	58.0%	33238	61.6%
Chemotherapy				
No	312	22.5%	20586	38.2%
Yes	1074	77.5%	33346	61.8%
Radiotherapy				
No	1021	73.7%	42283	78.4%
Yes	365	26.3%	11649	21.6%
Marital status				
Married	768	55.4%	32013	59.4%
Single	618	44.6%	21919	40.6%

3 All variables showed significant differences between VEOPC and older PC patients ($p < 0.001$).

Table 2 (on next page)

Multivariate SH analysis of CSS for the association between age and the prognosis of PDAC patients

We included all variables in the SH multivariate analysis. HR, Hazard ratio; 95%CI, 95%confidence index; Significant results ($p < 0.05$) were bolded.

1 **Table 2. Multivariate SH analysis of CSS for the association between age and the prognosis of PDAC patients**

Age (Years)	For the whole cohort		For patients with surgery		For patients without surgery	
	HR (95%CI)	p value	HR (95%CI)	p value	HR (95%CI)	p value
<45	Reference		Reference		Reference	
45-59	1.08 (1.02-1.14)	0.010	0.98 (0.86-1.12)	7.50E-01	1.10 (1.03-1.16)	0.003
60-69	1.10 (1.04-1.17)	<0.001	1.01 (0.89-1.15)	9.00E-01	1.13 (1.06-1.20)	<0.001
70-79	1.16 (1.09-1.23)	<0.001	1.04 (0.91-1.19)	5.70E-01	1.19 (1.12-1.26)	<0.001
>79	1.17 (1.10-1.24)	<0.001	1.00 (0.87-1.17)	9.60E-01	1.19 (1.12-1.27)	<0.001

2 We included all variables in the SH multivariate analysis. HR, Hazard ratio; 95%CI, 95%confidence index; Significant results ($p < 0.05$) were bolded.

Table 3(on next page)

Multivariate SH analyses of CSS for each variable in VEOPC patients

We included all variables in the SH multivariate analysis. HR, Hazard ratio; 95%CI, 95%confidence index; Significant results ($p < 0.05$) were bolded.

1 **Table 3. Multivariate SH analyses of CSS for each variable in VEOPC patients**

Characteristics	HR (95%CI)	p value
Sex		
Male	Reference	
Female	0.87 (0.77-0.98)	0.021
Race		
Caucasian	Reference	
African American	1.02 (0.84-1.23)	0.880
American Indian/Alaska Native	1.00 (0.67-1.50)	1.000
Asian or Pacific Islander	1.54 (1.30-1.83)	<0.001
Location		
Head of pancreas	Reference	
Body of pancreas	0.89 (0.73-1.08)	0.220
Tail of pancreas	1.07 (0.87-1.32)	0.530
Pancreatic duct	2.01 (1.34-3.02)	0.001
Other specified parts of pancreas	0.65 (0.33-1.29)	0.220
Overlapping lesion of pancreas	0.92 (0.72-1.18)	0.520
Pancreas, NOS	1.28 (1.03-1.60)	0.025
Surgery		
No surgery	Reference	
Local or partial pancreatectomy	0.40 (0.26-0.61)	<0.001
Local or partial pancreatectomy and duodenectomy	0.46 (0.36-0.60)	<0.001
Total pancreatectomy with or without gastrectomy or duodenectomy	0.41 (0.28-0.61)	<0.001
Pancreatectomy NOS or surgery NOS	0.77 (0.36-1.64)	0.490
Tumor size		
<=2cm	Reference	
2 to 4	1.07 (0.87-1.32)	0.510
4 to 6	1.11 (0.88-1.41)	0.360
>6	1.28 (0.94-1.74)	0.110
Unknown	1.01 (0.78-1.32)	0.930
LNR		
0	Reference	
<=0.2	1.39 (1.02-1.90)	0.037
0.2-0.4	1.80 (1.30-2.51)	<0.001
0.4-1	1.90 (1.37-2.64)	<0.001
No nodes were examined	1.33 (1.04-1.70)	0.021
T stage		
T1-T2	Reference	
T3-T4	1.16 (0.98-1.37)	0.082
Unknown	0.95 (0.73-1.22)	0.680

N stage		
NO	Reference	
N1	1.12 (0.95-1.31)	0.170
Unknown	1.34 (1.09-1.66)	0.005
M stage		
M0	Reference	
M1	1.48 (1.25-1.75)	<0.001
Unknown	1.09 (0.60-1.96)	0.780
Grade		
I-II	Reference	
III-IV	1.29 (1.07-1.56)	0.007
Unknown	1.09 (0.93-1.29)	0.290
Chemotherapy		
No	Reference	
Yes	0.63 (0.53-0.76)	<0.001
Radiotherapy		
No	Reference	
Yes	1.09 (0.94-1.25)	0.250
Marital status		
Married	Reference	
Single	1.15 (1.02-1.30)	0.021

- 2 We included all variables in the SH multivariate analysis. HR, Hazard ratio; 95%CI, 95%confidence index;
- 3 Significant results (p < 0.05) were bolded.

Figure 1

CIF plots of the association between age and the prognosis of PDAC patients

a-c, The CIF plots of the association between age and the prognosis of all cohort (a), patients with surgery (b) or without surgery (c).





Figure 2

Nomogram for predicting 6 months, 12 months, 18 months and 24 months cancer specific death of VEOPC patients

The nomogram is used by summing the points identified on the top scale for each independent variable and drawing a vertical line from the total points scale to the 6 months, 12 months, 18 months and 24 months cancer specific death to obtain the probability of survival. The total points projected to the bottom scale indicate the probability of the 6 months, 12 months, 18 months and 24 months cancer specific death. Sex: 1 = Male, 2 = Female; Race: 1 = Caucasian, 2 = African American, N = Other race; Location: 250 = Head of pancreas, 251 = Body of pancreas, 252 = Tail of pancreas, 253 = Pancreatic duct, 257 = Other specified parts of pancreas, 258 = Overlapping lesion of pancreas, 259 = Pancreas, NOS; Surgery, 0 = No surgery, 1 = Local or partial pancreatectomy, 2 = Local or partial pancreatectomy and duodenectomy, 3 = Total pancreatectomy with or without gastrectomy or duodenectomy, 4 = Pancreatectomy NOS or surgery NOS; Tumor size: 2 = ≤ 2 cm, 4 = 2cm to 4cm, 6 = 4cm to 6cm, 8 = > 6 cm, 9 = Unknown; LNR: 0 = 0, 1 = ≤ 0.2 , 2 = 0.2-0.4, 3 = > 0.4 , 4 = No nodes were examined; T stage: 1 = T1-T2, 2 = T3-T4, 9 = Unknown; N stage: 0 = NO, 1 = N1, 9 = Unknown; M stage: 0 = M0, 1 = M1, 9 = Unknown; Grade: 1 = I-II, 2 = III-IV, 9 = Unknown; Chemotherapy: 0 = none/unknown and 1 = yes; Radiotherapy: 0 = none/unknown or refused, 1 = beam radiation or combination of beam with implants or isotopes or radiation with method or source not specified or radioactive implants or radioisotopes and N = Recommended, unknown if administered; Marital status: 0 = married; 1 = widowed or single (never married or having a domestic partner) or divorced or separated. The asterisk means significant.



Total-points-to-outcome nomogram:




Figure 3

Calibration curves and AUC plots for internal validation of SH based nomograms

a-d, the calibration plots for 6 months (a), 12 months (b), 18 months (c) and 24 months (d) cancer specific death prediction of VEOPC patients; The x-axis shows the nomogram predicted probability while the y-axis is the actual survival estimated by the SH method, the black thick line overlaps the grey line indicating near perfect calibration; e, time dependent AUC plots of SH-based nomograms.




Figure 4

CIF plots based on the competing nomogram scores for the VEOPC patients

the whole cohort of VEOPC patients were divided into three equally distributed groups. The groups were named “High”, “Intermediate” and “Low” according to the levels of patients’ competing nomogram scores.

