

Evaluation of a new Argovit as an antiviral agent included in feed to protect the shrimp *Litopenaeus vannamei* against White Spot Syndrome Virus infection

Carlos R Romo Quiñonez¹, Ana R Álvarez Sánchez², Pindaro Álvarez-Ruiz³, Maria C Chávez Sánchez⁴, Nina Bogdanchikova⁵, Alexey Pestryakov⁶, Claudio H Mejia-Ruiz^{Corresp. 7}

¹ Laboratorio Biotecnología de Organismos Marinos, Programa de Acuicultura, Centro de Investigaciones Biológicas del Noroeste, La Paz, BCS, México

² Universidad Técnica Estatal de Quevedo, Quevedo, Los Rios, Ecuador

³ Departamento de Acuicultura, Centro Interdisciplinario de Investigación para el Desarrollo Integral Regional, I.P.N., Guasave, Sinaloa, México

⁴ Unidad de Acuicultura y Manejo Ambiental, Centro de Investigación en Alimentación y Desarrollo, Mazatlan, Sinaloa, México

⁵ Centro de Nanociencias y Nanotecnología, Universidad Nacional Autónoma de México, Ensenada, Baja California, México

⁶ Tomsk Polytechnical University, Tomsk, Russia

⁷ Laboratorio Biotecnología de Organismos Marinos, Programa de Acuicultura., Centro de Investigaciones Biológicas del Noroeste, La Paz, BCS, México

Corresponding Author: Claudio H Mejia-Ruiz

Email address: hmejia04@cibnor.mx

In this study, four experimental assays were conducted to evaluate the use of a new silver nanoparticle formulation named Argovit-4, which was prepared with slight modifications to enhance its biological activity against white spot syndrome virus (WSSV) in shrimp culture. The goals of these assays were to (1) determine the protective effect of Argovit-4 against WSSV, (2) determine whether Argovit-4 supplemented in feed exhibits toxicity towards shrimp, (3) determine whether Argovit-4 as antiviral additive in feed can prevent or delay/reduce WSSV-induced shrimp mortality, and (4) determine whether Argovit-4 supplemented in feed alters the early stages of the shrimp immune response. In bioassay 1, several viral inocula calibrated at 7 shrimp infectious doses (SID) were exposed to 40, 100, 200 and 1000 ng/SID of Ag+ and then intramuscularly injected into shrimp for 96 h. In assay 2, shrimp were fed Argovit-4 supplemented in feed at different concentrations (10, 100 and 1000 µg per gram of feed) for 192 h. In assay 3, shrimp were treated with Argovit-4 supplemented in feed at different concentrations and then challenged against WSSV for 192 h. In bioassay 4, quantitative real-time RT-qPCR was performed to measure the transcriptional responses of five immune-relevant genes in the haemocytes of experimental shrimp treated with Argovit-4 supplemented in feed at 0, 6, 12, 24 and 48 h. The intramuscularly injected Argovit-4 showed a dose-dependent effect ($p < 0.05$) on the cumulative shrimp mortality from 0-96 h post-infection. In the second bioassay, shrimp fed Argovit-4 supplemented in feed did not show signs of toxicity for the assayed doses over the 192-h experiment. The third and fourth bioassays showed that shrimp challenged with

WSSV at 1000 µg/g feed exhibited reduced mortality without altering the expression of some immune system-related genes according to the observed level of transcriptional. This study is the first show that the new Argovit-4 formulation has potential as an antiviral additive in feed against WSSV and demonstrates a practical therapeutic strategy to control WSSV and possibly other invertebrate pathogens in shrimp aquaculture.

Evaluation of a new Argovit as an antiviral agent included in feed to protect the shrimp *Litopenaeus vannamei* against White Spot Syndrome Virus infection

Carlos Rosendo Romo-Quiñonez¹, Ana Ruth Álvarez-Sánchez², Píndaro Alvarez-Ruiz^{3*}, María Cristina Chávez-Sánchez⁴, Nina Bogdanchikova⁵, Alexey Pestryakov⁶, Claudio Humberto Mejía-Ruiz^{1*}

1. Laboratorio Biotecnología de Organismos Marinos, Centro de Investigaciones Biológicas del Noroeste, S. C. La Paz, B.C.S. México.
2. Universidad Técnica Estatal de Quevedo. Quevedo, Los Ríos, Ecuador.
3. Departamento de Acuicultura, Centro Interdisciplinario de Investigación para el Desarrollo Integral Regional, I.P.N. Guasave, Sinaloa. México.
4. Unidad de Acuicultura y Manejo Ambiental, Centro de Investigación en Alimentación y Desarrollo, A.C. Mazatlán, Sinaloa. México.
5. Centro de Nanociencias y Nanotecnología, Universidad Nacional Autónoma de México, Ensenada, México.
6. Tomsk Polytechnic University, Tomsk, Russia.

*Corresponding Author 1:

Claudio Humberto Mejía-Ruiz,

IPN 195, Col. Playa Palo de Santa Rita Sur, La Paz, B.C.S., 23096, México

Email address: hmejia04@cibnor.mx

*Corresponding Author 2:

Píndaro Álvarez-Ruiz

E-mail: pindaroalvarez@hotmail.com

Abstract

In this study, four experimental assays were conducted to evaluate the use of a new silver nanoparticle formulation named Argovit-4, which was prepared with slight modifications to enhance its biological activity against white spot syndrome virus (WSSV) in shrimp culture. The

goals of these assays were to (1) determine the protective effect of Argovit-4 against WSSV, (2) determine whether Argovit-4 supplemented in feed exhibits toxicity towards shrimp, (3) determine whether Argovit-4 as antiviral additive in feed can prevent or delay/reduce WSSV-induced shrimp mortality, and (4) determine whether Argovit-4 supplemented in feed alters the early stages of the shrimp immune response. In bioassay 1, several viral inocula calibrated at SID_{50} (shrimp infectious doses 50% endpoint) were exposed to 40, 100, 200 and 1000 ng/ SID_{50} of Ag⁺ and then intramuscularly injected into shrimp for 96 h. In bioassay 2, shrimp were fed Argovit-4 supplemented in feed at different concentrations (10, 100 and 1000 µg per gram of feed) for 192 h. In bioassay 3, shrimp were treated with Argovit-4 supplemented in feed at different concentrations and then challenged against WSSV for 192 h. In bioassay 4, quantitative real-time RT-qPCR was performed to measure the transcriptional responses of five immune-relevant genes in haemocytes of experimental shrimp treated with Argovit-4 supplemented in feed at 0, 6, 12, 24 and 48 h. The intramuscularly injected Argovit-4 showed a dose-dependent effect ($p < 0.05$) on the cumulative shrimp mortality from 0-96 h post-infection. In the second bioassay, shrimp fed Argovit-4 supplemented in feed did not show signs of toxicity for the assayed doses over the 192-h experiment. The third and fourth bioassays showed that shrimp challenged with WSSV at 1000 µg/g feed exhibited reduced mortality without altering the expression of some immune system-related genes according to the observed level of transcriptional. This study is the first show that the new Argovit-4 formulation has potential as an antiviral additive in feed against WSSV and demonstrates a practical therapeutic strategy to control WSSV and possibly other invertebrate pathogens in shrimp aquaculture.

Keywords: Feed additive; AgNPs; Shrimp; WSSV; Gene expression

Introduction

The white spot syndrome virus (WSSV) is highly aggressive towards shrimp aquaculture worldwide, for which several studies have attempted to develop a treatment ([Álvarez-Sánchez et al., 2018](#)). This pathogen can cause 100% mortality in shrimp culture in just a few days, causing large economic losses, business closures and high unemployment rates ([Sánchez-Martínez, Aguirre-Guzmán & Mejía-Ruiz, 2007](#); [Bustillo, Escobedo & Sotelo, 2009](#)). Currently, nanotechnology has attracted a great deal of attention as an emerging field of research and

technological development that open the possibility of handling material at the nanometric level, with dimensions between 1 and 100 nm (Meneses-Marquez *et al.*, 2018). In this size range, nanoparticles show new physical and chemical properties that can be used in different scientific areas for the unprecedented study of phenomena that occur at the atomic and molecular level (Parveen, Misra & Sahoo. 2012; Esmaeillou *et al.*, 2017). Among the most interesting and promising nanomaterials are silver nanoparticles (AgNPs), which have shown broad-spectrum antiviral activity (Bello-Bello *et al.*, 2018; Galdiero *et al.*, 2011; Chris, Singh & Agarwal. 2018). Several *in vitro* studies have demonstrated the effect of AgNPs as an antiviral alternative against human viruses, including human immunodeficiency virus (Elechiguerra *et al.*, 2005; Lara *et al.*, 2010), H1N1 influenza A virus (Mori *et al.*, 2013), monkeypox virus (Rogers *et al.*, 2008), Tacaribe virus (Speshock *et al.*, 2010), hepatitis B virus (Leu *et al.*, 2008) and herpes simplex virus (Baram-Pinto *et al.*, 2009).

The mechanism of action of AgNPs as an antiviral agent has been studied against several enveloped viruses (Lara *et al.*, 2011; Bogdanchikova *et al.*, 2016), and it has suggested that AgNPs undergo preferential binding with viral envelope glycoproteins to inhibit viruses from binding to host cells (Elichiguerra *et al.*, 2005; Lara *et al.*, 2010). In addition, multiple studies have shown that nanoparticles can be recognized by the immune system and modulate the induction of immunostimulatory effects (Dacoba *et al.*, 2017; Boraschi *et al.*, 2017).

Recent advances have been made in aquaculture using AgNP-based formulations. In one study, Argovit® that was administered intramuscularly in shrimp *Litopenaeus vannamei* showed antiviral activity against WSSV (Juarez-Moreno *et al.*, 2017), a highly lethal and contagious pathogen. This result indicated that a single dose of silver nanoparticles was able to increase shrimp survival by up to 80% without toxic effects. In another study, Ochoa-Meza *et al.* (2019) evaluated the ability of silver nanoparticles coated with commercial non-toxic PVP (Argovit®) to promote the immune system response of shrimp infected with WSSV and observed a late immunostimulatory activity of two immune-related genes. In the same report, authors showed that a single dose of AgNP was able to increase the survival of the shrimp against severe infections of WSSV, even under adverse conditions in the presence of a high concentration of ferrous ions.

Crustaceans have an innate immune system, similar to other invertebrates, however, they lack a developed adaptive immune system. The innate immune system of shrimp consists of several

natural defence mechanisms: a microbial recognition system, a prophenoloxidase system (proPO system), a coagulation system, phagocytosis, encapsulation and the formation of nodules and oxygen reactive compounds (Kurtz, 2005; Ji, Xiao & Wang, 2009; Hirono, *et al.*, 2011). Several genes that participate in the shrimp immune system have been described in the species *L. vannamei*, some of which have been described in previous reports from our group (Flegel & Sritunyalucksana, 2010). To identify a potential mechanism of the antiviral and immunostimulatory activities of silver nanoparticles in shrimp, we selected five genes to study that are representative of different immunological routes. Two of the genes encode proteins (Rab6 and PAP) that participate in endocytosis and phagocytosis (Ye, Tang & Zhang 2012; Khimmakthong *et al.*, 2013). The genes encoding Crustin and PEN4 have been identified as participating in antiviral immunity against WSSV infections in *L. vannamei* (Sun *et al.*, 2017; Xiao *et al.*, 2018). Finally, the proPO gene activates the proPO system, which is involved in melanin formation in arthropods and other invertebrates (Sritunyalucksana & Soderhall, 2000) and results in the production of the melanin pigment that can often be seen as dark spots in the cuticle of arthropods and two antimicrobial peptides from defence system also in WSSV infections (Abad-Rosales *et al.*, 2019). These advances have demonstrated that AgNPs are effective against a wide spectrum of viruses, especially against the disease-causing WSSV, for which there have been no effective treatments until now (Verbruggen *et al.*, 2016). However, intramuscular administration of AgNPs is impractical, considering the large volumes of organisms to that would need to be treated. A practical alternative is the inclusion of AgNPs in balanced shrimp feed (Dananjaya *et al.*, 2016), guaranteeing a greater usefulness of AgNPs therapy.

Although the Argovit® formulation is no longer commercially available, new formulations have been developed with unique features (N. Bogdanchikova, Personal communication, December 2017). For instance, the Argovit-4 formulation retains essentially same the silver nanoparticle design of Argovit®. Thus, in this study, we investigated whether the AgNPs-based Argovit-4 could be used as an antiviral additive in feed to interfere with WSSV infection in shrimp.

Materials and Methods

Argovit-4 formulation

The silver nanoparticles Argovit-4, which replaced Argovit® (Novosibirsk, Russia; patent 2427380) was kindly donated by Dr. Nina Bogdanchikova from CNyN-UNAM, and this can be

requested by contacting Dr. Bodganchikova, (or it can be purchased at Bionag SAPI de CV, Tijuana, México. MC Martha Alvarado, (52) 664-317-99-21). The supplier's specifications indicate that Argovit-4 consists of spherical AgNPs with an average size of 35 ± 15 nm that are functionalized with non-toxic PVP (K30) in water. Table 1, which was modified from [Juarez-Moreno et al. \(2017\)](#), summarizes the physicochemical characterization of these formulations.

White spot syndrome viral stock

The WSSV strain (Guasave isolate, 2005) used in this study was kindly provided by the Biochemistry and Molecular Biology Laboratory from CIIDIR, Sinaloa, Mexico. The viral inoculum was prepared according to the methodology described by [Alvarez-Ruiz et al. \(2013\)](#). Muscular tissues from moribund shrimp were collected, weighed and homogenized in 10 volumes (W/V) of phosphate-buffered saline ($1 \times$ PBS; 137 mM NaCl, 2.7 mM KCl, 10 mM Na_2HPO_4 , and 2 mM KH_2PO_4) to pH 7.4. The homogenized tissue was centrifuged at $13,000 \times g$ (or g-force) for 20 min at 4°C , after which the supernatant was transferred to another microtube (1.6 mL) and centrifuged again at $13,000 \times g$ for 20 min at 4°C . The supernatant was passed through a $0.45 \mu\text{m}$ pore-sized filter and stored at -70°C .

Preparation of inoculum from WSSV-infected tissue

The experimental challenge with WSSV was conducted by the oral route using infected tissue. The infected tissue was prepared by intramuscularly injecting a batch of shrimp ($n=10$) of 10 ± 2 g with of 32 SID_{50} in $100 \mu\text{L}$ of PBS (assuming that 100% of organisms would be infected at 54 h post-infection). The shrimp with observable clinical signs of WSSV infection were individually separated and stored at -70°C until all of the shrimp became infected. Subsequently, after removing the cephalothorax, pleopods and exoskeleton the muscle tissue of the dead WSSV-infected shrimp was weighed and homogenized in 120 mL of marine water for every 24 g of tissue. The homogenized material was used as the inoculum used for the bioassay 4.

Shrimps

Batches of *L. vannamei* (4 ± 0.5 g) juveniles were obtained from a local farm in Guasave, Sinaloa, Mexico. The shrimps were transported to CIIDIR-IPN and placed in plastic tanks (1000-L capacity) with 500 L of filtered seawater at 30 PSU (Practical Salinity Units) with constant

aeration. The shrimps were fed *ad libitum* twice daily with Camaronina® commercial feed (Purina-Cargill LTD, Minnesota, USA) with 35% protein. Subsequently, the shrimps were verified to be free of WSSV, IHHNV and AHPND by endpoint PCR.

Experimental conditions

All experiments were performed in glass tanks (40-L capacity) containing 25 L of seawater (30 PSU) at $27 \pm 1^\circ\text{C}$ and continuous aeration with the shrimps acclimatized under these conditions for 24 h before starting the experiments. The experimental shrimp for bioassay 1 and 2 were fed Camaronina® at 4% of their body weight, while in bioassays 3, 4 and 5, the shrimp were fed feed supplemented with AgNPs. The uneaten feed and faecal material were removed by siphoning every 2 days with water exchange (50% volume) during all experimental assays.

In vivo viral titration and determination of minimal infectious dose by the intramuscular route

A batch of shrimp was used to titre the fresh inoculum using the intramuscular injection method described previously by [Alvarez-Ruiz et al. \(2013\)](#) and [Apun et al. \(2017\)](#). The inoculum was titrated by *in vivo* injecting 100 μL of serial dilutions from the original extract (10^{-3} , 10^{-4} , 10^{-5} and 10^{-6}) between the 3rd and 4th abdominal segments of shrimp ($n=5$ shrimp per dilution). The experimental assay was finished at 96 h post-infection (hpi). The viral titre was calculated using the method of [Reed & Muench \(1938\)](#), for the concentration of a test substance that produces an effect of interest in half of the test units (50% tissue culture infectious dose of a virus), which was described as $\text{SID}_{50}/\text{mL}$.

Preparation of Argovit-4 supplemented feed

Feed supplemented with AgNPs at different concentrations was made according to the modified method of [Gutiérrez-Dagnino et al. \(2015\)](#). Camaronina® 35% protein pellets (commercial feed for shrimp) were pulverized in a coffee grinder. Three homogeneous pastes supplemented with Argovit-4 AgNPs (10, 100 and 1000 $\mu\text{g/g}$ feed), sodium alginate (20 $\mu\text{g/g}$ feed; Sigma-Aldrich®, St. Louis, MO, USA), and distilled water (0.5 mL/g feed) were generated. In the control feed, AgNPs were replaced by distilled water. Subsequently, the pastes were manually pelletized with

a 50-mL disposable syringe (Millipore®, Merck, Darmstadt, Germany) and dried for 72 h at 4°C in darkness.

BIOASSAYS

Bioassay 1: Determination of the minimum WSSV infectious dose by intramuscular route

Bioassay 1 was designed to determine the effective dose of WSSV that is capable of infecting 100% of the shrimp (as determined by conspicuous clinical signs, such as a lack of appetite, erratic swimming and conspicuous atypical coloration). Three groups of shrimp (30 organisms per group, n=10 shrimp in three replicates) were intramuscularly inoculated between the 3rd and 4th abdominal segment with different WSSV infectious doses (7, 10 and 100 SID₅₀ in 100 µL of PBS). The control group was a mock inoculation with 100 µL of PBS. The mortality of the shrimp in every group was recorded and the bioassay finished was when all of the infected shrimp died.

Bioassay 2: Inhibition of pathogenicity by the application of a WSSV-Argovit-4 mixture

Several viral inoculums were calibrated at 7 SID₅₀ in 100 µL volumes and exposed at 40, 100, 200 and 1000 ng of Argovit-4. The viral inoculums and Argovit-4 (WSSV + AgNPs) mixtures were incubated for one hour on ice and then intramuscularly injected into 30 shrimp for every group (n=10 organisms per replicate) for each treatment. The positive control group was inoculated with WSSV (without AgNPs), while the control group was mock-treated with 100 µL of PBS. The experiment lasted up to 96 hpi, and every six hours the presence of dead organisms, clinical signs and the behaviour of the surviving shrimp were recorded.

Bioassay 3: Toxic effect of Argovit-4 supplemented in feed

The toxic effect of Argovit-4 supplemented in feed at different concentrations was tested first. The experiment consisted of four treatments with three replicates each (n=10 shrimp per replicate) that were organized as follows: (1) shrimps receiving feed without AgNPs (control group), (2) shrimps receiving AgNPs at 10 µg/g feed, (3) shrimps receiving AgNPs at 100 µg/g feed, (4) shrimps receiving AgNPs at 1000 µg/g feed. The shrimps were fed twice daily at 4% of

their body weight. The signs of toxicity (lethargic movement, empty gut or atypical colour change) and cumulative mortality were recorded daily. Previously a toxicity bioassay was carried out by intramuscular route in the shrimp for each of 5 different designs of Argovit, which did not occurred mortality and Argovit-4 was selected (data not shown).

Bioassay 4: Argovit-4 supplemented in feed against WSSV

For bioassay 4, the experimental design was performed in triplicate (n=10 shrimps by replicate). The shrimp were fed Argovit-4-supplemented feed at 0 and 12 h for the first and second feedings, respectively. In both cases, the feed weight corresponded to 4% of the shrimp body weight (4 g/organism). Subsequently, at 24 h, the shrimps were fed the WSSV inoculum (200 mg infected tissue per shrimp). Every shrimp received two daily rations of food corresponding to 4% of their body weight, and at 12 h post-treatment with AgNP in feed were challenged with 200 mg of infected tissue per shrimp for each replicate. The cumulative mortality was recorded twice a day until 192 h.

Bioassay 5: Immunostimulatory activity of Argovit-4 towards immune-related genes

To determine the immunostimulatory activity of Argovit-4 on shrimp *L. vannamei* juveniles, a bioassay was established in two glass tanks (40 L) with 25 L of seawater and continuous aeration. Two groups (n= 12 shrimp per tank) were treated with either feed supplemented with Argovit-4 (1000 µg/g) or control feed (feed without Argovit-4) at 4% of their body weight twice daily.

Haemolymph sampling

Shrimp haemolymph samples from each tank were collected in pools of three to obtain ~400 µL/shrimp every 0, 6, 12 and 48 h post-treatment. A 1 mL syringe (27 G×13 mm needle) was used for rinsing with 5% potassium oxalate in isotonic saline (850 mOsm/kg) to obtain haemolymph from the ventral sinus (base of the first abdominal segment ([Apun-Molina et al., 2017](#); [Guemez-Sorhouet et al., 2019](#))). The pools of haemolymph samples were centrifuged at 800 x g for 10 min at 4°C, after which the plasma was removed and the pellet was washed with 250 µL of cold PBS centrifugation as described above. The supernatant was removed, and the

haemocytes were suspended in microtubes containing 200 μ L of precooled TRIzol Reagent (TRI Reagent®, Molecular Research Center, Ohio, USA). The haemocyte samples were stored at -70°C until use.

Isolation of RNA and cDNA synthesis

Total RNA from the haemocyte and intestinal samples was extracted with TRIzol reagent according to the manufacturer's protocol. The RNA was quantified at 260 nm using a NanoDrop instrument (NanoDrop Technologies, Wilmington, DE) and then treated with DNase 1 (1 U/ μ L, Sigma-Aldrich®, St. Louis, MO, USA) to remove residual genomic DNA. cDNA synthesis was performed using the Improm II® Reverse Transcription System (Promega, Wisconsin, USA) with 500 ng of total RNA and random primers from the kit. The obtained cDNA was suspended in 80 μ L of ultrapure water and stored at -70°C until analysis.

Quantitative real-time PCR analysis

The transcriptional responses of genes encoding immune-related proteins (PEN4, Crustin, proPO, PAP and Rab6) from *L. vannamei* haemocytes were determined by quantitative real-time PCR (RT-qPCR) using a CFX96 system and CFX Manager version 3.0 (Bio-Rad Laboratories, Hercules, CA, USA). The expression of target genes was normalized to that of the β -actin gene as an endogenous control. Table 2 shows the sequences of specific primers used in this study. All amplifications were performed (in duplicate) in 15 μ L volume reactions and contained 7.5 μ L of PCR master-mix 2 \times (3.0 μ L of 5 \times Colorless GoTaq Flexi Buffer [Promega, Wisconsin, USA], 1.5 μ L of 25 mM MgCl₂ [Promega], 0.3 μ L of 10 mM dNTPs [Bioline, Taunton, MA, USA], 0.75 μ L of 20 \times EvaGreen® [Biotium, Fremont, CA, USA], 0.1 μ L of 5 U/ μ L GoTaq® Flexi DNA Polymerase [Promega, Wisconsin, USA], and 1.85 μ L of ultrapure water). Each master-mix aliquot was mixed with 0.35 μ L of each primer (10 μ M, Sigma-Aldrich®), 1.8 μ L of ultrapure water and 5 μ L of cDNA as template. The thermocycling program was as follows: an initial cycle of 95°C for 3 min, followed by 40 cycles of 95°C for 10 s, 58°C for 20 s, 72°C for 30 s and 79°C for 5 s (to acquire the fluorescence signal). After each reaction, a dissociation curve from 65 to 90°C was recorded at 0.5°C increments and examined for unique and specific products. The Pfaffl method was used to analyse the relative quantification data from the real-time PCR experiments (Pfaffl, 2003).

Statistical analysis

The percentage data of cumulative mortality (**Bioassays 2, 3 and 4**) were initially arcsine-transformed to perform the statistical analysis. Normality and homogeneity of variances assumptions were analysed by Kolmogorov-Smirnov and Levene's tests. Thereafter, Kruskal-Wallis ANOVA test (non-parametric ANOVA) followed by a multiple comparison test was used to determine significant differences between the experimental groups ($p < 0.05$). To determine the effect of Argovit-4 on the transcriptional response of five immune-relevant genes (**Bioassay 5**), the data were Log10-transformed to confirm its normality before statistical analysis. The data were analysed by Student's t-test to determine significant differences ($p < 0.05$). All bioassays were performed in triplicate and the results of cumulative mortality were expressed as mean \pm standard deviation. While, the immune-related gene expression results are reported as the mean values \pm standard error. Statistical analyses were performed using STATISTICA version 8.0 (StatSoft, Tulsa, OK).

Results

A 7 SID_{50} inoculum is the minimum effective dose to generate 100% shrimp mortality

In the challenge bioassay, the three assayed doses of WSSV resulted in 100% shrimp mortality at different time intervals. The shrimp group infected with 100 SID_{50} showed mortality after 30 hpi with values above 3%, reaching 100% of mortality at 54 hpi. In contrast, the shrimp group infected with 10 SID_{50} exhibited mortality after 36 hpi with values above 17%, reaching 100% at 60 hpi. Finally, in the shrimp group infected with 7 SID_{50} , mortality began after 36 hpi with values above 13%, reaching 100% at 72 hpi. The control group (PBS) did not show mortality during the bioassay ([Fig. 1](#)). The results revealed that 7 SID_{50} as the minimal dose capable causing 100% of shrimp to be infected with conspicuous clinical signs (lack of appetite, erratic swimming and atypical coloration).

Argovit-4/WSSV decreased the mortality of shrimp

The shrimp group treated with 1000 ng/SID₅₀ showed the lowest mortality (50%) at 60 hpi, whereas the shrimp groups treated with 40, 100 and 200 ng/SID₅₀ of Argovit-4 showed mortalities of 90, 83 and 70% at 60 hpi, respectively. The positive control showed 100% mortality at 72 hpi compared with 0% in the control group. Statistical analysis revealed that the susceptibility of the shrimp to WSSV mixed with Argovit-4 significantly decreased (Kruskal-Wallis test; $p=0.0169$) in a dose-dependent manner across the groups (Fig. 2).

Argovit-4 supplemented in feed is non-toxic

None of the assayed Argovit-4 doses supplemented in feed caused significant shrimp mortality (Kruskal-Wallis test; $p=0.0000$) and not affect the health or survival of shrimp, given that they did not show atypical behaviour (evaluated by observations for empty guts, lethargy, and erratic movement) by toxicity effects during the course of the bioassay at 192 h post-treatment.

Argovit-4 supplemented in feed has an antiviral effect against WSSV

The shrimp group exposed to infected WSSV tissue (without Argovit-4) showed a mortality of 43% at 84 hpi. The cumulative mortality in shrimp fed the 1000 µg/g doses was 3% after infection with WSSV at 192 h, followed by 10 and 16% for the 100 and 10 µg/g doses, respectively. In contrast, the uninfected control group did not show mortality. Interestingly, statistical analysis (Kruskal-Wallis test; $p=0.0001$) did not reveal a dose-dependent effect (Fig. 3). However, the results revealed an obvious antiviral effect with both doses assayed.

Argovit-4 does not induce an early immunostimulatory effect for immune-related genes

The expression analysis of genes encoding Crustin, PEN4, PAP and Rab6 showed no significant ($p<0.05$) changes in the haemocytes of shrimp treated with Argovit-4 supplemented in feed with respect to the control (Fig. 4A, 4B; 5A, 5B). Interestingly, only the gene encoding proPO was downregulated at 48 h (t-student test; $p=0.024$), which was possibly an indirect effect of low levels of AgNPs in the haemolymph (Fig. 6).

Discussion

Several studies have evaluated the effectiveness of AgNPs in aquatic diseases control because to

their antimicrobial potential against bacteria, virus and parasites (Barakat, El-Sayed & Gohar, 2016; Morales-Covarrubias *et al.*, 2016; Acedo-Valdez *et al.*, 2017; Pimentel-Acosta *et al.*, 2019). However, this is the first work that examine the efficacy of Argovit-4 (AgNPs-based formulation) as a potential antiviral additive in feed to protect the health of shrimp *L. vannamei* against WSSV.

In order to evaluate the protective effect of AgNPs on shrimp against WSSV, firstly, we determinate that 7 SID₅₀ was the minimum effective dose to generate 100% shrimp mortality according to WSSV inoculum used (**bioassay 1**) and then, we preincubated infectious WSSV inoculums (calibrated to 7 SID₅₀) with Argovit-4 at different concentrations, which were subsequently intramuscularly injected in shrimp (**bioassay 2**). Our results indicated that WSSV infectivity is drastically affected due to a delay in the onset of clinical signs and mortality in a manner dose-dependent ($p < 0.05$) (Figure 2), compared with that observed in the control (WSSV without Argovit-4). Some authors have observed that silver nanoparticles interaction with envelope glycoproteins of virus can inhibit post-entry stages during replication cycle (Rogers *et al.*, 2008; Lara *et al.*, 2010; Borrego *et al.*, 2016). Elechiguerra *et al.* (2005) suggested that AgNPs undergo specific interaction with HIV-1 due to their affinity to disulfide bonded regions (thiol groups provided in viral envelope protein domains). While, Lara *et al.* (2010) postulated that silver nanoparticles not only bind to the viral protein, but can also structurally modify it through the denaturing of disulfide-bound domain. Hence, we hypothesize that AgNPs could interact with WSSV envelope proteins (e.g. VP28; Tang *et al.*, 2007) and block the specific binding to membrane proteins of shrimp host cells. This can be seen in our results of the ability of AgNPs to decrease residual infectivity of viral particles after 60 minutes of incubation, as described by Lara *et al.* (2010). Nevertheless, further research is needed to elucidate the antiviral mechanism of silver nanoparticles against WSSV.

Recently, Juarez-Moreno *et al.* (2017) reported that prophylactic doses 50 ng/g Argovit® (AgNPs-based formulation) administered by intramuscular route in infected shrimp with WSSV evinced a rate survival of 80% without toxic effects after 96 h post infection. While, Ochoa-Meza *et al.* (2019) showed evidence that a single therapeutic dose 1.2 µg/g AgNPs can delay the onset of clinical signs and enhance the response of shrimp immune system without toxic effects in healthy shrimps. However, this results are not comparable with Juarez-Moreno *et al.* (2017) and Ochoa-Meza *et al.* (2019) in effectiveness, due to the different: (1) AgNP preparations used,

(2) weights of the shrimp and, (3) routes of exposure; considering that intramuscular administration is impractical in aquaculture due to the enormous quantity of organisms that are need to be treated (Kumar & Roy, 2017).

The administration by feeding represents the most practical prophylactic or therapeutic method for *in situ* shrimp culture (Alvarez-Sanchez *et al.*, 2017). Therefore, we prepared feed supplemented with Argovit-4 at 10, 100 and 1000 µg/g doses and determined their effects on shrimp health or mortality (**bioassay 3**). Our results showed no obvious toxicity effects on the behavioural activity of shrimp in the form of erratic swimming, changes in feeding rate, atypical body colour body, and increased mortality (1 shrimp death) in all treatments by the oral route. The lack of mortality and clinical signs can be explained by the structural design of Argovit® nanoparticles having been approved by sanitary international organizations for their use in veterinary and medical applications (Vazquez-Muñoz *et al.*, 2014; Bogdanchikova *et al.*, 2016; Borrego *et al.*, 2016).

Despite the potential of AgNPs as antimicrobial agent, in aquaculture industry, concerns remain regarding the potential negative impact to the marine and freshwater environment. However, to date, there have not been clear the *in vivo* toxicity of silver nanoparticles in marine environment, as the effects depend on AgNP properties (particle size, surface area, shape), concentration, AgNP colloidal stability, aggregation grade, sedimentation, and on the amount and species of products yielded from chemical interactions between AgNPs and other variables (Fabrega *et al.*, 2011; Sharma *et al.*, 2019; Pimentel-Acosta *et al.*, 2019),

It is important to clarify that Argovit-4, given its recent design, still does not present toxicity reports with which this AgNP should be better studied in terms of its safety if it is intended to apply to animals for human consumption and in turn study bioaccumulation in the same shrimp and in the ecotoxicological environment. We have presented in this research some results of AgNP exposure considering only the acute toxic response, that is, in the short term. This is because the antiviral activities observed are a consequence of the application of a single dose.

The main reason why a chronic toxicological study was not performed is precisely because of the observed antiviral effect. Fabrega *et al.* (2011), points out that invertebrates such as water flea *Daphnia pulex*, acute and chronic toxicity studies do not generate bioaccumulation, although other authors point out that in the event that AgNPs accumulate in the gills, they do not have harmful side effects (Xiang *et al.*, 2019), however, this could dependent of administered doses

(Monfared *et al.*, 2015). It is important to mention that more experiments should be carried out to consider the possibility of applying this nanoparticular design in the field. For example, bioassays that generate enough information to evaluate the bioaccumulation of flora and fauna accompanying the bodies of water where the shrimp are raised.

To determine the protective properties of Argovit-4 supplemented in feed against WSSV, the results of **bioassay 4** indicate that the 1000 µg/g treatment provided the best protective effect in shrimp against the inoculum of WSSV by the oral route than by injection (Fig. 3). This result was probably due to the high bioavailability of the silver nanoparticles in the digestive tract, which allowed the AgNPs-virus interactions to improve and promote entry by clathrin-mediated receptor routes (Coutiño, Ávila-Lagunes & Arroyo-Helguera, 2017). This is the first study that describes the use of AgNPs as an antiviral additive in feed against WSSV. In recent years, the use of metallic nanoparticles in aquaculture has been increased, for example in fish. For *Clarias gariepinus*, selenium nanoparticles (SeNPs) at a 680 µg/g dose was observed to improve feeds by increasing the proportion of fish food nutrients that pass across the gut tissue (Chris, Sing & Agarwal, 2018). For *Clarias batrachus*, it was also shown that iron nanoparticles (FeNPs) at a 400 µg/g dose is sufficient to ensure the growth and health of fish (Akter *et al.*, 2018).

Previous studies have demonstrated that because metallic nanoparticles are foreign elements to organisms, they can be recognized and internalized by immune system cells through different routes according to their size and shape (Jiang *et al.*, 2008; Coutiño, Ávila-Lagunes & Arroyo-Helguera, 2017; Dacoba *et al.*, 2017; Boraschi *et al.*, 2017). To test this possibility, and to determine the immunostimulatory effect in the early stages by assessing the expression of immune-relevant genes, **bioassay 5** was performed in which shrimp were fed Argovit-4 at a 1000 µg/g dose due to it being the therapeutic dose that best-protected shrimp against WSSV. Five genes encoding immune-related proteins were examined at different time intervals in shrimp haemocytes (PAP, Rab6, Crustin, PEN4, and proPO) by RT-qPCR analysis. PAP (Phagocytosis-Activating Protein) and Rab6 have been shown to be associated with phagocytosis in shrimp immune response (Deachamag *et al.*, 2006; Zhao *et al.*, 2011; Ye, Zong & Zhang, 2012; Ye, Tang & Zhang, 2012; Soto-Alcalá *et al.*, 2019). Crustin and PEN4 are antimicrobial peptides (AMPs) that participate in the humoral defence against pathogens or foreign particles (Wang *et al.*, 2010; Hirono *et al.*, 2011), while proPO is an important humoral effector in innate immunity, catalysing the upstream steps of melanization in shrimp and others invertebrates

(Gollas-Galván, Hernández-Lopez & Vargas-Albores, 1999; Wang *et al.*, 2010; Amparyup, Charoensapsri & Tassanakajon, 2013). The expression of the genes encoding PAP, Rab6, Crustin, and PEN4 in the treated shrimps showed no significant differences in the time intervals proposed with respect to the control (feed without Argovit-4). The expression of the gene encoding phagocytosis activating protein was not activated in samples from the silver nanoparticle-treated shrimp compared to that observed in the control shrimp samples, suggesting that the size of the particles is important, since this parameter facilitates their incorporation and internalization in haemocytes. This observation is reinforced by the same observed behaviour of the Rab6 gene, which participates in the endocytic pathway (Soto-Alcala *et al.*, 2019), since its expression was not altered at any time by the silver nanoparticles. On the other hand, peneaidins are a family of antibacterial proteins that have recently been shown to contribute to the arrest of WSSV-type viral particles. However, the expression of the PEN4 gene was not altered with respect to the time recorded from zero to 48 hours, which indicates that the AgNPs do not interact with the effectors of this gene. The same pattern was observed for the expression of the gene encoding Crustin, which has been shown to be activated in the first hours of WSSV infection (Xiao *et al.*, 2018; Sun, Wang & Zhu, 2017). Taken together, the patterns of the four assayed genes encoding PAP, Rab6, PEN3 and Crustin strongly suggests that the antiviral effect of the nanoparticles occurs by a different pathway than antibacterial peptides or endocytic phagocytosis.

The proPO system is one of the most studied in arthropods, where particles generated by cell destruction and attacks by pathogenic organisms are coated by melanin via the oxidation of phenols by the enzyme phenoloxidase, which are subsequently secreted as melanized corpuscles (Amparyup, Charoensapsri & Tassanakajon, 2013). The proPO gene encodes the phenoloxidase precursor enzyme, with *L. vannamei* encoding LvproPO1 and LvproPO2 (Wang *et al.*, 2010). Interestingly, proPO has been shown to be active during WSSV infection (Abad-Rosales *et al.*, 2019). In this study, proPO gene expression exhibited a significant decrease at 48 h compared with that observed in the control. Several authors have reported that during infection by WSSV, the virions are not covered by melanin due to the participation of at least two viral proteins identified in *P. monodon*. wsv164 interacts directly with PmproPO1 and PmproPO2 (Sangsuriya *et al.*, 2018), probably for an alternative route of infection. Similarly, the WSSV453 protein interacts with the proPO activating enzyme 2 (PmPPAE2), which could explain the observed

decrease in the expression of the proPO gene due to the presence of AgNPs, taking into account that no challenge against WSSV was performed in this experiment (Sutthangkul *et al.*, 2015; Sutthangkul *et al.*, 2017). Taken together, these results suggest that Argovits-4 does not induce the expression in of the assayed early stage immune-related genes, suggesting that the antiviral activity or immunostimulation by AgNP involves a different route to the endocytic pathway. A previous study by our group (Ochoa-Meza *et al.*, 2019) reported that the LGBP and Cu,ZnSOD genes are negatively deregulated by the presence of AgNPs (Argovit®), which suggested that there must be a path for the internalization of the nanoparticles and the activation of the immune system of haemocytes. However, such a path does not require the participation of any of the five genes analysed in this study.

Conclusions

In this study, we explored whether a new AgNP formulation (Argovit-4), which replaced Argovit®, has similar properties to protect and immunostimulant shrimp, allowing them to avoid infection by the white spot virus without causing toxic effects. In addition, we assessed if the expression of some genes immune system-related suggested a common means of activation.

1. Decreased or high concentrations of Argovit-4 nanoparticles do not generate mortality by causing toxicity towards shrimp.
2. The mixture of WSSV-Argovit-4 was capable of decreasing the mortality of shrimp by at least 50% by intramuscular administration.
3. Argovit-4 included in feed could be used to prevent infection by WSSV and can be considered a safe candidate to safeguard aquaculture systems in the future.
4. The zero expression of the PAP, Rab6, PEN4 and Crustin genes, suggest that the antiviral activity of Argovit-4 does not include any of the mechanisms by which such genes perform their function in the immune system of *L. vannamei*.
5. The late down regulation of proPO gene expression, suggests that the antiviral activity of Argovit-4, could to modify a different and unknown route to inhibit a WSSV infection.

Acknowledgements

For technical support: Rene R. Rebollar Prudente from the Biotechnological Marine Organism Laboratory at the Northwest Biological Research Center (CIBNOR), and Carina Gámez Jimenez, from the Biology and Biochemistry Laboratory of Centro de Investigaciones Interdisciplinarias para el Desarrollo Integral Regional del Instituto Politécnico Nacional (CIIDIR-IPN), for all the facilities and technical support during the experiments.

Funders

This study was supported by the National Council of Science and Technology (CONACyT), Mexico, through grant No. 258607 and the Tomsk Polytechnic University Competitiveness Enhancement Program project VIU-RSCBMT-65/2019. CONACYT CRQ-Scholarship: 493589.

Competing interest

The authors declare that there are no competing interests

Author contributions

Romo Quiñonez CR. PhD student. Performed all of the experiments.

Alvarez Sanchez AR. Participated in the experimental design and the discussion of the results.

Alvarez Ruiz P. Participated in the design and execution of the experiments and advised on the analysis of molecular techniques. Co-director of the PhD student.

Chavez Sanchez MC. Leader of the global project and participated in the discussion of results and the draft of the paper.

Bogdanchikova N. (nina@cnyn.unam.mx) Provided the AgNPs and participated in the writing of the paper, providing information regarding the chemical properties of Argovit-4.

Pestryakov A. Designed the new AgNPs (Argovit-4) and provided important data for its management.

Mejia Ruiz CH. Project leader in the management of AgNPs for their application in shrimp.

Participated in the experimental design, analysed results and the final manuscript. Principally responsible for the paper and co-director of the PhD student.

References

- 527 Abad-Rosales SM, Frías-Espericueta MG, Romero-Bernal OG, Lozano-Olvera R, Garcia-Gasca
528 SA, Montoya-Rodriguez L. and Voltolina D. 2019. White spot syndrome virus (WSSV)
529 infection and immunity responses in white shrimp (*Litopenaeus vannamei*) exposed to
530 sublethal levels of metals. *Aq. Res.* **50**:758–764. DOI: [10.1111/are.13932](https://doi.org/10.1111/are.13932)
- 531 Acedo-Valdez MR, Grijalva-Chon JM, Larios-Rodríguez E, Maldonado-Arce AD, Mendoza-
532 Cano F, Sanchez-Paz JA, Castro-Longoria R. 2017. Antibacterial effect of biosynthesized
533 silver nanoparticles in Pacific white shrimp *Litopenaeus vannamei* (Boone) infected with
534 necrotizing hepatopancreatitis bacterium (NHP-B). *Latin American Journal of Aquatic
535 Research* **45**(2):421–430. DOI: [10.3856/vol45-issue2-fulltext-17](https://doi.org/10.3856/vol45-issue2-fulltext-17)
- 536 Akter N, Alam J, Jewel AS, Khatun S, Akter S. 2018. Evaluation of dietary metallic iron
537 nanoparticles as feed additive for growth and physiology of Bagridae catfish *Clarias
538 batrachus* (Linnaeus, 1758). *International Journal of Fisheries and Aquatic Studies*
539 **6**(3):371–377.
- 540 Álvarez-Ruiz P, Mejía-Ruiz CH, Magallón-Barajas FJ, Escobedo-Bonilla CM. 2013. Silencing
541 Pacific white shrimp, *Litopenaeus vannamei* LvRab7 reduces mortality in brooders
542 challenged with white spot syndrome virus. *Aquac. Res.* **44**:772–782. DOI: [10.1111/j.1365-
543 2109.2011.03084.x](https://doi.org/10.1111/j.1365-2109.2011.03084.x)
- 544 Álvarez-Sánchez AR, Romo-Quinonez C, Rosas-Quijano R, Reyes AG, Barraza A, Magallón F,
545 Angulo C, Mejía CH. 2018. Production of specific dsRNA against white spot syndrome
546 virus in the yeast *Yarrowia lipolytica*. *Aquac. Res.* **49**:480–491. DOI: [10.1111/are.13479](https://doi.org/10.1111/are.13479)
- 547 Amparyup P, Charoensapsri W, Tassanakajon A. 2013. Prophenoloxidase system and its role in
548 shrimp immune responses against major pathogens. *Fish & Shellfish Immunology*
549 **34**(4):990–1001. DOI: [10.1016/j.fsi.2012.08.019](https://doi.org/10.1016/j.fsi.2012.08.019)
- 550 Apun-Molina JP, Robles-Romo A, Álvarez-Ruiz P, Santamaria-Miranda A, Arjona O, Racotta
551 IS. 2017. Influence of stocking density and exposure to white spot syndrome virus in
552 biological performance, metabolic, immune, and bioenergetics response of whiteleg shrimp
553 *Litopenaeus vannamei*. *Aquaculture* **47**:528–537. DOI: [10.1016/j.aquaculture.2017.06.027](https://doi.org/10.1016/j.aquaculture.2017.06.027)
- 554 Barakat KM, El-Sayed HS, Gohar YM. 2016. Protective effect of squilla chitosan–silver
555 nanoparticles for *Dicentrarchus labrax* larvae infected with *Vibrio anguillarum*.
556 *International Aquatic Research* **8**(2):179–189. DOI: [10.1007/s40071-016-0133-2](https://doi.org/10.1007/s40071-016-0133-2)

557 Baram-Pinto D, Shukla S, Perkas N, Gedanken A, Sarid R. 2009. Inhibition of herpes simplex
558 virus type 1 infection by silver nanoparticles capped with mercaptoethane sulfonate.
559 *Bioconjugate Chemistry* **20**(8):1497–1502. DOI: [10.1021/bc900215b](https://doi.org/10.1021/bc900215b)

560 Bello-Bello J, Spinoso-Castillo J, Arano-Avalos S, Martínez-Estrada E, Arellano-García M,
561 Pestryakov A, Toledano-Magaña Y, García-Ramos JC, Bogdanchikova N. 2018. Cytotoxic,
562 Genotoxic, and Polymorphism Effects on *Vanilla planifolia* Jacks ex Andrews after Long-
563 Term Exposure to Argovit® Silver Nanoparticles. *Nanomaterials* **8**(10):754. DOI:
564 [10.3390/nano8100754](https://doi.org/10.3390/nano8100754)

565 Boraschi D, Italiani P, Palomba R, Decuzzi P, Duschl A, Fadeel B, Moghimi SM. 2017.
566 Nanoparticles and innate immunity: new perspectives on host defence. *Seminars in*
567 *Immunology* **3**:33–51. DOI: [10.1016/j.smim.2017.08.013](https://doi.org/10.1016/j.smim.2017.08.013)

568 Bustillo M, Escobedo C, Sotelo R. 2009. Revisión de patogénesis y estrategias moleculares
569 contra el virus del síndrome de la mancha blanca en camarones peneidos. *Revista de*
570 *Biología Marina y Oceanografía* 1–11. DOI: [10.4067/S0718-19572009000100001](https://doi.org/10.4067/S0718-19572009000100001)

571 Chris UO, Singh NB, Agarwal A. 2018. Nanoparticles as feed supplement on Growth behaviour
572 of Cultured Catfish (*Clarias gariepinus*) fingerlings. *Materials Today: Proceedings*
573 **5**(3):9076–9081. DOI: [10.1016/j.matpr.2017.10.023](https://doi.org/10.1016/j.matpr.2017.10.023)

574 Coutiño EMR, Lagunes-Ávila L, Arroyo-Helguera O. 2017. Las nanopartículas de plata:
575 mecanismos de entrada, toxicidad y estrés oxidativo. *Revista de Educación Bioquímica*
576 **36**(2):39–54.

577 Dacoba TG, Olivera A, Torres D, Crecente-Campo J, Alonso MJ. 2017. Modulating the immune
578 system through nanotechnology. *Seminars in Immunology* **34**:78–102. DOI:
579 [10.1016/j.smim.2017.09.007](https://doi.org/10.1016/j.smim.2017.09.007)

580 Dananjaya SHS, Godahewa GI, Jayasooriya RGPT, Lee J, De Zoysa M. 2016. Antimicrobial
581 effects of chitosan silver nano composites (CAGNCs) on fish pathogenic *Aliivibrio* (*Vibrio*)
582 *salmonicida*. *Aquaculture* **450**:422–430. DOI: [10.1016/j.aquaculture.2015.08.023](https://doi.org/10.1016/j.aquaculture.2015.08.023)

583 Deachamag P, Intaraphad U, Phongdara A, Chotigeat W. 2006. Expression of a phagocytosis
584 activating protein (PAP) gene in immunized black tiger shrimp. *Aquaculture* **225**:165–175.
585 DOI: [10.1016/j.aquaculture.2006.01.010](https://doi.org/10.1016/j.aquaculture.2006.01.010)

Elechiguerra JL, Burt JL, Morones JR, Camacho-Bragado A, Gao X, Lara HH, Yacaman MJ. 2005. Interaction of silver nanoparticles with HIV-1. *Journal of Nanobiotechnology* **3**:1–10. DOI: [10.1186/1477-3155-3-6](https://doi.org/10.1186/1477-3155-3-6)

Esmacillou M, Zarrini G, Rezaee MA, Mojarad JS. 2017. Vancomycin Capped with Silver Nanoparticles as an Antibacterial Agent against Multi-Drug Resistance Bacteria. *Advanced Pharmaceutical Bulletin* **7**(3):479–483. DOI: [10.15171/apb.2017.058](https://doi.org/10.15171/apb.2017.058)

Fabrega, J., Luoma, S.N., Tyler, C.R., Galloway, T.S., Lead, J.R., 2011. Silver nanoparticles: behaviour and effects in the aquatic environment. *Environ. Int.* **37** (2): 517–531.

Flegel TW & Sritunyalucksana, K. 2010. Shrimp Molecular Responses to Viral Pathogens. *Marine Biotechnology* **13**(4):587–607. DOI: [10.1007/s10126-010-9287-x](https://doi.org/10.1007/s10126-010-9287-x).

Galdiero S, Falanga A, Vitiello M, Cantisani M, Marra V, Galdiero M. 2011. Silver nanoparticles as potential antiviral agents. *Molecules* **16**:8894–8918. DOI: [10.3390/molecules16108894](https://doi.org/10.3390/molecules16108894)

Gollas-Galván T, Hernández-López J, Vargas-Albores F. 1999. Prophenoloxidase from brown shrimp (*Penaeus californiensis*) hemocytes. *Comparative Biochemistry and Physiology Part-B* **122**(1):77–82. DOI: [10.1016/S0305-0491\(98\)10143-8](https://doi.org/10.1016/S0305-0491(98)10143-8)

González-Fernández A, Peleteiro-Olmedo M, Lozano-Fernández T, Simon-Vázquez R, Díaz-Freitas B. 2012. Nanotecnología y Sistema inmunitario. In: Monografías Dr. Antonio Esteve “Nuevas perspectivas en Inmunoterapia”, Ed Manel Juan. Barcelona: Fundación Dr. Antonio Esteve, 22–33.

Guemez-Sorhouet E, Villarreal H, Racotta IS, Naranjo J, Mercier L. 2019. Zootechnical and physiological responses of whiteleg shrimp (*Litopenaeus vannamei*) postlarvae reared in bioflocs and subjected to stress conditions during nursery phase. *Aquaculture Research* **1**–14. DOI: [10.1111/are.13994](https://doi.org/10.1111/are.13994)

Gutiérrez-Dagnino A, Luna-González A, Fierro-Coronado JA, Álvarez-Ruiz P, Flores-Miranda MC, Miranda-Saucedo S, Medina-Beltrán V, Escamilla-Montes R. 2015. Effect of inulin and fulvic acid on survival, growth, immune system, and WSSV prevalence in *Litopenaeus vannamei*. *Latin American Journal of Aquatic Research* **43**(5):912–921. DOI: [10.3856/vol43-issue5-fulltext-11](https://doi.org/10.3856/vol43-issue5-fulltext-11)

- Hirono I, Fagutao FF, Kondo H, Aoki, T. 2011. Uncovering the Mechanisms of Shrimp Innate Immune Response by RNA Interference. *Marine Biotechnology* **13**(4):622–628. DOI: [10.1007/s10126-010-9292-0](https://doi.org/10.1007/s10126-010-9292-0)
- Jiang W, Kim BYS, Rutka JT, Chan WCW. 2008. Nanoparticle-mediated cellular response is size-dependent. *Nature Nanotechnology* **3**(3):145–150. DOI: [10.1038/nnano.2008.30](https://doi.org/10.1038/nnano.2008.30)
- Ji P, Yao C. and Wang Z. 2009. Immune response and gene expression in shrimp (*Litopenaeus vannamei*) hemocytes and hepatopancreas against some pathogen-associated molecular patterns. *Fish & Shellfish Immunology* **27**(4):563–570. DOI: [10.1016/j.fsi.2009.08.001](https://doi.org/10.1016/j.fsi.2009.08.001)
- Kurtz, J. (2005). Specific memory within innate immune systems. *Trends in Immunology* **26**(4). DOI: [10.1016/j.it.2005.02.001](https://doi.org/10.1016/j.it.2005.02.001)
- Lara HH, Ayala-Nunez NV, Ixtapan-Turrent L, Rodriguez-Padilla C. 2010. Mode of antiviral action of silver nanoparticles against HIV-1. *J. Nanobiotech.* **8**:1. DOI: [10.1186/1477-3155-8-1](https://doi.org/10.1186/1477-3155-8-1)
- Lara HH, Garza-Treviño EN, Ixtapan-Turrent L, Singh DK. 2011. Silver nanoparticles are broad-spectrum bactericidal and virucidal compounds. *J. of Nanobiotech.* **30**(9):1–8. DOI: [10.1186/1477-3155-9-30](https://doi.org/10.1186/1477-3155-9-30)
- Lu L, Sun RW, Chen R, Hui CK, Ho CM, Luk JM, Lau GK, Che CM. 2008. Silver nanoparticles inhibit hepatitis B virus replication. *Antiviral Therapy* **13**:253–262.
- Khimmakthong U, Kongmee P, Deachamag P, Leggat U, Chotigeat W. 2013. Activation of an immune response in *Litopenaeus vannamei* by oral immunization with phagocytosis activating protein (PAP) DNA. *Fish & Shellfish Immunology* **34**(3):929–38. DOI: [10.1016/j.fsi.2013.01.004](https://doi.org/10.1016/j.fsi.2013.01.004)
- Kumar V, Roy S. 2017. Aquaculture Drugs: Sources, Active Ingredients, Pharmaceutic Preparations and Methods of Administration. *J Aquac Res Development* **8**:510. DOI: [10.4172/2155-9546.1000510](https://doi.org/10.4172/2155-9546.1000510)
- Martin GG, Quintero M, Quigley M, and Khosrovian H. 2000. Elimination of sequestered material from the gills of decapod crustaceans. *Journal of Crustacean Biology* **20**(2):209–217. DOI: [10.1163/20021975-99990032](https://doi.org/10.1163/20021975-99990032)
- Mejía-Ruiz CH, Vega-Peña S, Álvarez-Ruiz P, Escobedo-Bonilla CM. 2011. Double-stranded RNA against white spot syndrome virus (WSSV) *vp28* or *vp26* reduced susceptibility of *Litopenaeus vannamei* to WSSV, and survivors exhibited decreased susceptibility in

subsequent re-infections. *Journal of Invertebrate Pathology* **107**: 65e68. DOI: [10.1016/j.jip.2011.02.002](https://doi.org/10.1016/j.jip.2011.02.002)

Meneses-Márquez JC, Hamdan-Partida A, Monroy-Dosta MC, Castro-Mejía J, Bustos-Martínez JA. 2018. Silver nanoparticles applications (AgNPS) in aquaculture. *International Journal of Fisheries and Aquatic Studies* **6**:5–11.

Monfared AL, Bahrami AM, Hosseini E, Soltani S, Shaddel M. 2015. Effects of Nano-particles on Histo-pathological changes of the fish. *J Environ Health Sci Eng.* **13**: 62. DOI: [10.1186/s40201-015-0216-9](https://doi.org/10.1186/s40201-015-0216-9)

Morales-Covarrubias MS, García-Aguilar N, Bolan-Mejía MDC, Puello-Cruz AC. 2016. Evaluation of medicinal plants and colloidal silver efficiency against *Vibrio parahaemolyticus* infection in *Litopenaeus vannamei* cultured at low salinity. *Diseases of Aquatic Organisms* **122**(1):57–65. DOI: [10.3354/dao03060](https://doi.org/10.3354/dao03060)

Mori Y, Ono T, Miyahira Y, Nguyen VQ, Matsui T, Ishihara M. 2013. Antiviral activity of silver nanoparticle/chitosan composites against H1N1 influenza A virus. *Nanoscale Research Letters* **8**(1):93. DOI: [10.1186/1556-276X-8-93](https://doi.org/10.1186/1556-276X-8-93)

Parveen S, Misra R, Sahoo SK. 2012. Nanoparticles: a boon to drug delivery, therapeutics, diagnostics and imaging. *Nanomedicine* **8**:147-66. DOI: [10.1016/j.nano.2011.05.016](https://doi.org/10.1016/j.nano.2011.05.016)

Pfaffl MW. 2003. Quantification strategies in real-time PCR. In: A-Z of quantitative PCR. La Jolla, CA, USA: International University Line, 87–112.

Pimentel-Acosta CA, Morales-Serna FN, Chavez-Sanchez MC, Lara HH, Pestryakov A, Bogdanchikova N, Fajer-Avila EJ. 2019. Efficacy of silver nanoparticles against the adults and eggs of monogenean parasites of fish. *Parasitology Research* **118**(6):1741–1749. DOI: [10.1007/s00436-019-06315-9](https://doi.org/10.1007/s00436-019-06315-9)

Reed LJ, Muench H. 1938. A simple method of estimating fifty per cent endpoints. *Am. J. Epidemiol.* **27**(3):493–497. DOI: [10.1093/oxfordjournals.aje.a118408](https://doi.org/10.1093/oxfordjournals.aje.a118408)

Rogers JV, Parkinson CV, Choi YW, Speshock JL, Hussain SM. 2008. A preliminary assessment of silver nanoparticle inhibition of monkeypox virus plaque formation. *Nanoscale Research Letters* **3**(4):129–133. DOI: [10.1007/s11671-008-9128-2](https://doi.org/10.1007/s11671-008-9128-2)

Sanchez-Martínez JG, Aguirre-Guzman G, Mejía-Ruiz H. 2007. White Spot Syndrome Virus in cultured shrimp: A review. *Aquaculture Research* **38**(13):1339–1354. DOI: [10.1111/j.1365-2109.2007.01827.x](https://doi.org/10.1111/j.1365-2109.2007.01827.x)

- Sangsuriya P, Charoensapsri W, Sutthangkul J, Senapin S, Hirono I, Tassanakajon A, and Amparyup P. 2018. A novel white spot syndrome virus protein WSSV164 controls prophenoloxidases, PmpPOs in shrimp melanization cascade. *Dev. Comp. Imm.* **86**:109–117. DOI: [10.1016/j.dci.2018.05.005](https://doi.org/10.1016/j.dci.2018.05.005)
- Sharma VK, Sayes CM, Guo B, Pillai S, Parsons JG, Wang C, Yan B, Ma X. 2019. Interactions between silver nanoparticles and other metal nanoparticles under environmentally relevant conditions: a review. *Science of the Total Environment* **653**:1042–1051. DOI: [10.1016/j.scitotenv.2018.10.411](https://doi.org/10.1016/j.scitotenv.2018.10.411)
- Sritunyalucksana K, Soderhall K. 2000. The proPO and clotting system in crustaceans. *Aquaculture* **191**:53–69. DOI: [10.1016/S0044-8486\(00\)00411-7](https://doi.org/10.1016/S0044-8486(00)00411-7)
- Soto-Alcalá J, Álvarez-Ruiz P, Audelo-Naranjo JM, Esparza-Leal M, Luis-Villaseñor IE, Estrada-Godínez JA, Luna-González A, Gámez-Jiménez C, Diarte-Plata G. 2019. Transcriptional response of immune-related genes in *Litopenaeus vannamei* post-larvae cultured in recirculating aquaculture systems with and without biofloc. *Aq. Intern.* **27**:209–225 DOI: [10.1007/s10499-018-0317-4](https://doi.org/10.1007/s10499-018-0317-4)
- Speshock JL, Murdock RC, Braydich-Stolle LK, Schrand AM, Hussain SM. 2010. Interaction of silver nanoparticles with Tacaribe virus. *J. Nanobiotechnol.* **8**:19. DOI: [10.1186/1477-3155-8-19](https://doi.org/10.1186/1477-3155-8-19)
- Sun B, Wang Z, and F. Zhu. 2017. The crustin-like peptide plays opposite role in shrimp immune response to *Vibrio alginolyticus* and white spot syndrome virus (WSSV) infection. *Fish & Shellfish Immunology* **66**:487–496. DOI: [10.1016/j.fsi.2017.05.055](https://doi.org/10.1016/j.fsi.2017.05.055)
- Sutthangkul J, Amparyup P, Eum JH, Strand MR, Tassanakajon A. 2017. Anti-melanization mechanism of the white spot syndrome viral protein, WSSV453, via interaction with shrimp proPO-activating enzyme, PmpPPAE2. *Journal of General Virology.* **98**(4):69–778. DOI: [10.1099/jgv.0.000729](https://doi.org/10.1099/jgv.0.000729)
- Tang X, Wu J, Sivaraman J & Hew C. 2007. Crystal Structures of Major Envelope Proteins VP26 and VP28 from White Spot Syndrome Virus Shed Light on Their Evolutionary Relationship. *J. Virol.* **81**(12) 6709–6717. DOI: [10.1128/jvi.02505-06](https://doi.org/10.1128/jvi.02505-06)
- Tello-Olea M, Rosales-Mendoza S, Campa-Córdova AI, Palestino G, Luna-González A, Reyes-Becerril M, Velazquez E, Hernandez-Adame L, Angulo C. 2019. Gold nanoparticles (AuNP) exert immunostimulatory and protective effects in shrimp (*Litopenaeus vannamei*)

- 707 against *Vibrio parahaemolyticus*. *Fish & Shellfish Immunology* **84**:756–767. DOI:
708 [10.1016/j.fsi.2018.10.056](https://doi.org/10.1016/j.fsi.2018.10.056)
- 709 Vazquez-Muñoz R, Avalos-Borja M, Castro-Longoria E. 2014. Ultrastructural analysis of
710 candida albicans when exposed to silver nanoparticles. *PLoS ONE* **9**(10):1–10. DOI:
711 [10.1371/journal.pone.0108876](https://doi.org/10.1371/journal.pone.0108876)
- 712 Verbruggen B, Bickley LK, van Aerle R, Bateman KS, Stentiford GD, Santos EM, Tyler CR.
713 2016. Molecular mechanisms of white spot syndrome virus infection and perspectives on
714 treatments. *Viruses* **8**(1):1–29. DOI: [10.3390/v8010023](https://doi.org/10.3390/v8010023)
- 715 Xiang QQ, Wang D, Zhang JL, Ding CZ, Luo X, Tao J Ling J, Shea D and Chen LQ. 2019.
716 Effect of silver nanoparticles on gill membranes of common carp: Modification of fatty acid
717 profile, lipid peroxidation and membrane fluidity. *Environ. Poll.* On-line.
718 [DOI.ORG/10.1016/J.ENVPOL.2019.113504](https://doi.org/10.1016/j.envpol.2019.113504)
- 719 Xiao B, Fu Q, Niu S, Li H, Lü K, Wang S, Yin B, Weng S, Li C, He J. 2018. Penaeidins are a
720 novel family of antiviral effectors against WSSV in shrimp. *BioRxvi. Reprint Serv. Biol.*
721 DOI: [10.1101/467571](https://doi.org/10.1101/467571)
- 722 Ye T, Tang W, Zhang X. 2012. Involvement of Rab6 in the regulation of phagocytosis against
723 virus infection in invertebrates. *Journal of Proteome Research* **11**:4834–4846. DOI:
724 [10.1021/pr300274k](https://doi.org/10.1021/pr300274k)
- 725 Ye T, Zong R, Zhang X. 2012. The role of white spot syndrome virus (WSSV) VP466 protein in
726 shrimp antiviral phagocytosis. *Fish & Shellfish Immunology* **33**:350–358. DOI:
727 [10.1016/j.fsi.2012.05.017](https://doi.org/10.1016/j.fsi.2012.05.017)
- 728 Wang Y, Tseng CW, Lin HY, Chen I Chen YH, Chen YM, Chen TY, Yang HL. 2010. RNAi
729 knock-down of the *Litopenaeus vannamei* Toll gene (LvToll) significantly increases
730 mortality and reduces bacterial clearance after challenge with *Vibrio harveyi*. *Dev. Comp.*
731 *Immunol* **34**(1):49–58. DOI: [10.1016/j.dci.2009.08.003](https://doi.org/10.1016/j.dci.2009.08.003)
- 732 Zhao Z, Jianga C, Zhang X. 2011. Effects of immunostimulants targeting ran GTPase on
733 phagocytosis against virus infection in shrimp. *Fish & Shellfish Immunology* **31**:1013–1018.
734 DOI: [10.1016/j.fsi.2011.08.022](https://doi.org/10.1016/j.fsi.2011.08.022)

Figure 1

WSSV-Inoculum title

Cumulative mortality of shrimp *L. vannamei* injected with three different WSSV doses by intramuscular route. Shrimps infected with 100, 10 and 7 SID. (f) Control negative shrimp treated with PBS. The results are presented as the mean \pm SD.




Figure 2

Effect of mixture Argovit-4/WSSV of different doses injected by intramuscular route

Cumulative mortality of *L. vannamei* juveniles: 40, 100, 200, 1000 ng/(7)SID. (Control –) shrimp treated with PBS and were not infected; (Control +) shrimp treated with PBS and infected with WSSV inoculums. Statistical differences are represented by different letters (p<0.05). Points are presented as the mean \pm SD.




Figure 3

Effect of Argovit-4 supplemented in feed at different doses on cumulative mortality of *L. vannamei* juveniles

Shrimp nourished with Argovit-4 supplemented feed with concentration of silver: 10, 100, 1000 µg/g. (Control +) shrimp fed only with WSSV inoculum. (Control -) shrimp fed with feed lacking Argovit-4 additive. Procedure sequence: at 0 h the feeding with Argovit-4 supplemented feed was executed for the first time; at 12 h feeding with Argovit-4 supplemented feed was performed for the second time and at 24 h feeding with WSSV inoculum (200 mg infected tissue per shrimp) was performed. After that, every 12 h the shrimps were fed with commercial feed. Statistical differences are represented by different letters ($p < 0.05$). The results are presented as the mean \pm SD.




Figure 4

Relative expression of PEN4 and Crustin genes

The graphic show the RT-qPCR of; (A) Relative expression of PEN4 gene with no significant differences between AgNP treatment (black box) and control (white box); (B) Relative expression of Crustin gene, also no significant differences between AgNP treatment (black box) and control sample (white box). The data of the statistical analysis are represented in mean \pm standard error.




Figure 5

Relative expression of Rab6 and PAP genes.

The graphic show the RT-qPCR of; (A) Relative expression of Rab6 gene with no significant differences between AgNP treatment (black box) and control (white box); (B) Relative expression of PAP gene, also no significant differences between AgNP treatment (black box) and control sample (white box). The data of the statistical analysis are represented in mean \pm standard error.




Figure 6

Relative expression of proPO gene

The graphic show the RT-qPCR of the relative expression of proPO gene with no significant differences between AgNP treatment (black box) and control (white box) only at 48 hrs. The data of the statistical analysis are represented in mean \pm standard error (Analysis test in text).




Table 1(on next page)

Physicochemical Characteristics of silver nanoparticles

Physicochemical Characteristics of silver nanoparticles modified to Argovit-4 from Juarez-Moreno et al., 2017.

Properties	Mean
K- value	K30
PVP content	18.8 (% wt.)
Metallic silver content	1.2 (% wt.)
Silver nanoparticle morphology	spheroidal
Average diameter of metallic silver particles by TEM data (nm)	35±15
Size interval of metallic silver particles by TEM data	1 to 90 (nm)
Hydrodynamic diameter: metallic Ag with PVP	70 (nm)
Zeta potential	-15 (mV)
Surface plasmon resonance	420 (nm)
PVP structure by FTIR	Confirmed

1

2

3

4

Table 2(on next page)

Specific primers used

Specific primers used for PCR amplifications of immune system-related genes.

1

2

3

4

5

Genes	Primer sequences (5'-3')	Amplification Size (bp)	Reference
ProPO	Fw: CTGGGCCCCGGGAAGTCAAG Rv: GGTGAGCATGAAGAAGAGCTGGA	125	Soto-Alcalá et al., 2018
PEN 4	Fw: GCCCGTTACCCAAACCATC Rv: CCGTATCTGAAGCAGCAAAGTC	106	Wang et al., 2010
PAP	Fw: CGAAGTTCAGGTTGTGCGTG Rv: ACTGATGCACCATTGGCCTT	126	Soto-Alcalá et al., 2018
Rab 6	Fw: GTTCCGCAGCCTTATTCCCT Rv: ATCACTGCCTCGCTCTGTTC	133	Soto-Alcalá et al., 2018
Crustin	Fw: GAGGGTCAAGCCTACTGCTG Rv: ACTTATCGAGGCCAGCACAC	157	Wang et al., 2010
β-Actin	Fw: CCACGAGACCACCTACAAC Rv: AGCGAGGGCAGTGATTTC	142	Wang et al., 2007