

A taxonomic revision of the south-eastern dragon lizards of the *Smaug warreni* Boulenger species complex in southern Africa, with the description of a new species (Squamata: Cordylidae) (#40394)

1

First submission

Guidance from your Editor

Please submit by **17 Oct 2019** for the benefit of the authors (and your \$200 publishing discount).



Structure and Criteria

Please read the 'Structure and Criteria' page for general guidance.



Custom checks

Make sure you include the custom checks shown below, in your review.



Author notes

Have you read the author notes on the [guidance page](#)?



Raw data check

Review the raw data. Download from the [materials page](#).



Image check

Check that figures and images have not been inappropriately manipulated.

Privacy reminder: If uploading an annotated PDF, remove identifiable information to remain anonymous.

Files

Download and review all files from the [materials page](#).

14 Figure file(s)
5 Table file(s)
1 Raw data file(s)
1 Other file(s)

! Custom checks

Vertebrate animal usage checks

- ! Have you checked the authors [ethical approval statement](#)?
- ! Were the experiments necessary and ethical?
- ! Have you checked our [animal research policies](#)?

New species checks

- ! Have you checked our [new species policies](#)?
- ! Do you agree that it is a new species?
- ! Is it correctly described e.g. meets ICZN standard?



Structure and Criteria

Structure your review

The review form is divided into 5 sections. Please consider these when composing your review:

1. BASIC REPORTING
2. EXPERIMENTAL DESIGN
3. VALIDITY OF THE FINDINGS
4. General comments
5. Confidential notes to the editor

 You can also annotate this PDF and upload it as part of your review

When ready [submit online](#).

Editorial Criteria

Use these criteria points to structure your review. The full detailed editorial criteria is on your [guidance page](#).

BASIC REPORTING

-  Clear, unambiguous, professional English language used throughout.
-  Intro & background to show context. Literature well referenced & relevant.
-  Structure conforms to [PeerJ standards](#), discipline norm, or improved for clarity.
-  Figures are relevant, high quality, well labelled & described.
-  Raw data supplied (see [PeerJ policy](#)).

EXPERIMENTAL DESIGN

-  Original primary research within [Scope of the journal](#).
-  Research question well defined, relevant & meaningful. It is stated how the research fills an identified knowledge gap.
-  Rigorous investigation performed to a high technical & ethical standard.
-  Methods described with sufficient detail & information to replicate.

VALIDITY OF THE FINDINGS

-  Impact and novelty not assessed. Negative/inconclusive results accepted. *Meaningful* replication encouraged where rationale & benefit to literature is clearly stated.
-  All underlying data have been provided; they are robust, statistically sound, & controlled.
-  Speculation is welcome, but should be identified as such.
-  Conclusions are well stated, linked to original research question & limited to supporting results.

Standout reviewing tips

3



The best reviewers use these techniques

Tip

Support criticisms with evidence from the text or from other sources

Example

Smith et al (J of Methodology, 2005, V3, pp 123) have shown that the analysis you use in Lines 241-250 is not the most appropriate for this situation. Please explain why you used this method.

Give specific suggestions on how to improve the manuscript

Your introduction needs more detail. I suggest that you improve the description at lines 57- 86 to provide more justification for your study (specifically, you should expand upon the knowledge gap being filled).

Comment on language and grammar issues

The English language should be improved to ensure that an international audience can clearly understand your text. Some examples where the language could be improved include lines 23, 77, 121, 128 – the current phrasing makes comprehension difficult.

Organize by importance of the issues, and number your points

- 1. Your most important issue*
- 2. The next most important item*
- 3. ...*
- 4. The least important points*

Please provide constructive criticism, and avoid personal opinions

I thank you for providing the raw data, however your supplemental files need more descriptive metadata identifiers to be useful to future readers. Although your results are compelling, the data analysis should be improved in the following ways: AA, BB, CC

Comment on strengths (as well as weaknesses) of the manuscript

I commend the authors for their extensive data set, compiled over many years of detailed fieldwork. In addition, the manuscript is clearly written in professional, unambiguous language. If there is a weakness, it is in the statistical analysis (as I have noted above) which should be improved upon before Acceptance.

A taxonomic revision of the south-eastern dragon lizards of the *Smaug warreni* Boulenger species complex in southern Africa, with the description of a new species (Squamata: Cordylidae)

Michael F. Bates^{Corresp., Equal first author, 1, 2}, Edward L. Stanley^{Equal first author, 3}

¹ Department of Herpetology, National Museum, Bloemfontein, South Africa

² Department of Zoology and Entomology, University of the Free State, Bloemfontein, South Africa

³ Department of Herpetology, Florida Museum of Natural History, Gainesville, Florida, United States

Corresponding Author: Michael F. Bates

Email address: herp@nasmus.co.za

A recent molecular phylogeny of the large dragon lizards of the genus *Smaug* Stanley et al., 2011 recovered a south-eastern clade of relatively lightly-armoured, geographically-proximate species (*S. warreni* [Boulenger] and *S. barbertonensis* [Van Dam]) here referred to as the *S. warreni* species complex. Unexpectedly, *S. barbertonensis* was found to be paraphyletic, with individual sampled from northern Eswatini (formerly Swaziland) being more closely related to *S. warreni* than to *S. barbertonensis* from the type locality of Barberton in Mpumalanga Province, South Africa. Examination of voucher specimens used for the molecular analysis, as well as most other available museum material in this complex, indicated that the 'Eswatini' lineage—including populations in a small area on the northern Eswatini-Mpumalanga border, and northern KwaZulu-Natal Province in South Africa—was readily distinguishable from *S. barbertonensis* sensu stricto (and *S. warreni*) by its unique dorsal, lateral and ventral colour patterns. In order to further assess the taxonomic status of the three populations, a detailed morphological analysis was conducted. Multivariate analyses of scale counts and body dimensions indicated that the 'Eswatini' lineage and *S. warreni* were most similar. In particular, *S. barbertonensis* differed from the other two lineages by its generally lower numbers of transverse rows of dorsal scales, and a relatively wider head. High resolution Computed Tomography also revealed differences in cranial osteology between specimens from the three lineages. The 'Eswatini' lineage is described here as a new species, *Smaugswazicus* sp. nov., representing the eighth known species of dragon lizard. The new species appears to be near-endemic to Eswatini, with about 90% of its range located there. Our study indicates that *S. barbertonensis* sensu stricto is, therefore, a South African endemic restricted to an altitudinal band of about 300 m in the Barberton-Nelspruit-Khandizwe area of eastern

Mpumalanga Province, while *S. warreni* is endemic to the narrow Lebombo Mountain range of South Africa, Eswatini and Mozambique. We present a detailed distribution map for the three species, and a revised diagnostic key to the genus *Smaug*.



A taxonomic revision of the south-eastern dragon lizards of the *Smaug warreni* (Boulenger) species complex in southern Africa, with the description of a new species (Squamata: Cordylidae)

Michael F. Bates^{1,2}, Edward L. Stanley³

¹Department of Herpetology, National Museum, P.O. Box 266, Bloemfontein 9300, Free State Province, South Africa

²Department of Zoology and Entomology, University of the Free State, P.O. Box 339, Bloemfontein 9300, Free State Province, South Africa

³Department of Herpetology, Florida Museum of Natural History, Gainesville, Florida, 32611, USA

Corresponding author:

Michael F. Bates

36 Aliwal Street, Bloemfontein, South Africa

herp@nasmus.co.za

ABSTRACT



A recent molecular phylogeny of the large dragon lizards of the genus *Smaug* Stanley *et al.*, 2011 recovered a south-eastern clade of relatively lightly-armoured, geographically-proximate species (*S. warreni* [Boulenger] and *S. barbertonensis* [Van Dam]) here referred to as the *S. warreni* species complex. Unexpectedly, *S. barbertonensis* was found to be paraphyletic, with individuals sampled from northern Eswatini (formerly Swaziland) being more closely related to *S. warreni* than to *S. barbertonensis* from the type locality of Barberton in Mpumalanga Province, South Africa. Examination of voucher specimens used for the molecular analysis, as well as most other available museum material in this complex, indicated that the ‘Eswatini’ lineage—including populations in a small area on the northern Eswatini-Mpumalanga border, and northern KwaZulu-Natal Province in South Africa—was readily distinguishable from *S. barbertonensis* sensu stricto (and *S. warreni*) by its unique dorsal, lateral and ventral colour patterns. In order to further assess the taxonomic status of the three populations, a detailed morphological analysis was conducted. Multivariate analyses of scale counts and body dimensions indicated that the ‘Eswatini’ lineage and *S. warreni* were most similar. In particular, *S. barbertonensis* differed from the other two lineages by its generally lower numbers of transverse rows of dorsal scales, and a relatively wider head. High resolution Computed Tomography also revealed differences in cranial osteology between specimens from the three lineages. The ‘Eswatini’ lineage is described here as a new species, *Smaug swazicus* **sp. nov.**, representing the eighth known species of dragon lizard. The new species appears to be near-endemic to Eswatini, with about 90% of its range located there. Our study indicates that *S. barbertonensis* sensu stricto is therefore a South African endemic restricted to an altitudinal band of about 300 m in the Barberton–Nelspruit–Khandizwe area of eastern Mpumalanga Province, while *S. warreni* is endemic to the narrow Lebombo Mountain range of South Africa, Eswatini and Mozambique. We present a detailed distribution map for the three species, and a revised diagnostic key to the genus *Smaug*.

INTRODUCTION

Members of the *Smaug warreni* (Boulenger, 1908) species complex (*S. warreni*, *S. barbertonensis* [Van Dam, 1921], *S. depressus* [FitzSimons, 1930], *S. breyeri* [Van Dam, 1921], *S. vandami* [FitzSimons, 1930], *S. mossambicus* [FitzSimons, 1958] and *S. regius* [Broadley, 1962]) are large, robust and spinose girdled lizards (family Cordylidae) restricted to high-elevation regions of the north-eastern provinces of South Africa and Eswatini (formerly Swaziland), and the highlands of eastern Zimbabwe and adjacent Mozambique. Like most girdled lizards, members of the *S. warreni* complex are strictly rupicolous, inhabiting deep, horizontal or gently sloping crevices, often in shaded rocky outcrops (Jacobsen, 1989; Stanley & Bates, 2014). Due to their reliance on deep crevices they appear to be relatively substrate-specific, occurring in partially-vegetated boulder fields on gentle slopes.

The seven currently recognised taxa in the *S. warreni* complex are allopatric, occurring on separate mountain chains, and are distinguishable on the basis of differences in scalation and colour pattern (Jacobsen, 1989; Branch, 1998; Bates et al., 2014; Stanley & Bates, 2014). Despite these clear diagnoses, the group has a tortuous taxonomic history (see Stanley & Bates, 2014). For example, FitzSimons (1943) treated *Cordylus barbertonensis*, *C. b. depressus* and *C. breyeri* as subspecies of *C. warreni*, retained the subspecies *C. vandami perkoensis* (FitzSimons, 1930), and continued to recognise *C. laevigatus* (FitzSimons, 1933) as a valid species. Shortly thereafter, Loveridge (1944) revised the Cordylidae and treated all seven of the above taxa as subspecies of *Cordylus warreni*. FitzSimons (1958) later described *Cordylus warreni mossambicus*, and Broadley (1962) described *C. warreni regius*. *Cordylus warreni* was therefore considered a polytypic species with as many as nine subspecies (Branch, 1988). Jacobsen (1989) subsequently investigated the status of South African populations and on the basis of sympatry between *C. w. vandami* and *C. w. breyeri* at one locality, he recognised *vandami* as a full species. As a result of overlapping morphological character variation (scalation and colour pattern) he considered *C. w. perkoensis* a junior synonym of *C. vandami*, and *C. w. laevigatus* a junior synonym of *C. w. depressus*. Branch (1998) later followed Jacobsen's (1989) arrangement for South African and Eswatini (also spelled 'eSwatini') taxa, but also treated *C. breyeri*, *C. w. mossambicus* and *C. w. regius* as valid species (without providing reasons). Broadley (2006)

treated all taxa in the *C. warreni* complex (except *laevigatus* and *perkoensis*) as full species, but he too failed to provide justification for such action.

In a recent molecular study, Stanley *et al.* (2011) recovered the genus *Cordylus* as paraphyletic and allocated all members of the *C. warreni* complex, together with the large terrestrial species *C. giganteus* Smith, 1874, to a new genus, *Smaug*. A subsequent molecular phylogeny that focused on the *Smaug warreni* complex found that *S. warreni*, *S. barbertonensis*, *S. depressus*, *S. breyeri*, *S. vandami*, *S. mossambicus* and *S. regius* were all valid species (Stanley & Bates, 2014) (Fig. 1). The authors identified a south-eastern clade of three species-level taxa (hereafter referred to as the *S. warreni* species complex), comprising *S. warreni* and two lineages of *S. barbertonensis* from northern Eswatini and Mpumalanga Province, South Africa. The latter taxon was shown to be paraphyletic, with samples from northern Eswatini being more closely related to *S. warreni* than to topotypic *S. barbertonensis*. This led us to hypothesise that diagnosable morphological differences should exist between specimens referable to the three lineages.

FitzSimons (1943: 427) had in fact noted regional differences in colouration in specimens of *Cordylus warreni barbertonensis* as follows: “sides of body and tail with vertical barring of yellow”, “Lower surfaces brown, with irregularly scattered yellowish spots or short transverse bars” (Barberton, South Africa) versus “sides of body and tail with series of large yellow spots and narrow dark interspaces”, “lower surfaces yellowish-white, with irregular dark brown transverse bars on chest and belly, chin spotted with blackish and throat with vermiculate blackish markings” (Eswatini), but he did not suspect that this indicated separate taxonomic status for the two colour forms. Jacobsen (1989) examined 24 specimens of *C. w. barbertonensis* from Mpumalanga and the adjacent northern part of KwaZulu-Natal (formerly part of Transvaal), but did not distinguish different colour patterns.

In the present study it was found that specimens of the two ‘*S. barbertonensis*’ lineages had consistently different dorsal, lateral and ventral colour patterns, as well as other morphological differences. Populations from Eswatini and adjacent areas in Mpumalanga and

KwaZulu-Natal provinces in South Africa, initially referred to as '*Smaug* cf. *barbertonensis*' in this paper, are therefore described here as a new species.

MATERIALS AND METHODS

Study area

The study area comprises the South African provinces of Mpumalanga and (northern) KwaZulu-Natal, as well as Eswatini and adjacent parts of southern Mozambique. This area is bounded by latitudes 25°S and 28°S, and longitudes 30°30'E and 32°30'E.

Material examined

All available specimens in the Ditsong National Museum of Natural History, Pretoria (TM) and National Museum, Bloemfontein (NMB) were examined. Material of *Smaug* collected during *Jacobsen's* (1989) survey of the former Transvaal Province and *Boycott's* (1992) survey of Eswatini was, for the most part, deposited at Ditsong, and this includes the vast majority of museum material identified as *S. warreni* and *S. barbertonensis*. Some non-types of the new species referred to below are housed at American Museum of Natural History, New York (AMNH), Durban Natural Science Museum, Durban (DNSM), and Natural History Museum of Zimbabwe, Bulawayo (NMZB); and a few specimens of *S. warreni* are in the collections of AMNH and NMZB.

When collection co-ordinates and/or altitudes were not available in museum documentation, these were estimated using Google Earth Pro.

In addition to the data presented in this paper, comparative data consulted for the diagnoses of species and for the purposes of preparing a diagnostic key (see below) were obtained from specimens listed in Appendix 1 and *Boulenger* (1908), *Van Dam* (1921), *FitzSimons* (1930, 1933, 1943, 1958), *Loveridge* (1944), *Broadley* (1962, 1966), *De Waal* (1978), *Jacobsen* (1989) and *Mouton et al.* (2018).

Ethics approval

This project was approved by the National Museum Bloemfontein Ethics Clearance Committee (NMB ECC 2019/13).

External morphology

Head measurements (determined using vernier callipers): Length, measured from tip of snout to ear opening on right side; width, at widest point at about the level of the posterior borders of the parietals; depth, on right side from middle of posterior sublabial to highest point of posterior parietal. **Scalation** (examined using a binocular dissecting microscope, mostly a Nikon SMZ 745T): For the most part the morphological characters employed by FitzSimons (1943) were used, and in the same way, unless otherwise indicated. To avoid uncertainty, the following scale counts are described in detail: occipitals: large scales behind the posterior parietals, the outermost ones situated directly behind the elongated upper temporals; gular scales (often elongated and in longitudinal rows): counted transversely between posterior sublabials, the first row extending to the anterior end of the posterior sublabial; dorsal scale rows longitudinally: counted across the widest part of the body more-or-less midway between fore- and hindlimbs (scales of the most lateral rows are at least half the width of adjacent enlarged dorsals); dorsal scale rows transversely: counted from the first complete row behind the occipitals to the row that ends immediately anterior to the vent (when followed around to the ventral side); ventral scale rows longitudinally: counted across the widest part of the body, more-or-less midway between fore- and hindlimbs (lateral ventrals are rectangular or quadrangular, smooth or weakly keeled, flattened, and at least half the size of adjacent ventrals); ventral scale rows transversely: counted from the first row (which curves anteriorly) behind the posterior part of the forelimb insertion to the row (which curves posteriorly) immediately in front of the anterior part of the hindlimb insertion (i.e. scale rows between axilla and groin); lamellae under 4th toe of right foot were counted from the first scale entirely or largely [$> 60\%$] anterior to the junction between 3rd and 4th toes to the scale behind the claw, and incomplete lamellae (i.e. those that do not extend to either side) were excluded. **Sexing:** Males were identified by the presence of large femoral pores (usually with waxy plugs of secreted fluid) as well as differentiated femoral scales (genital glands). Females had minute pin-prick-like femoral pores without waxy plugs, and lacked differentiated femoral scales.

Osteological data

Osteological data was obtained from representative specimens of the *S. warreni* species complex via High Resolution X-ray Computed Tomography (HRCT). Specimens used were: *S. warreni* NMB R9292, AMNH-R-173381; *S. barbertonensis* NMB R9196 (topotype); *S. cf. barbertonensis* NMB R9201 (holotype of new species, see below), AMNH-R-173382. These specimens were scanned using a Phoenix v|tome|x S CT scanner at the American Museum of Natural History's Microscopy and Imaging Facility, and GE Inspection Technologies, LP Technical Solutions Center in San Carlos, California, or on a Phoenix v|tome|x M at the University of Florida's Nanoscale Research Facility. Each specimen was scanned twice: once to recover the full body, and a second higher resolution scan to focus on the cranial morphology. Current, voltage, and detector-time were modified to optimise the grayscale range, and specimens were scanned in sections to maximise resolution (Table S1). Raw data were processed using GE's proprietary datos|x software V.2.3 to produce a series of tomogram images which were then viewed, sectioned, measured and analysed using VG Studio Max 2.2 (Volume Graphics, Heidelberg, Germany). Individual skeletal elements and osteoderms were reconstructed separately for each scan, so as to facilitate osteological analysis. Tomograms and 3D mesh files for all datasets are available online at www.morphosource.org (see supplementary data for DOIs).

Statistical and multivariate analyses

Univariate analysis was conducted using Statistica v. 6. Principal component and linear discriminant analyses were run for three mensural characters (head length, width and height) and 12 meristic characters (highlighted in Table S2), taken from 78 adult museum specimens (>70 mm SVL), using the `prcomp` and `lda` commands in R `{stats}` and `{MASS}`. A second round of principal component analysis was performed using a reduced dataset comprising eight characters with the highest loading for the first two components (the three head measurements, number of gular scales, numbers of transverse and longitudinal rows of dorsal scales, numbers of transverse and longitudinal rows of ventral scales).

Species concept and species delimitation

We apply a lineage-based species concept whereby a species is represented by an independently evolving metapopulation lineage (see *Frost & Hillis, 1990; De Queiroz, 1998, 2007*). Genetic distinctness and morphological characters were the operational criteria for species delimitation.

The electronic version of this article in Portable Document Format (PDF) will represent a published work according to the International Commission on Zoological Nomenclature (ICZN), and hence the new names contained in the electronic version are effectively published under that Code from the electronic edition alone. This published work and the nomenclatural acts it contains have been registered in ZooBank, the online registration system for the ICZN. The ZooBank LSIDs (Life Science Identifiers) can be resolved and the associated information viewed through any standard web browser by appending the LSID to the prefix <http://zoobank.org/>. The LSID for this publication is: [urn:lsid:zoobank.org:pub:490BDD66-155F-423F-A4E9-DEAEEB024CC5]. The online version of this work is archived and available from the following digital repositories: PeerJ, PubMed Central and CLOCKSS.

RESULTS

Character analysis

Dorsal colour pattern. (Fig. 2) Specimens from all three clades recovered by *Stanley & Bates (2014)* are distinguishable on the basis of dorsal, lateral and ventral colour patterns. *Smaug warreni* has a medium to light brown (sometimes reddish-brown) dorsum with a series of 5–6 interrupted transverse bands between fore- and hindlimbs, each consisting of white or cream ocelli (spots or blotches) with dark (often black) borders. The dark-edges exaggerate the ocelli, but some specimens have only small pale markings which also lack heavy dark borders. The dorsum of *S. barbertonensis* is medium to dark brown (or even black), usually with 4–5 interrupted bands on the back formed mostly by transversely enlarged pale markings (rather than spots or blotches) with moderately dark edges. *Smaug cf. barbertonensis* is similar to the latter form, but there are usually 5–6 bands. However, in *S. barbertonensis* there is almost always a pale spot on the nape immediately posterior to the median occipitals, followed in close proximity by a distinct transverse band. In *S. cf. barbertonensis*, the spot on the nape is replaced by a pale band, followed after a distinct gap by another pale band (often divided medially) with a slightly

posteriorly-directed curvature. *Smaug warreni* often also has a pale band behind the occipitals, but it usually lacks the curved band that follows as in the case of the previous form.

Ventral colour pattern. (Fig. 2) In *S. warreni* the belly is generally white with the centre of each scale pale brown; the throat is white with a few small to medium-sized dark brown spots. In *S. barbertonensis* the belly is almost completely black or dark brown, with only a few pale markings on the sides; the throat is also dark, with occasional pale specks. In *S. cf. barbertonensis* the belly is white with 5–6 dark brown cross-bands, interrupted mid-ventrally by six longitudinal rows of brown scales; the throat is white with heavy, dark, reticulations.

Lateral colour pattern. (Figs 6, 9, 12) The flanks of *S. warreni* are often mostly cream with a few dark markings, but may consist of alternating light and dark vertical bands. In *S. barbertonensis* the flanks are primarily dark brown or even black, with a few narrow or moderate cream bands and/or spots/blotches. In contrast, the sides of the body in *S. cf. barbertonensis* consist of large cream spots or blotches on a dark background. In some cases the light patch behind the armpit is elongated (antero-posteriorly).

Scales at the edges of the ear openings. *Smaug barbertonensis* usually has generally elongated and spinose scales at the anterior edges of the ear openings (especially the central ones), whereas in most cases these scales are short and non-spinose in *S. cf. barbertonensis* and *S. warreni*.

Relative length of occipital scales. In all three forms in the complex there are usually six occipital scales, and the scales of the median pair are shorter and usually smaller than the others (although usually less distinctly so in *S. warreni*). In *S. warreni* the outer occipital is usually of similar size and shape to the adjacent inner occipital, but in *S. barbertonensis* and *S. cf. barbertonensis* the outer one is usually shorter and smaller. In *S. warreni* a small median occipital is common.

Quadrate variation. In *S. cf. barbertonensis* the quadrates have a pronounced ridge and concave region at the lateral edge of the adductor musculus mandibulae posterior origin, whereas in *S.*

barbertonensis and *S. warreni* the quadrates have a less pronounced ridge and a non-concave region (Fig. 3).

Scale counts. The three forms are similar in terms of scale counts (Table 1), but *S. barbertonensis* tends to have lower numbers of transverse dorsal scale rows than *S. warreni* and *S. cf. barbertonensis* (28–34 versus 31–41; Fig. 4A).

Head width. *Smaug barbertonensis*, when compared to both *S. warreni* and *S. cf. barbertonensis*, usually has a wider head relative to snout-vent length (SVL) (Fig. 4B).

Spinosity. A recent study by Mouton *et al.* (2018) that investigated the relationship between generation gland morphology and armour in the genus *Smaug* found that those species with multi-layer generation glands (*S. giganteus*, *S. breyeri*, *S. vandami*) had relatively long (basal) tail and occipital spines, while all other species (including *S. warreni*, *S. barbertonensis* and *S. cf. barbertonensis*) had two-layer glands and relatively short spines. The latter two forms were found to be slightly more spinose than *S. warreni* with regard to both tail and occipital spines.

Statistical analyses

Both principal components and linear discriminant analyses reveal clear separation in scale characters between *S. warreni* and *S. barbertonensis*, and *S. cf. barbertonensis* and *S. barbertonensis* (4% LDA mis-classification rate in both cases) (Fig. 4C–D, Table 2). *Smaug cf. barbertonensis* and *S. warreni* display similar pholidosis and head proportions and cannot be consistently sorted by these characters alone (25% LDA mis-classification rate). The first two principal components explain 30% of the variation in the total dataset and 44% of the total variation in the reduced dataset.

Systematics

Family Cordylidae

***Smaug swazicus* Bates & Stanley sp. nov.**

Swazi Dragon Lizard

Figs 5–7, Tables 3–4

urn:lsid:zoobank.org:act:A942675E-5E76-4FC9-AA8F-BFA7A4C131C7

Cordylus warreni barbertonenis (not Van Dam, 1921): FitzSimons, 1943: 426 (part: Hluti-Goedgegun, eSwatini); Branch, 1988: 164 (part) & 1998: 195 (part); Jacobsen, 1989 (part: Godlwayo; Nzulase; Farm Zwartkloof 60 HU); Adolphs, 1996: 15 (part); Bourquin, 2004: 96 (KwaZulu-Natal); Adolphs, 2006: 22 (part).

Smaug warreni barbertonensis (not Van Dam, 1921): Stanley *et al.*, 2011: 64 (part); Bates *et al.* 2014: 211 (part, including fig. on p. 211); Reissig, 2014: 190 (part, including figs 215– 217, 219).

Smaug sp. Stanley & Bates, 2014: 905.



Smaug cf. *barbertonenis* Mouton *et al.*, 2018: 464.

Holotype. NMB R9201, adult male (differentiated glandular femoral scales present) from Maguga Dam, Hhohho Region, Eswatini [26°04'32"S, 31°15'34"E; 2631AB; 562 m a.s.l.], collected by E.L. Stanley & J.M. da Silva, 31 October 2008.

Paratypes. Allotype: TM 78918, adult female (no differentiated femoral scales) from Nkomati Gorge, Malolotja Nature Reserve, Hhohho Region, Eswatini [26°03'15"S, 31°08'06"E; 2631AA; 640 m a.s.l.], collected by R.C. Boycott, 29 August 1993 (*Fig. 2*). Ten more paratypes: TM 83000, adult male, 1 km NW of Maguga Dam, Hhohho Region, Eswatini [26°04'04"S, 31°14'55"E; 2631AA; 618 m a.s.l.], R.C. Boycott, 25 March 1997; TM 83532, adult male, 5 km SE of Bhunya, Eswatini [26°32'16"S, 31°02'54"E; 2631CA; 960 m a.s.l.], R.C. Boycott, 28 June 2000; TM 42531, adult female, Mbutini Hills, 23 km N of Sepofaneni, Eswatini [26°31'34"S, 31°35'45"E;], W.D. Haacke, 3 September 1972; TM 51376, adult male, 15 km NW of Gilgal on route to Manzini, Lubombo district, Eswatini [2631DA], W.D. Haacke, 3 September 1972; TM 78931, juvenile, Nkomati Gorge, Malolotja Nature Reserve, Hhohho Region, Eswatini [26°03'14"S, 31°08'02"E; 2631AA; 669 m a.s.l.], R.C. Boycott, 14 September 1993; TM 78921, juvenile, Nkomati Valley, Hhohho Region, Eswatini [26°03'12"S, 31°14'24"E; 2631AA; 580 m a.s.l.], J. Linden, 31 October 1992; NMB R9194, adult male, Komati View Point, Malolotja Nature Reserve, Hhohho Region, Eswatini [26°04'29"S, 31°07'32"E; 2631AA; 1033 m a.s.l.],

E.L. Stanley & J.M. da Silva, 31 October 2008 (*Fig. 7*); NMB R9195, adult male, Komati View Point, Malolotja Nature Reserve, Hhohho Region, Eswatini [26°04'32"S, 31°08'01"E; 2631AA; 1052 m a.s.l.], E.L. Stanley & J.M. da Silva, 31 October 2008; NMB R9202, adult male from Maguga Dam, Hhohho Region, Eswatini [26°04'32"S, 31°15'35"E; 2631AB; 562 m a.s.l.], collected by E.L. Stanley & J.M. da Silva, 31 October 2008; TM 73290, adult female, Nzulase, Mpumalanga Province, South Africa [2531DC], N.H.G. Jacobsen, 29 March 1983.

Additional records (*material examined). SOUTH AFRICA: KwaZulu-Natal. Godlwayo Hill (27°20'S, 31°25'E; 750 m a.s.l.) TM 73290–1*, 73294*; Ithala Game Reserve (central point for mapping 27°30'S, 31°17'E) TM 51670*; Farm Zwartkloof 60HU (27°24'S, 31°33'E; ?420 m a.s.l.) TM 73285. ESWATINI: between Hluti and Goedgegun [now called Nhlangano] (no coordinates) TM 16827–9*, 16798*; same locality (27°12'20.1"S, 31°20'10.5"E, photographic record: T. Sparkes); 1 km NW of Maguga Dam wall (26°04'04"S, 31°14'55"E; 618 m a.s.l.) DNSM 1707 (identified as *Smaug barbertonensis* by R.C. Boycott), TM 83002*; 1 km SE of Maguga Dam wall (26°05'08"S, 31°16'20"E) DNSM 1710 (identified as *Smaug barbertonensis* by R.C. Boycott); Manzini, 25 km ESE of (26°31'S, 31°37'E) NMZB-UM 2026, 2529 (identified as *Cordylus warreni barbertonensis* by D.G. Broadley); Nwempisi Gorge, 12 km E of Mankayane (26°42'13.4"S, 31°11'49.1"E, sight record: R.C. Boycott). NO DATA: AMNH-R173382 (used for CT scanning).

Diagnosis. (includes 'additional material') Distinguished from all other cordylids by its unique dorsal, lateral and ventral colour patterns (see descriptions and figures). It also differs from the terrestrial *S. giganteus* by its smaller adult size (maximum SVL 145 mm versus 198 mm), and possession of six (occasionally four) moderate sized and weakly spinose occipitals, versus four (occasionally five) large and distinctly spinose occipitals. Differs from other species of *Smaug* as follows: from *S. vandami* by usually having six (versus usually four) occipitals; from *S. depressus* by having only 10–12 (versus 16–24) femoral pores per thigh in males; and from *S. breyeri* by having much less rugose head shields. It differs from *S. giganteus*, *S. breyeri* and *S. vandami* by having less spinose occipitals and tail spines, and two-layer (rather than multi-layer) generation glands.

Most similar to *S. barbertonensis* and *S. warreni*, but easily distinguishable by its colour pattern (back dark brown usually with 5–6 pale bands between fore- and hindlimbs, pale band on nape behind occipitals; flanks with large pale spots or blotches; venter pale with a dark central longitudinal band bordered on either side by dark transverse bands) compared to *S. barbertonensis* (back dark brown with 4–5 pale bands, pale spot or blotch on nape behind occipitals; flanks dark with narrow pale vertical markings; venter mostly dark brown or black) and *S. warreni* (back usually pale brown with 5–6 pale dark-edged bands, pale band on nape behind occipitals; flanks pale with brown markings; venter with brown markings on most scales); by usually having sort, non-spinose scales at the edges of the ear openings; and quadrates with a pronounced ridge and concave region at the lateral edge of the adductor musculus mandibulae posterior origin (no pronounced ridge or concave region in the other two species). Also differs as follows: outer occipitals shorter than the adjacent inner ones (of about equal length in *S. warreni*); head narrower than *S. barbertonensis* (head width/head length = 76–84% versus 80–92%); generally higher numbers of transverse dorsal scale rows (32–37 in 86% of specimens) than *S. barbertonensis* (29–32 in 81% of *S. barbertonensis*).

Description of holotype. NMB R9201. *External morphology:* Snout-vent length 138.8 mm, tail length (original) 187 mm, total length 325.8 mm, head length 40.2 mm, head width 31.2 mm, head depth 16.2 mm. Tail length/SVL = 1.35; head width/SVL = 22.4%; head width/head length = 77.5%; head depth/head length = 40.3%. Head strongly depressed, head shields rugose and moderately striated over parietal region. Frontonasal 1.05 times as wide as long, in contact with the rostral and loreals, separating the nasals, latter slightly swollen. Nostril – with a large inner flap attached posteriorly – situated in the posterior part of the nasal and in contact with the loreal and 1st supralabial. Prefrontals in contact at their inner angles. Frontal hexagonal, slightly widened anteriorly, anterior sides curved slightly inwards. Frontoparietals slightly broader than long. Posterior parietals larger than anterior ones; interparietal between four parietals, more sharply pointed anteriorly than posteriorly. Occipitals scales 6, well-developed, bluntly spinose, the outer ones shorter and smaller than the second ones, middle pair shortest and narrowest. Anterior upper temporals large, keeled at their lower edges. Gulars 23. Lateral temporals large, often bluntly keeled. Scales at anterior edge of ear opening (4 on left side of head, 5 right) projecting outwards as flattened and somewhat spatulate spines, the lowermost one narrow and

slender, the one above it distinctly spatulate and the largest. Supraoculars 4, the anterior one longest, the next (2nd) one broadest, posterior one the smallest. Supraciliaries 4, anterior one the longest. Lower eyelid opaque, consisting of about 10 small, vertically-elongated scales. Preocular at least twice the size of the loreal. Five large scales below the eye. Rostral 2.04 times wider than deep. Supralabials 6, 4th (longest) and 5th separated by a large suborbital shield (much narrower below than above). Mental 1.36 times as wide as long. Infralabials 6, 5th and 6th keeled, bordered below by five large sublabial shields. First pair of sublabials separated by an elongated scale in contact with the large mental and followed behind by two pairs of similar-sized scales, and numerous small elongated scales that increase in size until about the middle of the throat, but then reduce in size posteriorly. Sides of neck with irregular erect spines, the largest about twice as high as wide. Dorsal scales large, rugose, often striated, forming regular (but not always aligned) transverse series; four vertebral rows with smooth scales (probably due to rubbing against rocks), other dorsals keeled, but lateral scales keeled and spinose. Dorsals in 34 transverse series (from first row posterior to occipitals to row above vent) and 20 longitudinal rows. Ventrals smooth, mostly quadrangular, occasionally pentagonal near middle of venter posteriorly, mostly broader than long, two outer rows moderately keeled and weakly spinose, some scales of the 3rd row also weakly keeled, forming 14 longitudinal and 26 transverse series from axil to groin (with an additional seven rows to base of throat). A pair of slightly enlarged hexagonal preanal plates (slightly longer than wide), with smaller plates anteriorly and on the sides. Limbs above with large, keeled, spinose scales. Femoral pores 21 (10 left leg, 11 right). Differentiated femoral scales 46 (21 left, 25 right). Fourth toe on each foot with 16 subdigital lamellae. Tail with whorls of large, strongly keeled, spinose scales; each whorl separated by a smaller whorl of small, moderately keeled and weakly spinose scales; two upper lateral caudal scale rows consist of especially large and very strongly spinose scales (spines project backwards at angles of about 45°); subcaudal scales long, narrow, mostly pentagonal (occasionally rectangular) and moderately keeled.

Colour: Back dark brown to black with cream to yellow markings forming 5 interrupted transverse bands (with slightly dark borders in preservative) between fore-and hindlimbs, which continue along the tail, together with a band immediately behind the occipitals and another on the nape that is divided medially and curved slightly backwards. Belly cream with a brown

longitudinal band medially (six ventral plates wide) and short, broad, widely separated brown crossbands on either side between the limbs (at least four on the left, three on the right) which are often confluent with the darker parts of the back. The joining of these dark ventral and dorsal markings decorates the flanks with large cream-yellow spots/blotches. Top of limbs dark brown with numerous irregular cream to yellowish spots and blotches; underparts of limbs mostly cream with irregular brown markings, occasionally bands. Top of head brown with scattered irregular cream markings; throat mottled in brown and cream.

Osteodermal osteology: The dorsal and lateral sides of the trunk are covered in circular, well-separated osteoderms, dorsomedially unkeeled grading to well keeled and mucronate towards the sides. The nuchal osteoderms are small, becoming highly spined posterior to the tympanic opening. Ventral osteoderms are delicate and plate-like. The forelimbs are covered in keeled non-imbricate circular/ rhomboid osteoderms, while the hindlimbs are well armored, save for the ventral surface of the thigh, which lacks osteoderms. Osteoderms on the posterior part of the hindlimbs are heavily spinose. The caudal osteoderms are large, robust and arranged in imbricated whorls. Caudal osteoderms are feebly keeled and mucronate along the dorsal and ventral aspects, becoming more heavily spined laterally.

Postcranial skeleton: Tail complete, 26 presacral vertebrae, 32 caudal vertebrae. The haemapophyses of the first caudal osteoderms extend laterally to fuse to the posteroventral edge of the parapophysis, forming a biphiid rib. Four cervical, three sternal, two xiphisternal, 6 left and 7 right long asternal ribs with ossified costal cartilage, then 6 left and 5 right short asternal ribs and one very short pair of ribs immediately anterior to the sacral vertebrae. Cervical ribs 2–4 are distally flattened and biphiid, with the ventral processes more elongated. Pubis flattened and curved with a large, ventrally angled pectineal tubercle. Pubic symphysis flattened and triangular, separating the pubes entirely. Hyperischium and Hypoischium well developed. Ilium triangular in cross-section, with a feeble iliac tubercle. Sternal plate broad with no fontanelle. Interclavicle cruciform, clavicles flattened dorsally. Epicoracoid connects the scapular ray to the primary and secondary coracoid rays, but not to the anterior process of the scapular. Phalanges display a typical pattern of 2–3–4–5–3. Metatarsal 5 with elongated medial process at midbody.

443 *Cranial skeleton:* The scales of the dorsal and temporal regions of the skull and the ventrolateral
 444 aspects of the jaws are underlain with rugose osteoderms. These osteoderms fuse to the proximal
 445 parietal, frontal and postorbital bones, although the mesokinetic and metakinetic joints appear
 446 unobstructed and flexible. Lateral maxilla and anterior aspect of the premaxilla lack osteoderms.
 447 The parietal is pentagonal, with five osteoderms that underlie the parietal shields fused to its
 448 dorsal surface, and a bifid medioposterior process that extends either side of the sagittal crest of
 449 the supraoccipital. Three large osteoderms are fused to the frontal, which is unpaired and clasped
 450 by the parietal at its posteriolateral edge. The upper temporal fenestra is obscured anteriorly by a
 451 large osteoderm fused to the dorsal surface of the postorbital bone, posteriorly by an unfused
 452 rectangular osteoderm that overlies the squamosal. Premaxilla is unpaired and contains seven
 453 pleurodont teeth and five foramina, with a dorsal process that extends posteriorly to be clasped
 454 by the nasals, which themselves insert into the frontal. The maxilla is scinciform, with a deeply
 455 grooved crista dentalis, 9 left or 8 right lateral foramina, and 19 teeth. Teeth display pleurodont
 456 attachment and are unicuspid, with a slight concave surface where they connect with the
 457 mandibular teeth. No palpebral is present and the prefrontal connects directly to the anteriormost
 458 superorbital osteoderm. The jugal is triangular in cross-section and asymmetrically T-shaped,
 459 with a tapering anterior process and a broad, truncated posterior process that extends along and
 460 past the posterior edge of the maxilla. Lacrimal bone is small, flattened and oval. Pterygoids are
 461 edentate and extend back to connect with the quadrates, becoming C-shaped in cross-section
 462 posterior to the epipterygoid condyle. The squamosal is curved and blade-like, circular in cross-
 463 section anteriorly, becoming flattened posteriorly, where it articulates with the cephalic condyle
 464 of the quadrate and the braincase. Supratemporals are flattened, ovoid and not fused with the
 465 elongate paroccipital processes. The posterior aspect of the prootic not fully fused with the oto-
 466 occipital, resulting in a deep groove along the dorsal aspect of the para-occipital processes.
 467 Quadrates very broad with a pronounced ridge and concave region at the lateral edge of the
 468 adductor musculus mandibulae posterior origin. The supraoccipital has a strong sagittal crest that
 469 extends posteriorly to contact the ventral surface of the medioposterior process of the parietal.
 470 The prootic bears an extended alar process and a well-developed, rhomboid crista prootica, and
 471 a very weak supratrigeminal process. Basipterygoid processes are well developed and flattened.
 472 The lower jaw possesses a large adductor fossa, a highly flattened and medially extended
 473 retroarticular process, a medially open Meckelian canal that is closed posteriorly by a large

splénial, and a dentary with a strong subdental shelf; 21 mandibular teeth, and nine dentary foramina.

476

477 **Variation in paratypes** (including allotype TM 78918). *External morphology:* Tail length/SVL

478 1.12–1.44 (SVL: 129.4–143.8 mm, $N = 3$); head width/SVL = 21.6–23.9% in males (SVL:

479 129.4–145.0 mm, $N = 5$), 20.3–23.1% in females (SVL: 102.8–143.8 mm, $N = 4$); head

480 width/head length = 78.3–82.7% (SVL: 102.8–145.0 mm, $N = 9$); head depth/head length =

481 41.9–48.7% (SVL: 102.8–145.0 mm, $N = 9$). In TM 78918, shields on anterior part of head

482 smooth, weakly rugose on posterior part of head but without striations; in two juveniles: head

483 shields smooth (TM 78921) or weakly rugose without striations (TM 78931). Frontonasal

484 0.89–1.12 (0.94–1.05 in juveniles) times as wide as long. Nasal scale fragmented on left side in

485 TM 83000. Small infranasal present on both sides of head in TM 78918. Frontal with anterior

486 sides straight in TM 78921, strongly curved inwards in NMB R9202, separated from rostral by a

487 small rectangular scale in TM 42531. Prefrontals in narrow contact in TM 51376, anterior half of

488 prefrontals in contact in TM 78918 and 83532. Frontoparietals about as wide as long in TM

489 73290, 78918, 78931 and 83532. Interparietal sunken in NMB R9195, about as large as an

490 anterior parietal in two juveniles (TM 78921, 78931), triangular in TM 51376 and 73290, and as

491 pointed posteriorly as it is anteriorly in TM 83000 and 83532. Occipitals 6 but middle pair

492 separated by a small elongate scale in TM 78931, mostly very weakly spinose, all of about the

493 same size in TM 78918, outer scale and the one adjacent to it of similar length in NMB R9202

494 and TM 73290; in TM 83532 scales of the middle pair are shortest but of similar width to the

495 others, and wider than the outer occipitals; but the middle and outer scales may be similar in size

496 (TM 42531, 51376); middle occipitals the same size as second occipitals on either side in TM

497 78931; on left side of TM 83000 the outer occipital is about equal in size to the occipital adjacent

498 to it; in TM 51376 the inner occipitals are rugose only, not spinose. Gulars 22–28 (25 in

499 allotype). Posterior upper temporal scale keeled at its lower edge in TM 42531, 78918 and

500 78921; anterior and posterior upper temporals similar in size and keeled at the sides in TM

501 51376. Lateral temporals rugose only (not keeled) in NMB R9194 and TM 51376. Scales at

502 anterior edge of ear opening 4–6 (3rd from the top is tiny in TM 83532), lowermost spine often

503 not slender and similar to other small spines, but elongate and distinctly spiny in TM 42531. First

504 and 2nd (TM 78921, 78931) and 1st and 3rd (TM 78918) supraoculars about the same length,

2nd and 3rd on left side of NMB R9194 largely fused. Supraciliaries 5 (left side: NMB R9202, TM 42531; right: TM 83532), first and second supraciliaries about equal in length in TM 78931. Lower eyelid transparent in TM 73290 and TM 83532, usually consisting of several irregular scales (e.g. NMB R9195). Preocular about 1.5 times (TM 42531, 51376, 83532) and 1.75 times (TM 73290) larger than loreal. Six large scales below the eye in TM 42531, TM 73290 (left) and TM 78931 (left), and four in TM 83000; large suborbital shield divided in TM 73290. Rostral 2.14–2.81 (1.87–2.10 in juveniles) times wider than deep. Supralabials 7 on left side of head in TM 73290 and TM 78921; sixth (of six) in TM 51376 is granular and 2nd is fragmentary; 4th (of 6) distinctly keeled in TM 42531; 3rd and 4th fused in NMB R9202. Mental 1.17–1.66 (1.16 in juvenile TM 78931) times as wide as long. Fourth and 6th infralabial weakly keeled in TM 78931. Fifth and most posterior sublabial on either side of head rugose and keeled (TM 73290, 78918); 1st pair of sublabials in contact (NMB R9202; TM 78918, 78921, 78931, 83000, 83532), or separated by a narrow groove (TM 51376), large rectangular scale (NMB R9194), elongated triangular scale (NMB R9195), or separated posteriorly by a tiny pair of granules (TM 73290); 1st pair of sublabials followed by three (not two) pairs of smaller, slightly enlarged scales in TM 78931, and by one pair of distinctly enlarged scales in NMB R9202. Spines on sides of neck only about 1.5 times (not twice) as high as wide in juveniles (TM 78921, 78931) and TM 83000. Dorsal scales of TM 78918 and 78931 with short folds rather than distinct striations; two vertebral scale rows smooth in TM 42531, 4–6 rows smooth in TM 83532, 6–8 rows smooth in NMB R9195, none smooth in TM 83000, all vertebrals keeled in juveniles. Dorso-lateral and lateral scales usually keeled and spinose, but weakly spinose in juveniles. Dorsals in 32–41 (34 in allotype) transverse, and 21–26 (21 in allotype) longitudinal, rows. Ventrals occasionally pentagonal (TM 73290, 78918, 83532), longer than broad on anterior part of belly (TM 73290, 78918) or mostly square (NMB R9195, TM 51376 and 83000). All ventrals smooth in TM 51376; in NMB R9194 and TM 83532 only the outermost row of ventrals is moderately keeled and weakly spinose, with rows 2–3 very weakly keeled only; some scales of the 3rd row also very weakly spinose in NMB R9194 (including first inner row) and TM 73290, 78918, 83000; all three outer rows weakly keeled in NMB R9195. Ventrals in 23–29 (28 in allotype) transverse rows (6–9 additional rows on throat), and occasionally only 12 (NMB R9194, TM 51376) longitudinal rows. Enlarged hexagonal preanal plates 3 (TM 78918) or 4 (TM 83532); median preanal plates (pair) pentagonal in TM 42531, 51376, 78931 and 83000, heptagonal in TM

73290 (left side) which also has two extranumerary plates posterior to the large pair, and irregular in TM 78921; no enlarged plates anterior to median and lateral plates in TM 78931; enlarged median pair of plates in TM 42531, 51376, 78931 and 83000 much elongated, about twice as long as wide. Femoral pores 20–24 (10–12 on each thigh, 10 in allotype), appearing as small, shallow pits in females. Differentiated glandular femoral scales in males 19–61 (9–35 per thigh). Fourth toe with 16–19 (18 in allotype) subdigital lamellae.

Colour: Dorsum dark brown to black in preservative. Back with 4–6 (usually 5) interrupted transverse bands between fore- and hindlimbs, which are usually without dark borders after preservation in alcohol. In TM 73290 there is a roundish cream spot on the nape between the pale band behind the occipitals and the band on the neck. Belly with 5–6 brown crossbands on either side of the median band (prominent in the centre of the belly).

Variation in additional material. (All localities in KwaZulu-Natal; material examined only for the characters listed below.) *External morphology* ($N = 10$ unless otherwise indicated): Tail length/SVL 1.35–1.42 (SVL: 97.9–132.7 mm, $N = 3$); head width/SVL = 23.3–24.6% in males (SVL: 97.9–130.7 mm, $N = 3$), 21.4–22.7% in females (SVL: 123.9–132.7 mm, $N = 3$); head width/head length = 76.2–83.9% (SVL: 97.9–132.7 mm, $N = 6$); head depth/head length = 36.2–47.6% (SVL: 97.9–132.7 mm, $N = 6$). Preoculars 1; supraoculars 4 ($N = 9$); supraciliaries 4 (3 on right side in TM 73294; $N = 8$); postnasals 1 ($N = 9$); suboculars 4–5 (4 left, 5 right in TM 83002); supralabials (anterior to median subocular) usually 4 (5 in TM 16828; $N = 9$); infralabials 6 ($N = 9$); sublabials 5 (4 on left side in TM 16827; $N = 9$); occipitals 6 (additional small median scale in TM 16798 and 83002); gulars 24–29 ($N = 9$); scales at anterior edges of ear openings elongate and distinctly spinose (rather than short and blunt) in TM 16828; dorsal rows transversely 31–38; dorsal rows longitudinally 21–24; ventral rows transversely 25–27 ($N = 9$); ventral rows longitudinally usually 14, but 12 in TM 73285 and 773291; femoral pores 11–13 (males, $N = 4$), 10–13 (females, $N = 6$); differentiated femoral scales (generation glands) in males 19–29 per thigh ($N = 3$); lamellae under fourth toe 16–19.

Colour: Similar to holotype. Back with 5–6 (4 in TM 73285) interrupted transverse bands (with slightly dark borders in preservative). Belly with 5–6 brown crossbands on either side of the median band (prominent in the centre of the belly).

Size. Largest male (NMB R9194 [paratype], Komati View Point, Eswatini) $140.0 + 202 = 342$ mm, but NMB R9195 (paratype, Komati View Point) has SVL of 145.0 mm (tail broken/missing). Largest female (TM 78918 [allotype], Nkomati Gorge, Eswatini) $143.8 + 161 = 305$ mm.

Etymology. Named for the Kingdom of Eswatini, the country where most of the species' range is located. Both 'eSwatini' and 'Swaziland' derive from the word *iSwazi*, after the name of an early chief, *Mswati II*.

Distribution. Highveld and Middleveld regions of Eswatini, and adjacent areas in the South African provinces of (eastern) Mpumalanga (in Nkomazi municipality) and (northern) KwaZulu-Natal (in uPhongolo and Abaqulusi municipalities) (*Fig. 8*) at elevations of 462 to 1052 m a.s.l.

Natural history. Rupicolous, living in deep, horizontal (or gently sloping) crevices in granitic rock along hillsides, usually in the partial shade of trees (see also Jacobsen 1989). According to R C Boycott (in litt., 2019), rocky terrain in closed canopy bushveld is the preferred habitat in Eswatini. A specimen in Ithala Game Reserve in KwaZulu-Natal was photographed on a tree trunk (ReptileMAP, VM no. 152451).

Note. The photograph of a specimen of '*Smaug warreni barbertonensis*' from 'Barberton' in Bates et al. (2014) is in fact the same one used for *Fig. 7* in the current paper (i.e. NMB R9194, paratype of *S. swazicus* **sp. nov.**).

Smaug barbertonensis (Van Dam, 1921)

Barberton Dragon Lizard

Figs 9–11

Zonurus barbertonensis Van Dam, 1921: 240 (Barberton); Power, 1930: 14 & 17 (Barberton).
Zonurus barbertonensis barbertonensis FitzSimons, 1933 (by implication after describing
Zonurus barbertonensis depressus).
Cordylus warreni barbertonensis FitzSimons, 1943: 426 (part, Barberton and Nelspruit);
 Loveridge, 1944: 20 (Barberton); Branch, 1988: 164 (part), 1998: 195 (part); Jacobsen, 1989:
 590 (part: 5 km S of Nelspruit; Barberton Townlands 369JU; Broedershoek 129JU;
 Friedenheim 282JT; Karino to White River; Khandizwe; Nelspruit); Adolphs, 1996: 15 (part)
 & 2006: 22 (part);
Smaug warreni barbertonensis Stanley *et al.*, 2011 (part); Bates *et al.*, 2014: 211 (part, but
 excluding fig. on p. 211); Reissig, 2014: 190 (part, including fig. 218).
Smaug barbertonensis Stanley & Bates, 2014: 905; Mouton *et al.*, 2018: 463.

Diagnosis. Distinguished from all other cordylids by its unique combination of dorsal, lateral and ventral colour patterns (see descriptions and figures). It also differs from the terrestrial *S. giganteus* by its smaller adult size (maximum SVL 140 mm versus 198 mm) and possession of six (occasionally four) moderate sized and weakly spinose, versus four (occasionally five) large and distinctly spinose, occipitals. Differs from other species of *Smaug* as follows: from *S. vandami* by usually having six (versus usually four) occipitals; from *S. depressus* by having only 10–12 (versus 16–24) femoral pores in males; from *S. breyeri* by having much less rugose head shields. It differs from *S. giganteus*, *S. breyeri* and *S. vandami* by having less spinose occipitals and tail spines, and two-layer (rather than multi-layer) generation glands.

Most similar to *S. swazicus* **sp. nov.** and *S. warreni*, but easily distinguishable by its colour pattern (back dark brown with 4–5 pale bands between fore- and hindlimbs, pale spot or blotch on nape behind occipitals; flanks dark with narrow pale vertical markings; venter mostly dark brown or black) compared to *S. swazicus* **sp. nov.** (back dark brown usually with 5–6 pale bands, pale band on nape behind occipitals; flanks with large pale spots or blotches; venter pale with a dark central longitudinal band bordered on either side by dark transverse bands) and *S. warreni* (back usually pale brown with 5–6 pale dark-edged bands; flanks pale with brown markings; venter with brown markings on most scales); by usually having more elongate and spinose scales

at the edges of the ear openings (shorter and non-spinose in *S. swazicus* **sp. nov.** and *S. warreni*); and quadrates lacking a pronounced ridge and concave region at the lateral edge of the adductor musculus mandibular posterior origin (with a pronounced ridge and concave region in *S. swazicus* **sp. nov.**). Also differs as follows: outer occipitals shorter than the adjacent inner ones (of about equal length in *S. warreni*); head relatively wider than the other two species (head width/head length = 80–92% versus 73–84%); lower numbers of transverse dorsal scale rows (29–32 in 81%) compared to *S. swazicus* **sp. nov.** (32–37 in 86%) and *S. warreni* (32–38 in 92%); and lower numbers of longitudinal dorsal scale rows (20–24, mean 21.7) compared to *S. warreni* (22–28, mean 23.6).

Variation. ($N = 26$ unless otherwise indicated) *External morphology:* Tail length/SVL 1.21–1.42 (SVL: 109.6–134.0 mm, $N = 7$); head width/SVL = 22.5–25.7% in males (SVL: 111.0–134.0 mm, $N = 9$), 20.9–24.1% in females (SVL: 109.6–139.9 mm, $N = 11$); head width/head length = 80.4–92.0% (SVL: 109.6–139.9 mm, $N = 20$); head depth/head length = 38.2–48.2% (SVL: 109.6–139.9 mm, $N = 20$). Frontonasal in contact with the rostral and loreals, separating the nasals, latter slightly swollen; nostril – with a large inner flap attached posteriorly – situated in the posterior part of the nasal and in contact with the loreal and first supralabial ($N = 4$, topotypes). Scales at anterior edge of ear opening 4–6 on either side of head, projecting outwards as flattened, somewhat spatulate spines, the lowermost one narrow and slender, the one above it distinctly spatulate and the largest; scales generally elongate and somewhat spinose (more so than in *S. swazicus* and *S. warreni*), but short and blunt in TM 55789, 73292 and three juveniles (TM 4275, 26643, 73286). Preoculars 1; postoculars 1; supraoculars 4; supraciliaries usually 4 (5 on one side in three specimens); postnasals 1; suboculars usually 4–6 (often different on either side of head), but 7 on left side in NMB R9192); supralabials (anterior to median subocular) usually 4 (3 on one side in three specimens, and 5 on one side in two specimens); infralabials usually 5–6 (5 on left and 7 on right in TM 73281, and 6 left and 7 right in TM 55787); sublabials usually 5 (6 in NMB R9192); occipitals usually 6 (7 in TM 4468 and 4472, 8 in TM 73284; NMB R9192 has a single tiny median granule, TM 73293 has two such granules), outer occipitals shorter than those adjacent to them and those of the median pair the shortest ($N = 4$); gulars 23–31, but 20 in NMB R9193; dorsal scale rows transversely 28–34, longitudinally 20–24; ventral scale rows transversely 25–28, longitudinally usually 14 (16 in TM 73283);

femoral pores per thigh 8–11 (males, $N = 11$), 8–12 (females and juveniles, $N = 15$); differentiated femoral scales (generation glands) in males ≥ 100 mm SVL: 16–36 per thigh ($N = 10$), in juvenile males < 66 mm SVL: 18–22 per thigh ($N = 2$); lamellae under fourth toe 15–19.

Colour: Back dark brown to black with cream to yellow markings (mostly transversely enlarged) forming 4–5 (usually 4) interrupted transverse bands (usually with dark borders in preserved material) that continue onto the tail, together with a band on the nape and a cream spot, blotch or elongate marking (absent in TM 73286) immediately behind the median occipitals. Belly mostly brown (older material) to black (as in life) with a few cream patches or short ‘bands’ on either side, which are occasionally joined to the pale bands on the side of the back. The flanks are dark brown to black, usually with narrow, cream to yellowish, vertically elongated bars, occasionally spots (e.g. TM 51066). Top of limbs with numerous irregular cream to yellowish spots and blotches; underparts of limbs mostly cream with irregular brown markings, occasionally bands. Top of head brown or black with scattered irregular cream markings. Throat (including sublabials) mostly black or brown with occasional irregular scattered cream markings, but about half dark and half pale in TM 73283 and 73293, and mostly plain cream in TM 4275 (juvenile).

Size. Largest male (TM 73287, Broedershoek) $129.0 + 182.8 = 311.8$ mm, but TM 51066 (between Karino and White River) has SVL of 134.0 mm (tail broken/missing). Largest female (TM 4273: holotype, Barberton) $134.0 + 175 = 309$ mm, but TM 4468 (Barberton) has SVL of 139.9 mm (tail broken/missing).

Natural history. Rupicolous, living in deep, horizontal (or gently sloping) crevices in and between large granitic boulders, often in the partial shade of trees (see also Jacobsen 1989). For *S. ‘barbertonensis’*, FitzSimons (1943: 427) noted that the diet is similar to that of *S. warreni* (see below), but includes cetonid beetles and small land snails; usually five young are produced, and based on his examination of a series of females, fertilisation occurs in early spring, with young born at the end of summer. However, one of FitzSimons’ (1943) localities (‘Hluti–Goedgegun’, Eswatini) is referable to *S. swazicus* sp. nov., so it is not possible to know which species his data applies to. Computed Tomography scanning of NMB R9192 revealed four large embryos, and a large beetle in the stomach.

Distribution. Restricted to the Barberton, Nelspruit and Khandizwe areas of eastern Mpumalanga Province, South Africa (*Fig. 8*) at elevations of 724 to 1008 m a.s.l. An isolated record for this species at Farm: Jessievale, Ermelo district (2630AB, Bates *et al.* 2014) is in fact referable to *Cordylus vittifer* (re-examination by both authors of VM no. 1400 on ReptileMAP).

Localities. SOUTH AFRICA: Mpumalanga Province. Barberton (25°47'S, 31°03'E) TM 4273–5, 4468–9, 4471–2; Barberton army base – NMB R9191, 9196 (25°46'26"S, 31°03'20"E; 861 m a.s.l.), NMB R9192–3 (25°46'27"S, 31°03'21"E; 861 m a.s.l.); Barberton Townlands (25°47'S, 31°03'E) TM 73281, 73283–4; Broedershoek 129JU (25°27'S, 31°07'E, 753 m a.s.l.) TM 73286–7; Friedenheim 282JT (25°26'S, 30°59'E; 754 m a.s.l.) TM 55787; Karino and White River, between (2531AC) TM 51066; Khandizwe (25°28'S, 31°25'E; 724 m a.s.l.) TM 73292–3; Nelspruit, 5 km S of (25°32'S, 30°57'E; 824 m a.s.l.) TM 44873; Nelspruit, 14 km W of, on road to Machadodorp (2530BD) TM 26643; Nelspruit, 82 Ehmke Street, Extension 5 (*Fig. 11*); Nelspruit, Van Riebeeck Park (25°28'S, 30°59'E) TM 55788–9.

Notes. Van Dam's (1921) type locality of 'Barberton' does not indicate the exact locality at which the specimens were collected, so it is considered appropriate to treat the localities of 'Barberton Townlands' (Jacobsen 1989) and 'Barberton army base' as topotypic.

***Smaug warreni* (Boulenger, 1908)**

Lebombo Dragon Lizard

Figs 12–14

Zonurus warreni Boulenger, 1908: 232 (Ubombo); Hewitt, 1909: 36; Boulenger, 1910: 467 & 468; Power, 1930: 14 & 17; FitzSimons, 1930: 30; Lawrence, 1937: 111. *Cordylus warreni warreni* FitzSimons, 1943: 424 (Ubombo & Ingwavuma); Loveridge, 1944: 19 (Ubombo); Branch, 1988: 164, 1998: 195; Jacobsen, 1989: 586 (Duikershoek, Halfkroonspruit, Jozini Dam, Mananga, The Hippos); Adolphs, 1996: 15; Bourquin, 2004: 96 (KwaZulu-Natal); Adolphs, 2006: 22.

Smaug warreni warreni Stanley *et al.*, 2011; Bates *et al.*, 2014: 210; Reissig, 2014: 187.

Smaug warreni Stanley & Bates, 2014: 905; Mouton *et al.*, 2018: 463.

Diagnosis. Distinguished from all other cordylids by its unique combination of dorsal, lateral and ventral colour patterns (see descriptions and figures). It also differs from the terrestrial *S. giganteus* by its smaller adult size (maximum SVL 141 mm versus 198 mm) and possession of six (occasionally 7 or 8) moderate sized and weakly spinose, versus four (occasionally five) large and distinctly spinose, occipitals; and from other species of *Smaug* as follows: Differs from *S. vandami* by usually having six (versus usually four) occipitals. Differs from *S. depressus* by having only 8–12 (versus 16–24) femoral pores in males; and from *S. breyeri* by having much less rugose head shields. It differs from *S. giganteus*, *S. breyeri* and *S. vandami* by having less spinose occipitals and tail spines, and two-layer (rather than multi-layer) generation glands.

Most similar to *S. swazicus* **sp. nov.** and *S. barbertonensis* but easily distinguished by its colour pattern (back usually pale brown with 5–6 pale dark-edged bands between fore- and hindlimbs, pale band on nape behind occipitals; flanks pale with brown markings; venter with brown markings on scales) compared to *S. swazicus* **sp. nov.** (back dark brown usually with 5–6 pale bands, pale band on nape behind occipitals; flanks with large pale spots or blotches; venter pale with a dark central longitudinal band bordered on either side by dark transverse bands) and *S. barbertonensis* (back dark brown with 4–5 pale bands, pale spot or blotch on nape behind occipitals; flanks dark with narrow pale vertical markings; venter mostly dark brown or black); by usually having less spinose-like scales at the edges of the ear openings than *S. swazicus* **sp. nov.**; and quadrates without a pronounced ridge and concave region at the lateral edge of the adductor musculus mandibulae posterior origin (with a pronounced ridge and concave region in *S. swazicus* **sp. nov.**). Also differs as follows: outer occipitals and scales adjacent to them of about equal length (outer occipitals shorter than the adjacent inner ones in the other two species); head narrower than *S. barbertonensis* (head width/head length = 73–83% versus 80–92%); generally higher numbers of transverse dorsal scale rows (32–38 in 92% of specimens) than *S. barbertonensis* (29–32 in 81%); and greater numbers of longitudinal dorsal scale rows (22–28, mean 23.6) than *S. barbertonensis* (20–24, mean 21.7).

Variation. *External morphology* ($N = 39$ unless otherwise indicated): Tail length/SVL 1.06–1.43 (SVL: 105.0–128.9 mm, $N = 6$); head width/SVL = 22.2–24.2% in males (SVL: 105.3–127.3 mm, $N = 9$), 19.4–23.3% in females (SVL: 100.6–141.1 mm, $N = 21$); head width/head length = 72.5–83.1% (SVL: 100.6–141.1 mm, $N = 30$); head depth/head length = 35.0–47.2% (SVL: 100.6–141.1 mm, $N = 31$). Frontonasal in contact with the rostral and loreals (often in narrow contact, especially on right side in TM 50130), separating the nasals, latter slightly swollen; nostril – with a large inner flap attached posteriorly – situated in the posterior part of the nasal and in contact with the loreal and 1st supralabial (TM 50130: loreal in very narrow contact with nostril on left side of head and separated by first supralabial on right) ($N = 16$). Scales at anterior edge of ear opening usually 4–5 (3 on right side of head of TM 63567) projecting outwards as flattened, somewhat spatulate, spines, the lowermost one narrow and slender (especially so in TM 47449), the one above it distinctly spatulate and the largest; spines generally long and somewhat spinose in TM 50130, 70961 ($N = 16$). Preoculars 1; supraoculars 4 (2nd and 3rd on left side largely fused in NMB R10913); supraciliaries usually 4–5 (3 on left side of TM 78967, 4 left and 6 right in TM 13639, 6 on left side of 78969); postnasals 1; suboculars usually 4–5, occasionally 6; supralabials (anterior to median subocular) usually 4 (3 on one side in four specimens, 5 on one side of TM 78967); infralabials usually 6 (5 on both sides of TM 582, 5 on one side of TM 47449, 7 on both sides of TM 15320, 7 on one side in six specimens); sublabials 5; occipitals usually 6, but an additional – often narrow and much elongated – scale medially in 41% of specimens (median scale large in TM 50660, granular in TM 78966); outermost occipitals and those adjacent to them of similar size and length, but scales of the inner pair shorter and often smaller, except in NMB R10878 in which all occipitals are of similar length, although those of the inner pair are wider ($N = 16$); gulars 23–32 ($N = 37$); dorsal scale rows transversely 31–41, longitudinally 22–28; ventral scale rows transversely 23–27, longitudinally usually 14 (13 in NMB 9199, 12 in TM 2808 and 78969); femoral pores 7–13 (males [smallest is 86.2 mm SVL], $N = 11$), 8–13 (females and juveniles, $N = 27$); differentiated femoral scales (generation glands) in males 13–38 per thigh ($N = 10$); lamellae under fourth toe 15–20.

Colour: Back usually tan/khaki brown with irregular, black-bordered, cream or white blotches (ocelli) forming 5–6 slightly interrupted transverse bands between the legs that continue onto the tail, together with a band immediately behind the occipitals and another on the nape. Ocelli may

be in close proximity, creating the impression of distinct bands across the back. Occasional specimens have grey-brown backs with small, widely separated white spots lacking obvious black-borders (e.g. fig. 212 in *Reissig, 2014*). Belly white to cream, usually with numerous small, square, rectangular or irregular pale to dark brown markings; occasionally mostly without markings except for the sides (e.g. NMB R10898 and 10912) or with a large dark blotch on each ventral (NMB R9292). The flanks are mostly pale whitish, occasionally with some darker colouring and pale vertical bars. Top of limbs with numerous irregular cream to yellowish spots and blotches; underparts of limbs mostly cream with occasional scattered, irregular brown markings. Top of head tan/khaki brown with dark brown patches and scattered, irregular, cream markings (or small yellow speckles); throat mostly white to cream with varying amounts of darker markings (often extensive and bold).

Size. Largest male (TM 78963, Mananga Mountain, Mpumalanga, South Africa) $127.3 + 193.5$ [on museum tag] = 320.8 mm. Largest female (NMB R9292, Mananga Mountain) $128.9 + 182 = 311$ mm but TM 53869 (Lomahasha, Eswatini) has SVL of 141.1 mm (tail broken/missing).

Natural history. Diurnal and rupicolous, occurring in crevices between or under rocks on outcrops along the Lebombo mountains. According to *FitzSimons' (1943)*, ants, beetles, fossorial wasps, myriapods, frogs and lizards are eaten. *Loveridge (1944)* noted that for a sample from Ubombo, one specimen had eaten 32 *Eristalis* (drone fly) maggots, another lizard contained millipedes and ants, while a third had consumed a large grasshopper. Females give birth to 4–5 young in late summer (*FitzSimons, 1943*).

Distribution. Endemic to the Lebombo Mountains of eastern Eswatini, adjacent western Mozambique and South Africa (north-eastern Mpumalanga and north-eastern KwaZulu-Natal) (*Fig. 8*) at elevations of 82 to 745 m a.s.l.

Localities. (*specimens examined by D.G. Broadley – data not included in morphological analysis) MOZAMBIQUE: Estatuene ($26^{\circ}24'18''\text{S}$, $32^{\circ}04'42''\text{E}$) NMZB-UM 30510–3*; Meponduine ($25^{\circ}56'45''\text{S}$, $31^{\circ}58'44''\text{E}$) NMZB 30514*, 30562*; near Moambo close to Komati River ($25^{\circ}35'23''\text{S}$, $32^{\circ}14'47''\text{E}$; photo: L. Verburgt [pers. comm.], in *Reissig, 2014*: 188, fig.

812 213). SOUTH AFRICA: KwaZulu-Natal Province. Bhokweni (27°22'S, 32°03'E) TM 78969–
 813 72, 78974; Ingwavuma (27°08'S, 32°01'E) TM 15319–20; Mayaluka (26°54'52"S, 32°00'31"E)
 814 NMB R10878, 10911–2; Farm: Middlein 84 (27°21'03.0"S, 31°59'10.7"E; photographic record:
 815 S. Nielsen); Ubombo (27°34'S, 32°05'E) TM 582, 2808, 13639–41; NMZB–UM 1542*.
 816 Mpumalanga. Duikerhoek 489JU (25°42'S, 31°57'E) TM 78966; Halfkroon Spruit, Kruger
 817 National Park (2531BD) TM 78973, 'J6955'; Mananga Mountain (25°58'S, 31°52'E) TM
 818 78961–5; Mananga Mountain, 2 km SSW of Nsizwane (25°54'12"S, 31°52'12"E) NMB R9197–
 819 200, 9292; The Hippos 192JU (25°28'S, 31°57'30"E) TM 78967–8. ESWATINI: Lomahasha
 820 (25°59'S, 31°59'E) TM 50130–1, 53869, 63567; Lubombo foothills, 6 km E of Big Bend
 821 (26°47'S, 31°59'E) TM 70960–1; Siteki, S of (26°28'S, 31°56'E) TM 47449; Tshaneni
 822 (25°59'S, 31°46'E) TM 50660–1. NO DATA: AMNH 173381 (used for CT scanning).

823

824 **Notes.** *Boulenger's (1908)* description was based on two male specimens (i.e. syntypes) from
 825 Ubombo in the Lebombo Mountains of KwaZulu-Natal. His description includes a plate with a
 826 splendid drawing (by A.H. Searle) depicting a specimen with somewhat indistinct, narrow, dark
 827 crossbars on the back, each containing scattered pale spots. This illustration, together with
 828 *Boulenger's (1908: 233)* description: "Dark brown above, with small yellow black-edged spots
 829 forming more or less regular transverse series on the body; lower parts pale brown" characterise
 830 *S. warreni* (but back usually light brown, see *Fig. 14*). *Reissig (2014)* noted that the 'type
 831 specimen' is NHM 1946.8.8.1.



A revised diagnostic key to the genus *Smaug*

- 1a. Occipitals greatly enlarged, the outer ones strongly spinose and about twice as long as those of the median pair; dorsal scales strongly spinose; ventral scales imbricate; lamellae under fourth toe 10–12 *S. giganteus*
- 1b. Occipitals moderately to weakly enlarged, those of the outer pair somewhat larger or of similar size to the others; dorsal scales not strongly spinose; ventral scales non-imbricate; lamellae under fourth toe 14–20 2
- 2a. Occipitals of the outermost pair largest (and longest), innermost the smallest (and shortest). 3
- 2b. Occipitals of the outermost pair not the largest (or longest), innermost of similar size to other occipitals or slightly smaller (and shorter) 4
- 3a. Dorsum mostly plain brown, at most with occasional scattered pale markings; belly cream or brown; throat plain or with small brown spots; ventrals in 10–14 rows longitudinally *S. breyeri*
- 3b. Dorsum brown with transversely enlarged cream markings, at least at the sides of the back, but often extensively on the back and tail; belly dark with short pale transverse markings, especially towards the edges; throat pale with dark reticulations; ventrals in 12–16 rows longitudinally *S. vandami*
- 4a. Back with few or no pale markings; flanks mostly plain, and brightly coloured (red, orange or yellow) in males; first supralabial with distinct upward prolongation; dorsals in 22–30 rows longitudinally 5
- 4b. Back usually with distinct pale markings (except the ‘laevigata’ form of *S. depressus*); flanks of males with light and dark markings, and not brightly coloured; first supralabial with moderate or no upward prolongation; dorsals in 13–24 rows longitudinally 6

- 862 5a. Loreal large and not elongated, separated from nostril by upward prolongation of first
 863 supralabial; preocular usually widely separated from the nasal by the loreal; throat of male
 864 uniform dark brown *S. mossambicus*
- 865 5b. Loreal small and elongated, in contact with nostril; preocular large and usually in contact (or
 866 nearly so) with the nasal above the loreal; throat of male yellow or orange, with dark
 867 infuscations *S. regius*
- 868
- 869 6a. Back usually with distinct, small to moderate, scattered white spots or irregular markings, not
 870 forming crossbands, or completely plain grey ('laevigata' form); dorsals in 13–21 rows
 871 longitudinally; differentiated femoral scales (generation glands) in males 14–16
 872 *S. depressus*
- 873 6b. Back with distinct crossbands (usually interrupted) consisting of cream spots, blotches or
 874 transversely enlarged bars, often with dark edges; dorsals in 18–28 rows longitudinally;
 875 differentiated femoral scales (generation glands) in males 19–38 7
- 876
- 877 7a. Outer occipital of similar length to the occipital adjacent to it; small occipital often present
 878 between median pair; back medium to light brown with distinctly dark-edged pale spots or
 879 blotches forming crossbands; belly usually with centre of each scale brown (not mostly black
 880 or brown, or with brown crossbars interrupted by a median band); throat with small brown
 881 spots *S. warreni*
- 882 7b. Outer occipital usually shorter than the occipital adjacent to it; small occipital seldom present
 883 between median pair; back dark brown to black with pale markings, mostly in the form of
 884 narrow, transversely enlarged bars; throat black or with dark reticulations 8
- 885
- 886 8a. Back with 4–5 pale crossbands between the fore-and hindlimbs, with a pale spot on the nape
 887 behind the occipitals; throat mostly black; flanks dark with narrow, pale, vertical bars; belly
 888 mostly black, with a few pale markings at the sides; scales at anterior edges of ear openings
 889 elongated and spinose; dorsals in 28–34 (mostly ≤ 32) rows transversely
 890 *S. barbertonensis*
- 891 8b. Back usually with 5–6 pale crossbands between the fore-and hindlimbs, with a pale band on
 892 the nape behind the occipitals; throat pale with dark reticulations; flanks with large cream

spots and blotches; belly with brown crossbars interrupted by a dark median band; scales at anterior edges of ear openings short and blunt; dorsals in 31–41 (mostly ≥ 32) rows transversely *S. swazicus* **sp. nov.**

DISCUSSION

A molecular assessment of the genus *Smaug* by Stanley & Bates (2014) resulted in an unexpected finding – that populations in Eswatini, previously considered referable to *S. barbertonensis*, formed a distinct evolutionary lineage phylogenetically more closely allied to *S. warreni*, a species restricted to the narrow Lebombo mountain range. We subsequently noted distinct differences in colour pattern between vouchers of each of the three lineages referred to above, which we now collectively refer to as the *S. warreni* species complex. In the present study we therefore conducted a detailed morphological analysis of museum material of all populations identified as *S. warreni* and *S. barbertonensis*.



We found that specimens of the ‘Eswatini’ lineage—including populations in small areas on the northern Eswatini-Mpumalanga border, and in northern KwaZulu-Natal Province in South Africa—were readily distinguishable from *S. barbertonensis* sensu stricto (and *S. warreni*) by their unique dorsal, lateral and ventral colour patterns. FitzSimons (1943) had in fact noted differences in colour pattern between specimens of *S. barbertonensis* from the type locality of Barberton and specimens from ‘Hluti-Goedegun’ in Eswatini (now referred to the new species), but this had been regarded as merely representing regional variation.

In order to further assess the taxonomic status of the three populations, a detailed morphological analysis was conducted. Multivariate analyses of scale counts and body dimensions indicated that the ‘Eswatini’ lineage and *S. warreni* were most similar. In particular, *S. barbertonensis* differed from the other two lineages by its generally lower numbers of transverse rows of dorsal scales, more spinose scales at the anterior edges of the ear openings, and a relatively wider head. Also, the outer two occipital scales in *S. warreni* are of similar length, distinguishing it from the other two species which have the outer occipital usually slightly shorter than the adjacent inner occipital. High resolution Computed Tomography revealed differences in cranial osteology between specimens from all three lineages, with the

‘Eswatini’ lineage being remarkable in having a pronounced ridge and concave region at the lateral edge of the posterior origin of the adductor musculus mandibulae.

The ‘Eswatini’ lineage is described here as a new species, *Smaug swazicus* **sp. nov.** It appears to have a fairly widespread distribution in Eswatini west of the Lebombo Mountains, and a somewhat peripheral distribution in South Africa near its borders with Mpumalanga and KwaZulu-Natal provinces. We estimate that about 90% of its range is in Eswatini and suggest that it be considered near-endemic to that country. Recognition of the new species means that *S. barbertonensis* sensu stricto is a South African endemic restricted to an altitudinal band of about 300 m in the Barberton–Nelspruit–Khandizwe area of eastern Mpumalanga Province, while *S. warreni* is endemic to the narrow Lebombo Mountain range of South Africa, eSwatini and Mozambique.

The phylogenetic analysis of Stanley & Bates (2014) did not include samples from the southern part of the range (especially KwaZulu-Natal) of *S. swazicus* **sp. nov.**, but northern (including one locality in Mpumalanga) and southern material is morphologically indistinguishable, and so we provisionally treat *S. swazicus* **sp. nov.** as a single species.

The geographical break between *S. barbertonensis* and *S. swazicus* appears to correspond to the location of the ancient Makhonjwa mountain range, which lies directly south of Barberton. This range, also referred to as the Barberton Greenstone belt, is made up of some of the world’s oldest exposed rocks (3.6 Ga) which contain fossilised evidence of the earliest life on Earth (De Wit 2010). The time-calibrated phylogenetic analyses of the *S. warreni* species complex by Stanley & Bates (2014) indicates that the *S. warreni*–*S. swazicus* **sp. nov.** lineage diverged from *S. barbertonensis* during the late Miocene, around 7.5 million years ago. This is somewhat earlier than the most recent and extreme period of uplift of the eastern escarpment (Partridge & Maud, 1987), suggesting that populations on either side of the Makhonjwa mountains were isolated before that time. The population east of the Makhonjwa mountains split around 6.2 million years ago (Stanley & Bates, 2014), after which time *S. warreni* became closely associated with the narrow Lebombo mountain range.

Conservation implications

Due to their obligate rupicolous ecology, members of the *Smaug warreni* species complex are not subject to the same levels of habitat destruction as their terrestrial congener, *S. giganteus*. *Jacobsen (1989)* listed all species of *Smaug* (except the Vulnerable *S. giganteus*) as protected schedule 2 (Transvaal Nature Conservation Ordinance 12 of 1983), while *Bates et al. (2014)* and *Bates & Mouton (2018a)* reported their global conservation status as “Least Concern”, while recommending that further research is needed to assess the impact of tree removal from the habitat of *S. barbertonensis* (i.e. *S. barbertonensis* and *S. swazicus* **sp. nov.**) as crevices in the partial shade of trees are often selected for shelter (*Jacobsen, 1989*). In this regard Richard C. Boycott (2019, in litt.) noted that when he visited the locality ‘between Hluti and Goedgegun’ in Eswatini (as reported by FitzSimons 1943) a few years ago, *S. swazicus* **sp. nov.** was not present, possibly because all large trees along the rocky hillsides had disappeared, such that dappled shade was no longer available. The species appeared to have been replaced by skinks (*Trachylepis varia* [Peters] and *T. margaritifer* [Peters]). Part of the natural range of *S. swazicus* **sp. nov.** was inundated and thus lost to the species when the Maguga Dam in Eswatini was filled in 2002/3, although about 20 specimens were collected by Boycott and relocated downstream from the Dam as part of the Maguga Dam Comprehensive Mitigation Plan (R C Boycott, 2019, in litt.). The recognition of *S. swazicus* **sp. nov.** means that the range of *S. barbertonensis* sensu stricto now covers only about 180 km². Also, this species has been recorded within a narrow altitudinal band of only 300 m. Its conservation status should therefore be monitored. Despite being endemic to the narrow Lebombo mountain range, *S. warreni* apparently does not face any significant threats, and it is therefore also considered “Least Concern” (*Bates et al., 2014; Bates & Mouton, 2018a*). It appears to occur throughout the Lebombo range, from low to high altitudes (82 to 745 m a.s.l.). Using *IUCN (2012, 2017)* criteria, we suggest that all three species be regarded as Least Concern at this time.

CONCLUSIONS

Following the finding by *Stanley & Bates (2014)* that the south-eastern assemblage of populations referable to the *S. warreni* species complex comprised three distinct genetic lineages, we hypothesised that morphological differences should also exist between specimens referable to these lineages. Distinct differences were indeed identified between populations with regard to

colour pattern, scalation and cranial osteology, necessitating the description of a new species, *S. swazicus* **sp. nov.**, which appears to be near-endemic to eSwatini. This finding means that *S. barbertonensis* sensu stricto is endemic to South Africa, with a restricted range that may require monitoring in future to ensure that the species does not become threatened with extinction. Also, sampling of populations referable to *S. swazicus* **sp. nov.** in South Africa's KwaZulu-Natal Province is needed to investigate whether additional cryptic diversity exists in this species complex. *Smaug warreni* is endemic to the Lebombo range in South Africa, Eswatini and Mozambique. There are now **nine** known species of dragon lizards (*Smaug*).



ACKNOWLEDGEMENTS

We thank Lauretta Mahlangu (Ditsong Natural History Museum, Pretoria) for access to, and for loans of, *Smaug* material in her care; the late Donald Broadley for data on specimens in the collection of the Natural History Museum of Zimbabwe (Bulawayo); Richard Boycott for information about the distribution of this genus in Eswatini; and T. Busschau for the use of his photographs.

REFERENCES

- ADOLPHS, K. 1996. Bibliographie der Gürtelchsen und Schildchsen (Reptilia: Sauria: Cordylidae & Gerrhosauridae). Squamata Verlag, Sankt Augustin, 255 pp.
- ADOLPHS, K. 2006. Bibliotheca Cordyliformium. Squamata Verlag, Sankt Augustin, 303 pp.
- BATES, M. F., W. R. BRANCH, A. M. BAUER, M. BURGER, J. MARAIS, G. J. ALEXANDER, AND M. S. DE VILLIERS eds. 2014. Atlas and Red List of the Reptiles of South Africa, Lesotho and Swaziland. Suricata 1. Pretoria: South African National Biodiversity Institute.
- BATES, M.F. & MOUTON, P.L.F.N. 2018a. *Smaug barbertonensis*. The IUCN Red List of Threatened Species 201`8: e.T110167262A115679786.

<http://dx.doi.org/10.2305/IUCN.UK.2018-2.RLTS.T110167262A115679786.en>.

Downloaded: 2 January 2019.

BATES, M.F. & MOUTON, P.L.F.N. 2018b. *Smaug warreni*. The IUCN Red List of Threatened Species 2018: e.T110167218A115679533. <http://dx.doi.org/10.2305/IUCN.UK.2018-2.RLTS.T110167218A115679533.en>. Downloaded: 2 January 2019.

BOULENGER, G. A. 1908. On a collection of fresh-water fishes, batrachians, and reptiles from Natal and Zululand, with description of new species. *Annals of the Natal Government Museum (Pietermaritzburg)* 1:219–235.

BOULENGER, G. A. 1910. A Revised List of the South African Reptiles and Batrachians, with Synoptic Tables, special reference to the specimens in the South African Museum, and Descriptions of New Species. *Annals of the South African Museum* 5(9):455–538.

BOURQUIN, O. 2004. Reptiles (Reptilia) in KwaZulu-Natal: I – diversity and distribution. *Durban Museum Novitates* 29:57–103.

BRANCH, W. R. 1988. Field guide to snakes and other reptiles of southern Africa. Struik Publishers, Cape Town, 328 pp.

BRANCH, W. R. 1998. Field guide to snakes and other reptiles of southern Africa. Third edition. Struik Publishers, Cape Town, 399 pp.

BROADLEY, D. G. 1962. On some reptile collections from the North–Western and North–Eastern Districts of Southern Rhodesia 1958–61, with descriptions of four new lizards. *Occasional Papers of the National Museums of Southern Rhodesia* 3:787–843.

BROADLEY, D. G. 1966. The herpetology of south-east Africa. Unpublished Ph.D. thesis, University of Natal, Pietermaritzburg.

- BROADLEY, D. G. 2006. CITES Standard Reference for the Species of *Cordylus* (Cordylidae, Reptilia): CITES Nomenclature Committee for the 14th meeting of the CoP. 10.
- DE QUIEROZ, K. 1998. The General Lineage Concept of Species, Species Criteria, and the Process of Speciation. A Conceptual Unification and Terminological Recommendations. Pp. 55–75 in D. J. Howard, and S. H. Berlocher (Eds.). Endless forms: species and speciation. Oxford University Press, U.S.A.
- DE QUIEROZ, K. 2007. Species Concepts and Species Delimitation. Systematic Biology 56(6):879–886.
- DE WIT, M. J. 2010. The deep-time treasure chest of the Makhonjwa mountains. South African Journal of Science 106(5/6): 1–2.
- DE WAAL, S. W. P. 1978. The Squamata (Reptilia) of the Orange Free State, South Africa. Memoires van die Nasionale Museum, Bloemfontein 11:i–iii, 1–160.
- FITZSIMONS, V. 1930. Descriptions of new South African Reptilia and Batrachia, with distribution records of allied species in the Transvaal Museum Collection. Annals of the Transvaal Museum (Pretoria) 14:20–48.
- FITZSIMONS, V. 1933. Descriptions of five new lizards from the Transvaal and Southern Rhodesia. Annals of the Transvaal Museum (Pretoria) 15:273–280.
- FITZSIMONS, V. 1943. The lizards of South Africa. Memoirs of the Transvaal Museum (Pretoria) 1:1–528.
- FITZSIMONS, V. 1958. A new *Cordylus* from Gorongosa, Mocambique. Annals of the Natal Museum (Pietermaritzburg) 14:351–353.

- FROST, D. R. AND D. M. HILLIS 1990. Species in concept and practice: Herpetological applications. *Herpetologica* 46(1):87–104.
- HEWITT, J. 1909. Description of a New Species of *Platysaurus* and Notes on the Specific Characters of certain Species of *Zonuridae*, together with Synoptical Keys to all the known South African Species, and a résumé of our knowledge of their Distribution; and a Key to the known Genera of South African Lizards. *Annals of the Transvaal Museum* 2(1):29–40, 1 plt.
- IUCN [International Union for Conservation of Nature]. 2012. *IUCN Red List Categories and Criteria: Version 3.1*. Second edition. IUCN, Gland, Switzerland and Cambridge, UK.
- IUCN [International Union for Conservation of Nature] [Standards and Petitions Subcommittee]. 2017. Guidelines for Using the IUCN Red List Categories and Criteria. Version 13. Prepared by the Standards and Petitions Subcommittee. (Downloadable from: <http://www.iucnredlist.org/documents/RedListGuidelines.pdf>.)
- JACOBSEN, N. H. G. 1989. A herpetological survey of the Transvaal. Unpublished PhD thesis, University of Natal.
- LAWRENCE, R. F. 1937. The Girdle Tailed Lizard and its Mites. *The Cape Naturalist*, 1(4):107–113.
- LOVERIDGE, A. 1944. Revision of the African lizards of the family Cordylidae. *Bulletin of the Museum of Comparative Zoology at Harvard College*, 95(1):1–118, 12 pls. 
- MOUTON, P. LE F. N., A. FLEMMING, M. F. BATES, AND BROECKHUIZEN, C. 2018. The relationship between generation gland morphology and armour in Dragon Lizards (*Smaug*): A reassessment of ancestral states for the Cordylidae. *Amphibia–Reptilia* 39:457–470.

- PARTRIDGE, R. R., AND T. C. MAUD. 1987. Geomorphic evolution of southern Africa since the Mesozoic. *South African Journal of Geology* 90(2):179–208.
- POWER, J. H. 1930. On the South African Species of the genus *Zonurus*. *Annals of the Transvaal Museum*, 14(1):11–19, 2 pls.
- REISSIG, J. 2014. Girdled Lizards and their relatives: Natural History, Captive Care and Breeding. Edition Chimaira, Frankfurt am Main, 249 pp.
- SMITH, A. 1844. Illustrations of the Zoology of South Africa. Consisting chiefly of figures and descriptions of the objects of natural history collected during an expedition into the interior of South Africa, in the years 1834, 1835, and 1836. Fitted out by ‘The Cape of Good Hope Association for Exploring Central Africa’. London: Smith, Edler and Co.
- STANLEY, E. L., AND M. F. BATES 2014. Here be dragons: a phylogenetic and biogeographical study of the *Smaug warreni* species complex (Squamata: Cordylidae) in southern Africa. *Zoological Journal of the Linnean Society* 172:892–909.
- STANLEY, E. L., A. M. BAUER, T. R. JACKMAN, W. R. BRANCH, AND P. LE F. N. MOUTON 2011. Between a rock and a hard polytomy: rapid radiation in the rupicolous girdled lizards (Squamata: Cordylidae). *Molecular Phylogenetics and Evolution* 58:53–70.
- VAN DAM, G. 1921. Descriptions of new species of *Zonurus*, and notes on the species of *Zonurus* occurring in the Transvaal. *Annals of the Transvaal Museum (Pretoria)* 7:239–243.

Table 1(on next page)

Comparison of scalation data in the three species of the *Smaug warreni* species complex. Data for *S. cf. barbertonensis* is based on all type and additional material.

For ventral scale rows longitudinally, rare exceptions are indicated in parentheses, and superscripts indicate the relevant numbers of specimens. Femoral pores (males and females >95 mm SVL only) and differentiated femoral scales (generation glands, males >95 mm SVL only) are expressed as average number per thigh (left and right sides examined).

Table 1. Comparison of scalation data in the three species of the *Smaug warreni* complex. Data for *S. cf. barbertonensis* is based on all type and additional material. For ventral scale rows longitudinally, rare exceptions are indicated in parentheses, and superscripts indicate the relevant numbers of specimens. Femoral pores (males and females >95 mm SVL only) and differentiated femoral scales (generation glands, males >95 mm SVL only) are expressed as average number per thigh (left and right sides examined).

	Dorsal scale rows longitudinally	Dorsal scale rows transversely	Ventral scale rows longitudinally	Ventral scale rows transversely	Lamellae under 4th toe	Femoral pores (per thigh) (males and females)	Differentiated femoral scales in males (per thigh)	Gular scales between posterior sublinguals
<i>Smaug</i>	20–26	31–41	14	23–29	16–19	10–13	9.5–33	22–29
<i>cf. barbertonensis</i>	22.6 ± 1.53 <i>N</i> = 22	34.7 ± 2.37 <i>N</i> = 22	(12 ²)	26.3 ± 1.24 <i>N</i> = 21	17.6 ± 0.85 <i>N</i> = 22	11.2 ± 0.90 <i>N</i> = 17	23.2 ± 7.25 <i>N</i> = 9	25.7 ± 1.93 <i>N</i> = 20
<i>Smaug</i>	20–24	28–34	14	25–28	15–19	8.5–11.5	18.5–35.5	(20) 23–31
<i>barbertonensis</i>	21.7 ± 1.46 <i>N</i> = 26	30.5 ± 1.48 <i>N</i> = 26	(16 ¹)	26.4 ± 0.86 <i>N</i> = 26	17.7 ± 1.04 <i>N</i> = 26	9.9 ± 0.79 <i>N</i> = 20	25.9 ± 5.16 <i>N</i> = 9	27.0 ± 2.47 <i>N</i> = 26
<i>Smaug</i>	22–28	31–41	14	23–27	15–20	7.5–13	14.5–38	23–32
<i>warreni</i>	23.6 ± 1.42 <i>N</i> = 39	35.2 ± 2.11 <i>N</i> = 39	(12 ² , 13 ¹)	25.9 ± 1.00 <i>N</i> = 39	16.8 ± 1.20 <i>N</i> = 39	10.4 ± 1.30 <i>N</i> = 31	25.7 ± 6.81 <i>N</i> = 9	26.8 ± 1.94 <i>N</i> = 37

Table 2(on next page)

Results of linear discriminate analysis.

Table 2: Results of linear discriminate analysis.

Character	LD1	LD2
Head width	-0.7844609	-0.1758688
Head length	0.88220298	0.1207981
Head depth	-0.5397809	0.02969013
Supraciliaries	-0.3409638	0.64462659
Subocular	-0.3379355	-0.3798594
Supralabials	-1.2012326	-1.3092285
Infralabials	0.25093769	0.44987475
Sublabial	-2.1044475	2.13129379
Occipitals	-0.0857466	1.30120872
Gulars	-0.1238755	0.17796345
Dorsals transversely	0.42902426	-0.2617689
Dorsals longitudinally	0.15039435	0.34392291
Ventrals transversely	-0.2437863	0.04772589
Ventrals longitudinally	-0.3060718	0.95058204
Femoral pores	-0.0504546	-0.0592088
Differentiated femoral scales (generation glands)	0.00847055	0.00312826
Lamellae on 4th toe	-0.0151936	-0.7436493

Proportion of trace: LD1 = 0.593, LD2 = 0.407

Table 3(on next page)

Mensural data (mm) for the type series of *Smaug swazicus* sp. nov. (M = male, F = female, J = juvenile; r = regenerating tail).

Table 3. Mensural data (mm) for the type series of *Smaug swazicus* **sp. nov.** (M = male, F = female, J = juvenile; r = regenerating tail).

Museum number	Type status	Sex	Snout-vent length	Tail length	Total length	Head length	Head width	Head depth
NMB R9201	Holotype	M	138.76	187	325.76	40.19	31.15	16.19
TM 78918	Allotype	F	143.80	161	304.80	37.23	29.14	15.97
TM 83000	Paratype	M	129.35	150	279.35	34.86	27.99	14.85
TM 83532	Paratype	F	132.08			35.11	28.55	15.08
TM 42531	Paratype	F	102.77			29.23	23.71	12.25
TM 51376	Paratype	M	129.60			36.37	29.60	17.71
TM 78931	Paratype	J	70.14			21.00	16.30	8.00
TM 78921	Paratype	J	65.76			19.33	14.07	8.14
NMB R9194	Paratype	M	139.95	202	341.95	40.53	33.51	17.92
NMB R9195	Paratype	M	145.01			41.96	34.36	18.71
NMB R9202	Paratype	M	141.23	140r	281.23+	40.11	32.73	16.61
TM 73290	Paratype	F	140.11	116.40r	256.51+	37.49	30.97	17.86

Table 4(on next page)

Meristic data for the type series of *Smaug swazicus* sp. nov. (H = holotype, A = allotype, P = paratype); values on either side of a slash refer to the animal's left and right sides respectively.

Meristic data for the type series of *Smaug swazicus* **sp. nov.** (H = holotype, A = allotype, P = paratype); values on either side of a slash refer to the animal's left and right sides respectively.

Table 4. Meristic data for the type series of *Smaug swazicus* **sp. nov.** (H = holotype, A = allotype, P = paratype); values on either side of a slash refer to the animal's left and right sides respectively.

Museum number	Type status	Dorsal scale rows longitudinally	Dorsal scale rows transversely	Ventral scale rows longitudinally	Ventral scale rows transversely	Lamellae under 4 th toe	Femoral pores (left+right)	Differentiated femoral scales (left+right)	Gular scales (betwn poster sublinguals)
NMB R9201	H	20	34	14	26	16	10/11	21/25	23
TM 78918	A	21	34	14	28	18	10/10	0/0	25
TM 83000	P	22	34	14	23	16	10/10	10/9	28
TM 83532	P	25	41	14	29	18	11/10	0/0	27
TM 42531	P	22	37	14	25	17	10/10	0/0	28
TM 51376	P	26	36	12	26	18	?/11	?/24	26
TM 78931	P	21	35	14	26	17	12/12	10/16	25
TM 78921	P	23	32	14	25	18	11/11	12/12	24
NMB R9194	P	22	34	12	26	17	11/12	26/35	22
NMB R9195	P	25	35	14	27	17	12/11	33/?	25
NMB R9202	P	22	34	14	27	18	12/11	15/19	24
TM 73290	P	24	37	14	27	19	11/10	0/0	?

Figure 1

Time calibrated phylogram of six concatenated nuclear and mitochondrial genes for the genus *Smaug*.

Bootstrap support values (MP/ML) are shown above branches, and posterior probabilities below branches. (Modified from: Stanley & Bates 2014, fig. 3.)




Figure 2

Differences in colour pattern in the *Smaug warreni* species complex.

From left to right: Dorsal views of *Smaug barbertonensis* (NMB R9196, topotype), *S. cf. barbertonensis* (TM 78918, allotype, see below) and *S. warreni* (TM 63567); ventral views of *S. barbertonensis* (TM 55789), *S. cf. barbertonensis* (TM 78918) and *S. warreni* (TM 78969) (photos: M.F. Bates).



Figure 3

Quadrates variation in the *Smaug warreni* species complex.

(A) *S. warreni* NMB R9292. (B) *S. warreni* AMNH-R-173381. (C) *S. cf. barbertonensis* AMNH-R-173382. (D) *S. cf. barbertonensis* NMB R9201 (holotype, see below). (E) *S. barbertonensis* NMB R9196 (topotype). Diagnostic ridges on the quadrates of *S. cf. barbertonensis* are indicated using arrows.




Figure 4

Morphological variation in the *Smaug warreni* species complex.

A) Transverse rows of dorsal scales. (B) Scatterplot showing head length and head width (corrected by SVL). (C) Principal Component analysis of eight meristic and linear characters. (D) Linear Discriminate analysis of 15 meristic and linear characters.




Figure 5

Smaug swazicus sp. nov. Dorsal, ventral and lateral views of the head of the holotype (NMB R9201).




Figure 6

left Habitat at Malolotja National Park, Eswatini, vicinity of type locality of *Smaug swazicus* sp. nov.

upper right Dorsal colouration of live paratype of *S. swazicus* (NMB R9194). *lower right* Ventral colouration of the same specimen. (Photos: E.L. Stanley)



Figure 7

Cranial and postcranial osteology of *Smaug swazicus* sp. nov. (holotype, NMB R9201).

Body osteoderms are highlighted in blue.




Figure 8

Map showing recorded localities of *Smaug barbertonensis* (red) *S. swazicus* sp. nov. (green) and *S. warreni* (blue).

Filled circles denote museum records, while open circles show geo-referenced photo vouchers from University of Cape Town's Animal Demography Unit Virtual Museum. The type locality for each species is represented by a star.




Figure 9

Smaug barbertonensis. Dorsal, ventral and lateral views of the head of NMB R9191 (topotype).




Figure 10

left *Smaug barbertonensis* habitat, Nelspruit (Extension 5), Mpumalanga Province, South Africa.

upper right Dorsal colouration of *S. barbertonensis* from the latter locality. lower right Ventral colouration of *S. barbertonensis* (same specimen). (Photos: T. Busschau).



Figure 11

Cranial and postcranial osteology of *Smaug barbertonensis* (NMB R9196, topotype).

Body osteoderms are highlighted in blue.




Figure 12

Smaug warreni. Dorsal, ventral and lateral views of the head of TM 50130 (Lomahasha, Eswatini).




Figure 13

left: Smaug warreni habitat, northern Lebombo mountains, Mpumalanga Province, South Africa.

right: Dorsal colouration of S. warreni. (Photos: E.L. Stanley).



Figure 14

Cranial and postcranial osteology of *Smaug warreni* (NMB R9292).

Body ostoderms are highlighted in blue.

