

# Low-cost, low-input RNA-seq protocols perform nearly as well as high-input protocols

Peter A Combs, Michael B Eisen

Recently, a number of protocols extending RNA-sequencing to the single-cell regime have been published. However, we were concerned that the additional steps to deal with such minute quantities of input sample would introduce serious biases that would make analysis of the data using existing approaches invalid. In this study, we performed a critical evaluation of several of these low-volume RNA-seq protocols, and found that they performed slightly less well in metrics of interest to us than a more standard protocol, but with at least two orders of magnitude less sample required. We also explored a simple modification to one of these protocols that, for many samples, reduced the cost of library preparation to approximately \$20/sample.

# Low-cost, low-input RNA-seq protocols perform nearly as well as high-input protocols

Peter A. Combs

Graduate Program in Biophysics,  
University of California, Berkeley

Michael B. Eisen

Department of Molecular and Cell Biology  
and Howard Hughes Medical Institute,  
University of California, Berkeley, CA

January 6, 2015

## Abstract

Recently, a number of protocols extending RNA-sequencing to the single-cell regime have been published. However, we were concerned that the additional steps to deal with such minute quantities of input sample would introduce serious biases that would make analysis of the data using existing approaches invalid. In this study, we performed a careful evaluation of several of these low-volume RNA-seq protocols, and found that they performed slightly less well in metrics of interest to us than a more standard protocol, but with at least two orders of magnitude less sample required. We also explored a simple modification to one of these protocols that, for many samples, reduced the cost of library preparation to approximately \$20/sample.

## 1 Introduction

Second-generation sequencing of RNA (RNA-seq) has proven to be a sensitive and increasingly inexpensive approach for a number of different experiments, including annotating genes in genomes, quantifying gene expression levels in a broad range of sample types, and determining differential expression between samples. As technology improves, transcriptome profiling has been able to be applied to smaller and smaller samples, allowing for more powerful assays to determine transcriptional output. For instance, our lab has used RNA-seq on single *Drosophila* embryos to measure zygotic gene activation [21] and medium-resolution spatial patterning [5]. Further improvements will allow an even broader array of potential experiments on samples that were previously too small.

For instance, over the past few years, a number of groups have published descriptions of protocols to perform RNA-seq on single cells (typically mammalian cells) [29, 27, 28, 12, 16]. A number of studies, both from the original authors of the single-cell RNA-seq protocols and from others, have assessed various aspects of these protocols, both individually and competitively [2, 32, 22]. One particularly powerful use of these approaches is to sequence individual cells in bulk tissues, revealing different states and cellular identities [3, 30].

However, we felt that published descriptions of single-cell and other low-volume protocols did not adequately address whether a change in concentration of a given RNA between two samples would result in a proportional change in the FPKM (or any other measure of transcriptional activity) between those samples. While there are biases inherent to any protocol, we were concerned that direct amplification of the mRNA would select for PCR compatible genes in difficult to predict, and potentially non-linear ways. For many of the published applications of single cell RNA-seq, this is not likely a critical flaw, since the clustering approaches used are moderately robust to quantitative changes. However, to measure spatial and temporal activation of genes across an embryo, it is important that the output is monotonic with respect to concentration, and ideally linear. A linear response allows for more easily interpretable experimental results, without necessarily relying on complicated transformations of the data.

While it is possible to estimate absolute numbers of cellular RNAs from an RNA-seq experiment, doing so requires spike-ins of known concentration and estimates of total cellular RNA content [24, 20]. However, many RNA-seq experiments do not do these controls, nor are such controls strictly necessary under reasonable, though often untested, assumptions of approximately constant RNA content. While ultimately absolute concentrations will be necessary to fully predict properties such as noise tolerance of the regulatory circuits [10, 9], many current modeling efforts rely only on scaled concentration measurements, often derived from *in situ*-hybridization experiments [8, 15, 13]. Given that, we felt it was not important that different protocols should necessarily agree on any particular expression value for a given gene, nor are we fully convinced that absolute expression of any particular gene can truly reliably be predicted in a particular experiment.

In order to convince ourselves that data generated from limiting samples would be suitable for our purposes, we evaluated several protocols for performing RNA-seq on extremely small samples. We also investigated a simple modification to one of the protocols that reduced sample preparation cost per library by more than 2-fold. Finally, we evaluated the effect of read depth on quality of the data. This study provides a single, consistent comparison of these diverse approaches, and shows that in fact all data from the low-volume protocols we examined are usable in similar contexts to the earlier bulk approach.

## 2 Results

### 2.1 Experiment 1: Evaluation of Illumina TruSeq

In our hands, the Illumina TruSeq protocol has performed extremely reliably with samples on the scale of 100ng of total RNA, the manufacturer recommended lower limit of the protocol. However, attempts to create libraries from much smaller samples yielded low complexity libraries, corresponding to as much as 30-fold PCR duplication of fragments. Anecdotally, less than 5% of libraries made with at least 90ng of total RNA yielded abnormally low concentrations, which we observed correlated with low complexity (Data not shown). To determine the lower limit of input needed to reliably produce libraries, we attempted to make libraries from 40, 50, 60, 70, and 80 ng of *Drosophila* total RNA, each in triplicate.

Table 1: Total TruSeq cDNA library yields made with a given amount of input total RNA. Yields measured by Nanodrop of cDNA libraries resuspended in 25 $\mu$ L of EB. The italicized samples were unusually low, and when analyzed with a Bioanalyzer, showed abnormal size distribution of cDNA fragments.

| Amount Input RNA | Replicate A   | Replicate B | Replicate C |
|------------------|---------------|-------------|-------------|
| 40 ng            | <i>57 ng</i>  | 425 ng      | 672 ng      |
| 50 ng            | 435 ng        | 768 ng      | 755 ng      |
| 60 ng            | <i>115 ng</i> | 663 ng      | 668 ng      |
| 70 ng            | 300 ng        | 593 ng      | 653 ng      |
| 80 ng            | 468 ng        | 550 ng      | 840 ng      |

We considered the two libraries with lower than usual concentration to be failures. While a failure rate of approximately 1 in 3 might be acceptable for some purposes, we ultimately wanted to perform RNA sequencing on precious samples, where a failure in any one of a dozen or more libraries would necessitate regenerating all of the libraries. Furthermore, due to the low sample volumes involved (less than approximately 500pg of poly-adenylated mRNA), common laboratory equipment is not able to determine the particular point in the protocol where the failures occurred.

Thus, we consider 70 ng of total RNA to be the conservative lower limit to the protocol. While this is about 30% smaller than the manufacturer suggests, it is still several orders of magnitude larger than we needed it to be. We therefore considered using other small-volume and “single-cell” RNA-seq kits, which we had less experience with and less faith in the data.

## 2.2 Experiment 2: Competitive Comparison of Low-volume RNAseq protocols

We first sought to determine whether the low-volume RNAseq protocols available faithfully recapitulate linear changes in abundance of known inputs. We generated synthetic spike-ins by combining *D. melanogaster* and *D. virilis* total RNA in known, predefined proportions of 0, 5, 10, and 20% *D. virilis* RNA. For each of the low-volume protocols, we used 1ng of total RNA as input, whereas for the TruSeq protocol we used 100ng.

Although pre-defined mixes of spike-in controls have been developed and are commercially available [17], we felt it was important to ensure that a given protocol would function reproducibly with natural RNA, which almost certainly has a different distribution of 6-mers, which could conceivably affect random cDNA priming and other amplification effects. Furthermore, our spike-in sample more densely covers the approximately  $10^5$  fold coverage typical of RNA abundances. It should be noted, however, that our sample is not directly comparable to any other standards, nor is the material of known strandedness. We assumed that the majority of each sample is from standard annotated transcripts, but did not verify this prior to library construction and sequencing.

The different protocols had a variation in yield of libraries from between 6 fmole (approximately 3.6 trillion molecules) and 2,400 femtomoles, with the TruSeq a clear outlier at the high end of the range, and the other protocols all below 200 fmole (Table 2). While the number of PCR cycles in the final enrichment steps can be adjusted, all of these quantities are sufficient to generate hundreds of millions of reads—far more than is typically required for an RNA-seq experiment. We pooled the samples, attempting equimolar fractions in the final pool; however, due to a pooling error, we generated significantly more reads than intended for the TruSeq protocol, and correspondingly fewer in the other protocols. Unless otherwise noted, we therefore sub-sampled the mapped reads to the lowest number of mapped reads in any sample in order to provide a fair comparison between protocols.

We were interested in the fold-change of each *D. virilis* gene across the four samples, rather than the absolute abundance of any particular gene. Therefore, after mapping and gene quantification, we normalized the abundance  $A_{ij}$  of every gene  $i$  across the  $j = 4$  samples by a weighted average of the quantity  $Q_j$  of *D. virilis* in sample  $j$ , as show in equation 1. Thus, within a given gene, a linear fit of  $\hat{A}_{ij}$  vs  $Q_j$  should have a slope of one and an intercept of zero.

$$\hat{A}_{ij} = A_{ij} \div \frac{\sum_j Q_j A_{ij}}{\sum_j (Q_j)^2} \quad (1)$$

We filtered the *D. virilis* genes for those with at least 20 mapped fragments in the sample with 20% *D. virilis*, then calculated an independent linear regression for each of those genes. As expected, for every protocol, the mean slope was 1 ( $t$ -test,  $p < 5 \times 10^{-7}$  for all protocols). Similarly, the average intercepts for all protocols was 0 ( $t$ -test,  $p < 5 \times 10^{-7}$  for all protocols). Also

134 unsurprisingly, the TruSeq protocol had a noticeably higher mean correlation  
 135 coefficient ( $0.98 \pm 0.02$ ) than any of the other protocols ( $0.95 \pm 0.06$ ,  $0.92 \pm 0.09$ ,  
 136 and  $0.95 \pm 0.06$  for Clontech, TotalScript, and SMART-seq2, respectively). The  
 137 mean correlation coefficient was statistically and practically indistinguishable  
 138 between the Clontech samples and the SMART-seq2 samples ( $t$ -test  $p = .11$ ,  
 139 Figure 1).

140 While the TruSeq protocol clearly performed better than the low-volume  
 141 kits, we wondered how well an ideal RNA-seq protocol could perform. We sim-  
 142 ulated an experiment with known levels of *D. virilis* spike in and assuming a  
 143 multinomial distribution of read counts, and repeated the simulation 1,000 times  
 144 to estimate the distribution of relevant quality metrics. Surprisingly, the mean  
 145 correlation coefficient for the TruSeq protocol was higher than the mean corre-  
 146 lation coefficient of every repetition of the simulation, though indistinguishable  
 147 for practical purposes ( $0.984$  vs  $0.982$ ). The slopes were equally well clustered  
 148 around 1, with an interquartile range of  $0.0864$  for the TruSeq protocol com-  
 149 pared to  $0.0843$ , the mean of all simulations; 13% of simulations had a higher  
 150 IQR. We thus conclude that the major limiting factor for the TruSeq protocol  
 151 to generate a linear response in the data is likely the sequencing depth, whereas  
 152 the other protocols all contain additional bias.

153 Indeed, the only major differentiator we could find between the low-volume  
 154 protocols we measured was cost. For only a handful of libraries, the kit-based  
 155 all inclusive model of the Clontech and TotalScript kits could be a significant  
 156 benefit, allowing the purchase of only as much of the reagents as required. By  
 157 contrast, the Smart-seq2 protocol requires the a la carte purchase of a number  
 158 of reagents, some of which are not available or more expensive per unit for  
 159 smaller quantities. Furthermore, there could potentially be a “hang-  
 160 buns” problem, where reagents are sold in non-integer multiples of each other,  
 161 leading to leftovers. Many of these reagents are not single-purpose, however, so  
 162 leftovers could in principle be repurposed in other experiments.

Table 2: Summary of protocols used in experiments 2 and 3. Cost is estimated per sample assuming a large number of libraries at US catalog prices as of May 2014, and includes RNA extraction.

| Protocol                      | Shorthand | Cost per library |
|-------------------------------|-----------|------------------|
| TruSeq                        | TruS      | \$45             |
| Clontech                      | CT        | \$105            |
| TotalScript                   | TotS      | \$115            |
| Smart-seq2, standard protocol | SS        | \$55             |
| Smart-seq2, 2.5 fold dilution | SS—2.5x   | \$28             |
| Smart-seq2, 5 fold dilution   | SS—5x     | \$20             |

| Experiment | Protocol | % D. virilis | Yield (fmole) | Reads      | Mapped     |
|------------|----------|--------------|---------------|------------|------------|
| 2          | CT       | 0            | 6.5           | 3,803,843  | 3,374,520  |
| 2          | "        | 5            | 15.7          | 4,372,738  | 4,164,781  |
| 2          | "        | 10           | 47.4          | 10,013,087 | 9,527,023  |
| 2          | "        | 20           | 17.8          | 4,781,463  | 4,317,101  |
| 2          | TotS     | 0            | 176.8         | 3,281,134  | 2,930,058  |
| 2          | "        | 5            | 170.2         | 2,498,134  | 2,237,330  |
| 2          | "        | 10           | 102.5         | 5,777,523  | 5,424,366  |
| 2          | "        | 20           | 119.9         | 6,068,996  | 5,740,496  |
| 2          | TruS     | 0            | 2,401.0       | 67,560,511 | 64,024,881 |
| 2          | "        | 5            | 2,001.1       | 23,370,854 | 22,589,083 |
| 2          | "        | 10           | 2,174.2       | 39,454,390 | 38,093,763 |
| 2          | "        | 20           | 2,379.2       | 35,265,536 | 34,304,792 |
| 2          | SS2      | 0            | 34.3          | 2,439,518  | 2,297,087  |
| 2          | "        | 5            | 59.6          | 2,550,023  | 2,419,889  |
| 2          | "        | 10           | 67.9          | 2,534,628  | 2,444,568  |
| 2          | "        | 20           | 39.8          | 2,504,340  | 2,389,850  |
| 3          | SS2—2.5x | 0            | 104.4         | 15,769,915 | 14,393,959 |
| 3          | "        | 1            | 124.7         | 21,349,748 | 20,084,131 |
| 3          | "        | 5            | 113.0         | 17,047,120 | 16,329,641 |
| 3          | "        | 10           | 103.5         | 23,762,232 | 22,372,562 |
| 3          | "        | 20           | 123.8         | 20,809,781 | 20,041,548 |
| 3          | SS2—5x   | 0            | 59.4          | 19,214,155 | 17,324,598 |
| 3          | "        | 1            | 58.6          | 23,832,274 | 22,364,220 |
| 3          | "        | 5            | 65.4          | 18,149,452 | 17,157,450 |
| 3          | "        | 10           | 28.8          | 15,821,419 | 14,869,864 |
| 3          | "        | 20           | 57.2          | 22,466,345 | 21,620,603 |

Table 3: Sequencing summary statistics for samples. Protocols are the short-hands used in table 2. Reads indicates the total number of reads, and Mapped the total number of reads that mapped at least once to either genome. Experiments 2 and 3 were run in a single HiSeq lane each.




Figure 1: **Comparison of linearity between different RNA-seq protocols.** A) Normalized levels of gene expression  $\hat{A}$  across samples using the TruSeq protocol, where each line is for a different gene. B-E) Distributions of slopes, intercepts, and correlation coefficient for linear regressions of the abundance of each gene, as in panel A.



Table 4: Distribution of fit parameters. A simple linear fit,  $\hat{A}_{ij} = m \cdot Q_j + b$  was computed for each gene  $i$ , and a correlation coefficient  $r$  calculated. For brevity,  $\bar{x}$  is the mean of some variable  $x$ , and  $\sigma_x$  is its standard deviation.

| Protocol                      | $\bar{m} \pm \sigma_m$ | $b \pm \sigma_b$  | $\bar{r} \pm \sigma_r$ |
|-------------------------------|------------------------|-------------------|------------------------|
| TruSeq                        | 1.01 $\pm$ 0.0698      | -0.108 $\pm$ 1.05 | 0.98 $\pm$ 0.019       |
| Clontech                      | 1.01 $\pm$ 0.12        | -0.217 $\pm$ 1.79 | 0.95 $\pm$ 0.061       |
| TotalScript                   | 0.952 $\pm$ 0.129      | 0.715 $\pm$ 1.93  | 0.93 $\pm$ 0.094       |
| Smart-seq2                    | 1.03 $\pm$ 0.121       | -0.506 $\pm$ 1.82 | 0.95 $\pm$ 0.057       |
| Smart-seq2, 2.5 fold dilution | 0.996 $\pm$ 0.111      | 0.0623 $\pm$ 1.67 | 0.96 $\pm$ 0.053       |
| Smart-seq2, 5 fold dilution   | 1.01 $\pm$ 0.111       | -0.173 $\pm$ 1.66 | 0.96 $\pm$ 0.049       |

### 2.3 Experiment 3: Further modifications to the SMART-seq2 protocol

Although the SMART-seq2 was the cheapest of the protocols when amortized over a large number of samples, we wondered whether it could be performed even more cheaply without compromising data quality. This would enable us to include more biological replicates in the future experiments for which we are evaluating these protocols. In the original protocol, we noticed that roughly 60% of the cost came from the Nextera XT reagents. Thus, reducing the cost of tagmentation was the obvious goal to target.

We made additional libraries, again starting with 1ng of total RNA. We amplified a single set of spike-in samples with 0, 5, 10, and 20% *D. virilis* total RNA as in experiment 2, and made a single an additional sample with 1% *D. virilis* RNA. Starting at the point in the SMART-seq2 protocol where tagmentation was started, we performed reactions in volumes 2.5 $\times$  and 5 $\times$  smaller, using proportionally less cDNA as well. Due to the low total yield, we increased the number of enrichment cycles from 6 to 8 (see methods).

When normalized to the same number of reads as in experiment 2, the protocols with diluted Nextera reagents performed effectively identically: for instance, the mean correlation coefficients were in both cases  $0.96 \pm 0.05$  (Fig. 2 and Table 4). This is despite the additional cycles of enrichment, which improved yield.

Because we used a common set of pre-amplified cDNA samples that was performed in a distinct pre-amplification from experiment 2, we can estimate the contribution of that pre-amplification to the overall variation. If, in fact, the pre-amplification is a major contributor to the variation, then we would expect to find that the correlation between, for instance, the slopes of two runs of the same experiment with different pre-amplifications would be significantly lower than the correlation between the slopes of two runs using the same pre-amplified cDNA pools.

Unsurprisingly, the sets of samples that used the same preamplification were

more correlated with each other than with the set of samples that used a separate pre-amplification (Fig. 3). By analogy to dual-reporter expression studies[7], we term variation along the diagonal “extrinsic noise” ( $\eta_{ext} = \text{std}(m_1 + m_2)$ ), and variation perpendicular to the diagonal “intrinsic noise” ( $\eta_{int} = \text{std}(m_1 - m_2)$ ), being intrinsic to the pre-amplification step. Using that metric, the intrinsic noise is lower for the samples with the same pre-amplification ( $\eta_{int} = 0.09$ ) than for the samples with different pre-amplifications ( $\eta_{int} = 0.16$ ). Somewhat surprisingly, the extrinsic noise is higher for the samples with the same pre-amplification ( $\eta_{ext} = 0.20$  vs  $\eta_{ext} = 0.16$ ), perhaps due to the 2 additional cycles of PCR enrichment.




Figure 2: Distributions of slopes, intercepts, and correlation coefficients for experiment 3. Nextera XT reactions were reduced in volume by the indicated amount.

### 3 Discussion

When sample size is not the limiting factor, it is clear that using well-established protocols that involve minimal sequence-specific manipulation of the sample yields the best results, both in terms of reproducibility and linearity of response. However, if it is not practical to collect such relatively large samples, we believe that any of the “single-cell” protocols we have tested should perform similarly, and can be used as a drop-in replacement. While preamplification steps do introduce some detectable variance, it is not vastly detrimental to the data quality, and does not introduce obvious sequence-specific biases.

Such methods should be strongly preferred if it is feasible to collect a suitably homogenous sample. While bulk tissues may be a mixture of multiple distinct cell types, this may or may not affect the particular research question an RNAseq experiment is designed to answer. In our hands, the lower limit of reliable library construction using the Illumina TruSeq kit is approximately 70ng of total RNA; with non-homogenous samples, the practical limit is likely to be even lower. Although we believe there is significant user-to-user variation, it





Figure 3: **Estimating the source of preamplification noise.** Plotted are the estimated slopes for each gene between experiments. The blue, “Different pre-amplification” compares the 2.5× diluted and full sized reactions, whereas the green “same pre-amplification” points compare the 2.5× and 5× dilution samples, which used the same preamplified cDNA but different tagmentation reactions.

seems unreasonable to expect order-of-magnitude improvements are possible in techniques for precious samples. We suggest that this limit may be related to cDNA binding to tubes or purification beads, but since the quantities are lower than the detection threshold of many standard quality control approaches, we cannot directly verify this, nor do we believe that knowing the precise cause is likely to suggest remediation techniques.

Compared to the regimes these protocols were designed for, we used a relatively large amount of input RNA—1 ng of total RNA—corresponding to approximately 50 nuclei of a mid-blastula transition *Drosophila* embryo. Previous studies have shown that this amount of RNA is well above the level where stochastic variation in the number of mRNAs per cell will strongly affect the measured expression of a vast majority of genes [22]. It is nevertheless a small enough quantity to be experimentally relevant. For instance, we have previously dissected single embryos into approximately 12 sections, yielding approximately 10ng per section[5], and one could conceivably perform similar experiments on imaginal discs or antennal structures, which contain a similar amount of cells [19, 11].

One of the more striking results is that costs can be significantly reduced by simply performing smaller reactions, without noticeably degrading data quality. We do not suspect this will be true for arbitrarily small samples, such as from single cells. Instead, it is likely only true for samples near the high end of the

240 effective range of the protocol. We have not explored where this t breaks  
241 down, and strongly caution others to verify this independently using small pilot  
242 experiments before scaling up.

## 243 4 Methods

### 244 4.1 RNA Extraction, Library Preparation, and Sequenc- 245 ing

246 We performed RNA extraction in TRIzol (Life Technologies, Grand Island, NY)  
247 according to manufacturer instructions, except with a higher concentration of  
248 glycogen as carrier (20 ng) and a higher relative volume of TRIzol to the ex-  
249 pected material (1 mL, as in [21] and [5]). We quantified RNA concentrations  
250 using a fluorometric Qubit RNA HS assay (Life Technologies).

251 TruSeq libraries were prepared with the “TruSeq RNA Sample Preparation  
252 Kit v2” (Illumina Cat.#RS-122-2001) according to manufacturer instructions,  
253 except for the following modifications. All reactions were performed in half  
254 the volume of reagents. We find that this increases the effective concentration  
255 of RNA and cDNA. We performed all reactions and cleanups in 8-tube PCR  
256 strip tubes, which allowed us to reduce the volume of Resuspension Buffer to  
257 minimize volume left behind after each cleanup.

258 Clontech libraries were prepared with the “Low Input Library Prep Kit”  
259 (Clontech Cat.#634947). We generated cDNA by using TruSeq reagents until  
260 the cDNA synthesis step. Then, we used the Low Input Library Prep Kit to  
261 modify the cDNA into sequencing-competent libraries. We believe that a similar  
262 cDNA synthesis could be performed using oligo dT Dynabeads, RNA fragmen-  
263 tation reagents, and Superscript II (Life Technologies), for an approximate cost  
264 per sample of \$15.

265 TotalScript libraries were prepared with the “TotalScript RNA-Seq Kit” and  
266 “TotalScript Index Kit” (Epicentre Cat.#TSRNA1296 and TSIDX12910). We  
267 followed the manufacturer’s instructions, and used the oligo dT priming option.  
268 We performed the mixed priming option in parallel, which yielded approximately  
269 4-fold more library, but did not sequence them due to concerns of ribosomal  
270 contamination.

271 SMARTseq2 libraries were prepared according to the protocol in Picelli *et*  
272 *al.*(2014) [26]. Because we had already extracted and mixed the RNA, we began  
273 at step 5 with 3.7  $\mu$ L of dNTPs and 1  $\mu$ L of 37  $\mu$ M oligo dT primer, yielding the  
274 same concentration of primer and oligo as originally reported. We used 18 cycles  
275 for the preamplification PCR in step 14, added 1ng of cDNA to the Nextera XT  
276 reactions in step 28, and used 6 and 8 cycles for the final enrichment in step 33  
277 (experiments 2 and 3, respectively).

278 Libraries were quantified using a combination of Qubit High Sensitivity  
279 DNA (Life Technologies) and Bioanalyzer (Agilent Technologies, Sunnyvale,  
280 CA) readings, then pooled to equalize index concentration. Due to a pooling  
281 error in experiment 2, the TruSeq libraries were included at much higher abun-

dance. Pooled libraries were then submitted to the Vincent Coates Genome Sequencing Laboratory for 50bp single-end sequencing according to standard protocols for the Illumina HiSeq 2500. Bases were called using HiSeq Control Software v1.8 and Real Time Analysis v2.8.

## 4.2 Mapping and Quantification

Reads were mapped using STAR [6] to a combination of the FlyBase reference genome version 5.54 for *D. melanogaster* and *D. virilis* [23]. We randomly sampled the mapped reads to use an equal number in each sample compared. We used HTSeq (command line options `htseq-count --idattr='gene_name' --stranded=no --sorted=pos`) to count absolute read abundance per gene [1].

## 4.3 Simulation of Experiment 2

We wrote a Python script that simulated Experiment 2 assuming only uncorrelated counting noise in the number of reads per gene. The read counts from the sample with 20% *D. virilis* and the TruSeq protocol was used to generate the base probabilities. *D. virilis* gene probabilities were adjusted downwards, and the remaining probability was assigned evenly to the *D. melanogaster* genes. The SciPy function `stats.multinomial` was used to simulate read counts, assuming an equal number of reads as in the original experiment. Gene expression levels were normalized using equation 1, as in the actual experiment.

## 5 Acknowledgements

## 6 Additional Information and Declarations

### 6.1 Competing Interests

The authors declare no competing interests exist.

### 6.2 Author Contributions

Peter A. Combs conceived and designed the experiments, analyzed the data, and wrote the paper.

Michael B. Eisen conceived and designed the experiments and wrote the paper.

### 6.3 Data Deposition

We have deposited all reads in the NCBI GEO under the accession number GSE64673. The processed data is available at <http://eisenlab.org/lowvolume>. All custom analysis software is available at <https://github.com/eisenlab/SliceSeq>, and is primarily written in Python [31, 4, 14, 18, 25]. Commit `9fc810e7` was used to perform all analyses in this paper.

## 6.4 Funding

PAC is supported by National Institutes of Health training grant #T32 HG00047. MBE is a Howard Hughes Medical Institute Investigator. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

## References

- [1] S. Anders, P. T. Pyl, and W. Huber. HTSeq A Python framework to work with high-throughput sequencing data. Feb 2014.
- [2] Vipul Bhargava, Steven R Head, Phillip Ordoukhanian, Mark Mercola, and Shankar Subramaniam. Technical Variations in Low-Input RNA-seq Methodologies. *Scientific reports*, 4:3678, 2014.
- [3] Yosef Buganim, Dina A Faddah, Albert W Cheng, Elena Itskovich, Styliani Markoulaki, Kibibi Ganz, Sandy L Klemm, Alexander van Oudenaarden, and Rudolf Jaenisch. Single-Cell Expression Analyses during Cellular Reprogramming Reveal an Early Stochastic and a Late Hierarchic Phase. *Cell*, 150(6):1209–1222, sep 2012.
- [4] Peter J A Cock, Tiago Antao, Jeffrey T Chang, Brad A Chapman, Cymon J Cox, Andrew Dalke, Iddo Friedberg, Thomas Hamelryck, Frank Kauff, Bartek Wilczynski, and Michiel J L de Hoon. Biopython: freely available Python tools for computational molecular biology and bioinformatics. *Bioinformatics (Oxford, England)*, 25(11):1422–1423, jun 2009.
- [5] Peter A Combs and Michael B Eisen. Sequencing mRNA from cryo-sliced Drosophila embryos to determine genome-wide spatial patterns of gene expression. *PLoS ONE*, 8(8):e71820, 2013.
- [6] Alexander Dobin, Carrie A Davis, Felix Schlesinger, Jorg Drenkow, Chris Zaleski, Sonali Jha, Philippe Batut, Mark Chaisson, and Thomas R Gingeras. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics (Oxford, England)*, oct 2012.
- [7] Michael B Elowitz, Arnold J Levine, Eric D Siggia, and Peter S Swain. Stochastic gene expression in a single cell. *Science (New York, N.Y.)*, 297(5584):1183–1186, aug 2002.
- [8] Mayra Garcia, Marcos Nahmad, Gregory T Reeves, and Angelike Stathopoulos. Size-dependent regulation of dorsal-ventral patterning in the early Drosophila embryo. *Developmental Biology*, 381(1):286–299, sep 2013.
- [9] Thomas Gregor, William Bialek, Rob R de Ruyter van Steveninck, David W Tank, and Eric F Wieschaus. Diffusion and scaling during early

- embryonic pattern formation. *Proceedings of the National Academy of Sciences of the United States of America*, 102(51):18403–18407, dec 2005.
- [10] Thomas Gregor, David W Tank, Eric F Wieschaus, and William Bialek. Probing the limits to positional information. *Cell*, 130(1):153–164, jul 2007.
- [11] B S Hansson and S Anton. Function and morphology of the antennal lobe: new developments. *Annual review of entomology*, 45:203–231, 2000.
- [12] Tamar Hashimshony, Florian Wagner, Noa Sher, and Itai Yanai. CEL-Seq: Single-Cell RNA-Seq by Multiplexed Linear Amplification. *Cell reports*, aug 2012.
- [13] Xin He, Md Abul Hassan Samee, Charles Blatti, and Saurabh Sinha. Thermodynamics-based models of transcriptional regulation by enhancers: the roles of synergistic activation, cooperative binding and short-range repression. *PLoS Computational Biology*, 6(9), 2010.
- [14] John D Hunter. Matplotlib: A 2D graphics environment. *Computing In Science & Engineering*, 9(3):90–95, May-Jun 2007.
- [15] Garth R Ilesley, Jasmin Fisher, Rolf Apweiler, Angela H DePace, and Nicholas M Luscombe. Cellular resolution models for even skipped regulation in the entire Drosophila embryo. *eLife*, 2:e00522, 2013.
- [16] Saiful Islam, Una Kjällquist, Annalena Moliner, Pawel Zajac, Jian-Bing Fan, Peter Lönnerberg, and Sten Linnarsson. Characterization of the single-cell transcriptional landscape by highly multiplex RNA-seq. *Genome Research*, 21(7):1160–1167, jul 2011.
- [17] L. Jiang, F. Schlesinger, C. A. Davis, Y. Zhang, R. Li, M. Salit, T. R. Gingeras, and B. Oliver. Synthetic spike-in standards for RNA-seq experiments. *Genome Research*, 21(9):1543–1551, Sep 2011.
- [18] Eric Jones, Travis Oliphant, Pearu Peterson, and others. *SciPy: Open source scientific tools for Python*, 2001.
- [19] Ansgar Klebes, Brian Biehs, Francisco Cifuentes, and Thomas B Kornberg. Expression profiling of Drosophila imaginal discs. *Genome Biology*, 3(8):RESEARCH0038, jul 2002.
- [20] CharlesY. Lin, Jakob Lovén, PeterB. Rahl, RonaldM. Paranal, ChristopherB. Burge, JamesE. Bradner, TongIhn Lee, and RichardA. Young. Transcriptional Amplification in Tumor Cells with Elevated c-Myc. *Cell*, 151(1):56–67, Sep 2012.
- [21] Susan E Lott, Jacqueline E Villalta, Gary P Schroth, Shujun Luo, Leath A Tonkin, and Michael B Eisen. Noncanonical compensation of zygotic X transcription in early Drosophila melanogaster development revealed through single-embryo RNA-seq. *PLoS Biology*, 9(2):e1000590, 2011.

- 391 [22] Georgi K Marinov, Brian A Williams, Kenneth McCue, Gary P Schroth,  
392 Jason Gertz, Richard M Myers, and Barbara J Wold. From single-cell to  
393 cell-pool transcriptomes: stochasticity in gene expression and RNA splic-  
394 ing. *Genome Research*, 24(3):496–510, dec 2013.
- 395 [23] Peter McQuilton, Susan E St Pierre, Jim Thurmond, and FlyBase Con-  
396 sortium. FlyBase 101—the basics of navigating FlyBase. *Nucleic Acids*  
397 *Research*, 40(Database issue):D706–14, jan 2012.
- 398 [24] Ali Mortazavi, Brian A Williams, Kenneth McCue, Lorian Schaeffer, and  
399 Barbara Wold. Mapping and quantifying mammalian transcriptomes by  
400 RNA-Seq. *Nature Methods*, 5(7):621–628, jul 2008.
- 401 [25] F Perez and B E Granger. IPython: a system for interactive scientific  
402 computing. *Computing In Science & Engineering*, 2007.
- 403 [26] Simone Picelli, Omid R Faridani, Asa K Björklund, Gösta Winberg, Sven  
404 Sagasser, and Rickard Sandberg. Full-length RNA-seq from single cells  
405 using Smart-seq2. *Nature Protocols*, 9(1):171–181, jan 2014.
- 406 [27] Daniel Ramsköld, Shujun Luo, Yu-Chieh Wang, Robin Li, Qiaolin Deng,  
407 Omid R Faridani, Gregory A Daniels, Irina Khrebtukova, Jeanne F Loring,  
408 Louise C Laurent, Gary P Schroth, and Rickard Sandberg. Full-length  
409 mRNA-Seq from single-cell levels of RNA and individual circulating tumor  
410 cells. *Nature Biotechnology*, 30(8):777–782, aug 2012.
- 411 [28] Yohei Sasagawa, Itoshi Nikaido, Tetsutaro Hayashi, Hiroki Danno,  
412 Kenichiro D Uno, Takeshi Imai, and Hiroki R Ueda. Quartz-Seq: a highly  
413 reproducible and sensitive single-cell RNA sequencing method, reveals non-  
414 genetic gene-expression heterogeneity. *Genome Biology*, 14(4):R31, apr  
415 2013.
- 416 [29] Fuchou Tang, Catalin Barbacioru, Yangzhou Wang, Ellen Nordman,  
417 Clarence Lee, Nanlan Xu, Xiaohui Wang, John Bodeau, Brian B Tuch,  
418 Asim Siddiqui, Kaiqin Lao, and M Azim Surani. mRNA-Seq whole-  
419 transcriptome analysis of a single cell. *Nature Methods*, 6(5):377–382, may  
420 2009.
- 421 [30] Barbara Treutlein, Doug G Brownfield, Angela R Wu, Norma F Neff,  
422 Gary L Mantalas, F Hernan Espinoza, Tushar J Desai, Mark A Kras-  
423 now, and Stephen R Quake. Reconstructing lineage hierarchies of the dis-  
424 tal lung epithelium using single-cell RNA-seq. *Nature*, 509(7500):371–375,  
425 may 2014.
- 426 [31] G Van Rossum and F L Drake. *Python language reference manual*, 2003.
- 427 [32] Angela R Wu, Norma F Neff, Tomer Kalisky, Piero Dalerba, Barbara Treut-  
428 lein, Michael E Rothenberg, Francis M Mburu, Gary L Mantalas, Sopheak  
429 Sim, Michael F Clarke, and Stephen R Quake. Quantitative assessment



430 of single-cell RNA-sequencing methods. *Nature Methods*, 11(1):41–46, jan  
431 2014.