

Amendments to saline-sodic soils showed long-term effects on improving growth and yield of rice (*Oryza sativa* L.)

Dandan Zhao^{1,2}, Zhichun Wang^{Corresp., 1}, Fan Yang^{Corresp., 1}, Wendong Zhu^{1,2}, Fenghua An¹, Hongyuan Ma¹, Tibor Toth³, Xu Liao¹, Hongtao Yang¹, Lu Zhang¹

¹ Northeast Institute of Geography and Agroecology, Chinese Academy of Sciences (CAS), Changchun, China

² University of Chinese Academy of Sciences, Beijing, China

³ Research Institute for Soil Science and Agricultural Chemistry of the Hungarian Academy of Sciences, Budapest, Hungary

Corresponding Authors: Zhichun Wang, Fan Yang

Email address: wangzhichun@iga.ac.cn, yangfan@iga.ac.cn

Background. Saline-sodic soils are widely distributed in arid and semi-arid regions around the world. High levels of salt and sodium inhibit the growth and development of crops. However, there has been limited reports on both osmotic potential in soil solutions (OP_{ss}) and characteristics of Na^+ and K^+ absorption in rice in saline-sodic soils under various amendments application. **Methods.** A field experiment was conducted between 2009 and 2017 to analyze the influence of amendments addition to saline-sodic soils on rice growth and yield. Rice was grown in the soil with no amendment (CK), with desulfurization gypsum (DG), with sandy soil (SS), with farmyard manure (FM) and with the mixture of above amendments (M). The osmotic potential in soil solution, selective absorption of K^+ over Na^+ (SA), selective transport of K^+ over Na^+ (ST), the distribution of K^+ and Na^+ , and yield components in rice plants were investigated. **Results.** The results indicated that amendments application have positive effects on rice yield. The M treatment was the best among the tested amendments with the highest rice grain yield. M treatment increased the OP_{ss} values significantly to relieve the inhibition of the water uptake by plants.

Additionally, the M treatment significantly enhanced K^+ concentration and impeded Na^+ accumulation in shoots. SA values were reduced while ST values were increased for all amendments. In conclusion, a mixture of desulfurization gypsum, sandy soil and farmyard manure was the best treatment for the improvement of rice growth and yield in the Songnen Plain, northeast China.

Amendments to saline-sodic soils showed long-term effects on improving growth and yield of rice (*Oryza sativa* L.)

Dandan Zhao^{1,2}, Zhichun Wang^{1*}, Fan Yang^{1*}, Wendong Zhu^{1,2}, Fenghua An¹, Tibor Toth³, Xu Liao¹, Hongtao Yang¹, Lu Zhang¹, Hongyuan Ma¹

¹ Northeast Institute of Geography and Agroecology, Chinese Academy of Sciences (CAS), China, Changchun 130102, (China)

² University of Chinese Academy of Sciences, Beijing 100049, (China)

³ Research Institute for Soil Science and Agricultural Chemistry of the Hungarian Academy of Sciences, Herman Otto 15, H-1525 Budapest, (Hungary)

Corresponding Author:

Zhichun Wang, Fan Yang

Shengbei Street No. 4888, Changchun city, Jilin Province, 130102, China

Email address: wangzhichun@iga.ac.cn

Abstract

Background. Saline-sodic soils are widely distributed in arid and semi-arid regions around the world. High levels of salt and sodium inhibit the growth and development of crops. However, there has been limited reports on both osmotic potential in soil solutions (OP_{ss}) and characteristics of Na^+ and K^+ absorption in rice in saline-sodic soils under various amendments application.

Methods. A field experiment was conducted between 2009 and 2017 to analyze the influence of amendments addition to saline-sodic soils on rice growth and yield. Rice was grown in the soil with no amendment (CK), with desulfurization gypsum (DG), with sandy soil (SS), with farmyard manure (FM) and with the mixture of above amendments (M). The osmotic potential in soil solution, selective absorption of K^+ over Na^+ (SA), selective transport of K^+ over Na^+ (ST), the distribution of K^+ and Na^+ , and yield components in rice plants were investigated.

Results. The results indicated that amendments application have positive effects on rice yield. The M treatment was the best among the tested amendments with the highest rice grain yield. M treatment increased the OP_{ss} values significantly to relieve the inhibition of the water uptake by plants. Additionally, the M treatment significantly enhanced K^+ concentration and impeded Na^+ accumulation in shoots. SA values were reduced while ST values were increased for all amendments. In conclusion, a mixture of desulfurization gypsum, sandy soil and farmyard manure was the best treatment for the improvement of rice growth and yield in the Songnen Plain, northeast China.

Introduction

Soil salinity-sodicity is one of the main impediments for crop productivity and sustainability in arid and semiarid areas (Suarez, 2001; Qadir et al., 2006). Saline-sodic soils comprise approximately 3.67×10^7 ha, and Songnen Plain is one of the major saline-sodic areas in China (Yao et al., 2008; Yang et al., 2016). pH stress and Na^+ toxicity are the main causes of the degradation in saline-sodic soils (Gharaibeh et al., 2010). Efforts have been made to ameliorate saline-sodic soils including desulfurization gypsum, farmyard manure, sand, hydraulic engineering and phytoremediation (Qadir et al., 2007; Wang et al., 2012; Ahmad et al., 2013).

Desulfurization gypsum provides a sources of Ca^{2+} to replace exchangeable Na^+ , thereby improving the physical condition of the soil and increasing water infiltration (Oster, 1982; Wang et al., 2013; Wang & Yang, 2018). Manure application improves soil structure and alleviates soil sodicity (Yu et al., 2010). Sanding to saline-sodic soils changes soil compactness and reduces salt content (Wang et al., 2010). These amendments showed various improvements of saline-sodic soil properties in practice.

Crops respond to salinity and sodicity in two phases: (1) a continuous osmotic phase that occurs when the potential energy of the saline-sodic soil solution is lowered by its osmotic pressure, thus inhibiting the water uptake of plants; and (2) a slower ionic phase due to ion toxicity or ion imbalance as plants accumulate salt ions over a period of time (Munns & Tester, 2008). Most amendment studies focused on soil physiochemical properties (Chi et al., 2012; Zhao et al., 2018) rather than on the osmotic potential in the soil solution and the selective absorption of ions by plants, although they have important effects on crop

biomass (Wang et al., 2009).

Rice showed moderate sensitivity to salinity and sodicity (Maas & Hoffman, 1977). Kelly & Rengasamy (2006) showed that osmotic stress is one of the major factors in reducing crop yield. The decreasing the osmotic potential of the soil solution was inhibitory to the water uptake of plant roots (Duarte & De Souza, 2016). The survival of rice plants under saline-sodic conditions is correlated with Na^+ and K^+ accumulations in plant tissues (Song & Fujiyama, 1996). Yamanouchi et al. (1987) found that Na^+ concentrations in shoots are inversely correlated with the relative plant growth and yield. The susceptibility of rice plants to salinity and sodicity stress is due to the limited ability to restrict Na^+ transportation to shoots (Matsushita & Matoh, 1991). This Na^+ restricts K^+ uptake and K^+ is an essential macronutrient for the growth of plants and cannot be substituted by Na^+ (Bhandal & Malik, 1988). The ability of plants to keep a high cytoplasmic K^+/Na^+ ratio is one of the most important mechanisms of salt tolerance (Maathuis & Amtmann, 1999).

In this study, we measured the osmotic potential in the soil solution, characterized K^+ and Na^+ absorption of rice, K^+ and Na^+ concentrations in shoots and roots, selective absorption/transport for K^+ over Na^+ , distribution of K^+ , Na^+ in rice organs, and yield of rice under various soil treatments, including chemical treatment (desulfurization gypsum, DG), physical treatment (sandy soil, SS) and organic treatment (farm manure, FM) as well as mixed treatment (M) in saline-sodic soil for planting rice in field. We hypothesized that (1) amendments would increase the osmotic potential in soil solutions; (2) amendments would alter the ion selective absorption and selective transport in saline-sodic soils and (3) the grain yield of rice would be highest by M application according to the synergy among treatments when they applied together in the Songnen Plain, northeast China.

Materials & Methods

Location description

The study was conducted from 2009 to 2017 at Da'an Sodic Land Experiment Station (45°35'58"-45°36'28"N, 123°50'27"-123°51'31"E, 132.1), operated by the Chinese Academy of Sciences. The climate of this region is semi-humid to semi-arid continental monsoon. The annual mean air temperature is 4.7°C, and the mean annual precipitation of this area is approximately 400-500 mm, and 80% or more of the

precipitation occurs between May and September.

The soil at this study site is classified as clay loam ~~texture~~ with montmorillonite as a dominant mineral. The soil prior to the start of the experiment represents a typically severe saline-sodic soil with pH (1: 5 H₂O) of 10.47, electrical conductivity (EC) (1: 5 H₂O) of 2.36 mS cm⁻¹, soil organic C (SOC) of 2.80 g kg⁻¹ and exchangeable sodium percentage at 79.7% in the top 20-cm soil layer, which is considered to be the effective rooting zone. The main soluble cation was Na⁺, while the anions were HCO₃⁻ and CO₃²⁻. Based on the World Reference Base for Soil Resources (IUSS Working Group WRB, 2007), the main soil type was classified as solonetz.

Field design and treatments

The experiment was arranged in a random block design with three replicates of 20 m² for each plot. There were five treatments: (1) CK, without amendment application; (2) DG, amended with desulfurization gypsum (containing 93% CaSO₄·2H₂O) at 3 kg m⁻²; (3) SS, amended with sandy soil at 6 kg m⁻²; (4) FM, amended with 6 kg m⁻² farmyard manure (5) M, amended with the mixture of desulfurization gypsum, sandy soil and farmyard manure, the amounts of which are equal to those in the DG, SS and FM treatments. Some essential properties of the amendments used in the present study are presented in Table 1 (Luo et al., 2018). Plastic cloth buried between plots to a depth of 1 m soil separated plots to prevent disturbance of lateral movement of amendments, water and salt.

The soil amendments were only applied once before the start of this experiment in the late autumn, 2009. The soil amendments were mixed with the 0-20 cm soil layer by rotary cultivator and then irrigation was carried out after 24 h. The CK was also treated by the same method except for the amendment. Agronomic and fertilizer management practices for rice cultivation were the same in all plots and were in accordance with the prevalent system of agriculture in this area. Chemical fertilizers were broadcast over the soil annually at rates of 207 kg N ha⁻¹ (as urea containing 46% N), 78 kg P ha⁻¹ (as calcium super phosphate containing 12% P₂O₅) and 60 kg K ha⁻¹ (as potassium sulfate containing 45% K₂O). The soil was then plowed to mix the fertilizers into the subsoil.

The local rice cultivar (G19) was planted after wet plowing and sinking between May 20 to the end of May every year for the experiment. Rice seed was sown on normal soil in a greenhouse in early April for

nursing, and the 40-day seedlings were transplanted into the plots with a fixed planting spacing of 30×16.7 cm. Planting space of 30×16.7 cm is a common practice to avoid lodging, and cultivation of 3-5 seedlings per hill is recommended in saline-sodic soil in the Songnen plain (Wang et al, 2010). The depth of 3-7 cm standing water was maintained in the paddy through flood irrigation and runoff drainage during the growth stages of rice. The soils were all drained in the middle of September for harvest.

Measurements

K^+ and Na^+ concentrations in rice plant were measured by sampling three hills excluding the border hills from each plot on 20 days before harvesting. The selected rice hills were observed to be representative of the plot. The rice plants were separated into roots, leaves, sheaths and panicles. The roots were thoroughly washed with water to remove the soil particles. Clean roots were used for estimating Na^+ and K^+ concentrations. Plant samples were dried for 48h at $80^\circ C$ in an air-forced oven. Dried materials were finely grounded using a ball mill. They were then digested using an acid mixture [sulphuric acid: perchloric acid (H_2SO_4 : $HClO_4$ =4:1)] (Mori et al., 2011). K^+ and Na^+ concentrations were determined using an atomic absorption spectrometer (GGX-900). K^+ and Na^+ concentrations in the shoot were calculated from K^+ and Na^+ concentrations and dry weights of grains, leaves and sheaths, K^+ and Na^+ concentrations in the whole plant were calculated from K^+ and Na^+ concentrations and dry weights of grains, leaves, sheaths and roots.

At harvest in October, the following growth and yield data were determined in 2010, 2012, 2015 and 2017: plant height, panicle length, number of grains per panicle, 1000-grain weight and grain yield (Zeng et al., 2000).

To analyze the soil properties as affected by different amendments, soil sampling was performed after harvest of the rice in the November, 2017. All soil samples, obtained from each plot at six depths of 0-10 cm, 10-20 cm, 20-40 cm, 40-60 cm, 60-80 cm and 80-100 cm, . were dried at $105^\circ C$ for 24 h and passed through a 2-mm diameter sieve. Soil samples were analyzed for electric conductivity (EC in $dS\ m^{-1}$), soluble K^+ , Na^+ and Ca^{2+} using 1: 5 soil to water extracts as described by Sumner (1993).

The EC of 1: 5 soil to water extracts ($EC_{1:5}$) was determined by DDS-307 conductivity meter (Shanghai Precision Scientific Instrument Co., Ltd., China), the concentrations in $mmol/L$ of K^+ , Na^+ were determined using flame photometry (FP-6410) and the concentration of Ca^{2+} was measured by EDTA titration (Jackson,

140 1956).

141 The osmotic potential can serve as a good index for evaluating plant response to saline-sodic stress
142 (De Souza et al., 2012). In this experiment, we regard the 1: 5 soil to water extracts as soil solution, and the
143 osmotic potential in the soil solution (OP_{ss}) was calculated as follows:

$$144 \quad OP_{ss}=(-0.36)\times 10EC \text{ (Bohn et al., 2002)}$$

145 **SA and ST Calculation**

146 Selective absorption of K⁺ over Na⁺ (SA) represents the net capacity of a plant to absorb K⁺ relative
147 to Na⁺ from the shallow soil (0-40 cm); Selective transport of K⁺ over Na⁺ (ST) reflects the net capacity of
148 a plant to favour transport of K⁺ over Na⁺ from the root to shoot (Wang et al., 2004). In this study, SA and
149 ST values were calculated according to the following formula (Wang et al., 2002; Wang et al., 2004) using
150 data obtained from the experiments described earlier:

$$151 \quad SA= (K/Na \text{ in root dry weight})/ (\text{soil } K/Na \text{ at } 0\text{-}40 \text{ cm depth})$$

$$152 \quad ST= (K/Na \text{ in shoot dry weight})/ (K/Na \text{ in root dry weight})$$

153 **Statistical analysis**

154 Data were statistically evaluated by analysis of variance (ANOVA). The differences between mean
155 values were evaluated by the Duncan test. All analyses were done with the SPSS 20.0 software (New York,
156 USA). A *P*=0.05 value was used to designate statistically significant results.

157 **Results**

158 **Effect of amendments application on osmotic potential in soil solution**

159 The osmotic potential in the soil solution (OP_{ss}) was increased by amendments application compared
160 to the control. The amplitude of variation of OP_{ss} was from -4.39 bars in the 80-100 cm soil layer under CK
161 treatment to -1.04 bars in the 10-20 cm soil layer under M treatment. In the 0-40 cm soil layer, amendments
162 application generally increased the OP_{ss} values in the following order: M>DG>SS>FM>CK (Figure 1). In
163 the 0-10 cm soil layer, the M, DG, SS and FM treatment increased the OP_{ss} by 53.8%, 40.1%, 29.1% and
164 12.2% compared to the CK treatment, respectively. In the same soil layer, the highest OP_{ss} was observed
165 for M, which means that the ability to reduce the salt concentration of soil solution is strongest, followed
166 by DG.

Effect of amendments application on Na⁺ and K⁺ concentrations in rice shoots and roots

The Na⁺ concentration in shoots of rice plants varied with different amendments applied in the saline-sodic soil (Figure 2A). Rice shoots of plants in M treatment showed the lowest Na⁺ concentration of 0.91 mg/g dry weight, and the Na⁺ concentrations in FM, CK, SS and DG treatments were 4.4%, 7.7%, 8.8% and 11.0% higher than that in M treatment, respectively. The difference in Na⁺ concentration between DG and M treatments was significant. The mean root Na⁺ concentration was highest in the CK treatment, and 0.8%, 7.1%, 9.2% and 15.1% lower in M, SS, FM and DG treatments, respectively. However, the differences on Na⁺ concentration in rice root among amendment treatment and CK were non-significant. Among amendments treatment and CK were non-significant (Figure 2B). Amendments application significantly enhanced K⁺ concentration in rice shoots compared to the control treatment, with the highest K⁺ concentration found for DG (Figure 2C). The K⁺ concentration in rice roots with M, SS, FM treatments were lower than that with the control treatment. The lowest K⁺ concentration was observed for FM, which was 16.8% lower than that of CK (Figure 2D).

Selective absorption and transport of K⁺ over Na⁺ in rice plant

Compared to the CK treatment, the M treatment significantly decreased the SA value of the rice by 74.8% (Figure 3A). However, the M treatment significantly increased the ST value of the rice compared to the CK treatment, which was 1.5 times more than the ST value of the CK treatment (Figure 3B). Amendment application hindered the uptake of K⁺ over Na⁺ from soil to root (SA) compared with CK (Figure 3A), which is probably a consequence of rice physiological adjustment. Amendment application enhanced the uptake of K⁺ over Na⁺ from root to shoot (ST) compared with CK (Figure 3B). This was attributed to strong selective transport of K⁺ over Na⁺ under amendment application.

The mean Na⁺ concentration in the soil extracts decreased from a maximum (6.68 mmol/L) in the CK treatment to 3.16, 4.35, 5.11 and 6.60 mmol/L with M, DG, SS and FM treatments, respectively. The differences among M, DG and CK were significant (Figure 4A). Amendment application slightly enhanced the K⁺ concentration in the soil extracts compared to CK (Figure 4B). The Ca²⁺ concentration in the soil extracts were higher for treatments with amendments than the one without, and differences among the 4

different amendments were not significant (Figure 4C).

Characteristics of distribution of Na⁺ and K⁺ in rice with different amendments application

There were little differences in Na⁺ concentration in the whole rice plants among different treatments in the saline-sodic soils in this experiment (Figure 5A). K⁺ concentration in the whole plant was significantly enhanced after amendment application, but the differences between the 4 treatments with amendments were non-significant (Figure 5B). Na⁺ absorbed by the whole plant was almost the same with and without amendments, which was different from the observations on rice organs (Figure 5A and Table 2). The Na⁺ concentrations in rice roots and grains both decreased when applying amendments in the saline-sodic soils; which were contrary to the rise of K⁺ concentrations in sheaths and leaves (Table 2). Compared to the control treatment, DG, SS, FM and M treatments increased the K⁺ concentrations in rice sheaths by 57.2%, 54.9%, 44.1% and 25.5%, respectively.

For the distribution of ions in rice organs, there was a higher proportion of the total K⁺ in leaves. More Na⁺ was found in roots. The order of accumulation of Na⁺ in various organs was roots>leaves>sheaths>grains (Table 2). The order is imposed by the fact that the root system retains more Na⁺ and prevents Na⁺ from being transported to the aboveground organs in saline-sodic soils, resulting in higher K⁺ proportion in leaves, sheaths and grains. This was also illustrated as being beneficial to normal metabolic activity (Borsani, 2001; Ahmad & Jabeen, 2005).

Relationship between OP_{ss}, selective absorption and yield of rice

The grain yield of the four years of 2010, 2012, 2015 and 2017 were taken as representative of the trend of rice yield from 2009 to 2017 (Figure 6). In terms of grain yield, the M was the best treatment, next is the DG treatment. The grain yield of rice with amendments application were significantly higher than without amendments application except in 2015.

Amendment treatments significantly enhanced the grain yield of rice compared to the control in 2017 (Table 3). The differences, however, among different amendments were not significant at $P<0.05$. Soil amendment application generally increased the 1000-grain weight in the following order: FM>M>DG>SS>CK (Table 3). Additionally, the FM and M treatments significantly increased the 1000-

grain weight to 1.16 and 1.13 times more than the CK treatment, respectively (Table 3). Compared to the CK treatment, the SS treatment considerably enhanced the number of grains per panicle (Table 3). There was no significant difference on rice height and panicle length between various treatments (Table 3).

Significant positive correlations were found between OP_{ss} in the 0-20 cm soil layer and the 1000-grain weight ($R^2=0.992$, $P<0.001$, Table 4). Significant negative correlations were found between SA and rice grain yield ($R^2=0.925$, $P<0.05$, Table 4) and between SA and 1000-grain weight of rice ($R^2=0.884$, $P=0.047$, Table 4). There was no significant correlation between either SA or ST and other growth parameters and yield of rice (Table 4).

Discussion

Characteristics of Na^+ and K^+ absorption in rice

Compared with the CK treatment, the selectivity absorption of K^+ over Na^+ (SA) decreased significantly with the M application in this study. When the M applied, osmotic stress and Na^+ toxicity were significantly decreased leading to better plant growth in saline-sodic soils (Swarup, 1982; Yuncai et al., 2005; Luo et al., 2018; Shi et al., 2019). Similar to our results, previous studies have shown that plants accumulate excessive Na^+ in their shoots under stress caused by high salinity-sodicity (Roy & Mishra, 2014), and Na^+ concentration in shoots increased significantly with a surge in soil salinity-sodicity (Syed & Abdur, 2017).

Adding amendments reduces the salinity-sodicity stress of plants growing in the amended soil (Chaganti & Crohn, 2015). Therefore, the rice planted in the CK plot was under a higher external salinity-sodicity stress. As a result, the SA value of rice plants with CK was higher than those with amendments application and maintained a high cytosolic K^+/Na^+ ratio. This is thought to be one of the most important mechanisms of salt tolerance exhibited by plants (Gorham, 1990; Dubcovsky et al., 1996; Munns & James, 2008; Munns et al., 2010).

Effects of Ca^{2+} on SA and ST values

The competition of K^+ and Na^+ entry into plants can result in significant adverse effects on plants' growth, where concentrations of Na^+ often exceed those of K^+ (Tester & Davenport, 2003). Therefore, the maintenance of a high K^+/Na^+ ratios in plants is essential (Maathuis & Amtmann, 1999). Amendments of

Ca²⁺ promoted K⁺ over Na⁺ absorption, resulting in the enhancement of selectivity absorption of K⁺ over Na⁺ (Alama et al., 2002). Ca²⁺ can replace Na⁺ in plants which restores cell wall stability and plasma membrane integrity (Zhang et al., 2010; Wu & Wang, 2012). Although alleviation of Na⁺ toxicity by supplemental Ca²⁺ was confirmed, the responses varied with different plant species. Under similar saline-sodic conditions, amendments of Ca²⁺ were found to obviously increased K⁺/Na⁺ selectivity of both roots and shoots (SA and ST values) in *Medicago sativa* (Al-Khateeb, 2006) and *Cornus sericea* (Renault & Affifi, 2009). This is contrast to Wang et al. (2007) who reported that amendments of Ca²⁺ had no influence on SA and ST values of *Suaeda maritima*. In addition, the responses of Ca²⁺ to Na⁺ also varied with osmotic potential in soil solution in the same plant species. In rice, Ca²⁺ did not have significant effects on selective absorption and selective transport of K⁺ over Na⁺ of plants when subjected to low osmotic potential in soil solution (Yeo & Flowers, 2010). This is consistent with the results obtained in the present study: there were not significant differences among CK, DG, SS and FM treatment on SA and ST values (Figure 3). In contrast, M application significantly decreased roots Na⁺ absorption and increased shoots K⁺ accumulation in rice. It is proposed that the presence of Ca²⁺ could enhance K⁺/Na⁺ selectivity and regulate ion homeostasis in rice under low saline-sidicity condition.

In rice, a minority of the ions reaching the shoots of plant are the consequence of leakage along the transpirational bypass flow to the xylem and Ca²⁺ application can reduce the bypass flow of rice (Faiyue et al., 2010; Anil et al., 2005). This reduction in the bypass flow is positively related with the concomitant reduction in the shoot Na⁺ uptake (Anil et al. 2005). In addition, a majority of the ions reaching the shoots of rice should be transported via the symplast pathway. Therefore, Ca²⁺ plays important role in regulating apoplast and symplast pathways involved in Na⁺ transport.

Yield of rice

Transient salinity affects the plants' absorption of available water, which results in a reduction in plant yield (Rengasamy, 2010a, 2010b). However, application of amendments to saline-sodic soils can alleviate the salinity-sodicity stress on plants (Irshad et al., 2002). Amendments application in our study enhanced the OP_{ss} values, and then decreased osmotic pressure of the soil solution. This ultimately increased the plant growth and yield of rice in the saline-sodic soils. Applying a small amount of calcium thus was shown to

enhance the plants' salt tolerance (Cramer, 1992).

The grain yield of rice was higher with amendment application compared to that with CK treatment. There were significant negative correlations between SA and rice grain yield ($R^2=0.925$, $P<0.05$, Table 4), a possible explanation for this result is that the low SA values of plant was partly ascribed to its relative sensitivity to root which increased the Na^+ accumulation in root and therefore decreased the SA values (Leland & Eugene, 1999; Ren et al., 2005; Tsialtas & Maslari, 2009).

DG, SS and FM application are known to improve the root environment and increase rice yield (Abrishamkesh et al., 2015). In this study, we found that the mixture of DG, SS and FM application reduced the absorption of Na^+ in rice shoots ~~significantly~~ and led to the highest rice grain yields, which may be due to the synergistic effect of these three amendments. However, the ~~each amendment~~ contribution to the rice yield needs to be quantified in future studies.

Conclusions

In this field experiment, the amendments application significantly increased the yield of rice. In particular, the M treatment was the best among the tested amendment treatments with the highest rice grain yield in the saline-sodic soils although the differences between amendments treatment were not significant. Relative to the CK treatment, the FM and M treatments significantly enhanced the 1000-grain weight and the SS treatment significantly improved the number of grains per panicle. The amendments treatment increased the OP_{ss} significantly, thus relieving the inhibition of water uptake by plants. In addition, a positive effect of amendments application on reducing Na^+ accumulation and increasing the uptake of K^+ of rice shoot was observed. Amendments application increased ST values and decreased SA values. Moreover, there existed an ion regionalization distribution in rice plant; there was a higher K^+ proportion in leaves and a higher Na^+ proportion in roots. Collectively, the mixture of desulfurization gypsum, sandy soil and farmyard manure provided excellent results for increasing the yield of rice in the saline-sodic soils in the Songnen Plain, northeast China.

Acknowledgements

We thank Da'an Sodic Land Experiment Station of China for providing experimental plot.

References

- 302 **Abrishamkesh S, Gorji M, Asadi H. 2015.** Effects of rice husk biochar application on the properties of alkaline soil
303 and lentil growth. *Plant, Soil and Environment* **61**: 475–482.
- 304 **Ahmad R, Jabeen R. 2005.** Foliar spray of mineral elements antagonistic to sodium a technique to induce salt
305 tolerance in plants growing under saline conditions. *Pakistan Journal of Botany* **37**: 913–920.
- 306 **Ahmad S, Ghafoor A, Akhtar ME, Khan MZ. 2013.** Ionic displacement and reclamation of saline-sodic soils using
307 chemical amendments and crop rotation. *Land Degradation and Development* **24**: 170–178.
- 308 **Alama S, Huqb SMI, Kawai S, Islam A. 2002.** Effects of applying calcium salts to coastal saline soils on growth
309 and mineral nutrition of rice varieties. *Journal of Plant Nutrition* **25**: 561–575.
- 310 **Al-khateeb SA. 2006.** Effect of calcium/sodium ratio on growth and ion relations of alfalfa (*Medicago sativa* L.)
311 seedling s growth under saline condition. *Journal of Agronomy* **5**: 175–181.
- 312 **Anil VS, Kirishnamurthy P, Kuruvilla S, Suchartha K, Thomas G, Mathew MK. 2005.** Regulation of the uptake
313 and distribution of Na⁺ in shoots of rice (*Oryza sativa*) variety Pokkali: role of Ca²⁺ in salt tolerance response.
314 *Physiologia Plantarum* **124**: 451–464.
- 315 **Bhandal IS, Malik CP. 1988.** Potassium estimation, uptake, and its role in the physiology and metabolism of
316 flowering plants. *International Review of Cytology* **110**: 205–254.
- 317 **Bohn HL, Myer RA, O'Connor GA. 2002.** Soil chemistry. John Wiley & Sons.
- 318 **Borsani O. 2001.** Identification of two loci in tomato reveals distinct mechanisms for salt tolerance. *The Plant Cell*
319 *Online* **13**: 873–888.
- 320 **Chaganti VN, Crohn DM. 2015.** Evaluating the relative contribution of physiochemical and biological factors in
321 ameliorating a saline-sodic soil amended with composts and biochar and leached with reclaimed water. *Geoderma*
322 **259-260**: 45–55.
- 323 **Chi CM, Zhao CW, Sun XJ, Wang ZC. 2012.** Reclamation of saline-sodic soil properties and improvement of rice
324 (*Oriza sativa* L.) growth and yield using desulfurized gypsum in the west of Songnen Plain, northeast China.
325 *Geoderma* **187**: 24–30.
- 326 **Cramer GR. 1992.** Kinetics of maize leaf elongation. II. Responses of a Na-excluding cultivar and a Na-including
327 cultivar to varying Na/Ca salinities. *Journal of Experimental Botany* **43**: 857–864.
- 328 **De Souza ER, Freire MBGDS, da Cunha KPV, do Nascimento CWA, Ruiz HA, Lins CMTSouza ERD, Freire**

- 329 **MBGDS, Karina Patrícia Vieira da Cunha. 2012.** Biomass, anatomical changes and osmotic potential in *Atriplex*
330 *nummularia* Lindl. cultivated in sodic saline soil under water stress. *Environmental and Experimental Botany* **82**:
331 20–27.
- 332 **Duarte HHF, De Souza ER. 2016.** Soil water potentials and *Capsicum annuum* L. under salinity. *Revista Brasileira*
333 *de Ciência do solo* **40**: 1-11.
- 334 **Dubcovsky J, Santa MG, Epatein E, Luo MC, Dvorak J. 1996.** Mapping of the K⁺/Na⁺ discrimination *locusKna1*
335 in wheat. *Theoretical and Applied Genetics* **92**: 448–454.
- 336 **Faiyue B, Vijayalakshmi C, Nawaz S, Nagato Y, Taketa S, Ichii M, Azzawi MJA, Flowers TJ. 2010.** Studies on
337 sodium bypass flow in lateral rootless mutants lrt1 and lrt2, and crown rootless mutant crl1 of rice (*Oryza sativa*
338 L.). *Plant Cell and Environment* **33**: 687–701.
- 339 **Gharaibeh MA, Eltaif NI, Shra’Ah SH. 2010.** Reclamation of a calcareous saline-sodic soil using phosphoric acid
340 and by-product gypsum. *Soil Use and Management* **26**: 141–148.
- 341 **Gorham J. 1990.** Salt tolerance in the Triticeae: K/Na discrimination in Aegilops species. *Journal of Experimental*
342 *Botany* **41**: 1095–1101.
- 343 **Irshad M, Honna T, Eneji AE, Yamamoto S. 2002.** Wheat response to nitrogen source under saline conditions.
344 *Journal of Plant Nutrition* **25**: 2603–2612.
- 345 **Jackson ML. 1956.** Soil chemical analysis-advanced course: Published by the author. Madison, Wisconsin: 895.
- 346 **Kelly J, Rengasamy P. 2006.** Diagnosis and management of soil constraints: transient salinity, sodicity and alkalinity.
347 The University of Adelaide and Grain Research and Development Corporation, Australia.
- 348 **Leland E, Eugene V. 1999.** Crop response and management of salt-affected soils. *Handbook of plant and crop stress*
349 169.
- 350 **Luo SS, Wang SJ, Lei T, Shi SH, Xu SQ, Yang F, Li XJ, Wang ZC, Tian CJ. 2018.** Aggregate-related changes in
351 soil microbial communities under different ameliorant applications in saline-sodic soils. *Geoderma* **329**: 108–117.
- 352 **Maas EV, Hoffman GJ. 1977.** Crop salt tolerance-current assessment. *Journal of the irrigation and drainage division*
353 **103**: 115–134.
- 354 **Maathuis FJ, Amtmann A. 1999.** K⁺ nutrition and Na⁺ toxicity: the basis of cellular K⁺/Na⁺ ratios. *Annals of Botany*
355 **84**: 123–133.

- Matsushita N, Matoh T. 1991.** Characterization of Na⁺ exclusion mechanisms of salt-tolerant reed plants in comparison with salt-sensitive rice plants. *Physiologia Plantarum* **83**: 170–176.
- Mori S, Suzuki K, Oda R, Higuchi K, Maeda U, Yoshida M, Tadano T. 2011.** Characteristics of Na and K absorption in *Suaeda salsa* (L.) Pall. *Soil Science and Plant Nutrition* **57**: 377–386.
- Munns R, James RA. 2008.** Mechanisms of salinity tolerance. *Annual Review of Plant Biology* **59**: 651–681.
- Munns R, Tester M. 2008.** Mechanism of salinity of salt tolerance. *Annual Review of Plant Biology* **59**: 651–681.
- Munns R, Wallace PA, Teakle NL, Colmer TD. 2010.** Measuring soluble ion concentrations (Na⁺, K⁺, Cl⁻) in salt-treated plants. In *plant Stress Tolerance, Methods and Protocols*. New York, USA, Human press, Springer Science.
- Oster JD. 1982.** Gypsum usage in irrigated agriculture: a review. *Fertilizer Research* **3**: 73–89.
- Qadir M, Noble A, Schubert S. 2006.** Sodicty-induced land degradation and its sustainable management: Problems and prospects. *Land Degradation & Development* **17**: 661–676.
- Qadir M, Schubert S, Badia D, Sharma BR, Qureshi AS, Murtaza G. 2007.** Amelioration and nutrient management strategies for sodic and alkali soils. *CAB Reviews Perspectives in Agriculture. Veterinary Science Nutrition and Natural Resources* **21**: 1–13.
- Ren ZH, Gao JP, Li LG. 2005.** A rice quantitative trait locus for salt tolerance encodes a sodium transporter. *Nature genetics* **37**: 1141.
- Renault S, Affifi M. 2009.** Improving NaCl resistance of red-osier dogwood: role of CaCl₂ and CaSO₄. *Plant and soil* **315**: 123–133.
- Rengasamy P. 2010a.** Osmotic and ionic effects of various electrolytes on the growth of wheat. *Australian Journal of Soil Research* **48**: 120–124.
- Rengasamy P. 2010b.** Soil processes affecting crop production in salt-affected soils. *Functional Plant Biology* **37**: 613–620.
- Roy C, Mishra R. 2014.** Impact of NaCl stress on the physiology of four cultivars of *S. lycopersicum*. *Research in Plant Biology* **4**: 09–20.
- Shi SH, Tian L, Nasir F, Bahadur A, Batool A, Luo SS, Yang F, Wang ZC, Tian CJ. 2019.** Response of microbial communities and enzyme activities to amendments in saline-alkaline soils. *Applied Soil Ecology* **135**: 16–24.
- Song JQ, Fujiyama H. 1996.** Difference in response of rice and tomato subjected to sodium salinization to the

addition of calcium. *Soil Science and Plant Nutrition* **42**: 503–510.

Suarez DL. 2001. Sodic soil reclamation: Modelling and field study. *Soil Research* **39**: 1225–1246.

Sumner ME. 1993. Sodic soils: new perspectives. *Australian Journal of Soil Research* **31**: 683–750.

Swarup A. 1982. Availability of ions, zinc and phosphorus in submerged sodic soil as affected by amendments during the growth period of rice crop. *Plant and Soil* **66**: 37–43.

Syed GA, Abdur R. 2017. The influence of salinity and drought stress on sodium, potassium and proline content of *Solanum lycopersicum* L. cv. Rio granded. *Pakistan Journal of Botany* **49**: 1–9.

Tester M, Davenport R. 2003. Na⁺ tolerance and Na⁺ transport in higher plants. *Annals of Botany* **91**: 503–527.

Tsialtas JT, Maslari N. 2009. Selective absorption of K over Na in sugar beet cultivars and its relationship with yield and quality in two contrasting environments of central Greece. *Journal of agronomy and crop science* **195**: 384–392.

Wang CM, Zhang JL, Liu XS, Li Z, Wu GQ, Cai JY. 2009. *Puccinellia tenuiflora* maintains a low Na⁺ level under salinity by limiting unidirectional Na⁺ influx resulting in a high selectivity for K⁺ over Na⁺. *Plant, Cell and Environment* **32**: 486–496.

Wang J, Bai Z, Yang P. 2013. Effect of byproducts of flue gas desulfurization on the soluble salts composition and chemical properties of sodic soils. *Plos one* **8**: e71011.

Wang J, Bai Z, Yang P. 2012. Sodic soil properties and sunflower growth as affected by byproducts of flue gas desulfurization. *Plos one* **7**: e52437.

Wang J, Yang P. 2018. Potential flue gas desulfurization gypsum utilization in agriculture: A comprehensive review. *Renewable and Sustainable Energy Reviews* **82**: 1969–1978.

Wang MM, Liang ZW, Wang ZC, Huang LH, Ma HY, Liu M, Gu X. 2010. Effect of sand application and flushing during the sensitive stages on rice biomass allocation and yield in a saline-sodic soil. *Journal of Food, Agriculture and Environment* **8**: 692–697.

Wang MM, Liang ZW, Yang F, Ma HY, Huang LH, Liu M. 2010. Effects of number of seedlings per hill on rice biomass partitioning and yield in a saline-sodic soil. *Journal of Food Agriculture and Environment* **8**: 628–633.

Wang S, Wan C, Wang Y, Chen H, Zhou Z, Fu Y, Sosebee RE. 2004. The characteristics of Na⁺, K⁺ and free proline distribution in several drought resistant plants of the Alxa Desert, China. *Journal of Arid Environments* **56**:

525–539.

Wang S, Zheng W, Ren J, Zhang C. 2002. Selectivity of various types of salt-resistant plants for K^+ over Na^+ . *Journal of Arid Environment* **52**: 457–472.

Wang SM, Zhang JL, Flower TJ. 2007. Low-affinity Na^+ uptake in the halophyte *Suaeda maritima*. *Plant Physiology* **145**: 559–571.

Wang SM, Zhao GO, Gao YS, Tang ZC, Zhang CL. 2004. *Puccinellia tenuiflora* exhibits stronger selectivity for K^+ over Na^+ than wheat. *Journal of Plant Nutrition* **27**: 1841–1857.

Wu GQ, Wang SM. 2012. Calcium regulates K^+/Na^+ homeostasis in rice (*Oryza sativa* L.) under saline conditions. *Plant Soil and Environment* **58**: 121–127.

Yamanouchi M, Maeda Y, Nagai T. 1987. Relationship between the salt accumulation in the shoots and the degree of salinity tolerance in rice cultivars. *Soil Science Plant Nutrition* **58**: 591–594.

Yang F, An FH, Ma HY, Wang ZC, Zhou X, Liu Z. 2016. Variations on soil salinity and sodicity and its driving factors analysis under microtopography in different hydrological conditions. *Water* **8**: 227.

Yao RJ. 2008. Development and prospect of the research on salt-affected soils in China. *Acta Pedologica Sinica* **45**: 837–845 (in Chinese).

Yeo AR, Flowers TJ. 2010. The Absence of an effect of the Na/Ca ratio on sodium chloride uptake by rice (*Oryza sativa* L.). *New Phytologist* **99**: 81–90.

Yu JB, Wang ZC, Meixner FX, Yang F, Wu HF, Chen XB. 2010. Biogeochemical characterizations and reclamation strategies of saline sodic soil in northeastern China. *CLEAN-Soil, Air, Water* **38**: 1010–1016.

Yunca H, Schmidhalter U. 2005. Drought and salinity: A comparison of the effects of drought and salinity. *Journal of Plant and Nutrition and Soil Science* **168**: 247–273.

Zeng L, Shannon MC. 2000. Salinity effects on seedling growth and yield components of rice. *Crop Science* **40**: 996.

Zhang JL, Flowers TJ, Wang SM. 2010. Mechanisms of sodium uptake by roots of higher plants. *Plant and Soil* **326**: 45.

Zhao YG, Wang SJ, Li Y, Liu J, Zhuo YQ, Chen HX, Wang J, Xu LZ., Sun ZT. 2018. Extensive reclamation of saline-sodic soils with flue gas desulfurization gypsum on the Songnen Plain, Northeast China. *Geoderma* **321**: 52–60.

Tables

Table 1. Properties of the amendments used in the present study.

Table 2 Na^+ and K^+ concentrations and K^+/Na^+ ratios in different organs of rice plant with various treatments.

Table 3. Effects of amendments application on growth and yield of rice plant in 2017.

Table 4. Correlation coefficients among OP, SA, ST values and different growth and yield of rice in 2017.

Figures

Figure 1. Osmotic potential of the 1: 5 soil water extract at various soil profile depths with different amendments application. CK, control, without amendments application; DG, desulfurization gypsum; SS, sandy soil; FM, farmyard manure; M, mixture of desulfurization gypsum, sandy soil and farmyard manure. Bars represent the standard error of the mean of three replications.

Figure 2. Na^+ and K^+ concentrations in different parts of rice plants with various treatments. CK, control, without amendments application; DG, desulfurization gypsum; SS, sandy soil; FM, farmyard manure; M, mixture of desulfurization gypsum, sandy soil and farmyard manure. Bars represent the standard error of the mean of three replications. Different letters denote means that are significantly different from each other ($P<0.05$).

Figure 3. Selective absorption (SA) and selective transport (ST) of rice with various treatments. CK, control, without amendments application; DG, desulfurization gypsum; SS, sandy soil; FM, farmyard manure; M, mixture of desulfurization gypsum, sandy soil and farmyard manure. Bars represent the standard error of the mean of three replications. Different letters denote means that

are significantly different from each other ($P<0.05$).

Figure 4. Na^+ , K^+ and Ca^{2+} concentrations in the 1: 5 soil water extract (0-40 cm) with various treatments. CK, control, without amendments application; DG, desulfurization gypsum; SS, sandy soil; FM, farmyard manure; M, mixture of desulfurization gypsum, sandy soil and farmyard manure. Bars represent the standard error of the mean of three replications. Different letters denote means that are significantly different from each other ($P<0.05$).

Figure 5. Na^+ and K^+ concentrations in the whole rice plant with various treatments. CK, control, without amendments application; DG, desulfurization gypsum; SS, sandy soil; FM, farmyard manure; M, mixture of desulfurization gypsum, sandy soil and farmyard manure. Bars represent the standard error of the mean of three replications. Different letters denote means that are significantly different from each other ($P<0.05$).

Figure 6. The trend of grain yield. CK, control, without amendments application; DG, desulfurization gypsum; SS, sandy soil; FM, farmyard manure; M, mixture of desulfurization gypsum, sandy soil and farmyard manure. Bars represent the standard error of the mean of three replications.

Figure 1

Osmotic potential of the 1: 5 soil water extract at various soil profile depths with different amendments application.

CK, control, without amendments application; DG, desulfurization gypsum; SS, sandy soil; FM, farmyard manure; M, mixture of desulfurization gypsum, sandy soil and farmyard manure.

Bars represent the standard error of the mean of three replications.




Figure 2

Na⁺ and K⁺ concentrations in different parts of rice plants with various treatments.

CK, control, without amendments application; DG, desulfurization gypsum; SS, sandy soil; FM, farmyard manure; M, mixture of desulfurization gypsum, sandy soil and farmyard manure.

Bars represent the standard error of the mean of three replications. Different letters denote means that are significantly different from each other ($P < 0.05$).




Figure 3

Selective absorption (SA) and selective transport (ST) of rice with various treatments.

CK, control, without amendments application; DG, desulfurization gypsum; SS, sandy soil; FM, farmyard manure; M, mixture of desulfurization gypsum, sandy soil and farmyard manure.

Bars represent the standard error of the mean of three replications. Different letters denote means that are significantly different from each other ($P<0.05$).




Figure 4

Na⁺, K⁺ and Ca²⁺ concentrations in the 1: 5 soil water extract (0-40 cm) with various treatments.

CK, control, without amendments application; DG, desulfurization gypsum; SS, sandy soil; FM, farmyard manure; M, mixture of desulfurization gypsum, sandy soil and farmyard manure. Bars represent the standard error of the mean of three replications. Different letters denote means that are significantly different from each other ($P<0.05$).




Figure 5

Na⁺ and K⁺ concentrations in the whole rice plant with various treatments.

CK, control, without amendments application; DG, desulfurization gypsum; SS, sandy soil; FM, farmyard manure; M, mixture of desulfurization gypsum, sandy soil and farmyard manure.

Bars represent the standard error of the mean of three replications. Different letters denote means that are significantly different from each other ($P < 0.05$).




Figure 6

The trend of grain yield.

CK, control, without amendments application; DG, desulfurization gypsum; SS, sandy soil; FM, farmyard manure; M, mixture of desulfurization gypsum, sandy soil and farmyard manure. Bars represent the standard error of the mean of three replications.




Table 1(on next page)

Properties of the amendments used in the present study.

Note: EC, electrical conductivity; SOC, soil organic carbon.

Table 1. Properties of the amendments used in the present study.

Property	Desulfurization gypsum	Sandy soil	Farm manure
pH	7.62	8.92	8.30
EC (dS/m)	34.20	0.78	-
SOC (g/kg)	-	4.23	263.30
K ⁺ (g/kg)	1.00	0.001	13.60
Na ⁺ (g/kg)	1.59	0.008	4.11
Ca ²⁺ (g/kg)	265.30	0.10	7.49
Mg ²⁺ (g/kg)	1.68	0.01	10.20

Note: EC, electrical conductivity; SOC, soil organic carbon.

Table 2 (on next page)

Na⁺ and K⁺ concentrations and K⁺/Na⁺ ratios in different organs of rice plant with various treatments.

Note: the small letters after data in a column for each treatment indicate that ion contents were significantly different at $P=0.05$. CK, control, without amendments application; DG, desulfurization gypsum; SS, sandy soil; FM, farmyard manure; M, mixture of desulfurization gypsum, sandy soil and farmyard manure.

Table 2. Na⁺ and K⁺ concentrations and K⁺/Na⁺ ratios in different organs of rice plant with various treatments.

Treatment	Organ	Na ⁺ (mg/g DW)	K ⁺ (mg/g DW)	K ⁺ /Na ⁺
CK	Grain	0.53c	2.75ab	7.07a
	Leaf	1.38b	3.61a	2.57b
	Sheath	1.06b	1.99b	1.90b
	Root	2.38a	3.01a	1.36b
DG	Grain	0.25d	2.62b	10.61a
	Leaf	1.63b	5.06a	3.18b
	Sheath	1.30c	4.42a	3.40b
	Root	2.02a	3.04b	1.53c
SS	Grain	0.27d	2.69c	10.11a
	Leaf	1.59b	4.93a	3.11b
	Sheath	1.19c	3.56b	2.93b
	Root	2.21a	2.57c	1.28c
FM	Grain	0.19d	2.60b	15.15b
	Leaf	1.61b	5.19a	3.23b
	Sheath	1.13c	2.67b	2.40a
	Root	2.16a	2.51b	1.25b
M	Grain	0.24c	2.61b	12.59a
	Leaf	1.27b	4.62a	3.85b
	Sheath	1.31b	4.65a	3.54b
	Root	2.36a	2.80b	1.21c

Note: the small letters after data in a column for each treatment indicate that ion contents were significantly different at $P=0.05$. CK, control, without amendments application; DG, desulfurization gypsum; SS, sandy soil; FM, farmyard manure; M, mixture of desulfurization gypsum, sandy soil and farmyard manure.

Table 3(on next page)

Effects of amendments application on growth and yield of rice plant in 2017.

Note: mean value and its standard error (SE) are reported. Different letters denote means that are significantly different from each other ($P<0.05$). CK, control, without amendments application; DG, desulfurization gypsum; SS, sandy soil; FM, farmyard manure; M, mixture of desulfurization gypsum, sandy soil and farmyard manure.

Table 3. Effects of amendments application on growth and yield of rice plant in 2017.

Treatment	Height (cm)	Panicle length (cm)	Number of grains per panicle	1000-grain weight (g)	Grain yield (kg/ha)
CK	89.0±3.2a	14.3±0.4a	59.3±3.9b	19.4±0.6b	4130±1349.2b
DG	87.8±3.4a	15.3±0.4a	78.5±9.5ab	21.7±0.6ab	6030±209.9a
SS	89.8±2.1a	15.3±0.6a	86.7±5.2a	20.7±0.8ab	5560±79.4a
FM	94.2±1.1a	14.3±0.3a	68.9±8.4ab	22.4±0.7a	5790±209.9a
M	92.5±2.6a	14.3±0.3a	78.2±4.6ab	21.9±0.9a	6030±209.9a

Note: mean value and its standard error (SE) are reported. Different letters denote means that are significantly different from each other ($P<0.05$). CK, control, without amendments application; DG, desulfurization gypsum; SS, sandy soil; FM, farmyard manure; M, mixture of desulfurization gypsum, sandy soil and farmyard manure.

Table 4(on next page)

Correlation coefficients among OP, SA, ST values and different growth and yield of rice in 2017.

Note: * and ** denote correlation at the 0.05 and 0.01 levels of significance, respectively.

Table 4. Correlation coefficients among OP, SA, ST values and different growth and yield of rice in 2017.

	OP _{ss} (bars)	SA (selective absorption)	ST value (selective transport)	Height (cm)	Panicle length (cm)	Number of grains per panicle	1000-grain weight (g)
SA (selective absorption)	-0.857						
ST (selective transport)	0.628	-0.879*					
Height (cm)	0.695	-0.589	0.278				
Panicle length (cm)	-0.146	-0.205	0.391	-0.62			
Number of grains per panicle	0.276	-0.492	0.319	-0.08	0.727		
1000-grain weight (g)	0.992**	-0.884*	0.671	0.619	-0.024	0.375	
Grain yield (kg/ha)	0.789	-0.925*	0.821	0.303	0.477	0.714	0.855

Note: * and ** denote correlation at the 0.05 and 0.01 levels of significance, respectively.