

Temporal changes in the most effective pollinator of a bromeliad pollinated by bees and hummingbirds

Roberta Luisa Barbosa Leal¹, Marina M Moreira^{1,2}, Alessandra R Pinto³, Julia O Ferreira¹, Miguel Rodriguez-Girones⁴, Leandro Freitas^{Corresp. 1}

¹ Jardim Botânico do Rio de Janeiro, Rio de Janeiro, Rio de Janeiro, Brazil

² Centro de Ciências Agrárias, Universidade Federal do Espírito Santo, Alegre, Espírito Santo, Brazil

³ Programa de Pós-Graduação em Ecologia, Universidade Federal do Rio de Janeiro, Rio de Janeiro, Rio de Janeiro, Brazil

⁴ Estación Experimental de Zonas Áridas, Almería, Spain

Corresponding Author: Leandro Freitas

Email address: lfreytas.jbot@gmail.com

A generalist pollination system may be characterized through the interaction of a plant species with two or more functional groups of pollinators. The spatiotemporal variation of the most effective pollinator is the factor most frequently advocated to explain the emergence and maintenance of generalist pollination systems. There are few studies merging variation in floral visitor assemblages and the efficacy of pollination by different functional groups. Thus, there are gaps in our knowledge about the variation in time of pollinator efficacy and frequency of generalist species. In this study, we evaluated the pollination efficacy of the floral visitors of *Edmundoa lindenii* (Bromeliaceae) and their frequency of visits across four reproductive events. We analysed the frequency of the three groups of floral visitors (large bees, small bees, and hummingbirds) through focal observations in the reproductive events of 2015, 2016, 2017, and 2018. We evaluated the pollination efficacy (fecundity after one visit) through selective exposure treatments and the breeding system by manual pollinations. We tested if the reproductive success after natural pollination varied between the reproductive events, and also calculated the pollen limitation index. *Edmundoa lindenii* is a self-incompatible and parthenocarpic species, requiring the action of pollinators for sexual reproduction. Hummingbirds had higher efficacy than large bees, and small bees acted only as pollen larcenists. The relative frequency of the groups of floral visitors varied between the reproductive events. Pollen limitation has occurred only in the reproductive event of 2017, when visits by hummingbirds were scarce and reproductive success after natural pollination was the lowest. We conclude that hummingbirds and large bees were the main and the secondary pollinators of *E. lindenii*, respectively, and that temporal variations in the pollinator assemblages had effects on its reproductive success. Despite their lower pollination efficacy, large bees ensured seed set when hummingbirds failed. Thus, we provide

evidence that variable pollination environments may favour generalization, even under differential effectiveness of pollinator groups if secondary pollinators provide reproductive assurance.

Temporal changes in the most effective pollinator of a bromeliad pollinated by bees and hummingbirds

Roberta Luísa Barbosa Leal¹, Marina Muniz Moreira^{1,2}, Alessandra Ribeiro Pinto³, Julia de Oliveira Ferreira¹, Miguel Angel Rodríguez-Gironés⁴, Leandro Freitas¹

¹ Jardim Botânico do Rio de Janeiro, Rio de Janeiro, RJ, Brazil

² Universidade Federal do Espírito Santo, Centro de Ciências Agrárias, Alegre, ES, Brazil

³ Universidade Federal do Rio de Janeiro, Programa de Pós-Graduação em Ecologia, Rio de Janeiro, RJ, Brazil

⁴ Estación Experimental de Zonas Áridas, Almería, Spain

Corresponding Author:

Leandro Freitas

Rua Pacheco Leão, 915, Jardim Botânico, Rio de Janeiro, Brasil.

Email address: lfreytas.jbot@gmail.com

ABSTRACT

Background

A generalist pollination system may be characterized through the interaction of a plant species with two or more functional groups of pollinators. The spatiotemporal variation of the most effective pollinator is the factor most frequently advocated to explain the emergence and maintenance of generalist pollination systems. There are few studies merging variation in floral visitor assemblages and the efficacy of pollination by different functional groups. Thus, there are gaps in our knowledge about the variation in time of pollinator efficacy and frequency of generalist species. In this study, we evaluated the pollination efficacy of the floral visitors of *Edmundoa lindenii* (Bromeliaceae) and their frequency of visits across four reproductive events.

Methods

We analysed the frequency of the three groups of floral visitors (large bees, small bees, and hummingbirds) through focal observations in the reproductive events of 2015, 2016, 2017, and

2018. We evaluated the pollination efficacy (fecundity after one visit) through selective exposure treatments and the breeding system by manual pollinations. We tested if the reproductive success after natural pollination varied between the reproductive events, and also calculated the pollen limitation index.

Results

Edmundoa lindenii is a self-incompatible and parthenocarpic species, requiring the action of pollinators for sexual reproduction. Hummingbirds had higher efficacy than large bees, and small bees acted only as pollen larcenists. The relative frequency of the groups of floral visitors varied between the reproductive events. Pollen limitation has occurred only in the reproductive event of 2017, when visits by hummingbirds were scarce and reproductive success after natural pollination was the lowest.

INTRODUCTION

In most plants, flowers are visited by a diverse assemblage of animals, which characterizes generalist pollination systems (Waser *et al.*, 1996). Based on the behavior and morphophysiological traits, floral visitors can be arranged in different functional groups of pollinators, which may differ in their contribution to the plant reproductive success (Fenster *et al.*, 2004). This difference in contribution occurs as these pollinators can vary in frequency of visits and ability to transfer pollen (Shuttleworth & Johnson, 2008; Ollerton, 2017). Studies encompassing generalist pollination systems mainly report floral visitor assemblages and visitation rates (*e.g.*, Thompson, 2001; Freitas & Sazima, 2006; Scrok & Varassin, 2011); despite not all visitors are actual pollinators (Armbruster, Fenster & Dudash, 2000; Ollerton, 2017). Therefore, studies that evaluate the pollination efficacy (for instance, measuring seed set after one visit, Freitas, 2013) of different functional groups of pollinators are necessary to better understand pollinators' relevance on plants with a generalist pollination system.

Based on the principle of “the most effective pollinator,” a plant having more than one functional group of pollinators could be interpreted as an intermediary stage in the shift from one specialized pollinator to another (Stebbins, 1970). An alternative explanation is that generalist systems are not transient and may be favored in certain scenarios, for example, under unpredictable pollination environments (Herrera, 1988; 1996; Waser *et al.*, 1996; Armbruster *et al.*, 2000; Gómez & Zamora, 2006; Ollerton *et al.*, 2007). Accordingly, the spatiotemporal

variation of the most effective pollinator is the factor most frequently advocated to explain the emergence and maintenance of generalist pollination systems. Several studies have explored temporal variation in composition and frequency of floral visitors (*Fenster & Dudash, 2001; Ivey, Martinez & Wyatt, 2003; Zych et al., 2018*). Other studies have quantified the effectiveness (i.e., the product of efficacy times visitation rate, after *Herrera, 1987; Freitas 2013*) of different pollinator species or functional groups (e.g., *Amorim, Galetto & Sazima, 2013; Salas-Arcos, Lara & Ornelas, 2017*). However, information about floral visitor assemblages and their effectiveness on pollination over multiple reproductive events is restricted to a few systems (*Herrera, 1987; 1988; Larsson, 2005; Wiggam & Ferguson, 2005*). Thus, there are gaps in knowledge about the variation of pollinators in generalist plants and their effectiveness over time.

Generalized pollination systems have ecological and evolutionary dimensions (*Armbruster et al., 2000*), therefore the effect of several pollinators in the process of evolutionary generalization depends on the selective pressures exerted by those floral visitors. In this sense, differences in pollination efficacy (sensu *Freitas, 2013*) among functional groups may be enhanced if the variations in the pollination environment affect the plant reproductive success. For instance, pollen limitation (PL), the lower fruit and/or seed production due to inadequate pollen receipt, is widespread in angiosperms (*Ashman et al., 2004; Knight et al., 2005*), and similarly to the pollinator effectiveness, its magnitude varies at several scales (*Bennett et al., 2018*). However, how temporal variations in the pollination environment and PL levels are related is a fundamental but poorly understood aspect to a better understanding of the mechanisms that lead to the maintenance of generalized pollination systems (see *Koski et al., 2018*).

Bromeliaceae is the largest family of predominantly Neotropical angiosperms (*Givnish, 2017*) and hummingbirds are the main group of pollinators of them (*Benzing, 2000*). However, several species are known to have mixed pollination systems, for example involving hummingbirds and bees or bats (*Givnish et al., 2014*). *Edmundoa lindenii* (Regel) Leme bears flowers that are visited by both hummingbirds and bees in montane tropical forests, however we do not know the role of those groups on the plant sexual reproduction over time. Here, we evaluated the efficacy of three groups of floral visitors and their frequency of visits over four reproductive events. In particular, we addressed the following questions: (1) Do hummingbirds,

large bees and small stingless bees pollinate this species, considering the high divergence of traits between them? (2) Is the pollination efficacy of hummingbirds higher than large bees, since hummingbirds pollinate most species of Bromeliaceae? (3) Are the relative and gross frequencies of floral visits by each group equivalent over four reproductive events, since clear environmental variations were not noted in those years? (4) However, if they are not equivalent, are those variations in frequency related to the reproductive success in natural conditions and the occurrence of pollen limitation?

MATERIALS AND METHODS

Study site and species

This study was conducted in an area covered by montane Atlantic Forest, located in the Serra dos Órgãos National Park (PARNASO), Rio de Janeiro state, Brazil (22° 52'- 22° 54' S and 42° 09'- 45° 06' W, ca. 960 m a.s.l.) among four reproductive events (from 2014 to 2018). The average annual rainfall at the study site is 2,436 mm, with the rainiest period between December and March and colder and drier months from June to August. The mean annual temperature is 18.6 °C, with minimum and maximum monthly temperatures of 13.7 °C and 22.9 °C (climate data for 2015 to 2018 from the meteorological station located inside the PARNASO). The field research reported here was performed using the required permit (SISBIO No. 34882, No. 432793).

Edmundoa lindenii (Regel) Leme (Bromeliaceae - Bromelioideae) is a terrestrial, saxicolous, or epiphyte herb, endemic to the Atlantic Forest in south and southeastern Brazil (Martinelli *et al.*, 2008; BFG, 2015). In the study area, this species blooms between December and February and produces fruits between March and April; its flowers are visited by bees and hummingbirds (R.L.B. Leal *et al.*, pers. obs.).

Floral biology

We measured inflorescences of *E. lindenii* (n = 16 individuals) directly in the field with a digital calliper, considering the following traits: scape length, inflorescence diameter, and bract length. Flowers (n = 73) from 28 individuals were collected in the field, stored in 70% alcohol, and measured in the laboratory with a digital calliper considering the following structures: corolla tube length (i.e., from septal nectary to the opening of the corolla) and the width of the corolla tube opening. We counted the number of ovules in 25 flowers (n = 15 individuals).

To analyse the colour quantitatively, we measured the spectral reflectance of petals, sepals, and bracts. For this, 12 flowers ($n = 6$ individuals) were collected in the field, stored in thermal bags containing moist paper, and brought to the laboratory, where they were immediately measured (Lunau *et al.*, 2011). We measured the reflectance using an USB2000 spectrophotometer (OceanOptics, Inc., Dunedin, FL, USA) coupled with a deuterium–halogen light source (DH-2000; OceanOptics, Inc., Ostfildern, Germany), with a light emission range between 215 and 1700 nm. We took all reflectance measurements at a 45° angle in relation to the plant structure, and we used barium sulphate as the white standard and black paper as the black standard (Chittka & Kevan, 2005).

We used the logarithm version of the receptor noise-limited model (Vorobyev *et al.*, 2001) to compare the colours of the petals, sepals, and bracts. Chromatic distances were calculated according to the trichromatic formulation for bees and the tetrachromatic formulation for hummingbirds (Vorobyev & Osorio, 1998). We modelled spectral sensitivity curves using data from *Sephanoides sephaniodes* (Herrera *et al.*, 2008) to estimate hummingbird colour distances, and from *Bombus terrestris* for bees (Telles & Rodríguez-Gironés, 2015). In all cases, we used standard daylight illumination (D65 – Wyszecki & Stiles, 1982). Using these models, we determined the spectral location of each structure in a colour space for each pollinator.

The distance between two points in a colour space provides an approximation of the perceived colour difference (Endler & Mielke, 2005). We evaluated colour distances between sepals, petals, and bracts. Using the receptor noise-limited model, we estimated that two colours were discriminable if their distance was greater than 0.27 units for bees (Telles & Rodríguez-Gironés, 2015) and 1.0 for hummingbirds (Vorobyev *et al.*, 1998). For representation, we also calculated the colour loci of the flower colours in the respective colour space models: the colour hexagon for bees (Chittka 1992) and the colour tetrahedron for hummingbirds (Vorobyev *et al.*, 1998).

Nectar

We measured the nectar volume in flowers previously bagged in bud stage, with a graduated microliter syringe (Hamilton, Nevada, USA), and the concentration with hand-held refractometer (Bellingham + Stanley Eclipse, UK). To evaluate nectar production during anthesis, without the effect of nectar removal, 36 flowers ($n = 10$ individuals) previously bagged at the bud stage were

measured once after anthesis onset. In total, we performed measurements at four different times of the day: 7:00 am (n = 10 flowers); 8:30 am (n = 10); 10:00 am (n = 10); and 11:30 am (n = 6). To evaluate if nectar removal stimulates its secretion, 24 flowers were submitted to four treatments (n = 6 individuals, four flowers per individual – one flower per treatment per individual): R = nectar was measured once at 11:00 am; R1 = nectar was measured twice (at 10:00 am and 11:30 am); R2 = nectar was measured three times (at 8:30 am, 10:00 am and 11:30 am); and R3 nectar was measured four times (at 7:00 am, 8:30 am, 10:00 am and 11:30 am). Thus, for the treatments R1, R2, and R3, nectar was remeasured on the same flowers. We calculated the total amount of sugar (mg) per flower by multiplying nectar volume (μL) by its corrected concentration (mg/μL) according to *Dafni, Kevan & Husband (2005)*.

Breeding system and pollen limitation

We evaluated the breeding system and pollen limitation (PL) through manual pollination treatments. Floral buds of different individuals were previously bagged with “voile” bags, and the flowers submitted to the following treatments: (1) spontaneous self-pollination - 20 flowers from 8 individuals, were bagged and not manipulated in 2016; (2) hand self-pollination - 49 flowers from 20 individuals, were supplemented manually with pollen from the same flower and bagged in 2016; (3) hand cross-pollination - 130 flowers from 47 individuals, were supplemented with pollen from other individual, located at least 10 m away, and then bagged. We conducted the cross-pollination treatment in the years 2016 (26 flowers), 2017 (48 flowers), and 2018 (56 flowers). (4) Pollination under natural conditions - 131 flowers from 36 individuals were marked and kept unbagged. We evaluated the flowers from natural pollination in 2016 (49 flowers), 2017 (20 flowers), and 2018 (62 flowers). At the end of the experiments we collected the fruits and counted the number of seeds per fruit in the laboratory. We calculated the index of pollen limitation as $IPL = 1 - P_n/P_c$ where P_n is the proportion of seed-bearing fruits multiplied by the mean number of seeds per fruit of flowers exposed to natural pollination, and P_c is the proportion of seed-bearing fruits multiplied by the mean number of seeds per fruit of flowers after hand cross-pollination (adapted from *Lloyd and Schoen, 1992; Larson and Barrett, 2000*). We used as response variable of reproductive success in all analyses such an estimate combining fruits bearing seeds and the number of seeds, which is appropriate because *E. lindenii* is parthenocarpic (i.e., flowers develop into fruits independent of pollination). Values of $IPL \leq 0.2$ indicate absence of PL, whereas $IPL > 0.8$ indicates strong PL (*Freitas, Wolowski & Sigiliano,*

2010). We assessed the self-incompatibility by the index of incompatibility (ISI), a relative measure of the seeds produced after self- and cross-pollination (Zapata and Arroyo, 1978). Species with $ISI < 0.30$ may be classified as self-incompatible (Ramirez & Brito, 1990).

Frequency and efficacy of floral visitors

We performed focal observations (sensu Dafni et al., 2005) to evaluate the identity of floral visitors and their frequency of visits, by censuses of 30-min per individual ($n = 190$ individuals) between 6:00 am and 12:00 am, totalizing 184 hours of observation. Observations were done in four reproductive events, in the years 2015 (43.5 hours, $n = 50$ individuals), 2016 (39.0 hours, $n = 42$), 2017 (51.0 hours, $n = 48$), and 2018 (50.5 hours, $n = 50$). Images and videos were captured during the visits to evaluate the foraging behaviour and the floral resources obtained. The visits were identified as legitimate or illegitimate by the expected mode of pollination, considering the shape and arrangement of the flower parts (sensu Irwin et al., 2010; Freitas, 2018). Specimens of insects were collected for posterior identification. We grouped the floral visitors into three functional groups based on identity, body size and foraging behaviour, as following: hummingbirds, large bees (length ≥ 10 mm), and small bees (< 10 mm).

We evaluated the efficacy of the three functional groups of floral visitors through experiments of selective exposition. In this experiment, flowers previously bagged at the bud stage were exposed to a single visit by any visitor of the three functional groups, as follows: small bees (48 flowers in 2016), large bees (20 flowers in 2017 and 60 in 2018), and hummingbirds (20 flowers in 2017 and 65 in 2018). Here we also considered the proportion of seed-bearing fruits multiplied by the mean number of seeds per fruit as the response variable of the pollination efficacy by each functional group.

Data analyses

We performed all the analyses in R version 3.4.4 (R Development Core Team, 2018). We evaluated the production of nectar during anthesis and the effect of nectar removal in nectar secretion by analyses of variance (one-way ANOVA), using the function *aov*. We assessed the differences between treatments (time of anthesis and number of removals) by Tukey HSD post-hoc test, using the function *TukeyHSD*.

We conducted a linear model to evaluate if the reproductive success after natural pollination varied between three reproductive events (2016, 2017, and 2018). Prior to analyses, we cubic-root transformed values of the response variable to meet the normality assumptions. We used the reproductive events (three levels: 2016, 2017, and 2018) as fixed effect. We established the model using the function *lm* and we tested the model assumptions by visual inspection of the residuals using the *qqnorm* function. We calculated the significance of each term in the model using the function *Anova* (Type II) from the *car* package (Kuznetsova et al., 2017) and the differences between levels of categorical factors using the function *lsmeans* from *lsmeans* package (Lenth, 2016).

To evaluate whether hummingbirds and large bees differ in their efficacy, we conducted a linear model. Prior to analyses, we cubic-root transformed values of the response variable to meet the normality assumptions. We used the functional group of pollinators (two levels: hummingbirds and large bees) and the year when the treatments were conducted (two levels: 2017 and 2018) as fixed effects. We established the model using the function *lm* and we tested the model assumptions by visual inspection of the residuals using the *qqnorm* function. We calculated the significance of each term in the model using the function *Anova* (Type II) from the *car* package (Kuznetsova et al., 2017) and the differences between levels of categorical factors using the *lsmeans* package (Lenth, 2016). We did not compare the efficacy of small bees as no seeds were produced after their visits.

To evaluate if the frequency of visits varies between the four reproductive events and the group of floral visitors, we conducted a log-linear model for categorical data (Wood, 2017). We used the relative frequency of visits as a response variable, and the reproductive events (four levels: 2015, 2016, 2017, and 2018) and the functional groups (three levels: hummingbirds, large bees, and small bees) as fixed effects. We established the model using the *glm* function with the family set to *poisson* (Lindsey, 1997). We tested the model assumptions by visual inspection of the residuals. We calculated the significance of each term in the model using the *anova* function (test = *Chisq*) and the differences between levels of categorical factors using the *emmeans* function available in the *emmeans* package.

RESULTS

Floral biology

The flowers of *E. lindenii* are grouped in a compound corymboid inflorescence with *ca.* 100–150 flowers, inserted in the leaf rosette (*Fig. 1*). Inflorescence diameter reached 121.32 ± 17.01 mm and scape length 296.87 ± 23.86 mm (mean \pm SD throughout the text). The flowers are hermaphrodite, with the androecium presenting six stamens included in the corolla and anthers with longitudinal dehiscence (*Fig. 1*). The gynoecium is also included in the corolla and the style ends in a three-lobed stigma (*Fig. 1*). The inferior and trilocular ovary contained 197.9 ± 54.12 ovules. The length of bracts and sepals was 55.15 ± 6.99 mm and 26.0 ± 4.0 mm, respectively. The corolla is tubular (length: 17.95 ± 2.92 mm) with a narrow opening (3.11 ± 1.17 mm). The flowers have diurnal anthesis, characterized by the presence of exposed pollen grains and receptive stigma. The anthesis began at around 06:00 am and lasted for about six hours when corolla closed. The secretion of nectar started in the beginning of anthesis, and it did not increase over time ($F = 0.44$; $df = 3$; $p = 0.726$; *Fig. 2A*). However, the removal of nectar stimulated new secretion ($F = 6.632$, $df = 3$, $p = 0.00273$, *Fig. 2B*).

The bracts reflect red wavelengths, whereas the corolla is UV-reflecting white, and the sepals are UV-absorbing white (*Fig. 3*). The colour of petals, sepals, and bracts, as well as open or closed flowers, is distinguishable by bees and hummingbirds. Flower colour was 1–7 times above the discrimination criteria (0.27) for bee vision (petals-sepals 4 ± 1 , bracts-sepals 4 ± 2) and 5–15 times above the discrimination criteria (1.0) for hummingbirds (petals-sepals 8 ± 2 , bracts-sepals 12 ± 4 , bracts-petals 15 ± 7).

Breeding system and pollen limitation

Edmundoa lindenii is self-incompatible ($ISI = 0.08$; *Table 1*) and parthenocarpic (*Table 1*), requiring the action of pollinators for sexual reproduction. We observed that the reproductive success after natural pollination varied between reproductive events (ANOVA: $F=12.9$, $df = 2$ $p < 0.001$). The lowest reproductive success occurred in the reproductive event of 2017 (contrasts: **2016-2017**: $t = 5.050$; $df = 103$; $p < 0.001$; **2016-2018**: $t = 2.100$; $df = 103$; $p = 0.095$; **2017-2018**: $t = -3.840$; $df = 103$; $p < 0.001$; *Fig. 4*). Pollen limitation was expressive only in 2017 (PL index: 2016 = -0.21, 2017 = 0.70, 2018 = -0.003).

Floral visitors and temporal variation

The flowers of *E. lindenii* were visited by 11 species of animals belonging to three functional groups (hummingbirds, large bees, and small bees; *Table 2*). Hummingbirds were the group with the highest species richness with seven species (*Table 2; Fig. 1*). The small bee *T. spinipes* was the only visitor that conducted illegitimate visits, resulting in damage of corolla and/or anthers by chewing. Hummingbirds and large bees foraged for nectar acting as legitimate visitors, and small bees collected pollen.

The pollinator functional group affected the reproductive success of *E. lindenii* (*Table 3*) and hummingbirds had higher efficacy than large bees (contrast: $t = 3.015$ $df = 148$, $p = 0.003$; *Table 4; Fig. 5*). Small bees did not act as pollinators, as none of the flowers they visited produced seeds. The frequency of visits varied between functional groups of floral visitors with significant interaction between functional group and reproductive event (*Table 4; Fig. 6*).

DISCUSSION

The frequency of floral visitors' groups (hummingbirds, large bees, and small bees) varied between reproductive events of *E. lindenii*, and this variation resulted in a reduction in the natural pollination in one year, when pollen limitation was also recorded. The existence of year-to-year changes in the composition of floral visitors has been found in several systems (*Schemske & Horvitz, 1984; Traveset & Sáez, 1997; Price et al., 2005; Olesen et al., 2008; Petanidou et al., 2008*), while in others, pollination efficacy between different years was studied (*Fishbein & Venable, 1996; Stoepler et al., 2012*). However, there are fewer studies that consider both plant reproductive success and variations in the pollinator assemblages along time (*Herrera, 1990; Fleming et al., 2001; Salas-Arcos et al., 2017*). Thus, through observational and experimental approaches, we have shown that the reproductive success of a generalist species responded to temporal variation of its assemblage of pollinators.

Overall hummingbirds had higher efficacy and visitation frequency than large bees, so they could be pointed as the main pollinators of *E. lindenii*. Hummingbirds also act as the main pollinators in other species of Bromeliaceae (*Schmid et al., 2011; Magalhães et al., 2018*). However, large bees did not always act as secondary pollinators of *E. lindenii*: they were the main pollinators when the frequency of visits by hummingbirds was low. Pollinators belonging to different functional groups may vary in their pollination effectiveness, exerting pressures toward specialization (*Stebbins, 1970; Rosas-Guerrero et al. 2014*). Thus, evolution of a

generalized pollination system is expected when different pollinators play the same role as selective agents or if the less effective pollinators provide reproductive assurance, offsetting fluctuations of the most effective pollinators. This could be the case for large bees and *E. lindenii*, as they can ensure sexual reproduction when hummingbirds fail. In short, our results are consistent with the hypothesis that the maintenance of generalist pollination is related to the existence of variable pollination environments (Waser *et al.*, 1996) and support that combined measurements of reproductive success and pollinator assemblages along time and space as an interesting approach in this regard (see Gómez & Zamora, 2006 for additional suggestions). Curiously, there was no evidence of declining populations of hummingbirds in the study area in 2017, the causes of the drastic reduction of visits to the flowers of *E. lindenii* by hummingbirds that year are completely unknown.

Despite the differences on effectiveness, flowers of *E. lindenii* are attractive to both hummingbirds and bees. Floral visitors identify and select flowers using a variety of characteristics, including size and color contrasts (Papiorek *et al.*, 2016), and hummingbirds and bees can detect different color spectra (Chittka & Waser, 1997; Vorobyev & Osorio, 1998). The floral traits of *E. lindenii* are detectable by both groups of floral visitors. Petals had UV reflection, sepals absorbed UV, and bracts were red. These results correspond to the expected pattern for attraction of bees and hummingbirds, as bees have a spectrum of vision that includes UV wavelengths, around 300–400 nm (Kevan, Chittka & Dyer, 2001), and hummingbirds are known for their preference for red-coloured flowers that mostly are UV-absorbent (Lunau *et al.* 2011). However, trade-offs between selective pressures exerted by different pollinators may occur if they differ in preference for floral traits (Gervasi & Schiestl, 2017). As pointed out above, we may speculate that the temporal fluctuation in the visitor frequencies of *E. lindenii* may reduce the probability of pollinators exerting consistent selective pressures on its floral traits (Schemske & Horvitz, 1984; Gómez & Zamora, 2006). In this case, attractive and non-restrictive flowers for large bees are important for reproductive assurance in reproductive events with a low frequency of hummingbirds, thus it would not be expected a shift to specialized hummingbird-flowers. However, further experiments are needed to support that selection toward pollination specialization by hummingbirds is restrained in this system.

In addition to visual signals, the nectar of *E. lindenii* is accessible to hummingbirds and large bees, thus its flowers are attractive and legitimately accessible to the two groups. The

dynamics of nectar production influences the behaviour of pollinators during visits to the flowers (*Parachnowitsch, Manson & Sletvold, 2018*). Although nectar production of *E. lindenii* did not increase over time, its removal stimulated further secretion. Similar results have been found in other bromeliads, including some species visited by hummingbirds and bees (*Galetto & Bernardello, 1992; Ordano & Ornelas, 2004*). Nectar secretion after withdrawal may favor repeated visits to the flower. This is consistent with our results of the pollination efficacy experiment in *E. lindenii*. Specifically, seed set after one visit by a large bee or hummingbird was lower than seed set of flowers exposed to pollinators during the whole anthesis, indicating that more than one visit is necessary to achieve maximum fecundity in this species. At last, nectar secretion after removals and repeated visits by pollinators could be related to the male component of the reproductive success (see *Ordano & Ornelas, 2004*).

The population of *E. lindenii* at PARNASO was self-incompatible, but self-compatibility has been registered in other populations of this species (*Matallana et al., 2010*). Variations between self-incompatibility and self-compatibility within species are common in plant evolution and may indicate transitions between reproductive systems (*Igic, Lande & Kohn, 2008*). Studies have shown that compatibility barriers can be broken by genetic changes (such as mutations) (e.g., *Sassa et al., 1997*), physiological factors, elevated temperatures, and stress (e.g., *Tezuka et al., 1997*), allowing for self-pollination. Moreover, breeding systems may be related to the degree of pollination generalization, linking shifts in pollination and incompatibility systems. For instance, *Wessinger and Kelly (2018)* found a relationship between self-compatibility and attributes related to the attraction of hummingbirds, including red flowers and loss of floral aroma and UV-absorbing pigments. In contrast, we found self-incompatibility and UV-absorbing sepals in *E. lindenii* despite higher frequency of visits by hummingbirds. Self-incompatibility in this species seems to work as a barrier to autogamous pollination, since small bees access the anthers, make long visits to the flower, and manipulate the pollen. In fact, pollinators usually do not operate independently of herbivores (florivores in this case), which may generate a trade-off between the fitness functions by each kind of organism (*Ashman, 2002; Gómez & Zamorra, 2006; Gélvez-Zúñiga et al., 2018*).

CONCLUSION

Our results allow us to conclude that hummingbirds and large bees were the main and the secondary pollinators of *E. lindenii*, respectively. Moreover, small bees had a negative effect on its reproduction. These results could indicate a higher degree of specialization of this system than the apparent generalization considering floral visitor composition (see Padyšáková *et al.*, 2013), which would be in accordance with the most effective pollinator principle (Stebbins, 1970). However, temporal variations in the pollinator assemblages had effects on reproductive success of *E. lindenii*, leading to the occurrence of pollen limitation when visits by hummingbirds were scarce. Despite their lower pollination efficacy, large bees ensured seed set when hummingbirds failed. Plant pollination generalization has been associated with similar effectiveness by the different pollinators (Waser *et al.*, 1996). However, we provide evidence that variable pollination environments may favour generalization, even under differential effectiveness of pollinator groups if secondary pollinators provide reproductive assurance. This reinforces the idea of different mechanisms driving the evolution of generalized pollination systems (Schiestl, Balmer & Gervasi, 2018).

ACKNOWLEDGMENTS

The authors thank Gabriel C. Rocha for field and data analysing assistance; Cristovão Albuquerque (Publicase, Brazil) for language editing; Mariano Ordano for critical review; and the staff of the Serra dos Órgãos National Park for logistics support and for allowing access to Park facilities.

REFERENCES

- Amorim FW, Galetto L, Sazima M. 2013. Beyond the pollination syndrome: nectar ecology and the role of diurnal and nocturnal pollinators in the reproductive success of *Inga sessilis* (Fabaceae). *Plant biology* 5(2): 317-327 DOI: 10.1111/j.1438-8677.2012.00643.x
- Armbruster S, Fenster C, Dudash M. 2000. Pollination 'principles' revisited: specialization, pollination syndromes, and the evolution of flowers. In: Totland Ø, ed. *The Scandinavian Association for Pollination Ecology honours Knut Fægri*. Oslo: Det Norske Videnskaps-Akademi, 179-200.

- Ashman T. 2002. The role of herbivores in the evolution of separate sexes from hermaphroditism. *Ecology* 83: 1175-1184 DOI: 10.1890/0012-9658(2002)083[1175:trohit]2.0.co;2
- Ashman T, Knight TM, Steets JA, Amarasekare P, Burd M, Campbell DR, Dudash MR, Johnston MO, Mazer SJ, Mitchell RJ, Morgan MT, Wilson WG. 2004. Pollen limitation of plant reproduction: ecological and evolutionary causes and consequences. *Ecology*, 85: 2408-2421 DOI:10.1890/03-8024
- Bates D, Maechler M, Bolker B, Walker S. 2015. lme4: Linear mixed-effects models using Eigen and S4. R package version 1.1–7. 2014.
- BFG. 2015. Growing knowledge: an overview of seed plant diversity in Brazil. *Rodriguésia* 66: 1085-1113 DOI: 10.1590/2175-7860201566411
- Bennett JM, Steets JA, Burns JH, Durka W, Vamosi JC, Arceo-Gomez G, Burd M, Burkle LA, Ellis AG, Freitas L, Li J, Rodger JG, Wolowski M, Xia J, Ashman TL, Knight TM. 2018. GloPL, a global data base on pollen limitation of plant reproduction. *Scientific Data* 5: 180249 DOI: 10.1038/sdata.2018.249
- Chittka, L. 1992. The colour hexagon: a chromaticity diagram based on photoreceptor excitations as a generalized representation of colour opponency. *Journal of Comparative Physiology A* 170(5): 533-543 DOI: 10.1007/bf00199331
- Chittka L, Waser NM. 1997. Why red flowers are not invisible to bees. *Israel Journal of Plant Sciences* 45(2-3), 169-183 DOI: 10.1080/07929978.1997.10676682
- Chittka L, Kevan PG. 2005. Flower colour as advertisement. In Dafni A, Kevan PG, Husband BC, eds. *Practical Pollination Biology*. Cambridge, Canada: Enviroquest, 157–196.
- Dafni, A., Kevan, P. G., & Husband, B. C. 2005. Practical pollination biology. *Practical pollination biology*. Haifa: Institute of Evolution, University of Haifa
- Fenster CB, Armbruster WS, Wilson P, Dudash MR, Thomson JD. 2004. Pollination syndromes and floral specialization. *Annual Review of Ecology, Evolution, and Systematics* 35: 375-403 DOI: 10.1146/annurev.ecolsys.34.011802.132347
- Fenster CB, Dudash MR. 2001. Spatiotemporal variation in the role of hummingbirds as pollinators of *Silene virginica*. *Ecology* 82(3): 844-851 DOI: 10.2307/2680202
- Fishbein M, Venable DL. 1996. Diversity and temporal change in the effective pollinators of *Asclepias tuberosa*. *Ecology* 77(4): 1061-1073 DOI: 10.2307/2265576

- 429 Fleming TH, Sahley, CT, Holland JN, Nason JD, Hamrick JL. 2001. Sonoran desert columnar
430 cacti and the evolution of generalized pollination systems. *Ecological Monographs* 71(4)
431 511-530 DOI: 10.1890/0012-9615(2001)071[0511:sdccat]2.0.co;2
- 432 Freitas L. 2013. Concepts of pollinator performance: is a simple approach necessary to achieve a
433 standardized terminology? *Brazilian Journal of Botany* 36: 3-8 DOI: 10.1007/s40415-013-
434 0005-6
- 435 Freitas L. 2018. Precisamos falar sobre o uso impróprio de recursos florais. *Rodriguésia* 68(4):
436 2223-2228 DOI: 10.1590/2175-7860201869446
- 437 Freitas L, Sazima M. 2006. Pollination biology in a tropical high-altitude grassland in Brazil:
438 interactions at the community level. *Annals of the Missouri Botanical Garden* 93: 465-516
439 DOI: 10.3417/0026-6493(2007)93[465:PBIATH]2.0.CO;2
- 440 Freitas L, Wolowski M, Sigiliano MI. 2010. Ocorrência de limitação polínica em plantas de
441 Mata Atlântica. *Oecologia Australis* 14(1): 251-265 DOI: 10.4257/oeco.2010.1401.15
- 442 Galetto, L., & Bernardello, L. M. 1992. Extrafloral nectaries that attract ants in Bromeliaceae:
443 structure and nectar composition. *Canadian Journal of Botany* 70(6): 1101-1106 DOI:
444 10.1139/b92-136
- 445 Gélvez-Zúñiga I, Teixido AL, Neves AC, Fernandes GW. 2018. Floral antagonists counteract
446 pollinator-mediated selection on attractiveness traits in the hummingbird-pollinated *Collaea*
447 *cipoensis* (Fabaceae). *Biotropica* 50: 797-804 DOI:10.1111/btp.12574
- 448 Gervasi DD, Schiestl FP. 2017. Real-time divergent evolution in plants driven by pollinators.
449 *Nature Communications* 8(1), 14691 DOI: 10.1038/ncomms14691
- 450 Givnish TJ. 2017. A New World of plants. *Science* 358 (6370): 1535-1536 DOI:
451 10.1126/science.aar4191
- 452 Givnish TJ, Barfuss MH, Van Ee B, Riina R, Schulte K, Horres R, Gonsiska PA, Jabaily RS,
453 Crayn DM, Smith JA, Winter K, Brown GK, Evans TM, Holst BK, Luther H, Till W, Zizka
454 G, Berry PE, Sytsma KJ. 2014. Adaptive radiation, correlated and contingent evolution, and
455 net species diversification in Bromeliaceae. *Molecular Phylogenetics and Evolution* 71: 55–
456 78 DOI: 10.1016/j.ympev.2013.10.010
- 457 Gómez JM, Zamora R. 2006. Ecological factors that promote the evolution of generalization in
458 pollination systems. In Waser N, Ollerton J. eds. *Plant-pollinator interactions: from*
459 *specialization to generalization*. Chicago: University of Chicago Press, 145-165.

- Herrera CM. 1987. Components of pollinator "quality": comparative analysis of a diverse insect assemblage. *Oikos* 50: 79-90 DOI: 10.2307/3565403
- Herrera CM. 1988. Variation in mutualisms - the spatio-temporal mosaic of a pollinator assemblage. *Biological Journal of the Linnean Society* 35: 95-125 DOI: 10.1111/j.1095-8312.1988.tb00461.x
- Herrera CM. 1990. Daily patterns of pollinator activity, differential pollinating effectiveness, and floral resource availability, in a summer-flowering Mediterranean shrub. *Oikos* 58(3): 277-288 DOI: 10.2307/3545218
- Herrera CM. 1996. Floral traits and plant adaptation to insect pollinators: a devil's advocate approach. In Lloyd DG., Barrett SCH. eds. *Floral biology: studies on floral evolution in animal-pollinated plants*. New York: Chapman & Hall, 65-87.
- Herrera G, Zagal JC, Diaz M, Fernández MJ, Vielma A, Cure M, Martinez J, Bozinovic F, Palacios AG. 2008. Spectral sensitivities of photoreceptors and their role in colour discrimination in the green-backed firecrown hummingbird (*Sephanoides sephaniodes*). *Journal of Comparative Physiology A* 194(9): 785-794 DOI: 10.1007/s00359-008-0349-8.
- Igic B, Lande R, Kohn JR. 2008. Loss of self-incompatibility and its evolutionary consequences. *International Journal of Plant Sciences* 169(1): 93-104 DOI: 10.1086/523362
- Ivey CT, Martinez P, Wyatt R. 2003. Variation in pollinator effectiveness in swamp milkweed, *Asclepias incarnata* (Apocynaceae). *American Journal of Botany* 90(2): 214-225 DOI: 10.3732/ajb.90.2.214
- Irwin RE, Bronstein JL, Manson JS, Richardson L. 2010. Nectar robbing: ecological and evolutionary perspectives. *Annual Review of Ecology, Evolution, and Systematics* 41(1): 271-292 DOI: 10.1146/annurev.ecolsys.110308.120330
- Kevan PG, Chittka L, Dyer AG. 2001. Limits to the salience of ultraviolet: lessons from colour vision in bees and birds. *Journal of Experimental Biology* 204(14): 2571-2580. Available at <https://jeb.biologists.org/content/jexbio/204/14/2571.full.pdf> (accessed 28 Sept 2019)
- Kuznetsova A, Brockhoff PB, Christensen RHB. 2017. lmerTest Package: tests in linear mixed effects models. *Journal of Statistical Software* 82(13): 1-26 DOI: 10.18637/jss.v082.i13
- Larson BM, Barrett SC. 2000. A comparative analysis of pollen limitation in flowering plants. *Biological Journal of the Linnean Society* 69(4): 503-520 DOI: 10.1111/j.1095-8312.2000.tb01221.x

- 491 Larsson M. 2005. Higher pollinator effectiveness by specialist than generalist flower-visitors of
492 unspecialized *Knautia arvensis* (Dipsacaceae). *Oecologia* 146: 394-403 DOI:
493 10.1007/s00442-005-0217-y
- 494 Lenth RV. 2016. Least-squares means: the R package lsmeans. *Journal of Statistical Software*
495 69(1): 1-33 DOI: 10.18637/jss.v069.i01
- 496 Lindsey JK. 1997. *Applying generalized linear models*. New York: Springer.
- 497 Lloyd DG, Schoen DJ. 1992. Self- and cross-fertilization in plants. I. Functional dimensions.
498 *International Journal of Plant Sciences* 153: 358-369. Available at
499 <http://www.jstor.org/stable/2995676> (accessed 20 Oct 2019).
- 500 Lunau K, Papiorek S, Eltz T, Sazima M. 2011. Avoidance of achromatic colours by bees
501 provides a private niche for hummingbirds. *Journal of Experimental Biology* 214(9): 1607-
502 1612 DOI: 10.1242/jeb.052688
- 503 Magalhães AFP, Maruyama PK, Tavares LAF, Martins RL. 2018. The relative importance of
504 hummingbirds as pollinators in two bromeliads with contrasting floral specializations and
505 breeding systems. *Botanical Journal of the Linnean Society* 188(3): 316-326 DOI:
506 10.1093/botlinnean/boy050
- 507 Martinelli G, Vieira CM, Gonzalez M, Leitman P, Piratininga A, Costa AF, Forzza RC 2008.
508 Bromeliaceae da Mata Atlântica brasileira: lista de espécies, distribuição e conservação.
509 *Rodriguésia* 59(1): 209-258 DOI: doi.org/10.1590/2175-7860200859114
- 510 Matallana G, Godinho MAS, Guilherme FAG, Belisario M, Coser TS, Wendt T. 2010. Breeding
511 systems of Bromeliaceae species: evolution of selfing in the context of sympatric
512 occurrence. *Plant Systematics and Evolution* 289(1-2): 57-65 DOI: 10.1007/s00606-010-
513 0332-z
- 514 Olesen JM, Bascompte J, Elberling H, Jordano P. 2008. Temporal dynamics in a pollination
515 network. *Ecology* 89(6): 1573-1582 DOI: 10.1890/07-0451.1
- 516 Ollerton J. 2017. Pollinator diversity: distribution, ecological function, and conservation. *Annual*
517 *Review of Ecology, Evolution, and Systematics* 48: 353-376 DOI: 10.1146/annurev-ecolsys-
518 110316-022919
- 519 Ordano M, Ornelas JF. 2004. Generous-like flowers: nectar production in two epiphytic
520 bromeliads and a meta-analysis of removal effects. *Oecologia* 140(3): 495-505 DOI:
521 10.1007/s00442-004-1597-0

- 522 Padyšáková E, Bartoš M, Tropek R, Janeček S. 2013. Generalization versus specialization in
523 pollination systems: visitors, thieves, and pollinators of *Hypoestes aristate* (Acanthaceae).
524 *PLoS One* 8(4):e59299 DOI:10.1371/journal.pone.0059299
- 525 Papiorek S, Junker RR, Alves-dos-Santos I, Melo GA, Amaral-Neto LP, Sazima M, Wolowski
526 M, Freitas L, Lunau K. 2016. Bees, birds and yellow flowers: pollinator-dependent
527 convergent evolution of UV patterns. *Plant Biology* 18(1): 46-55 DOI: 10.1111/plb.12322
- 528 Parachnowitsch AL., Manson JS, Sletvold N. 2018. Evolutionary ecology of nectar. *Annals of*
529 *Botany* 123(2): 247-261 DOI: 10.1093/aob/mcy132
- 530 Petanidou T, Kallimanis AS, Tzanopoulos J., Sgardelis SP, Pantis JD. 2008. Long-term
531 observation of a pollination network: fluctuation in species and interactions, relative
532 invariance of network structure and implications for estimates of specialization. *Ecology*
533 *Letters* 11(6): 564-575 DOI: 10.1111/j.1461-0248.2008.01170.x
- 534 Price MV, Waser NM, Irwin RE, Campbell DR, Brody AK. 2005. Temporal and spatial
535 variation in pollination of a montane herb: a seven-year study. *Ecology* 86(8): 2106-2116
536 DOI: 10.1890/04-1274
- 537 Ramirez N, Brito Y. 1990. Reproductive biology of a tropical palm swamp community in the
538 Venezuelan llanos. *American Journal of Botany* 77(10): 1260-1271 DOI: 10.1002/j.1537-
539 2197.1990.tb11378.x
- 540 Rosas-Guerrero V, Aguilar R, Martén-Rodríguez S, Ashworth L, Lopezaraiza-Mikel M, Bastida
541 JM, Quesada M. 2014. A quantitative review of pollination syndromes: do floral traits
542 predict effective pollinators? *Ecology Letters* 17: 388-400. DOI: 10.1111/ele.12224
- 543 Salas-Arcos L, Lara C, Ornelas JF. 2017. Reproductive biology and nectar secretion dynamics of
544 *Penstemon gentianoides* (Plantaginaceae): a perennial herb with a mixed pollination system?
545 *PeerJ* 5: e3636 DOI: 10.7717/peerj.3636
- 546 Sassa H, Hirano H, Nishio T, Koba T. 1997. Style-specific self-compatible mutation caused by
547 deletion of the S-RNase gene in Japanese pear (*Pyrus serotina*). *The Plant Journal* 12(1):
548 223-227 DOI: 10.1046/j.1365-313X.1997.12010223.x
- 549 Schemske DW, Horvitz CC. 1984. Variation among floral visitors in pollination ability: a
550 precondition for mutualism specialization. *Science* 225(4661): 519-521 DOI:
551 10.1126/science.225.4661.519

- Schiestl FP, Balmer A, Gervasi DD. 2018. Real-time evolution supports a unique trajectory for generalized pollination. *Evolution* 72: 2653-2668. DOI:10.1111/evo.13611
- Schmid S, Schmid VS, Zillikens A, Harter-Marques B, Steiner J. 2011. Bimodal pollination system of the bromeliad *Aechmea nudicaulis* involving hummingbirds and bees. *Plant Biology* 13: 41–50 DOI: 10.1111/j.1438-8677.2010.00348.x
- Scrok GJ, Varassin IG. 2011. Reproductive biology and pollination of *Aechmea distichantha* Lem. (Bromeliaceae). *Acta Botanica Brasilica* 25(3): 571-576 DOI: 10.1590/s0102-33062011000300009
- Shuttleworth A, Johnson SD. 2008. Bimodal pollination by wasps and beetles in the African milkweed *Xysmalobium undulatum*. *Biotropica* 40: 568-574 DOI: 10.1111/j.1744-7429.2008.00418.x
- Stebbins GL. 1970. Adaptive radiation of reproductive characteristics in angiosperms, I: Pollination mechanisms. *Annual Review of Ecology and Systematics* 1: 307-326 DOI: 10.1146/annurev.es.01.110170.001515
- Stoepler TM, Edge A, Steel A, O'Quinn RL, Fishbein M. 2012. Differential pollinator effectiveness and importance in a milkweed (*Asclepias*, Apocynaceae) hybrid zone. *American Journal of Botany* 99(3): 448-458 DOI: 10.3732/ajb.1100272
- R Core Team 2018. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. Available at <https://www.R-project.org/>.
- Telles FJ, Rodríguez-Gironés MA. 2015. Insect vision models under scrutiny: what bumblebees (*Bombus terrestris terrestris* L.) can still tell us. *The Science of Nature* 102(1-2): 4 DOI: 10.1007/s00114-014-1256-1
- Tezuka T, Tsuruhara A, Suzuki H, Takahashi SY. 1997. A connection between the self-incompatibility mechanism and the stress response in lily. *Plant and Cell Physiology* 38: 107-112. DOI: 10.1093/oxfordjournals.pcp.a029139
- Thompson JD. 2001. How do visitation patterns vary among pollinators in relation to floral display and floral design in a generalist pollination system? *Oecologia* 126(3): 386-394 DOI: 10.1007/s004420000531
- Traveset A, Sáez E. 1997. Pollination of *Euphorbia dendroides* by lizards and insects: spatio-temporal variation in patterns of flower visitation. *Oecologia* 111(2): 241-248 DOI: 10.1007/PL00008816

- 583 Vorobyev M, Osorio D. 1998. Receptor noise as a determinant of colour thresholds. *Proceedings*
584 *of the Royal Society of London. Series B: Biological Sciences* 265(1394): 351-358 DOI:
585 10.1098/rspb.1998.0302
- 586 Vorobyev MD, Osorio D, Bennett ATD, Marshall NJ, Cuthill IC. 1998. Tetrachromacy, oil
587 droplets and bird plumage colours. *Journal of Comparative Physiology A*. 183: 621–633
588 DOI: 10.1007/s003590050286
- 589 Vorobyev MD, Brandt R, Peitsch D, Laughlin SB, Menzel R. 2001. Colour thresholds and
590 receptor noise: behaviour and physiology compared. *Vision Research* 41:639–53. DOI:
591 10.1016/S0042-6989(00)00288-1
- 592 Waser NM, Chittka L, Price MV, Williams NM, Ollerton J. 1996. Generalization in pollination
593 systems, and why it matters. *Ecology* 77: 1043-1060 DOI: 10.2307/2265575.
- 594 Wessinger CA, Kelly JK. 2018. Selfing can facilitate transitions between pollination syndromes.
595 *American Naturalist* 191(5): 582-594. DOI: 10.1086/696856
- 596 Wiggam S, Ferguson CJ. 2005. Pollinator importance and temporal variation in a population of
597 *Phlox divaricata* L. (Polemoniaceae). *American Midland Naturalist* 154(1): 42-54 DOI:
598 10.1674/0003-0031(2005)154[0042:PIATVI]2.0.CO;2
- 599 Wood SN. 2017. Generalized additive models: an introduction with R. Boca Raton: Chapman
600 and Hall.
- 601 Zapata TR, Arroyo MTK. 1978. Plant reproductive ecology of a secondary deciduous tropical
602 forest in Venezuela. *Biotropica* 10(3): 221-230 DOI: 10.2307/2387907
- 603 Zych M, Junker RR, Nepi M, Stpiczyńska M, Stolarska B, Roguz K. 2018. Spatiotemporal
604 variation in the pollination systems of a supergeneralist plant: is *Angelica sylvestris*
605 (Apiaceae) locally adapted to its most effective pollinators? *Annals of Botany* 123(2): 415-
606 428 DOI: 10.1093/aob/mcy140

Figure 1

fig1 - *Edmundoa lindenii* flowers were visited by three functional groups: hummingbirds, large bees, and small bees.

(A) Inflorescence and flowers; (B) Visit by the hummingbird *Amazilia fimbriata*; (C) visit by the large bees *Bombus morio*; (D) visit by the small bee *Trigona spinipes*. All observations were made in the montane Atlantic Forest at Serra do Órgãos National Park, southeastern Brazil.




Figure 2

Nectar production in *Edmundoa lindenii* flowers did not increase during the anthesis, but the nectar removal stimulated new secretion.

(A) Nectar production along the anthesis at four different times of the day: 7:00h; 8:30h; 10:00h; and 11:30h. (B) Nectar production after experimental removal of nectar: R = no removal, sampled one time at 11:00h; R1 = one removal, sampled two times at 10:00h and 11:30h; R2 = two removals, sampled three times at 8:30h, 10:00h and 11:30h; and R3 = three removals, sampled four times at 7:00h, 8:30h, 10:00h and 11:30h. The horizontal line in the both boxplots represents the median values, the upper and lower sides of the box represent the corresponding quartiles, and vertical lines are minimum and maximum values of the data range. Dots are outliers. Different letters indicate statistical significance between pairs of years ($p < 0.05$) by ANOVA post-hoc test (TukeyHSD).





Figure 3

fig3 - Attractive structures of *Edmundoa lindenii* include red bracts, UV- reflecting white petals, and UV-absorbing white sepals, and can be detected by bees and hummingbirds.

(A) The reflectance spectra of attractive structures in *E. lindenii* inflorescences. For each structure, the colored line represents the mean reflectance and the corresponding color shading represents the standard deviation. Red = bract reflectance; blue = sepal reflectance; green = petal reflectance. (B) Hexagon model for bee vision based on the photoreceptors of *Bombus terrestris*. (C) Tetrahedron model for bird vision based in the photoreceptors of *Sephanoides sephaniodes*. In both models, the gray point represents achromatic center, the red point represents mean loci for bracts, the blue point indicates the mean loci for sepals, and the green point represents mean loci for petals.




Figure 4

fig4 - The reproductive event of 2017 presented the lower reproductive success by natural pollination.

Reproductive success after natural pollination among three reproductive events (2016, 2017, and 2018). Seeds = proportion of fruits with seeds formed multiplied by the number of seeds in each fruit. Different letters show statistically significant difference ($p < 0.05$).




Figure 5

fig5A - Hummingbirds had higher efficacy than large bees in two reproductive events.

Pollinator efficacy in the reproductive events of 2017 (A) and 2018 (B). Seeds = proportion of fruits with seeds formed multiplied by the number of seeds in each fruit. Different letters show statistically significant difference ($p < 0.05$).




Figure 6

fig6 - Relative frequency of visits by each functional group in *Edmundoa lindenii* flowers varied among the four reproductive events.

The relative frequency of visits in the reproductive events of 2015, 2016, 2017 and 2018. HB = hummingbirds; LB = large bees; SB = small bees.




Table 1 (on next page)

The population of *Edmundoa lindenii* in PARNASO is self-incompatible and parthenocarpic.

Fruit and seeds production after hand pollination treatments at Serra do Órgãos National Park, southeastern Brazil. Reproductive success = proportion of fruits with seeds multiplied by the number of seeds.

Table 1:
The population of *Edmundoa lindenii* in PARNASO is self-incompatible and parthenocarpic.
 Fruit and seeds production after hand pollination treatments at Serra do Órgãos National Park, southeastern Brazil. Reproductive success = proportion of fruits with seeds multiplied by the number of seeds.

Treatments	Flowers (n)	Fruits with seeds (n)	Fruit set	Seeds (mean \pm sd)	Reproductive success
Spontaneous self-pollination	20	1	0.05	26	1.30
Manual self-pollination	49	4	0.08	126.00 \pm 58.17	10.08 \pm 4.65
Cross-pollination total	130	127	0.98	116.11 \pm 58.97	113.64 \pm 61.27
Cross-pollination 2016	26	23	0.89	126.57 \pm 53.66	99.65 \pm 57.91
Cross-pollination 2017	48	48	1.00	138.44 \pm 63.47	138.44 \pm 63.47
Cross-pollination 2018	56	56	1.00	98.88 \pm 54.65	98.88 \pm 54.65
Natural conditions total	131	108	0.82	119.08 \pm 66.31	101.98 \pm 58.43
Natural conditions 2016	49	45	0.92	131.19 \pm 55.58	120.69 \pm 51.13
Natural conditions 2017	20	11	0.55	76.73 \pm 73.66	42.20 \pm 40.51
Natural conditions 2018	62	52	0.84	118.04 \pm 70.23	99.15 \pm 58.99

Table 2 (on next page)

Floral visitors and the resources they have taken in *Edmundoa lindenii* flowers.

All records were made in 2015 to 2018 at Serra do Órgãos National Park, southeastern Brazil.

Functional groups: HB = hummingbirds, LB = large bees, SB = small bees. Type of resources:

P = pollen, N = nectar.

Table 2:
Floral visitors and the resources they have taken in *Edmundoa lindenii* flowers.
 All records were made in 2015 to 2018 at Serra do Órgãos National Park, southeastern Brazil.
 Functional groups: HB = hummingbirds, LB = large bees, SB = small bees. Type of resources: P = pollen, N = nectar.

Family	Species	Functional group	Resource
Trochilidae	<i>Amazilia lactea</i> (Lesson, 1832)	HB	N
	<i>Amazilia versicolor</i> (Vieillot, 1818)	HB	N
	<i>Leucocholoris albicollis</i> (Viellot, 1818)	HB	N
	<i>Phaethornis eurynome</i> (Lesson, 1832)	HB	N
	<i>Ramphodon naevius</i> (Dumont, 1818)	HB	N
	<i>Thalurania glaucopis</i> (Gmelin, 1788)	HB	N
	<i>Amazilia fimbriata</i> (Gmelin, 1788)	HB	N
Apidae	<i>Bombus morio</i> (Swederus, 1787)	LB	N/P
	<i>Bombus brasiliensis</i> (Lepeletier, 1835)	LB	N/P
	<i>Euglossa</i> sp.	LB	N
	<i>Trigona spinipes</i> (Fabricius, 1793)	SB	P

Table 3(on next page)

The pollination efficacy of hummingbirds was higher than efficacy of large bees in *Edmundoa lindenii*.

ANOVA results on the pollination efficacy of functional groups of pollinators of *E. lindenii*, measured by seed set after a single visit to the flower at Serra do Órgãos National Park, southeastern Brazil.

Table 3:
The pollination efficacy of hummingbirds was higher than efficacy of large bees in
Edmundoa lindenii.
 ANOVA results on the pollination efficacy of functional groups of pollinators of *E. lindenii*,
 measured by seed set after a single visit to the flower at Serra do Órgãos National Park,
 southeastern Brazil.

Effects	DF	F	p
Reproductive event (year)	2	0.15	0.860
Functional group of pollinators	1	71.07	<0.001
Reproductive event : functional group	1	2.60	0.078

Table 4(on next page)

The reproductive events and the functional groups of flower visitors of *Edmundoa lindonii* had an effect on the relative frequency of visits.

ANOVA results on the relative frequency of floral visits by functional groups during the reproductive events of 2015, 2016, 2017 and 2018 at Serra do Órgãos National Park, southeastern Brazil.

Table 4:
The reproductive events and the functional groups of flower visitors of *Edmundoa lindenii* had an effect on the relative frequency of visits.
 ANOVA results on the relative frequency of floral visits by functional groups during the reproductive events of 2015, 2016, 2017 and 2018 at Serra do Órgãos National Park, southeastern Brazil.

Effects	DF	Deviance	P (> Chi)
Reproductive events (year)	3	0.020	0.999
Functional group of floral visitors	2	26.392	<0.001
Reproductive events : functional group	6	244.562	<0.001