

Trypophobia as an urbanized emotion: Comparative research in ethnic minority regions of China

Siqi Zhu ^{Corresp., 1}, **Kyoshiro Sasaki** ^{2, 3, 4}, **Yue Jiang** ⁵, **Kun Qian** ⁶, **Yuki Yamada** ^{Corresp. 3}

¹ Graduate School of Human-Environment Studies, Kyushu University, Fukuoka, Japan

² Faculty of Science and Engineering, Waseda University, Tokyo, Japan

³ Faculty of Arts and Science, Kyushu University, Fukuoka, Japan

⁴ Japan Society for the Promotion of Science, Tokyo, Japan

⁵ School of Ethnology and Sociology, Yunnan University, Kunming, China

⁶ Institute of Decision Science for a Sustainable Society, Kyushu University, Fukuoka, Japan

Corresponding Authors: Siqi Zhu, Yuki Yamada

Email address: knmoonless@gmail.com, yamadayuk@gmail.com

Trypophobia is a strong emotion of disgust evoked by clusters of holes or round objects (e.g., lotus seed pod). It has become increasingly popular and been studied since 2010s, mainly in the West and Japan. Considering this, trypophobia might be a modern emotion and hence, urbanization possibly plays key roles in trypophobia. To address this issue, we compared the degree of trypophobia between urban and less urban people in China. In an experiment, we asked participants about their degree of discomfort from trypophobic images. The results showed that trypophobia occurred in both groups, although the effect size was larger in urban than less urban people. Moreover, post-experimental interviews and post-hoc analyses revealed that older people in less urban area did not experience as much trypophobia. Our findings suggest that trypophobia links to urbanization and age-related properties.

Trypophobia as an urbanized emotion: Comparative research in ethnic minority regions of China

Siqi Zhu^{1*}, Kyoshiro Sasaki^{2,3,4}, Yue Jiang⁵, Kun Qian⁶, and Yuki Yamada^{2**}

¹ Graduate School of Human-Environment Studies, Kyushu University, Fukuoka, Japan

² Faculty of Arts and Science, Kyushu University, Fukuoka, Japan

³ Faculty of Science and Engineering, Waseda University, Tokyo, Japan

⁴ Japan Society for the Promotion of Science, Tokyo, Japan

⁵ School of Ethnology and Sociology, Yunnan University, Kunming, China

⁶ Institute of Decision Science for a Sustainable Society, Kyushu University, Fukuoka, Japan

Word count: 3068

*Corresponding author 1: Siqi Zhu

Address: Graduate School of Human-Environment Studies, Kyushu University,

744 Motooka, Nishi-ku, Fukuoka 819-0395, Japan

E-mail: knmoonless@gmail.com

**Corresponding author 2: Yuki Yamada

Address: Faculty of Arts and Science, Kyushu University,

744 Motooka, Nishi-ku, Fukuoka 819-0395, Japan

E-mail: yamadayuk@gmail.com

Abstract

Trypophobia is a strong emotion of disgust evoked by clusters of holes or round objects (e.g., lotus seed pod). It has become increasingly popular and been studied since 2010s, mainly in the West and Japan. Considering this, trypophobia might be a modern emotion and hence, urbanization possibly plays key roles in trypophobia. To address this issue, we compared the degree of trypophobia between urban and less urban people in China. In an experiment, we asked participants about their degree of discomfort from trypophobic images. The results showed that trypophobia occurred in both groups, although the effect size was larger in urban than less urban people. Moreover, post-experimental interviews and post-hoc analyses revealed that older people in less urban area did not experience as much trypophobia. Our findings suggest that trypophobia links to urbanization and age-related properties.

Keywords: emotion; cognition; cultural differences; vision; disgust

37

38

Introduction

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

Clusters of holes or round objects can induce strong feelings of disgust. This phenomenon is called *tryphobia* (Abbasi, 2011; Cole & Wilkins, 2013). It increasingly garnered public attention via the Internet and has then been investigated since the 2010s (Abbasi, 2011; Cole & Wilkins, 2013). Accordingly, it seems to be a relatively new emotional phenomenon. Previous studies have revealed the relationship between tryphobia and visual processing. Cole and Wilkins (2013) performed a spectral analysis of tryphobic images (e.g., lotus seed pod) and neutral images (e.g., golf cup) and showed that these images had high contrast energy at midrange spatial frequency in comparison to the neutral images. Their study indicated that spatial frequency information at midrange was involved with tryphobia. Another study revealed that low- and mid-spatial frequency information contributed to tryphobia (Sasaki, Yamada, Kuroki, & Miura, 2017). Based on these findings, tryphobic objects may be processed rapidly and unconsciously; indeed, there is empirical evidence supporting this (Sasaki, Watanabe, & Yamada, 2018; Shirai & Ogawa, 2019). Taken together, early visual processing plays key roles in tryphobia.

The question remains, however: does early visual processing alone contribute to tryphobia? Several studies have discussed cognitive factors of tryphobia. Recent studies have developed the Tryphobia Questionnaire (TQ) and investigated the relationship between tryphobia and personal traits (Chaya, Xue, Uto, Yao, & Yamada, 2016; Le, Cole, & Wilkins, 2015; Imaizumi, Furuno, Hibino, & Koyama, 2016a). In particular, Imaizumi and his colleagues showed that core disgust (i.e., emotional avoidance of pathogen infection) sensitivity and positively predicted the TQ score (Imaizumi, Furuno, Hibino, & Koyama, 2016b). This

finding indicates that disgust contributes to tryphobia, and other findings also support this idea (Kupfer & Le, 2018; Vlok-Barnard & Stein, 2017). Based on these studies, Yamada and Sasaki (2017) proposed the “involuntary protection against dermatosis” (IPAD) hypothesis. According to this hypothesis, tryphobic objects evoke unpleasant emotions because their appearance is associated with dermatosis and, as a result, an avoidance reaction to pathogens is induced. Indeed, Yamada and Sasaki also provided evidence supporting the IPAD hypothesis; the history of skin problems involves tryphobic discomfort. Thus, disgust toward infectious pathogens should involve tryphobia.

Studies on tryphobia have rapidly increased, as we mentioned above. However, these studies were mainly conducted in the West and in Japan. Additionally, numerous cross-cultural studies on emotions show cultural differences in emotional processing (Grossmann, Ellsworth, & Hong, 2012; Grossmann, Karasawa, Kan, & Kitayama, 2014; Hot, Saito, Mandai, Kobayashi, & Sequeira, 2006; Kitayama, Mesquita, & Karasawa, 2006; Masuda, Ellsworth, Mesquita, Leu, Tanida, & Van de Veerdonk, 2008; Tanaka, Koizumi, Imai, Hiramatsu, Hiramoto, & de Gelder, 2010), although these studies mainly addressed general positive and negative emotions. Recently, given technological advances, it has become possible for behavioral data to be collected from less-urbanized areas, which was previously difficult (Takahashi, Oishi, & Shimada, 2018). Considering that tryphobia has become popular and been studied since about 2010, mainly in Western and Japanese culture areas (Abbasi, 2011; Cole & Wilkins, 2013), it might be a modern emotion that possibly involves urbanization, and people in less-urbanized areas might not experience tryphobia. To address this issue, we examined whether tryphobia occurs in less-urbanized areas in China.

In the current study, we focused on two regions in Southwest China, Yunnan Province and Guangxi Zhuang Autonomous Region. Yunnan province has mountains at its southwestern and southern part, including the Ailao and Wuliang Mountains along the southern coast of the Honghe River. In these mountainous areas, several ethnic minorities live a less-urbanized life. For example, Hani people, a mountainous ethnic group in the frontier of the country, are distributed along the Ailao and Wuliang Mountains. Dai people also live in the Honghe region, but are more concentrated in valleys or the relatively flat regions of the Honghe River basin. Guangxi Zhuang Autonomous Region is located at the southeast edge of the Yunnan-Guizhou Plateau. Yao people live in rural areas of the northwest and northeast mountainous and hilly areas. Thus, the present study examined whether people in the less urbanized areas of the Honghe region of Yunnan (i.e., Hani and Dai people) and Qibainong mountainous area in northwest of Guangxi (i.e., Yao people) experience trypophobia. As the control group, we also asked Chinese people in an urbanized area in the Yunnan Province, China and in Fukuoka City, Japan, if they experienced trypophobia. We hypothesized that trypophobia would be more salient as urbanization progressed. Therefore, we predicted that less urban people would experience weaker discomfort from the trypophobic images than would urban people and that, in less urban people, there would be no or small difference in the degree of discomfort between the neutral and trypophobic images.

Methods

Ethics statement

The present study received approval from the psychological research ethics committee of the Faculty of Human-Environment Studies at Kyushu University (approval number: 2018-

002). The experiment was conducted according to the guidelines laid down in the Helsinki declaration. Informed consent was verbally obtained from all participants because some were pre-literate. Participants had the right to withdraw from the experiment at any time without providing any reason.

Participants

We performed a two-way mixed-design analysis of variance (ANOVA) with the image type (tryophobic and neutral) as a within-participant factor and the participant group (less urban and urban people) as a between-participant factor. The required sample sizes were calculated using G*power (Faul, Erdfelder, Lang, & Buchner, 2007). We mainly intended to test whether there is an interaction effect between the image type and the participant group. Thus, we performed a preliminary test to detect the interaction effect ($\alpha = .05$, $1 - \beta = .80$, $f = .25$), and then estimated the sample size to be 34 (i.e., 17 participants per exposure condition). However, a previous study suggests that at least 20 participants are required per group to avoid Type I errors (Simmons, Nelson, & Simonsohn, 2011). Thus, we set the minimal sample size to 40 participants (i.e., 20 per exposure condition) for the statistical analysis.

Thirty-four ethnic minorities living in less-urbanized areas (21 males and 13 females, mean age \pm SEM = 48.9 ± 3.34 years) and 34 students in urban areas (11 males and 23 females, mean age \pm SEM = 23.6 ± 0.36 years) participated in the experiment¹. The ethnic minorities of the less urban group consisted of 10 Hani, 10 Dai, and 14 Yao people. Dai participants' data were collected from the Qimaba town of the Huayao Dai minorities, while those of Hani participants were collected from Hani villages around Lüchun County. Both fields were in Honghe Hani and Yi Autonomous Prefecture of Yunnan Province, China. We also collected data of Yao people from the Qibainong Area of the Yao villages in Guangxi Zhuang

Autonomous Region, China. The data of urban students were collected from the campus of Yunnan University in Kunming, China and Kyushu University in Fukuoka, Japan. This urban group consisted of Han (the ethnic majority in China), Yi, Jingpo, and Zhuang peoples. Though the last three are relative minority groups among Yunnan and Kyushu University students, they are sufficient to be regarded as a control group because they had lived in a city for a long time and were urbanized to a large extent. All of them were naive to the purpose of this experiment and reported having normal vision.

Apparatus and stimuli

The stimuli were presented on a laptop (Dell Inspiron 7460). The resolution was 1920×1080 pixels, and the refresh rate was 100 Hz. The presentation of the stimuli and the collection of data were controlled by a computer. The stimuli were generated using PsychoPy3 (Peirce, 2007) and included 20 tryphobic and 20 neutral images (512×512 pixels) used in previous studies (Le et al., 2015; Sasaki et al., 2017; Yamada & Sasaki, 2017).

Procedure

The participants initiated each trial by pressing the spacebar on a computer keyboard. After the fixation mark was presented for 500 ms, the image stimulus appeared. They had to erase the stimulus by pressing a key when they thought that they had observed the stimulus enough. The rating scales were then presented. The participants were asked to evaluate the degree of discomfort for each image on a nine-point scale that ranged from 1 (strong discomfort) to 9 (strong comfort). Each participant performed 40 trials: two image types (tryphobic and neutral) \times 20 images. Trials were randomized for each participant. Afterward, a post-experimental interview was performed. The interview began with the question of whether participants felt uncomfortable when they saw images of a single hole or clusters of

holes. If they answered “Yes,” the subsequent question was about the part that made them feel uncomfortable. If they answered “No,” the subsequent question was about the feeling they actually experienced when they looked at these pictures and why they felt it.

Analysis

We calculated the average rating scores of the tryphobic and neutral images for each participant. We performed a two-way mixed ANOVA on the rating score with the image type as a within-participant factor and the participant group as a between-participant factor. When the interaction between the image type and participant group was significant, we performed a test of the simple main effects. We set the significance level at $\alpha = .05$ and reported η_p^2 s.

Results

The results are shown in Figure 1. The ANOVA revealed that the main effects of the image type and the participant group were significant (image type: $F(1, 66) = 106.31, p < .001, \eta_p^2 = .62$; participant group: $F(1, 66) = 8.92, p = .004, \eta_p^2 = .12$). Moreover, the interaction was significant ($F(1, 66) = 31.04, p < .001, \eta_p^2 = .32$). The simple main effect of the image type was significant in both the participant groups (less urban people: $F(1, 33) = 16.05, p < .001, \eta_p^2 = .33$; urban people: $F(1, 33) = 96.99, p < .001, \eta_p^2 = .75$). Furthermore, the simple main effect of the participant group was significant for the tryphobic image ($F(1, 66) = 19.29, p < .001, \eta_p^2 = .23$) while it was not significant in the neutral image ($F(1, 66) = 0.24, p = .63, \eta_p^2 = .004$).

After the experiment, participants verbally reported their impressions. Except for simple answers such as “yes” or “uncomfortable” to the questions about whether participants felt uncomfortable, a part of the other detailed reports are shown in Table 1. The full interview reports are available at the data repository (<https://osf.io/wvu8z/>).

--Figure 1 and Table 1 around here--

Discussion

The present study demonstrated that trypophobia occurs in Chinese ethnic minorities in less urban as well as urbanized people. As mentioned above, we hypothesized that urbanization is one of the factors contributing to trypophobia, and to examine this hypothesis, a comparison of the effect sizes of trypophobia on less urban and urban people would be informative. Based on the results for each group, a smaller effect size was found for less urban ($\eta_p^2 = .33$) than for urban people ($\eta_p^2 = .75$). Briefly, trypophobia was salient in urban people in comparison with less urban people. Additionally, the rating score of the trypophobic image was significantly lower in urban than in less urban people, while there was no significant difference in the neutral image scores. Thus, the urban people experienced more discomfort from the trypophobic, but not neutral, images than did the less urban people. These results suggest that trypophobia comes from urbanization.

Does urbanization alone regulate trypophobic responses? This may not be true. There are several differences in attributes and living environments between less urban and urban people; one possibility is a difference in age. In actuality, the participants' age was significantly higher in the less urban than in the urban people (Welch's *t*-test: $t(33.75) = 7.54$, $p < .001$, Cohen's $d = 1.83$). Did the participants' age also link to trypophobia? According to the post-experimental interview, the older less urban people do not experience much trypophobia. Moreover, the post-hoc tests showed that the differences in the rating scores (i.e., subtracting the scores of the trypophobic image from those of the neutral ones) significantly correlated with

the participants' age ($r = -.37$, $p = .002$), indicating that trypophobia weakened with participants' higher age. A similar (but considerably weak) pattern was found in a previous study (Imaizumi et al., 2016a), which used the TQ with Japanese people. Taken together, although we could not clearly conclude which factor contingent to urbanization involves trypophobia, age-related properties are possibly crucial ones.

The most interesting result of the present study was that, as mentioned earlier, even people living in less-urbanized regions strongly experienced trypophobia, suggesting its high prevalence. Here, one question arises: since when did this phenomenon exist? The study of trypophobia has been rapidly increasing since 2000. To the best of our knowledge, the oldest case of a fear of a hole, which seems to be very similar to trypophobia, was described in 1998 (Rufo, 1998). Our urbanization hypothesis can help to answer this question: trypophobia is a phenomenon that became stronger with urbanization, so it is likely that it was not strong enough to be discovered until global urbanization fully progressed. Furthermore, the rapid spread of trypophobia may be related to the development of image processing and information communication technologies (e.g., the Internet): someone experienced trypophobia, then posted this experience on a website (e.g., blog or forum), and as a result, trypophobia spread over the world². These occurrences played a critical role in the creation, processing, and dissemination of trypophobic images. In addition, because of the digital divide (Norris, 2001), the development produced a gap in familiarity with information technology between regions and generations, which supports our urbanized-trypophobia hypothesis: if there is any positive correlation between age and trypophobia intensity, this may be explained by the digital divide.

Because of differences in hygienic conditions, the severity of infectious diseases is significantly higher in less-urban regions than in urban areas (Paddock, 2014; World Health

Organization, 2002). Are these difference in the severity of infectious diseases associated with the present findings? Based on the IPAD hypothesis (Yamada & Sasaki, 2017), tryphobia is assumed to be an avoidance reaction to infectious pathogens because the appearance of tryphobic images is associated with skin diseases. This theory would predict that people in less-urban areas would experience tryphobia more strongly than would people in urban areas; however, this was not the case. Therefore, the difference in the severity of infectious diseases cannot simply account for the difference in tryphobia between urban and less-urban areas. The present findings give rise to a new possibility: infection habituation. People in less-urban areas are frequently in contact with infectious pathogens (Paddock, 2014; World Health Organization, 2002) and as a result, are habituated to them. Such habituation might reduce the fear of infectious diseases and, in turn, weaken the avoidance response to apparent sources of infection, thereby attenuating tryphobia. Further experimentation on this point is thus warranted.

The present study has some limitations worth discussing. First, our experiment was conducted using a laptop computer, but the middle-aged and older people in less-urban area were embarrassed about their performances using such digital experimental device. The difficulty in operation possibly caused increased variance in the data. Moreover, the embarrassment may have somewhat contaminated and biased the emotional responses of comfort in the minority sample, although the neutral condition did not differ between the groups. Second, less-urban participants included those who did not understand Mandarin Chinese, and hence, the experimental instruction sometimes required translation by experimentally naive local guides who used local ethnic languages. This translation may have changed nuances, affecting the results. In future research, these problems may be solved

through non-digital methods such as paper experiments and by inviting local psychological scientists to conduct experiments. Third, there were ethnic differences between urbanized and less-urbanized areas and these differences might have potential effects on our results that have not yet been discovered. Although there is no reasonable hypothesis on the differences in tryphobia among ethnic groups at this time and thus it is less likely that the ethnic differences contaminated our results, controlling these differences might be desirable in future studies.

Fourth, all participants reported that their visual acuity was normal, but their visual function was not actually measured and could not be confirmed. This may also affect the evaluation results. Indeed, provocative visual patterns in the modern urban environments possibly involve visual stress (Wilkins, Penacchio, & Leonards, 2018), which is linked to tryphobia (Imaizumi et al., 2016b). Moreover, as Sasaki and colleagues (2017) revealed, tryphobia is a phenomenon that is dependent on spatial frequency. According to their study, high-frequency components only minorly contributed to the strength of tryphobia. Differences in visual function have often been argued about in studies on cultural differences in visual perception (e.g., Ahluwalia, 1978; Berry, Poortinga, Breugelmans, Chasiotis, & Sam, 2012; Jahoda, 1971). If less-urban participants who participated in this study had visual characteristics biased toward processing of high-frequency components, it is possible that tryphobia as seen in the present results is underestimated.

Last but not least, through this research in the Honghe and Qibainong Areas, we found that the development of tryphobia in less-urban areas, which had been experimented with previously, greatly exceeded the initial expectations. These areas can still have some unique cultural symbols of ethnic minorities, such as residents wearing traditional ethnic costumes and holding traditional ceremonies during festivals. However, in recent times, the wooden buildings

have been transformed into brick buildings and most residents skillfully use mobile phones and electronic payments. These phenomena show that urbanization and modernization are accelerating even in the minority areas along the border. Therefore, the boundaries between urban and less-urban areas in China are gradually dissolving. Nevertheless, the results of the present study also seem to tell us that even in the course of such transformations, the regional differences in terms of trypanophobia still exist. Thus, testing trypanophobia in some more primitive ethnic settlements, such as with the Dulong in Nujiang, Yunnan, or ethnic minority areas in Thailand, Laos, and Myanmar, concerning the difference between these peripheral regions and urban areas is worth exploring in the future.

Conclusions

The present study aimed at testing whether trypanophobia occurs in people in rural areas, especially in Chinese ethnic minorities. Our hypothesis was that people in rural areas experienced weaker trypanophobia than people in urban areas. Consistent with this hypothesis, weaker trypanophobia occurred in rural people, suggesting that trypanophobia is a relatively new emotion that has emerged through urbanization. Future research using more peoples secluded from modern civilizations is needed.

284

285

References

286

Abbasi, J. (2011, July 25). Is tryphobia a real phobia? *Popular Science*. Retrieved from <http://www.popsci.com/tryphobia>

287

288

Ahluwalia, A. (1978). An intra-cultural investigation of susceptibility to 'perspective' and 'non-

289

perspective' spatial illusions. *British Journal of Psychology*, 69(2), 233-241.

290

Berry, J. W., Poortinga, Y. H., Breugelmans, S. M., Chasiotis, A., & Sam, D. L. (2012). *Cross-*

291

cultural psychology: Research and applications. Cambridge: Cambridge University

292

Press. Chicago.

293

Chaya, K., Xue, Y., Uto, Y., Yao, Q., & Yamada, Y. (2016). Fear of eyes: Triadic relation

294

among social anxiety, tryphobia, and dis- comfort for eye cluster. *PeerJ*, 4:e1942.

295

Cole, G. G., & Wilkins, A. J. (2013). Fear of holes. *Psychological Science*, 24, 1980–1985.

296

Faul, F., Erdfelder, E., Lang, A-G., & Buchner, A. (2007). G*Power 3: a flexible statistical

297

power analysis program for the social, behavioral, and biomedical sciences. *Behavior*

298

Research Methods, 39, 175–191.

299

Grossmann, I., Ellsworth, P. C., & Hong, Y. Y. (2012). Culture, attention, and emotion. *Journal*

300

of Experimental Psychology: General, 141, 31–36.

301

Grossmann, I., Karasawa, M., Kan, C., & Kitayama, S. (2014). A cultural perspective on

302

emotional experiences across the life span. *Emotion*, 14, 679–692.

303

Hot, P., Saito, Y., Mandai, O., Kobayashi, T., & Sequeira, H. (2006). An ERP investigation of

304

emotional processing in European and Japanese individuals. *Brain Research*, 1122,

305

171–178.

- 306 Imaizumi, S., Furuno, M., Hibino, H., & Koyama, S. (2016a). Development of the Japanese
307 Version of Trypophobia Questionnaire. *The Japanese Journal of Personality*, 25, 171–
308 173. (in Japanese)
- 309 Imaizumi, S., Furuno, M., Hibino, H., & Koyama, S. (2016b). Trypophobia is predicted by
310 disgust sensitivity, empathic traits, and visual discomfort. *SpringerPlus*, 5, 1–5.
- 311 Jahoda, G. (1971). Retinal pigmentation, illusion susceptibility and space perception.
312 *International Journal of Psychology*, 6(3), 199-207.
- 313 Kitayama, S., Mesquita, B., & Karasawa, M. (2006). Cultural affordances and emotional
314 experience: Socially engaging and disengaging emotions in Japan and the United
315 States. *Journal of Personality and Social Psychology*, 91, 890–903.
- 316 Kupfer, T. R., & Le, A. T. (2018). Disgusting clusters: Trypophobia as an overgeneralised
317 disease avoidance response. *Cognition and Emotion*, 32, 729–741.
- 318 Le, A. T., Cole, G. G., & Wilkins, A. J. (2015). Assessment of tryphobia and an analysis of
319 its visual precipitation. *Quarterly Journal of Experimental Psychology*, 68, 2304–
320 2322.
- 321 Masuda, T., Ellsworth, P. C., Mesquita, B., Leu, J., Tanida, S., & Van de Veerdonk, E. (2008).
322 Placing the face in context: Cultural differences in the perception of facial emotion.
323 *Journal of Personality and Social Psychology*, 94, 365–381.
- 324 Peirce, J. W. (2007). PsychoPy–psychophysics software in python. *Journal of Neuroscience*
325 *Methods*, 162, 8–13.
- 326 Norris, P. (2001). *Digital divide: Civic engagement, information poverty, and the Internet*
327 *worldwide*. Cambridge: Cambridge University Press. Chicago.

- Paddock, C. (2014, November 21). Water and sanitation in rural areas are 'seriously underfunded'. *Medical News Today*. Retrieved from <https://www.medicalnewstoday.com/articles/285848.php>.
- Rufo, M. (1998). La petite fille qui avait peur des trous. *Soins Pédiatrie/Puériculture*, 182, 3.
- Sasaki, K., Watanabe, K., & Yamada, Y. (2018). Invisible but unpleasant: Unconscious emotional processing of tryphobic objects. *PsyArXiv*, psyarxiv.com/5xshq
- Sasaki, K., Yamada, Y., Kuroki, D., & Miura, K. (2017). Trypophobic discomfort is spatial frequency dependent. *Advances in Cognitive Psychology*, 13, 224–231.
- Shirai, R., & Ogawa, H. (2019). Trypophobic images gain preferential access to early visual processes. *Consciousness and Cognition*, 67, 56–68.
- Simmons, J. P., Nelson, L. D., & Simonsohn, U. (2011). False-positive psychology: Undisclosed flexibility in data collection and analysis allows presenting anything as significant. *Psychological Science*, 22, 1359–1366.
- Takahashi, K., Oishi, T., & Shimada, M. (2017). Is 😊 smiling? Cross-cultural study on recognition of emoticon’s emotion. *Journal of Cross-Cultural Psychology*, 48, 1578–1586.
- Tanaka, A., Koizumi, A., Imai, H., Hiramatsu, S., Hiramoto, E., & de Gelder, B. (2010). I feel your voice: Cultural differences in the multisensory perception of emotion. *Psychological Science*, 21, 1259–1262.
- Vlok-Barnard, M., & Stein, D. J. (2017). Trypophobia: An investigation of clinical features. *Revista Brasileira de Psiquiatria*, 39, 337-341.

Wilkins, A.J. Penacchio, O. and Leonards, U. (2018). The built environment and its patterns: a view from the vision sciences. *Journal of Sustainable Design and Applied Research*, 6, 41-48.

World Health Organization. (2002). *The world health report 2002: Reducing risks, promoting healthy life*. World Health Organization.

Yamada, Y., & Sasaki, K. (2017). Involuntary protection against dermatosis: A preliminary observation on trypophobia. *BMC Research Notes*, 10:658.

357

358

Acknowledgements

359 This research was supported by JSPS KAKENHI (15H05709, 16H01866, 17H00875,
360 17H06342, 17J05236, 18H04199, 18K12015, and 19K14482).

361

362

Author Contributions

363 Contributed to conception and design: ZS, KS, QK, YY

364 Contributed to acquisition of data: ZS, JY

365 Contributed to analysis and interpretation of data: ZS, KS, QK, YY

366 Drafted and/or revised the article: ZS, KS, JY, QK, YY

367 Approved the submitted version for publication: ZS, KS, JY, QK, YY

368

369

Competing Interests

370 The authors declare no competing financial interests.

371

372

Data Availability

373 The datasets are available at <https://osf.io/wvu8z/>

374

375

Footnotes

376 ¹Initially, 20 ethnic minorities living in less-urbanized areas and 22 students from Yunnan
377 University (i.e., urban areas) participated in the experiment based on the power analysis. We
378 obtained similar results to that of the main text. According to the reviewer's suggestion, we

added additional data from 14 less urban people from Guangxi and 12 urban people from Kyushu University.

²Reviewer 2 also experienced a similar case.

Figure Legends

Figure 1. The results of the experiment. The error bars indicate standard errors of the mean.

Table 1. Post-experimental interview reports.

Table 1 (on next page)

Post-experimental interviews

1

Area	Ethnicity	Age	Gender	Report
Less urban	Dai	10	Female	The porous images made me feel dizzy.
		41	Female	I felt that many holes were dirty.
		40	Female	I felt a little nauseous and scared of the porous images.
		64	Female	No matter which image I saw, I felt good and did not feel dizzy or uncomfortable.
		72	Male	I did not feel uncomfortable. I evaluated the impression of the images based on whether things inside the images were practically useful or whether they were ornamentally valuable.
		57	Male	It did not matter how many holes the images contained. If the brightness of the image was dark, I felt uncomfortable. On the other hand, its brightness was high, which made me comfortable.
	Hani	42	Female	The images did not induce discomfort regardless of whether these contained many holes. I prefer plaid clothes.
		79	Male	I did not feel uncomfortable. I actually felt that the porous images were better-looking.
		30	Female	The image containing high-dense holes made me dizzy and thus I did not want to look at them.
		55	Male	The images containing fewer holes were good.
Urban	Yi	25	Female	I was very disgusted.
		20	Male	I felt nothing special about the images.
	Jingpo	25	Female	I hated the clusters of small holes, although the holes on the food were good.
	Han	21	Female	I did not feel good.
	Han	18	Male	The porous images were eye-catching, while I felt nothing special about the other images.
	Han	23	Female	When I saw the porous image, I felt as if my scalp was numb and my hair stood up.
	Han	26	Male	The holes would be disgusting if they were randomly arranged and thus looked worm-eaten and like acne. However, if the holes were regularly aligned like beehives, they were good.
	Han	22	Female	It did not matter if the porous images contained artificial objects. However, they were disgusting if they contained too many holes.
	Han	23	Male	Holes on natural things made me nauseous. In particular, when objects in the holes were similar to acne, I felt disgust. The dirty images also made me nauseous.
	Han	23	Male	When lots of holes were regularly aligned, they were disgusting.
	Han	24	Female	I felt nothing special about the images. When I saw the mushroom, I even wanted to eat it.

2

Figure 1

Figure 1.

The results of the experiment. The error bars indicate standard errors of the mean.

