

Comparative transcriptomics characterized the distinct biosynthetic abilities of terpenoid and paeoniflorin biosynthesis in herbaceous peony strains

Baowei Lu¹, Liangjing Cao², Qian Gao², Xuan Wang³, Yongjian Yang¹, Pengming Liu¹, Baoliang Yang¹, Tong Chen⁴, Xin-Chang Li², Qinghua Chen¹, Fengxia An^{Corresp., 1}, Jun Liu^{Corresp. 2}

¹ Bozhou University, Bozhou, Anhui, China

² National Key Facility for Crop Resources and Genetic Improvement, Institute of Crop Science, Chinese Academy of Agricultural Sciences, Beijing, China

³ Department of Biological Sciences, Xian Jiaotong-Liverpool University, Suzhou, China

⁴ National Resource Center for Chinese Materia Medica, Academy of Chinese Medical Sciences, Beijing, China

Corresponding Authors: Fengxia An, Jun Liu

Email address: An_fengxia@163.com, liujun@caas.cn

The herbaceous peony (*Paeonia lactiflora* Pall.) is a perennial flowering plant of the Paeoniaceae species that is widely cultivated for medical and ornamental uses. The monoterpene glucoside paeoniflorin and its derivatives are the active compounds of the *P. lactiflora* roots. However, the gene regulation pathways associated with monoterpene and paeoniflorin biosynthesis in *P. lactiflora* are still unclear. Here, we selected three genotypes of *P. lactiflora* with distinct morphologic features and chemical compositions that were a result of long-term reproductive isolation. We performed an RNA-sequencing experiment to profile the transcriptome changes of the shoots and roots. Using *de novo* assembly analysis, we identified 36,264 unigenes, including 521 genes responsible for encoding transcription factors. We also identified 28,925 unigenes that were differentially expressed in different organs and/or genotypes. Pathway enrichment analysis showed that the *P. lactiflora* unigenes were significantly overrepresented in several secondary metabolite biosynthesis pathways. We identified and profiled 33 genes responsible for encoding the enzymescontrolling the major catalytic reactions in the terpenoid backbone and in monoterpene biosynthesis. Our study identified the candidate genes in the terpenoid biosynthesis pathways, providing useful information for metabolic engineering of *P. lactiflora* intended for pharmaceutical uses and facilitating the development of strategies to improve marker-assist *P. lactiflora* in the future.

Comparative Transcriptomics Characterized the Distinct Biosynthetic Abilities of Terpenoid and Paeoniflorin Biosynthesis in Herbaceous Peony Strains

Baowei Lu^{1,3}, Liangjing Cao², Qian Gao², Xuan Wang^{2,3}, Yongjian Yang¹, Pengming Liu¹, Baoliang Yang¹, Tong Chen⁴, Xin-Chang Li², Qinghua Chen¹, Fengxia An^{1*}and Jun Liu^{2*}

¹Bozhou University, Anhui Province, 236800, China

²National Key Facility for Crop Resources and Genetic Improvement, Institute of Crop Science, Chinese Academy of Agricultural Sciences, Beijing 100081, China

³Department of Biological Sciences, Xian Jiaotong-Liverpool University, Suzhou 215123, China

⁴National Resource Center for Chinese Materia Medica, Academy of Chinese Medical Sciences, Beijing 100700, China

Corresponding Author:

FengxiaAn (An_fengxia@163.com) or Jun Liu (liujun@caas.cn)

Abstract

The herbaceous peony (*Paeonia lactiflora* Pall.) is a perennial flowering plant of the Paeoniaceae species that is widely cultivated for medical and ornamental uses. The monoterpene glucoside paeoniflorin and its derivatives are the active compounds of the *P. lactiflora* roots. However, the gene regulation pathways associated with monoterpene and paeoniflorin biosynthesis in *P. lactiflora* are still unclear. Here, we selected three genotypes of *P. lactiflora* with distinct morphologic features and chemical compositions that were a result of long-term reproductive isolation. We performed an RNA-sequencing experiment to profile the transcriptome changes of the shoots and roots. Using *de novo* assembly analysis, we identified 36,264 unigenes, including 521 genes responsible for encoding transcription factors. We also identified 28,925 unigenes that were differentially expressed among organs and/or genotypes. Pathway enrichment analysis showed that the *P. lactiflora* unigenes were significantly overrepresented in several secondary metabolite biosynthesis pathways. We identified and profiled 33 genes responsible for encoding the enzymes controlling the major catalytic reactions in the terpenoid backbone and in monoterpene biosynthesis. Our study identified the candidate genes in the terpenoid biosynthesis pathways, providing useful information for metabolic engineering of *P. lactiflora* intended for pharmaceutical uses and facilitating the development of strategies to improve marker-assist *P. lactiflora* breeding in the future.

57

58 Introduction

59 The herbaceous peony (*Paeonialactiflora* Pall.) is a flowering plant in the family Paeoniaceae, which is
60 native to Central and eastern Asia (Zhao et al. 2018; Zhao et al. 2017). Its dried root is harvested without the
61 bark in the autumn from plants that are between 3-5 years of age; this harvested material is named Radix
62 Paeoniae Alba or Baishao and is a well-known Chinese herb, used for over 2000 years (He & Dai 2011; Zha
63 et al. 2012). A water/ethanol extract of Radix Paeoniae Alba, now known as Total Glucosides of Peony (TGP),
64 was originally used in the treatment of typhoid (Li et al. 2011). Subsequently, TGP has been widely
65 prescribed for fever, rheumatoid arthritis, hepatitis, muscle cramping and spasms, systemic lupus
66 erythematosus, and dysmenorrhea (Fan et al. 2012; He & Dai 2011; Ji et al. 2013; Mao et al. 2012; Nam et al.
67 2013).

68 Paeoniflorin (C₂₃H₂₈O₁₁, molecular weight = 480.45) is the major medicinal component in *P. lactiflora*
69 roots. In vitro and in vivo studies in animal models have confirmed that TGP, paeoniflorin,
70 benzoylpaeoniflorin, galloylpaeoniflorin and their derivatives, are medicinally active compounds with multiple
71 pharmacological effects (Fan et al. 2012; He & Dai 2011; Zhou & Wink 2018). TGP can inhibit acute and
72 subacute inflammation, an effect which is potentially mediated by the suppression of prostaglandin E₂,
73 leukotriene B₄, and nitric oxide, as well as the intracellular calcium ion concentration (He & Dai 2011; Xu et
74 al. 2016). TGP has been known to protect cells against Ca²⁺ overload and oxidative stress (Zhang et al. 2017).
75 Moreover, the components of TGP, as important immunomodulatory effectors, can regulate the proliferation
76 and apoptosis of lymphocytes and balance the production of proinflammatory cytokines in a dose-dependent
77 manner (He & Dai 2011; Hu et al. 2018). In addition, paeoniflorin and its derivatives were shown to inhibit
78 tumor growth and macrophage-mediated lung metastases (Ou et al. 2011; Wu et al. 2015).

79 Paeoniflorin is a monoterpene glucoside that is biosynthesized from geranyl-pyrophosphate (GPP). GPP
80 is produced via a conversion from the universal terpenoid precursor, Isopentenyl pyrophosphate (IPP). In
81 plants and bacteria, IPP is produced from the two terpene biosynthesis pathways, the mevalonate pathway
82 (MVA), and the 1-deoxy-d-xylulose-5-phosphate/ methyl-erythritol-4-phosphate (DXP/MEP) pathway
83 (Kanehisa et al. 2012; Ren et al. 2009; Xie et al. 2011). The MVA pathway reactions take place in the cytosol

and are catalyzed by enzymes including hydroxyl methylglutaryl-CoA synthase, acetyl-CoA C-acetyltransferase, HMG-CoA reductase, mevalonate kinase, and phosphomevalonate kinase. The DXP/MEP pathway is catalyzed in the plastids. Pyruvate and glyceraldehyde 3-phosphate are converted by 1-deoxy-D-xylulose-5-phosphate synthase and 1-deoxy-D-xylulose-5-phosphate reductoisomerase to 1-deoxy-D-xylulose 5-phosphate and 2-C-methyl-D-erythritol 4-phosphate, respectively. The products are subsequently catalyzed by 4-diphosphocytidyl-2-C-methyl-D-erythritol (CDP-ME) synthase, CDP-ME kinase, and 2-C-methyl-D-erythritol 2,4-cyclodiphosphate synthase to mediate the formation of 2-C-methyl-D-erythritol 2,4-cyclopyrophosphate, which is then converted to (E)-4-hydroxy-3-methyl-but-2-enyl pyrophosphate (HMB-PP) by HMB-PP synthase. HMB-PP is converted to IPP and dimethylallyl pyrophosphate (DMAPP) by HMB-PP reductase. IPP and DMAPP are condensed by geranyl pyrophosphate synthase to produce GPP. In addition to producing a monoterpene, GPP is also a precursor to sesquiterpenes and diterpenes. The conversion from GPP to alpha-terpineol is critical for producing the monoterpene, which is catalyzed by (-)-alpha-terpineol synthase (EC 4.2.3.111, RLC1). Paeoniflorin can be modified by benzoic acid and gallic acid to produce benzoylpaeoniflorin and galloylpaeoniflorin, respectively. Benzoic acid and gallic acid are catalyzed by 3-deoxy-7-phosphoheptulonate synthase, 3-dehydroquinate synthase, and 3-dehydroquinate dehydratase/shikimate dehydrogenase.

With a long history of domestication and selection, the *P. lactiflora* strains used for medical purposes contain high levels of paeoniflorin and are nearly completely infertile due to embryo abortion in their traditional planting regions, like the Bozhou area. Strains have been reproduced through the vegetative propagation of shoots for hundreds of years. Thus these *P. lactiflora* accessions are reproductively isolated and may serve as suitable resources in the investigation of the genetic and molecular basis of the paeoniflorin biosynthesis pathways. Using sequence homology to search for the known sequences and domains, several studies identified a group of genes involved in paeoniflorin biosynthesis. For example, the previous study identified 24 genes, including 8 with full-length cDNA sequences and revealed transcriptional and phylogenetic associations with paeoniflorin biosynthesis (Yuan et al. 2013). However, the gene alias in the paeoniflorin biosynthesis pathways and their expression patterns have not been profiled in *P. lactiflora* strains on a genome-wide basis.

High throughput sequencing technologies have revolutionized genomic and transcriptomic studies. Improved algorithms are now available for *de novo* reassembly of the transcriptome of a non-model plant species without a valid reference genome sequence (Luo et al. 2017b). In this study, we assembled the transcriptome of roots and shoots derived from the 3 strains of *P. lactiflora* using *de novo* assembly analysis. By aligning the assembled genes with public databases, we globally annotated 34,203 unigenes in *P. lactiflora*. These genes may have potential functions. For instance, our analysis identified 521 transcription factor genes. Moreover, we profiled gene expression levels and identified a group of tissue- and/or strain-differential expressed genes using differential expression analysis. Of these genes, we identified the candidate genes associated with the nearly completed terpenoid backbone biosynthesis pathway. We verified the expression pattern of a selective group of candidate genes using the qRT-PCR assay. Our study provides a valuable dataset for updating our understanding of the gene regulatory network underlying paeoniflorin biosynthesis in *P. lactiflora*.

Materials and Methods

Plant materials for RNA-Seq

The *P. lactiflora* Pu-Bang, Xian-Tiao, and Guan-Shang strains were conserved and cultivated under field conditions at Bozhou University, Bozhou, China. The shoots and roots of 3-year old plants were isolated. To avoid circadian effects, we harvested all the tissues in the afternoon of the same day. The samples with three biological replicates were frozen in liquid nitrogen immediately after harvesting. The isolated samples and purified RNA were stored at -80 °C.

RNA extraction, library construction, and Illumina sequencing

The total RNA of individual samples was extracted and purified with the RNeasy® Plant Mini Kit (QIAGEN, Germany). RNA concentration was measured using a Nanodrop 2100 spectrophotometer. RNA Integrity values were checked using an Agilent Bioanalyzer. The samples with a RIN score >8.5 were used for library construction (Liu et al. 2014). The sequencing libraries were generated using a NEB Next Ultra RNA Library Prep Kit for Illumina (New England Biosystems), following the manufacturer's recommendations. Library sequencing was performed on a Hi-Seq X10 system with 150-cycle paired-end sequencing protocol (Illumina).

Bioinformatics analysis of RNA-seq datasets

Referring the methods used in recent studies with modification (Bedre et al. 2016; Lu et al. 2018), we assembled the transcriptome using Trinity (Haas et al. 2013). We aligned read sequences using HISAT2 (Kim et al. 2015). Fragments per kilobase of exon per million fragments mapped of assembled transcripts (FPKM) were calculated and normalized using DESeq2 with global normalization parameters (Anders et al. 2014; Love et al. 2014; Quinn & Chang 2016; Zhang et al. 2014). The sequences of the assembled unigenes were annotated by Trinotate (Haas et al. 2013). Coding regions of unigenes were predicted using Transdecoder (Haas et al. 2013). BLAST v2.7.1 was performed to determine the sequence homology (e-value cutoff of $1e^{-5}$) to UniProt/SwissProt, HMMR v3.1b2, EggNOG v4.5.1, and metabolic pathways were analyzed using the Kyoto Encyclopedia of Genes and Genomes database (Kanehisa et al. 2015). For the unigene with annotated homolog, we used the criterion of FPKM more than 0.8 to filter out the lowly expressed genes; for those without homolog, we used FPKM more than 1 as the cut-off criterion. Differential expression analysis was carried out using DESeq2 (Anders & Huber 2010). Genes with normalized fold-change greater than 2, significance *P*-value less than 0.05, and Benjamini-Hochberg false discovery rate less than 0.1 were considered to be differentially expressed genes.

Quantitative detection of paeoniflorin

We referred to the previous method in order to measure paeoniflorin in samples (Yuan et al. 2013). The dried samples (0.50 g) were weighed and extracted with 50 mL of 50% aqueous methanol with ultrasonication for 30 min. The extracted samples were diluted with 50 mL 50% aqueous methanol and filtered with a 0.45- μ m Millipore filter membrane (Millipore, MA, USA) at 25 °C. We used the Agilent 1200 LC Series (Agilent Technologies, Palo Alto, CA, USA) High Performance Liquid Chromatography (HPLC) system to measure the paeoniflorin abundance. The wavelength was set at 230 nm with a flow rate of 1.0 mL/min at a temperature of 25 °C. Standard compounds were purchased from the National Institutes for Food and Drug Control and the linearity of the standard compounds was checked at seven concentration solutions.

Quantitative RT-PCR

A total of 1 to 2 μ g RNA samples were treated by DNase I (RNeasy plant mini kit) and were reverse transcribed with oligo (dT) primer and SuperScript III (Invitrogen). cDNA samples were analyzed using quantitative PCR with SYBR Premix Ex Taq (Takara) and a Biorad CFX96 real-time PCR system. The

P. lactiflora Actin transcript sequence (Available at <https://www.ncbi.nlm.nih.gov/nuccore/JN105299>) was used as endogenous reference genes to design primers to normalize the expression levels among samples (Qi et al. 2018; Yuan et al. 2013). The qRT-PCR reactions were carried out with three- step cycles (95°C denaturation, 60°C annealing, 72°C amplification, 95°C denaturation, 60°C annealing and 95°C denaturation) and 40 cycles of amplification. We used three technical replicates to produce the average expression levels of the genes relative to that of the reference gene using the $2^{-\Delta\Delta CT}$ method (Livak & Schmittgen 2001). The primers are listed in Supplemental Table 1.

Accession numbers

The fastq-formatted RNA-seq datasets and the sampling information (accession number CRA001881, <https://bigd.big.ac.cn/gsa/browse/CRA001881>) are publically available on the Genome Sequence Archive database (BIG.Members. 2018; Wang et al. 2017).

Results

RNA-Seq and *de novo* assembly of the *P. lactiflora* transcriptome

The *P. lactiflora* Pu-Bang (PB) and Xian-Tiao (XT) accessions are the most widely used herbaceous peony stains for medical uses due to their high levels of paeoniflorin, which is derived mainly from their roots; whereas the Guan-Shang (GS) accession contains less paeoniflorin and is usually cultivated for ornamental uses. The morphological features of the 1- and 3-year old plants of the 3 strains were shown in Figure 1, respectively. We determined the paeoniflorin levels of the isolated root samples using High Performance Liquid Chromatography (HPLC). The results confirmed the accumulated levels of paeoniflorin in PB and XT roots compared with those of GS (Supplemental Fig. 1).

To systematically identify genes and explore the gene expression network underlying paeoniflorin biosynthesis in *P. lactiflora*, we purified the RNA samples with three biological replicates derived from the shoots and roots of 3-year old PB, XT, and GS plants, and carried out the RNA-sequencing analysis using the Illumina paired-end 150 bp protocol. After filtering out the low quality reads, we obtained 775.73 million reads in total (Supplemental table 2). Using Trinity (Haas et al. 2013), we performed *de novo* transcriptome assembly and obtained 36,264 unigenes encoding 72,910 transcripts with 986 nt contig N50 length and 42.7% average GC content. We also checked the completeness of *de novo* assembled Unigenes by aligning the

sequences with SwissPort database (TheUniProtConsortium. 2017). The results showed that 7,131 (50%) out of the 14,165 aligned sequences have more than 40% coverage of known transcripts (Supplemental table 3 & 4).

Functional annotation of expressed genes in the three *P. lactiflora* accessions

Due to embryo abortion and vegetative propagation, the *P. lactiflora* accessions have been undergoing reproductive isolation with a long cultivation history in the Bozhou region and were thought to be genetically distinct (Zhou et al. 2002). However, the genomic evidence supporting this point is still lacking. We measured the expression levels of unigenes by calculating normalized Fragments Per Kilobase of exon per million fragments Mapped (FPKM). The unigenes with an FPKM value higher than 1 in at least one were used to perform hierarchical clustering analysis based on the Pearson correlation efficiency. We analyzed the hierarchical structure of the gene expression levels on a genome-wide basis (Fig. 2A). Most of the biological replicate samples belonged to the same clusters. Principal component analysis results also showed a similar result confirming a high level of reproducibility of the biological replicate samples (Fig. 2B). However, the tissues and strains were distributed in different cluster clades. It was noted that all the clades derived from PB and XT were separated from those of GS, suggesting that the strains for medicinal uses are genetically divergent from the strains used for ornamental purposes, possibly due to the selection and reproductive isolation among the strains.

Next, we predicted the protein-coding potential for the unigenes using the Transdecoder and searched for the annotation for the unigenes by aligning the assembled transcripts and predicted peptide sequences to the protein sequences annotated by Swiss-Prot, TrEMBL, Pfam, and KEGG databases using Trinotate, BLASTP, and BLASTX (Boutet et al. 2016; Camon et al. 2003; El-Gebali et al. 2019; Haas et al. 2013; Kanehisa et al. 2017). In total, we identified 34,203 unigenes containing significant matches to the annotated genes/proteins in at least one database (Fig. 3B). Of them, 28,083 (82%) were reproducibly detected by at least 2 data resources. The annotation information, predicted protein sequences, and FASTA-formatted sequences of these genes were provided in supplemental materials that could serve as a reference annotation for future studies (Supplemental table 3).

Transcription factors (TFs) with DNA-binding domains are the major regulators controlling the activity and specificity of the gene transcription process. We predicted genes encoding TFs in our assembled unigene dataset and identified 521 TF-encoding unigenes belonging to 32 TF families (Fig. 3C). Of these, AP2/ERF, bHLH, FAR1, bZIP, and HB are the most abundant TF genes. The detailed information of the TF genes was provided in supplemental Table 5.

Identification of tissue- and/or strain-differentially expressed unigenes

Next, we searched for the differentially expressed unigenes. For each strain, we compared shoots and roots and extracted lists of root/shoot specific upregulated genes (fold change of expression level > 2 and false discovery rate < 0.05). Our analysis identified 10,125 up-regulated unigenes in roots and 11,911 in shoots. There were 1906 to 3737 unigenes specifically up-regulated in the roots and/or shoots of each strain; whereas only 332 (3%) and 886 (7%) unigenes were up-regulated in the roots and shoots of all the three *P. lactiflora* strains, respectively (Fig. 4). This result showed that a large number of the tissue-specific genes were also strain-specific, which is consistent with the fact that the three *P. lactiflora* strains have been undergoing genetic separation and selection during the last several centuries.

We performed the correlation analysis between the paeoniflorin levels and the expression levels of all the genes in shoots and roots of the three strains (supplemental table 3). Our analysis identified 3952 genes with the Pearson correlation coefficients higher than 0.6. Among the 161 gene with Pearson correlation coefficients higher than 0.9, we found several genes encoding transcription factors, E3 ubiquitin-protein ligase and oxidation-reduction related process, such as GT-2, ABCF4, SKP2A, FREE, GOR, TLP, RMA1H1, HAT5, XBAT31 and DELLA, suggesting transcriptional and post-translational regulation may be highly associated with paeoniflorin accumulation.

Identification of genes in terpenoid and paeoniflorin biosynthesis pathway

To further dissect the regulation pathways of the unigenes, we analyzed the gene list enrichment using the Kyoto Encyclopedia of Genes and Genomes (KEGG) datasets and KOBAS3.0 with hypergeometric testing and Benjamini and Hochberg correction (Kanehisa et al. 2012; Xie et al. 2011). In total, we identified 71 significantly enriched KEGG pathways in *P. lactiflora* (Supplemental Table 6). The list included several

KEGG terms of secondary metabolite biosynthesis, such as: Terpenoid backbone biosynthesis (P -value < 0.0152), Glycerophospholipid metabolism (P -value < 0.0001), Inositol phosphate metabolism (P -value < 0.0001), Pyruvate metabolism (P -value < 0.0002), Seleno compound metabolism (P -value < 0.0298), Ascorbate and aldarate metabolism (P -value < 0.0172) and Butanoate metabolism (P -value < 0.0352). *P. lactiflora* are generally known to have abundant secondary metabolites (Li et al. 2016a; Liu et al. 2017; Ma et al. 2016). Our transcriptome and pathway enrichment results are consistent with the metabolism profiling studies.

The previous studies have identified 19 EST sequences in the terpenoid backbone biosynthesis pathways in *P. lactiflora*, including 7 with full-length cDNA sequences (Yuan et al. 2013). However, the genes in the terpenoid backbone biosynthesis pathway have not been globally profiled and the enzyme catalyzing the initiation step from GPP to monoterpenoid biosynthesis has not been identified in *P. lactiflora* yet. In our datasets, we identified 32 genes with full-length CDSs encoding the enzymes controlling the major catalytic reactions in MVA and MEP pathways (Fig. 5). Compared with the previous study (Yuan et al. 2013), the genes encoding 10 previously reported enzymes were also identified in our study (Fig. 5). Moreover, we identified the unigene (E_H33980_c1_g4, *RLCI*) encoding (-)-alpha-terpineol synthase (EC 4.2.3.111) that can catalyze the conversion from GPP to alpha-terpineol (Kulkarni et al. 2013b), a monoterpene precursor of paeoniflorin. Our analysis identified the genes encoding the enzymes that almost completely catalyzed the reactions from the glycolysis products to the terpenoid backbone and monoterpenoid biosynthesis in *P. lactiflora*.

The high accumulation of paeoniflorin in XT and/or BP roots suggests some genes in the paeoniflorin biosynthesis pathway may be highly expressed in XT and/or PB roots. We analyzed the expression patterns of the aforementioned 33 unigenes (Fig 6). In XT roots, The expression levels of *HMGS*, *MK*, *PMK*, *GGR*, *FPSI* were higher than those in GS roots; whereas the expression levels of *PMK*, *MVD*, *GGR* in PB roots were higher than those in GS roots. The other genes did not clearly reveal XT- and/or PB-root differential expression patterns. Note that XT and PB presented the distinct expression pattern of these genes, which was consistent with the fact that the two strains were genetically isolated with a long history (Zhou et al. 2002). To confirm the expression levels determined by the RNA-seq experiment, we used a quantitative Real-time PCR (qRT-PCR) assay to measure the expression levels of several unigenes (Supplemental Fig. 2). These results

indicated that the different biosynthetic abilities of terpenoid biosynthesis among the strains may be largely contributed to by several master genes, also suggesting that the heterosis in crosses of PB and XT should be considered as a potential approach to breed *P. lactiflora* strains with high paeoniflorin content.

Discussion

The herbaceous peony has a large and complex genome. It was estimated that the genome size of the tree peony, a closely related species of *P. lactiflora*, is about 12.5 Gbp. Thus, assembling the high-quality genome of *P. lactiflora* remains a challenge. Recently, transcriptome studies on a large number of plant species with complex genomes, such as smooth cordgrass (*Spartina alterniflora*) (Bedre et al. 2016), buckwheat (Lu et al. 2018), and *Caragana korshinskii* (Li et al. 2016b), have been carried out. Transcriptome assembly is currently a feasible and cost-efficient technology to globally identify genes in the *P. lactiflora* genome because genome information is unavailable (Luo et al. 2017a; Luo et al. 2017b). In our study, we identified 36,264 unigenes. Using the homolog alignment analysis, we identified a large number of these assembled unigenes encoded in known regulatory domains/motifs, suggesting their molecular functions. Of these, we identified 521 transcription factor genes belonging to 32 families and profiled their expression patterns. Our study identified a large number of previously un-annotated genes in the *P. lactiflora* genome, which could supply valuable molecular information for future functional studies.

Plant terpenoids are widely used as traditional herbal remedies and for their aromatic qualities. Terpenoids are highly abundant in several accessions of *P. lactiflora*, suggesting a high potential for terpenoid biosynthesis in them. However, the terpenoid biosynthetic pathways are not yet fully understood in *P. lactiflora*. Previous studies have identified the 19 EST sequences in the terpenoid backbone biosynthesis pathway in *P. lactiflora* (Yuan et al. 2013). Paeoniflorin is accumulating in roots of 3-year-old plants and lacking in leaves and its accumulation levels are highly differed in different strains (Li et al. 2011; Yuan et al. 2013). The tissue- and strain-specific accumulation pattern of paeoniflorin facilitates us to explore the molecular basis of paeoniflorin biosynthesis pathways. In our study, we carry out a transcriptome analysis and identify 32 unigenes in the pathway and profiled their expression patterns that covered most of the reactions in

the terpenoid backbone biosynthesis pathway. The expression levels of *AACT1* and several transcription factor genes showed specific accumulation in XT/PB roots; whereas most of the genes in the terpene backbone biosynthesis showed differential expression patterns between strains rather between tissues. This result raised two testable hypotheses: 1) The several genes with *AACT1* expression patterns are master genes in the pathway; 2) The root-specific expression patterns of these genes associated with paeoniflorin levels could be found in other strains. The gene sequences and their expression patterns identified in our study could serve as a reference dataset for follow up studies to test their expression specificities in more *P. lactiflora* strains associated with paeoniflorin levels. With this data, the other unidentified gene members and the complete terpenoid backbone biosynthesis pathway in *P. lactiflora* strains can be deciphered in the future, using homologous cloning, phylogenetic analysis, and catalytic kinetics experiments.

Our transcriptome profiling analysis uncovered the tissue- and/or strain-differential expression patterns in the three *P. lactiflora* strains, suggesting the genes in the terpenoid biosynthesis pathway were diversified during the selection process. These results could supply valuable molecular information for future functional studies in *P. lactiflora*. Geranyldiphosphate synthases and farnesyldiphosphate synthase catalyze the important branch-point reactions from GPP to monoterpenes and sesquiterpenes. The catalytic activities of the two enzymes are sensitive to temperature and metal ion concentration (Kulkarni et al. 2013a). Under high temperature and Mg^{2+} rich condition, the monoterpene concentration was significant higher compared with those of sesquiterpenes. In our experiment, we identified several unigenes encoding Geranyldiphosphate synthases and farnesyldiphosphate synthase in *P. lactiflora*. Their catalytic activities could be investigated under different temperature and metal ion conditions in follow-up studies.

It has been known that *P. lactiflora* is highly sensitive to the photoperiod and temperature changes. The accessions grown in the Bozhou region have been undergoing reproductive isolation for a long period. Thus, the genetic background of each individual might be fixed. In our study, we identified a large number of genes in *P. lactiflora* and found gene expression patterns in the terpenoid pathway are highly diversified among different accessions. With the sequences of these genes, the phylogenetic structures and population genetics

backgrounds of germplasm resources could be further investigated to elucidate the domestication and selection process of *P. lactiflora*.

Conclusion

Using the RNA-sequencing protocol, we assembled the exon structures of 36,264 unigenes in the shoots and roots of three 3-year-old *P. lactiflorais* accessions, including 28,925 differentially expressed genes. We systematically annotated their molecular functions by aligning their sequences with those from multiple data resources. 521 transcription factor genes were identified. Of 3,952 genes, the Pearson Correlation Coefficients between the expression levels and the paeoniflorin levels were higher than 0.6. We identified 32 genes encoding the enzymes controlling the major catalytic reactions in MVA and MEP pathways and one gene encoding (-)-alpha-terpineol synthase. These genes contributed nearly complete terpenoid backbone biosynthesis pathways. By profiling gene expression patterns associated with MVP, MEP pathways, and I-PP to monoterpene conversation reactions, we uncovered several unigenes that were highly expressed in the roots of high-paeoniflorin accessions, indicating that the different biosynthetic abilities of terpenoid biosynthesis among the accessions may be largely contributed to by several master genes.

Author Contributions

F. A., B. L., and J. L. designed the experiment. B. L., Y. Y., P. L., Q. C., Y. Z. and B. Y. sampled the biological materials and extracted the RNA and sequenced the libraries. T. C., F. A., J.L. and Y. W. analyzed the datasets. B.L. J.L. and Y.W. prepared the figures and wrote the manuscript. All authors reviewed and approved the final version of the manuscript.

REFERENCE

- Anders S, and Huber W. 2010. Differential expression analysis for sequence count data. *Genome biology* 11:R106. 10.1186/gb-2010-11-10-r106
- Anders S, Pyl PT, and Huber W. 2014. HTSeq—a Python framework to work with high-throughput sequencing data. *Bioinformatics*:btu638.
- Bedre R, Mangu VR, Srivastava S, Sanchez LE, and Baisakh N. 2016. Transcriptome analysis of smooth cordgrass (*Spartina alterniflora* Loisel), a monocot halophyte, reveals candidate genes involved in its adaptation to salinity. *BMC genomics* 17:657-657. 10.1186/s12864-016-3017-3
- BIG.Members. 2018. Database Resources of the BIG Data Center in 2018. *Nucleic acids research* 46:D14-D20. 10.1093/nar/gkx897

- Boutet E, Lieberherr D, Tognolli M, Schneider M, Bansal P, Bridge AJ, Poux S, Bougueleret L, and Xenarios I. 2016. UniProtKB/Swiss-Prot, the Manually Annotated Section of the UniProt KnowledgeBase: How to Use the Entry View. In: Edwards D, ed. *Plant Bioinformatics: Methods and Protocols*. New York, NY: Springer New York, 23-54.
- Camon E, Barrell D, Brooksbank C, Magrane M, and Apweiler R. 2003. The Gene Ontology Annotation (GOA) Project--Application of GO in SWISS-PROT, TrEMBL and InterPro. *Comparative and functional genomics* 4:71-74. 10.1002/cfg.235
- El-Gebali S, Mistry J, Bateman A, Eddy SR, Luciani A, Potter SC, Qureshi M, Richardson LJ, Salazar GA, Smart A, Sonnhammer ELL, Hirsh L, Paladin L, Piovesan D, Tosatto SCE, and Finn RD. 2019. The Pfam protein families database in 2019. *Nucleic Acids Research* 47:D427-D432. 10.1093/nar/gky995
- Fan YF, Xie Y, Liu L, Ho HM, Wong YF, Liu ZQ, and Zhou H. 2012. Paeoniflorin reduced acute toxicity of aconitine in rats is associated with the pharmacokinetic alteration of aconitine. *Journal of ethnopharmacology* 141:701-708. 10.1016/j.jep.2011.09.005
- Haas BJ, Papanicolaou A, Yassour M, Grabherr M, Blood PD, Bowden J, Couger MB, Eccles D, Li B, Lieber M, MacManes MD, Ott M, Orvis J, Pochet N, Strozzi F, Weeks N, Westerman R, William T, Dewey CN, Henschel R, LeDuc RD, Friedman N, and Regev A. 2013. De novo transcript sequence reconstruction from RNA-seq using the Trinity platform for reference generation and analysis. *Nature protocols* 8:1494-1512. 10.1038/nprot.2013.084
- He DY, and Dai SM. 2011. Anti-inflammatory and immunomodulatory effects of paeonia lactiflora pall., a traditional chinese herbal medicine. *Frontiers in pharmacology* 2:10. 10.3389/fphar.2011.00010
- Hu B, Xu G, Zhang X, Xu L, Zhou H, Ma Z, Shen X, Zhu J, and Shen R. 2018. Paeoniflorin Attenuates Inflammatory Pain by Inhibiting Microglial Activation and Akt-NF-kappaB Signaling in the Central Nervous System. *Cellular physiology and biochemistry : international journal of experimental cellular physiology, biochemistry, and pharmacology* 47:842-850. 10.1159/000490076
- Ji Y, Wang T, Wei ZF, Lu GX, Jiang SD, Xia YF, and Dai Y. 2013. Paeoniflorin, the main active constituent of Paeonia lactiflora roots, attenuates bleomycin-induced pulmonary fibrosis in mice by suppressing the synthesis of type I collagen. *Journal of ethnopharmacology* 149:825-832. 10.1016/j.jep.2013.08.017
- Kanehisa M, Furumichi M, Tanabe M, Sato Y, and Morishima K. 2017. KEGG: new perspectives on genomes, pathways, diseases and drugs. *Nucleic Acids Research* 45:D353-D361. 10.1093/nar/gkw1092
- Kanehisa M, Goto S, Sato Y, Furumichi M, and Tanabe M. 2012. KEGG for integration and interpretation of large-scale molecular data sets. *Nucleic acids research* 40:D109-114. 10.1093/nar/gkr988
- Kanehisa M, Sato Y, Kawashima M, Furumichi M, and Tanabe M. 2015. KEGG as a reference resource for gene and protein annotation. *Nucleic acids research* 44:D457-D462.
- Kim D, Langmead B, and Salzberg SL. 2015. HISAT: a fast spliced aligner with low memory requirements. *Nat Methods* 12:357-360. 10.1038/nmeth.3317
- nmeth.3317 [pii]
- Kulkarni R, Pandit S, Chidley H, Nagel R, Schmidt A, Gershenzon J, Pujari K, Giri A, and Gupta V. 2013a. Characterization of three novel isoprenyl diphosphate synthases from the terpenoid rich mango fruit. *Plant physiology and biochemistry : PPB* 71:121-131. 10.1016/j.plaphy.2013.07.006
- Kulkarni R, Pandit S, Chidley H, Nagel R, Schmidt A, Gershenzon J, Pujari K, Giri A, and Gupta V. 2013b. Characterization of three novel isoprenyl diphosphate synthases from the terpenoid rich mango fruit.

- 398 *Plant Physiology and Biochemistry* 71:121-131. <https://doi.org/10.1016/j.plaphy.2013.07.006>
- 399 Li B, Bhandari DR, Rompp A, and Spengler B. 2016a. High-resolution MALDI mass spectrometry imaging of
- 400 gallotannins and monoterpene glucosides in the root of *Paeonia lactiflora*. *Scientific reports* 6:36074.
- 401 10.1038/srep36074
- 402 Li J, Chen CX, and Shen YH. 2011. Effects of total glucosides from paeony (*Paeonia lactiflora* Pall) roots on
- 403 experimental atherosclerosis in rats. *Journal of ethnopharmacology* 135:469-475.
- 404 10.1016/j.jep.2011.03.045
- 405 Li S, Fan C, Li Y, Zhang J, Sun J, Chen Y, Tian C, Su X, Lu M, Liang C, and Hu Z. 2016b. Effects of drought and salt-
- 406 stresses on gene expression in *Caragana korshinskii* seedlings revealed by RNA-seq. *BMC genomics*
- 407 17:200-200. 10.1186/s12864-016-2562-0
- 408 Liu D, Wang D, Qin Z, Zhang D, Yin L, Wu L, Colasanti J, Li A, and Mao L. 2014. The SEPALLATA MADS - box protein
- 409 SLMBP21 forms protein complexes with JOINTLESS and MACROCALYX as a transcription activator for
- 410 development of the tomato flower abscission zone. *The Plant Journal* 77:284-296.
- 411 Liu P, Xu Y, Yan H, Chen J, Shang E-X, Qian D-W, Jiang S, and Duan J-A. 2017. Characterization of molecular
- 412 signature of the roots of *Paeonia lactiflora* during growth. *Chinese Journal of Natural Medicines* 15:785-
- 413 793. [https://doi.org/10.1016/S1875-5364\(17\)30110-3](https://doi.org/10.1016/S1875-5364(17)30110-3)
- 414 Livak KJ, and Schmittgen TD. 2001. Analysis of relative gene expression data using real-time quantitative PCR and
- 415 the 2- $\Delta\Delta CT$ method. *methods* 25:402-408.
- 416 Love MI, Huber W, and Anders S. 2014. Moderated estimation of fold change and dispersion for RNA-seq data with
- 417 DESeq2. *Genome biology* 15:550.
- 418 Lu Q-H, Wang Y-Q, Song J-N, and Yang H-B. 2018. Transcriptomic identification of salt-related genes and de novo
- 419 assembly in common buckwheat (*F. esculentum*). *Plant Physiology and Biochemistry* 127:299-309.
- 420 <https://doi.org/10.1016/j.plaphy.2018.02.001>
- 421 Luo J, Duan J, Huo D, Shi Q, Niu L, and Zhang Y. 2017a. Transcriptomic Analysis Reveals Transcription Factors
- 422 Related to Leaf Anthocyanin Biosynthesis in *Paeonia qiui*. *Molecules* 22. 10.3390/molecules22122186
- 423 Luo J, Shi Q, Niu L, and Zhang Y. 2017b. Transcriptomic Analysis of Leaf in Tree Peony Reveals Differentially
- 424 Expressed Pigments Genes. *Molecules* 22. 10.3390/molecules22020324
- 425 Ma X, Chi YH, Niu M, Zhu Y, Zhao YL, Chen Z, Wang JB, Zhang CE, Li JY, Wang LF, Gong M, Wei SZ, Chen C, Zhang L,
- 426 Wu MQ, and Xiao XH. 2016. Metabolomics Coupled with Multivariate Data and Pathway Analysis on
- 427 Potential Biomarkers in Cholestasis and Intervention Effect of *Paeonia lactiflora* Pall. *Frontiers in*
- 428 *pharmacology* 7:14. 10.3389/fphar.2016.00014
- 429 Mao QQ, Ip SP, Xian YF, Hu Z, and Che CT. 2012. Anti-depressant-like effect of peony: a mini-review.
- 430 *Pharmaceutical biology* 50:72-77. 10.3109/13880209.2011.602696
- 431 Nam KN, Yae CG, Hong JW, Cho DH, Lee JH, and Lee EH. 2013. Paeoniflorin, a monoterpene glycoside, attenuates
- 432 lipopolysaccharide-induced neuronal injury and brain microglial inflammatory response. *Biotechnology*
- 433 *letters* 35:1183-1189. 10.1007/s10529-013-1192-8
- 434 Ou TT, Wu CH, Hsu JD, Chyau CC, Lee HJ, and Wang CJ. 2011. *Paeonia lactiflora* Pall inhibits bladder cancer growth
- 435 involving phosphorylation of Chk2 in vitro and in vivo. *Journal of ethnopharmacology* 135:162-172.
- 436 10.1016/j.jep.2011.03.011
- 437 Qi Z, Zhang Z, Wang Z, Yu J, Qin H, Mao X, Jiang H, Xin D, Yin Z, Zhu R, Liu C, Yu W, Hu Z, Wu X, Liu J, and Chen Q.
- 438 2018. Meta-analysis and transcriptome profiling reveal hub genes for soybean seed storage composition

- during seed development. *Plant, cell & environment*. 10.1111/pce.13175
- Quinn JJ, and Chang HY. 2016. Unique features of long non-coding RNA biogenesis and function. *Nat Rev Genet* 17:47-62. 10.1038/nrg.2015.10
- Ren ML, Zhang X, Ding R, Dai Y, Tu FJ, Cheng YY, and Yao XS. 2009. Two new monoterpene glucosides from *Paeonia lactiflora* Pall. *Journal of Asian natural products research* 11:670-674. 10.1080/10286020902980087
- TheUniProtConsortium. 2017. UniProt: the universal protein knowledgebase. *Nucleic acids research* 45:D158-D169. 10.1093/nar/gkw1099
- Wang Y, Song F, Zhu J, Zhang S, Yang Y, Chen T, Tang B, Dong L, Ding N, Zhang Q, Bai Z, Dong X, Chen H, Sun M, Zhai S, Sun Y, Yu L, Lan L, Xiao J, Fang X, Lei H, Zhang Z, and Zhao W. 2017. GSA: Genome Sequence Archive^{sup}. *Genomics, proteomics & bioinformatics* 15:14-18. 10.1016/j.gpb.2017.01.001
- Wu Q, Chen GL, Li YJ, Chen Y, and Lin FZ. 2015. Paeoniflorin inhibits macrophage-mediated lung cancer metastasis. *Chinese journal of natural medicines* 13:925-932. 10.1016/S1875-5364(15)30098-4
- Xie C, Mao X, Huang J, Ding Y, Wu J, Dong S, Kong L, Gao G, Li CY, and Wei L. 2011. KOBAS 2.0: a web server for annotation and identification of enriched pathways and diseases. *Nucleic acids research* 39:W316-322. 10.1093/nar/gkr483
- Xu W, Zhao Y, Qin Y, Ge B, Gong W, Wu Y, Li X, Xu P, and Xue M. 2016. Enhancement of Exposure and Reduction of Elimination for Paeoniflorin or Albiflorin via Co-Administration with Total Peony Glucosides and Hypoxic Pharmacokinetics Comparison. *Molecules* 21. 10.3390/molecules21070874
- Yuan Y, Yu J, Jiang C, Li M, Lin S, Wang X, and Huang L. 2013. Functional diversity of genes for the biosynthesis of paeoniflorin and its derivatives in *Paeonia*. *International Journal of Molecular Sciences* 14:18502-18519. 10.3390/ijms140918502
- Zha LP, Cheng ME, and Peng HS. 2012. Identification of ages and determination of paeoniflorin in roots of *Paeonia lactiflora* Pall. From four producing areas based on growth rings. *Microscopy research and technique* 75:1191-1196. 10.1002/jemt.22048
- Zhang Y, Qiao L, Xu W, Wang X, Li H, Chu K, and Lin Y. 2017. Paeoniflorin Attenuates Cerebral Ischemia-Induced Injury by Regulating Ca(2+)/CaMKII/CREB Signaling Pathway. *Molecules* 22. 10.3390/molecules22030359
- Zhang YC, Liao JY, Li ZY, Yu Y, Zhang JP, Li QF, Qu LH, Shu WS, and Chen YQ. 2014. Genome-wide screening and functional analysis identify a large number of long noncoding RNAs involved in the sexual reproduction of rice. *Genome Biol* 15:512. 10.1186/s13059-014-0512-1
- Zhao D, Wang R, Liu D, Wu Y, Sun J, and Tao J. 2018. Melatonin and Expression of Tryptophan Decarboxylase Gene (TDC) in Herbaceous Peony (*Paeonia lactiflora* Pall.) Flowers. *Molecules* 23. 10.3390/molecules23051164
- Zhao D, Wei M, Shi M, Hao Z, and Tao J. 2017. Identification and comparative profiling of miRNAs in herbaceous peony (*Paeonia lactiflora* Pall.) with red/yellow bicoloured flowers. *Scientific reports* 7:44926. 10.1038/srep44926
- Zhou H, Hu S, Guo B, Feng X, Yan Y, and Li JJYxbApS. 2002. A study on genetic variation between wild and cultivated populations of *Paeonia lactiflora* Pall. 37:383-388.
- Zhou JX, and Wink M. 2018. Reversal of Multidrug Resistance in Human Colon Cancer and Human Leukemia Cells by Three Plant Extracts and Their Major Secondary Metabolites. *Medicines* 5. 10.3390/medicines5040123

Figure 1

The developmental features of the three *P. lactifloras* strains

(A-C) shows the 30-day-old plants of Xian-Tiao (XT), Pu-Bang (PB), and Guan-Shang (GS) strains, which were grown from the shoots (D-F) isolated from 3-year-old plants. (G-I) and (J-L) show the leaves and roots of 3 year-old plants, respectively.




Figure 2

Paeoniflorin levels in the three *P.lactiflora* accessions

Bars show standard deviations of five biological replicates by HPLC system.




Figure 3

Functional annotation of the assembled unigenes

(A) The length distribution of the assembled unigenes. (B) Functional annotation of unigenes using the 4 databases. (C) Transcription factor genes identified in *P. lactiflora*.





Figure 4

Identification of the differentially expressed genes

The differentially expressed genes in roots (A) and shoots (B) of the three *P. lactiflora* accessions.

A Up-regulated genes in roots



B Up-regulated genes in shoots




Figure 5

The assembled genes encoding enzymes in terpenoid backbone and monoterpene biosynthesis pathways

The identified *P. lactiflora* enzymes are highlighted by red rectangles. The identified enzymes with cDNA and/or Expressed Sequence Tag evidence are indicated by arrows.




Figure 6

The expression levels of the genes encoding enzymes in terpenoid backbone and monoterpene biosynthesis pathways in shoot and root samples of the three *P. lactiflora* accessions

The heatmap plot presents the mean value of expression levels of the three biological replicates. The unigene ID, enzyme id and symbols are given according to their expression levels. S and R give shoots and roots respectively.

