

Linking diffuse radiation and ecosystem productivity of a desert steppe ecosystem

Cheng Li^{1,2}, Xin Jia^{1,2}, Jingyong Ma^{1,2}, Peng Liu^{1,2}, Ruizhi Yang^{1,2}, Yujie Bai^{1,2}, Muhammad Hayat^{1,2}, Jinglan Liu³ and Tianshan Zha^{1,2}

¹Yanchi Research Station, School of Soil and Water Conservation, Beijing Forestry University, Beijing, China

²Key Laboratory of State Forestry Administration on Soil and Water Conservation, Beijing Forestry University, Beijing, China

³School of Ecology and Nature Conservation, Beijing Forestry University, Beijing, China

ABSTRACT

Radiation components have distinct effects on photosynthesis. In the desert steppe ecosystem, the influence of diffuse radiation on carbon fixation has not been thoroughly explored. We examined this diffusion and its effect on ecosystem productivity was examined during the growing season from 2014 to 2015 on the basis of eddy covariance measurements of CO₂ exchange in a desert steppe ecosystem in northwest China. Our results indicated that the gross ecosystem production (GEP) and diffuse photosynthetically active radiation (PAR_{dif}) peaked when the clearness index (CI) was around 0.5. The maximum canopy photosynthesis (P_{max}) under cloudy skies (CI < 0.7) was 23.7% greater than under clear skies (CI ≥ 0.7). When the skies became cloudy in the desert steppe ecosystem, PAR_{dif} had a greater effect on GEP. Additionally, lower vapor pressure deficits (VPD ≤ 1 kPa), lower air temperatures (T_a ≤ 20 °C), and non-stressed water conditions (REW ≥ 0.4) were more conducive for enhanced ecosystem photosynthesis under cloudy skies than under clear skies. This may be due to the comprehensive effects of VPD and T_a on stomatal conductance. We concluded that cloudiness can influence diffuse radiation components and that diffuse radiation can increase the ecosystem production of desert steppe ecosystems in northwest China.

Submitted 15 August 2019

Accepted 2 April 2020

Published 5 May 2020

Corresponding author

Jinglan Liu, liujl66@bjfu.edu.cn

Academic editor

Leonardo Montagnani

Additional Information and
Declarations can be found on
page 14

DOI 10.7717/peerj.9043

© Copyright
2020 Li et al.

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

Subjects Ecosystem Science, Biosphere Interactions, Climate Change Biology, Biogeochemistry, Environmental Impacts

Keywords Diffuse radiation, Clearness index, Gross ecosystem production, Desert steppe ecosystem

INTRODUCTION

Gross ecosystem production (GEP), the overall photosynthetic fixation of carbon per unit of space and time (*Monteith, 1972; Monteith, 1977*), is the determining factor in biogeochemical cycles of terrestrial ecosystems (*Lin et al., 2017*). The primary influence on daytime carbon uptake in terrestrial ecosystems during the growing season is solar radiation (*Guan et al., 2006; Bai et al., 2012*). The incoming solar radiation's quantity and quality composition can affect GEP (*Gu et al., 2003; Bai et al., 2012*). Previous studies have indicated that atmospheric cloud content changes can affect solar radiation on the ground surface and can balance the direct and diffuse solar radiation components (*Gu et al., 2003; Niyogi et al., 2007; Urban et al., 2007; Zhang et al., 2011a; Zhang et al.,*

2011b; Gao et al., 2018). Since plant canopies can more efficiently use diffuse radiation for photosynthesis than direct radiation, increased diffuse radiation under cloudy sky conditions can significantly enhance GEP (Goudriaan, 1977; Gu et al., 2002; Farquhar, Roderick & Pinatubo, 2003; Gu et al., 2003; Alton, North & Los, 2007b; Mercado et al., 2009; Tong et al., 2017). Under cloudy sky conditions, the total and direct radiation received by the canopy decreases, while the amount of diffuse radiation increases. The increased diffuse radiation more evenly distributes radiation throughout the canopy (Roderick et al., 2001; He et al., 2013), improving leaf photosynthesis (Urban et al., 2007; Knohl & Baldocchi, 2008; Mercado et al., 2009; Still et al., 2009). Therefore, canopy photosynthesis under cloudy skies with more diffuse radiation was more effective than under clear skies (Gu et al., 2002; Farquhar, Roderick & Pinatubo, 2003; Alton, North & Los, 2007b; Mercado et al., 2009; Tong et al., 2017). How photosynthesis responds to different sky conditions in forests (Gu et al., 1999; Rocha et al., 2004; Dengel & Grace, 2010; Zhang et al., 2010; Urban et al., 2012; Xu et al., 2018; Park et al., 2018) and grassland ecosystems (Jing et al., 2010; Zhang et al., 2011b; Bai et al., 2012) has been researched in previous studies. Due to smaller leaf area index values and the weaker photosynthesis abilities for low canopy ecosystems, the positive effects of diffuse radiation may be reduced (Letts, Lafleur & Roulet, 2005) or eliminated (Kanniah, Beringer & Hutley, 2013). However, other studies have found that diffuse radiation enhances the carbon sink in grassland ecosystems (Zhang et al., 2011b; Bai et al., 2012). A previous study of 23 sampled sites (including grasslands, farmlands, and forests) found that light use efficiency can be improved under high diffuse radiation condition, but it was not significantly different across vegetation types in the great plains of the southern United States (Wang, Dickinson & Liang, 2008). Ultimately, our understanding of diffuse radiation on ecosystem carbon fixation remains limited because the effects differ across vegetation types. Additionally, diffuse radiation interacts with other environmental factors, such as temperature and moisture conditions (Cheng et al., 2015).

With increases in climate change and variability, China has seen a rising trend in diffuse radiation (Mercado et al., 2009; Ren et al., 2013). Previous studies have shown that China's diffuse radiation increased by $7.03 \text{ MJ m}^{-2} \text{ yr}^{-1}$ annually from 1981 to 2010 (Ren et al., 2013). Over the past 15 years, Northwest China's cloud coverage has reduced, but the cloud optical depth has increased (Ren et al., 2013). The question now is how diffuse radiation affects the GEP of the desert steppe ecosystem, and whether cloudiness positively or negatively affects the GEP.

We hypothesized that diffuse radiation positively affects a steppe ecosystem's GEP under different environmental conditions. We used the eddy covariance technique to measure the CO_2 exchange and to examine the effect of diffuse radiation on the GEP in a desert steppe ecosystem at the southern edge of the Mu Us desert from 2014 to 2015. Our research aimed to determine the relationship between diffuse radiation and canopy productivity and to examine whether cloud-induced changes in radiation components enhance canopy productivity. Our objectives were to: (1) quantify the effect of cloudiness on the light response process, (2) examine how GEP changes with diffuse radiation, and (3) explore the relationship between environmental factors and GEP under different sky conditions.




Figure 1 The distribution of the study flux site. The location of the study flux site (A) and (B), and vegetation condition (C). Map data ©2019 Google.

Full-size DOI: 10.7717/peerj.9043/fig-1

MATERIAL AND METHODS

Site description

Our research was conducted in a desert steppe ecosystem on the southern edge of the Mu Us desert ($37^{\circ}42'31''\text{N}$, $107^{\circ}13'47\text{E}$, 1,560 m above sea level) in Ningxia, Northwest China (Fig. 1). The long-term mean air temperature (1954–2004) was 8.1°C and the area is frost-free for an average of 165 days a year (Wang et al., 2014). Annual precipitation is 287 mm, 62% of which falls between July and September (Feng et al., 2013; Jia et al., 2014). Annual incoming shortwave radiation is $1.4 \times 10^5 \text{ J cm}^{-2}$ (Gong et al., 2016), and mean annual potential evapotranspiration is 2,024 mm (Jia et al., 2014). The site's ground is flat with a homogeneous underlying surface. Vegetation is dominated by C_3 herbaceous plants: *Leymus secalinus*, *Pennisetum flaccidum*, *Stipa breviflora*, *Cleistogenes squarrosa*, and a scattered encroachment of the C_3 shrub species *Artemisia ordosica*.

CO_2 flux and meteorological measurements

We used an eddy covariance (EC) system, which we mounted at a 4.2 m height above the ground, to measure the CO_2 exchange between vegetation and atmosphere (Xie et al., 2015). The EC system consisted of a closed-path infrared gas analyzer (IRGA; model LI-7200, LI-COR Biosciences, Lincoln, NE, USA) and a sonic anemometer (WindMasterTM Pro, Gill Instruments Ltd, Lyminster, England). We used the gas analyzer to measure fluctuations in CO_2 and water vapor concentrations. The anemometer was used to measure fluctuations in wind speed, direction, and sonic temperature. We calibrated the LI-7200 every three months using 99.99% nitrogen gas to calibrate zeros for both CO_2 and water vapor, and a

650 ppm CO₂ standard and a dew point generator (LI-610, LI-COR Inc., USA) to calibrate the span for CO₂ and water vapor, respectively. Wind speed, sonic virtual temperature, CO₂ concentrations, and H₂O concentrations were sampled at 10 Hz, and the data were stored in a data logger (LI-7550, LI-COR Inc., USA). Additionally, due to the short length of the canopy (1.4 m), the canopy's CO₂ storage was not used to estimate net ecosystem production (NEP) (Zhang *et al.*, 2007). The CO₂ storage term tended to be close to zero when using daily and annual time scales (Baldocchi, 2003; Jia *et al.*, 2014).

Simultaneously, we measured above-canopy meteorological variables at the top of the tower each half hour. Incident photosynthetically active radiation (PAR in $\mu\text{mol m}^{-2} \text{s}^{-1}$) was measured using LI-190SA quantum sensors (LI-COR Biosciences, Lincoln, NE, USA). Air temperature (T_a in °C) and relative humidity (RH) were measured using a thermohygrometer (HMP155A, Vaisala, Finland). The vapor pressure deficit (VPD) was calculated accordingly from T_a and RH. Precipitation (mm) was determined using a tipping bucket rain gauge (TE-525 MM, Campbell Scientific, UT, USA). Soil temperature (T_s in °C) and volumetric soil water content (SWC) profiles at 10 cm depths below ground adjacent to the EC tower were monitored using ECH₂O-5TE sensors (Decagon Devices Inc., Pullman, WA, USA). We stored all meteorological variables in data loggers (CR200X and CR3000, Campbell Scientific Inc., USA).

Data processing

We used the post-processing software Eddypro 4.0.0 to compute flux covariance from the raw data. The mean covariance between fluctuations in vertical wind speed and CO₂ concentration was determined by the net ecosystem CO₂ exchange (NEE, $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$) flux between the ecosystem and atmosphere. Downward fluxes values were considered as negative and upward fluxes values as positive. Our calculations included a 3-D coordinate rotation, spike detection, and checks for instantaneous records exceeding realistic absolute limits. In addition, we used air density fluctuations to correct the CO₂ fluxes (Webb, Pearman & Leuning, 1980). Nighttime data with friction velocities below the 0.18 m s⁻¹ threshold were removed to eliminate underestimating stable stratification (Jia *et al.*, 2014). The friction velocity threshold was estimated following the China FLUX standard method (Zhu *et al.*, 2006).

We chose a gap-filling model based on the magnitude and bias of residuals and the stability of model-parameter estimates (Xie *et al.*, 2016). Small gaps (≤ 2 h) were filled using linear interpolation, while larger gaps were filled during the night (PAR < 5 $\mu\text{mol m}^{-2} \text{ s}^{-1}$) according to a Q₁₀-model (Xie *et al.*, 2015):

$$NEE_{\text{night}} = Re_{10} \times Q_{10}^{(T_s - 10)/10} \quad (1)$$

where NEE_{night} is the nighttime NEE, T_s is the soil temperature at a 10-cm depth, Re_{10} is ecosystem respiration at $T_s = 10$ °C, and Q_{10} is the temperature sensitivity of Re .

Daytime ecosystem respiration (Re) was estimated from the nighttime NEE-based calibration of Eq. (1) using monthly values, assuming consistent temperature sensitivity between nighttime and daytime exchanges. NEP was estimated as—NEE. Once daytime

estimates of Re were available, gaps in daytime NEE were filled with:

$$NEE = Re - \frac{\alpha' \times PAR \times P_{max'}}{\alpha' \times PAR + P_{max'}} \quad (2)$$

where α' is the apparent quantum yield ($\mu\text{mol CO}_2 \mu\text{mol PAR}^{-1}$), PAR is photosynthetically active radiation in $\mu\text{mol m}^{-2} \text{s}^{-1}$, and $P_{max'}$ is the maximum apparent photosynthetic capacity of the canopy ($\mu\text{mol CO}_2 \text{m}^{-2} \text{s}^{-1}$). Off-season Re was considered as 24 h NEE fluxes. GEP was estimated as:

$$GEP = Re + NEP \quad (3)$$

The gaps in vapor pressure deficit (VPD, kPa) were filled from the gap-filled T_a and RH records above the canopy. T_s and T_a value gaps were filled using the mean diurnal variation (MDV) method (Jia et al., 2016), while missing PAR values were empirically filled with half-hourly PAR data from a meteorological tower about 3 km east. Other 30-min gaps <1.5 h in meteorological variables with underlying diurnal cycles were filled using linear interpolation, and longer gaps (i.e., gaps ≥ 1.5 h) were filled using the Mean Diurnal Variation method (Falge et al., 2001).

Relative extractable soil water (REW) was calculated using Eq. (4) below, with REW <0.4 and REW ≥ 0.4 representing soil water stress and non-stress, respectively (Granier et al., 1999).

$$REW = \frac{SWC - SWC_{min}}{SWC_{max} - SWC_{min}} \quad (4)$$

where SWC_{max} and SWC_{min} are the maximum and minimum soil water content measured at 10 cm depths, respectively.

Defining diffuse PAR and sky conditions

Diffuse PAR was calculated based on the clearness index (CI) and solar elevation angle (β) (Gu et al., 1999). The corresponding equations were as follows:

$$PAR_{dif} = PAR_{tot} \frac{[1 + 0.3(1 - q^2)]q}{1 + (1 - q^2)\cos^2(90^\circ - \beta)\cos^3\beta} \quad (5a)$$

$$q = (S_f/S_e)/CI \quad (5b)$$

When $0 \leq CI \leq 0.3$, constraint: $S_f/S_e \leq CI$,

$$S_f/S_e = CI(1.020 - 0.254CI + 0.0123 \sin\beta) \quad (5c)$$

When $0.3 < CI \leq 0.78$, constraint: $0.1CI \leq S_f/S_e \leq 0.97CI$,

$$S_f/S_e = CI(1.400 - 1.749CI + 0.177\sin\beta) \quad (5d)$$

When $0.78 < CI$, constraint: $S_f/S_e \geq CI$,

$$S_f/S_e = CI(0.486CI - 0.182 \sin\beta) \quad (5e)$$

where PAR_{tot} is the total PAR, PAR_{dif} is the diffuse PAR, and S_f is the total diffuse radiation received by the horizontal plane of the Earth's surface (W m^{-2}).

CI is defined as the ratio of the global solar radiation (S) received at the Earth's surface to the extraterrestrial irradiance (S_e) in a plane parallel to the Earth's surface ([Gu et al., 1999](#)), such that:

$$CI = \frac{S}{S_e} \quad (5f)$$

$$S_e = S_{sc}[1 + 0.033\cos(360t_d/365)] \sin\beta \quad (5g)$$

$$\sin\beta = \sin\varphi * \sin\delta + \cos\varphi * \cos\delta * \cos\omega \quad (5h)$$

where S_{sc} is the solar constant ($1,368 \text{ W m}^{-2}$), t_d is the day of the year, β is the solar elevation angle, φ is the degree of latitude, δ is the declination of the sun, and ω is the time angle.

CI has been commonly used to indicate sky conditions ([Rocha et al., 2004](#); [Letts, Lafleur & Roulet, 2005](#); [Urban et al., 2007](#); [Alton, 2008](#); [Knobl & Baldocchi, 2008](#)). To analyze how diffuse PAR responds to GEP under different sky conditions, we defined the two sky conditions as clear sky ($CI \geq 0.7$) and cloudy sky ($CI < 0.7$). The cloudy sky condition was divided into thin clouds ($0.3 < CI < 0.7$) and thick clouds ($CI \leq 0.3$) ([Chen, Shen & Kato, 2009](#)).

Diffuse light response parameters

We estimated photosynthetic parameters from light-response curves as described by the Michaelis–Menten equation below:

$$GEP = \frac{\alpha \times P_{max} \times PAR_{dif}}{\alpha \times PAR_{dif} + P_{max}} \quad (6)$$

where GEP is gross ecosystem production, PAR_{dif} is diffuse PAR, α is ecosystem apparent quantum yield as an initial slope of light response curve of photosynthesis, and P_{max} is maximum photosynthesis.

Statistical analyses

We fit [Eq. \(6\)](#) using GEP data collected each half hour from June to August to evaluate the variations in light response. The regressions were conducted on bin-averaged data using a PAR interval of $50 \mu\text{mol m}^{-2} \text{ s}^{-1}$. In order to test the dependency of the GEP_{day} -PAR relationship on different environmental factors, and to exclude the influences of plant phenology, we compiled GEP_{day} data from the mid-growing season into multiple groups. The daytime data from 10:00 to 16:00 in the mid-growing season (June–August, inclusive) was used for this analysis because plant photosynthesis is most active during this period, and the influence of the solar altitude angle on CI could be eliminated ([Bai et al., 2012](#); [Prior, Eamus & Duff, 1997](#)).

In order to further examine how GEP responds to changes in the diffuse PAR, environmental conditions were separated into three T_a classes ($T_a < 20 \text{ }^\circ\text{C}$, $20 < T_a \leq 25 \text{ }^\circ\text{C}$, and $T_a > 25 \text{ }^\circ\text{C}$), three VPD classes (VPD $< 1 \text{ kPa}$, $1 < \text{VPD} \leq 2 \text{ kPa}$, and VPD $> 2 \text{ kPa}$) ([Zhang et al., 2011a](#)), and two water conditions.

We used path analysis in AMOS (version 24.0, Chicago, IL, USA) to examine how environmental factor interactive controls affect GEP under different sky conditions. All

selected regression curves were statistically significant ($p < 0.05$) and based on several related studies (Zhang et al., 2010; Oliphant et al., 2011; Bai et al., 2012; Kanniah, Beringer & Hutley, 2013) that had been conducted using SPSS software (2012, ver. 22.0; SPSS Inc., USA). The standardized path coefficient (r), an analogy of the correlation coefficient, was used to quantify the effect of one variable on another (Shipley, 2002). The final model had a high goodness-of-fit when the good fit index (GFI) was greater than 0.9 and the root mean square error of approximation (RMSEA) was less than 0.05 (Kline, 2011).

RESULTS

Environmental factors and CI

Figures 2A and 2B show seasonal variations in the daily mean air temperature (T_a) and vapor pressure deficit (VPD). Seasonal patterns in T_a and VPD were similar across different years, peaking during the mid-growing seasons. Relative extractable soil water (REW) at 10-cm depths, daily sum precipitation, and cumulative precipitation showed clear seasonal patterns (Figs. 2C and 2D). The total rainfall at the middle of the growing seasons was 185.5 mm and 87.1 mm in 2014 and 2015, respectively, with one big rain event generating more than 20 mm day⁻¹ in 2014. Due to uneven rainfall from year to year, plants experienced 23 days of soil water stress (REW < 0.4) in 2014 and 86 days of soil water stress in 2015 (Figs. 2C and 2D).

Seasonal CI variations are shown in Figs. 2E and 2F. 2014 had a similar seasonal pattern as 2015. Changes in incident PAR daily totals were consistent with those of diffuse and direct PAR, with an annual total of 393.2 MJ m⁻² for incident PAR in 2014, 398.1 MJ m⁻² for incident PAR in 2015, 168.8 MJ m⁻² for diffuse PAR in 2014, and 167.0 MJ m⁻² for diffuse PAR in 2015 (Figs. 2G and 2H).

Figure 3 shows the effect of cloudiness on T_a , VPD, REW, and PAR (including total, direct, and diffuse) measurements during the studied period. T_a and VPD increased linearly with an increase in CI (Figs. 3A, 3B), while no obvious relationship was observed between REW and CI (Fig. 3C). Total PAR increased linearly with an increase in CI (Fig. 3D). As the sky became clearer (CI > 0.3), direct radiation increased exponentially with CI (Fig. 3D). The diffuse PAR increased as CI increased, peaking at a CI of 0.5 under thin cloud conditions, then decreased as CI continued to increase (Fig. 3D). When CI exceeded 0.8, the diffuse radiation increased.

Light response parameters and GEP under different sky conditions

Light response curves under different sky and environmental conditions are shown in Fig. 4A. We found significant quadratic relationships between the GEP and diffuse radiation under sunny and thin cloud conditions, and an approximately linear relationship under thick cloud conditions. There was a higher potential P_{\max} under cloudy conditions with a larger proportion of diffuse radiation than under clear sky conditions. Under clear sky conditions, we observed photo-inhibition phenomena (Fig. 4A). The GEP peaked when CI fluctuated around 0.5 (Fig. 4B).

The light-response curve α and P_{\max} parameters are listed in Table 1. As the sky became cloudy, the P_{\max} became larger and the α value became smaller. The P_{\max} was 39% and




Figure 2 The seasonal variations of environmental factors. (A) and (B) Daily mean air temperature (T_a) at 4.2 m above ground and vapor pressure deficit (VPD), (C) and (D) relative extractable soil water (REW) at 10 cm depth and daily-integrated and cumulative precipitation, (E) and (F) clearness index (CI), (G) and (H) daily-integrated incident total, diffuse and direct photosynthetically active radiation (PAR) in 2014 and 2015 growing season (May–October).

Full-size DOI: [10.7717/peerj.9043/fig-2](https://doi.org/10.7717/peerj.9043/fig-2)

68% greater under thick clouds than under thin clouds and clear skies, respectively. The α value was 400% and 68% greater under clear skies than under thick clouds and thin clouds, respectively. The P_{\max} was 23.7% greater under cloudy skies ($CI < 0.7$) than under clear skies ($CI \geq 0.7$), and the α value was 122% greater under clear skies than under cloudy skies (Table 1). As the T_a and VPD rose, the P_{\max} gradually became smaller (Table 1).

Under similar T_a , VPD, and water content classes, the P_{\max} values were significantly greater under cloudy conditions than under clear skies (Fig. 5). Compared to clear sky conditions, the cloudy condition P_{\max} values were 46.5%, 20.7%, and 30.4% higher under the three T_a classes; 5.3%, 20.9%, and 21% higher under the three VPD classes; 33% and 13% higher under the two water stressed conditions, respectively. Under cloudy sky conditions, the P_{\max} value under $T_a \leq 20$ °C was 19% higher than that under $20 < T_a \leq 25$ °C and 64% higher than that under $T_a > 25$ °C, respectively. Under cloudy sky conditions, the P_{\max} value was 2.3% higher under $VPD \leq 1$ kPa than under $1 < VPD \leq 2$ kPa, and 24% higher than under $VPD > 2$ kPa. In cloudy sky conditions, the P_{\max} was




Figure 3 The relationships between clearness index and the environmental factors. Relationships between clearness index and (A) air temperature (T_a) at 4.2 m above ground, (B) vapor pressure deficit (VPD), (C) relative extractable soil water (REW) at 10 cm depths, and (D) total incident photosynthetically active radiation (PAR), diffuse PAR and direct PAR. Data used are half-hourly values between 10 am and 4 pm in mid-growing season (June–August) in 2014 and 2015. Half-hourly data was bin-averaged by CI increment of 0.1. Bars indicate standard errors.

Full-size [DOI: 10.7717/peerj.9043/fig-3](https://doi.org/10.7717/peerj.9043/fig-3)




Figure 4 The relationships between different sky conditions and the gross ecosystem production. Daytime gross ecosystem production (GEP) as a function of (A) diffuse photosynthetically active radiation (PAR_{dif}) and of (B) the clearness index (CI) during mid-growing season (June–August) in 2014 and 2015. Half-hourly GEP was bin-averaged (A) by PAR increment of $50 \mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ and (B) by CI increment of 0.1. Bars indicate standard errors.

Full-size [DOI: 10.7717/peerj.9043/fig-4](https://doi.org/10.7717/peerj.9043/fig-4)

Table 1 The light-response curve parameters under different environment conditions and different sky conditions. Parameter values describing the response of daytime gross ecosystem production (GEP) to incident diffuse photosynthetically active radiation (PAR_{dif}) and incident total photosynthetically active radiation (PAR_{tot}) (the last column) between 10 am and 4 pm during mid-growing season (June–August) under different environmental conditions in 2014 and 2015.

Treatment	α $\mu\text{mol CO}_2$ $\mu\text{mol photon}^{-1}$	P_{max} $\mu\text{mol m}^{-2} \text{s}^{-1}$	Adj.R ²
$VPD \leq 1$	0.0162	2.86	0.59
$1 < VPD \leq 2$	0.0311	2.34	0.39
$VPD > 2$	0.0142	1.99	0.75
$T_a \leq 20$	0.0141	3.11	0.88
$20 < T_a \leq 25$	0.0209	2.46	0.54
$T_a > 25$	0.0148	2.27	0.56
$REW < 0.4$	0.0184	2.22	0.64
$REW \geq 0.4$	0.0214	2.37	0.60
Cloudy sky	0.0134	2.38	0.92
Thin clouds	0.0178	2.30	0.79
Thick clouds	0.0059	3.23	0.89
Clear sky	0.0299	1.93	0.30
Diffuse radiation	0.0102	2.608	0.85
Total radiation	0.0084	2.197	0.74

3.2% higher under non water-stressed conditions than under water-stressed conditions. The results indicated that lower T_a , lower VPD, and non-stressed REW accelerated the ecosystem photosynthetic potential under cloudy skies.

Direct and indirect influences of environmental factors on GEP

Based on the path coefficient analysis, our findings further confirmed the direct and indirect influences of PAR_{dif} on GEP. When the sky became cloudy, the PAR_{dif} more positively influenced GEP (Fig. 6). Additionally, the PAR_{dir} positively affected GEP under cloudy conditions and negatively affected GEP under clear sky conditions. In most cases, VPD and T_a negatively affected GEP. The influence of VPD on GEP was more negative under cloudy conditions than under clear skies. T_a was a weak influence on GEP under cloudy sky conditions. The indirect effect of T_a on GEP was negative, indicating that a T_a increase would indirectly lead to a GEP decrease via an increasing VPD. REW demonstrated positive effects on GEP under clear skies but no obvious effects under cloudy skies.

DISCUSSION

Factors influencing light response parameters

We concluded that the desert steppe had a higher P_{max} under cloudy conditions. Our findings were similar to those in previous studies on crop (Tong *et al.*, 2014) and grassland ecosystems (Jing *et al.*, 2010; Zhang *et al.*, 2011b; Bai *et al.*, 2012; Huang *et al.*, 2014). However, the P_{max} for a tropical savanna ecosystem was found to be highest under clear sky conditions (Kanniah, Beringer & Hutley, 2013). This may be due to a relative lower radiation intensity under cloudy conditions compared with sunny conditions in the




Figure 5 The maximum ecosystem photosynthesis P_{\max} under different environment conditions. Maximum ecosystem photosynthesis (P_{\max} in $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$) in clear skies and cloudy skies under various environmental conditions. Data were fitted using analysis of covariance (ANCOVA). Values above the bars were P_{\max} under different condition. T_{a1} , T_{a2} and T_{a3} represent air temperature $T_a \leq 20^\circ\text{C}$, $20 < T_a \leq 25^\circ\text{C}$ and $T_a > 25^\circ\text{C}$, respectively. VPD1, VPD2 and VPD3 represent vapor pressure deficit $\text{VPD} \leq 1 \text{ kPa}$, $1 < \text{VPD} \leq 2 \text{ kPa}$ and $\text{VPD} > 2 \text{ kPa}$, respectively. REW1 and REW2 represent relative extractable soil water content $\text{REW} < 0.4$ and $\text{REW} \geq 0.4$, respectively.

Full-size DOI: 10.7717/peerj.9043/fig-5




Figure 6 The path analysis diagram. Direct and indirect effects of environmental factors on gross ecosystem production (GEP) under clear sky and cloudy sky conditions during mid-growing season (June–August) in 2014 and 2015. Values in the path figure are standardized path coefficients (PV: -1 to 1). Dashed solid line with $\text{PV} < 0$ denotes negative correlation and solid line with $\text{PV} > 0$ denotes positive correlation.

Full-size DOI: 10.7717/peerj.9043/fig-6

tropical savanna. The α value in this study is consistent with those in forest ecosystem studies, where the cloudy sky appeared larger than the clear sky (Hollinger *et al.*, 1994; Rocha *et al.*, 2004; Dengel & Grace, 2010; Zhang *et al.*, 2010). This result indicates that the desert steppe ecosystem has a higher potential light use efficiency.

We found higher P_{\max} values under cloudy rather than sunny skies for two reasons. First, radiation composition and quality and light distribution variations within an ecosystem influenced the canopy's photosynthetic productivity under cloudy conditions (Knohl &

Baldocchi, 2008). Second, cloudy skies lead to a reduction in T_a and VPD, which increase canopy stomatal conductance and enhance photosynthetic rates (*Huxman et al., 2003; Xu et al., 2018*).

However, the results of this research are not consistent with other previous studies (*Letts, Lafleur & Roulet, 2005; Niyogi et al., 2007*) that did not find an increase in the photosynthesis of ecosystems with low leaf area indexes, such as grasslands and shrublands, under cloudy conditions. The explanation for this may be that a decrease in GEP under cloudy conditions caused by total radiation reduction is greater than an increase in GEP caused by diffuse fertilization. We attribute this inconsistency to environmental differences in the studied ecosystems, such as local thermal, water, and light conditions (*Zhang et al., 2011b*).

Interplay of factors on GEP under different sky conditions

Different sky conditions caused radiation composition and other environmental factor variations, leading to corresponding effects on GEP. Our path analysis results showed that in the desert steppe ecosystem under cloudy skies the PAR_{dif} more positively influenced GEP, while T_a and VPD suppressed GEP in most cases (*Fig. 6*). The existence of clouds may be both a cause and consequence of solar radiation, temperature, moisture, precipitation, and other atmospheric factors (*Gu et al., 1999; Zhang et al., 2011a; Zhang et al., 2011b*). These factors have direct and indirect effects on the biophysical processes of ecosystem productivity.

Our study showed that the P_{max} increased under lower T_a and VPD conditions. However, under the same environmental factor limitations, VPD reduced P_{max} more significantly than T_a under cloudy conditions (*Fig. 5*). The increase in T_a and VPD limited the GEP regardless of sky conditions. Under cloudy sky conditions, T_a and VPD had a secondary direct effect on GEP. A decrease in T_a can cause decreased ecosystem respiration (*Gu et al., 1999; Urban et al., 2007*). An increase in T_a might result in increased VPD, leading to the indirect inhibition of GEP. Increases in VPD can limit GEP under varying environmental conditions, and leaf guard cell's behavior is highly controlled by the inside and outside vapor pressure balance (*Farquhar & Sharkey, 1982*). On the other hand, higher VPD levels affect stomatal closure, thus controlling photosynthetic rates (*Zhou et al., 2013; Goodrich et al., 2015*).

Under clear sky conditions, sunlit canopy leaves showed an obvious photo-inhibition phenomenon (*Fig. 4A*), similar to that of an open shrub-land ecosystem (*Keenan et al., 2019*). Higher T_a and VPD under clear sky conditions (*Figs. 3A, 3B*) may cause the stomatal closure of leaf guard cells that are protecting themselves against high radiation (*Zhou et al., 2013; Goodrich et al., 2015*). Previous studies have shown that strong UV radiation, especially ultraviolet-B (UV-B), can inhibit photosynthesis (*Ekelund, 2000; Correia et al., 2005; Sangtarash et al., 2009; Li et al., 2010*). Plants that live under strong UV can regulate pigmentation, different enzyme mechanisms, and photosynthesis in order to protect themselves from high radiation (*Franklin & Forster, 1997; Ekelund, 2000*).




Figure 7 Daytime gross ecosystem production (GEP) as a function of diffuse photosynthetically active radiation (PAR_{dif}) and of total photosynthetically active radiation (PAR_{tot}) during mid-growing season (June–August) in 2014 and 2015. Half-hourly GEP was bin-averaged (a) by PAR increment of $50 \mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$. Bars indicate standard errors. Model parameters were fitted separately to diffuse and total radiation data, and the optimized model curves are shown as lines.

Full-size DOI: 10.7717/peerj.9043/fig-7

Potential impact of diffuse radiation on canopy productivity

Increases in diffuse radiation play an important role in ecosystem GEP enhancement. The GEP peaked when the CI was around 0.5 (Fig. 4B) and when PAR_{dif} reached its maximum (Fig. 3D). The results revealed that vegetation reached its light saturation when PAR_{dif} reached its maximum. This CI range was consistent with those found in other studies on semi-arid steppes (Zhang *et al.*, 2011b), grasslands (Bai *et al.*, 2012), croplands (Tong *et al.*, 2014), and forest ecosystems (Gu *et al.*, 1999), but was higher than the values obtained in studies on short grass in semi-arid regions (Jing *et al.*, 2010). We also observed that the light response curves differed under diffuse radiation and total radiation conditions (Fig. 7). The light-response curve α and P_{max} parameters are listed in Table 1. Two light response curves had similar trends, but the curves rose more steeply with light intensities under diffuse radiation. This result is consistent with previous studies (Oliphant *et al.*, 2011; Williams *et al.*, 2014). In the desert steppe ecosystem, a higher α value meant a higher potential light use efficiency, and a higher P_{max} meant a higher potential ecosystem production under diffuse radiation than under total radiation conditions. Those results indicated that the enhancement of canopy photosynthesis by diffuse radiation received by an ecosystem is more pronounced under cloudy conditions (Gu *et al.*, 1999; Alton *et al.*, 2007a; Oliveira *et al.*, 2007). The effect of radiation changes, associated with an increase in clouds or scattered aerosols, on photosynthesis depends on the balance between total PAR reduction (which tends to reduce photosynthesis) and increased PAR diffusion (which tends to increase photosynthesis) (Mercado *et al.*, 2009).

Due to a lack of measured PAR_{dif} data, our data was obtained using model calculations. We corrected the time zones according to Beijing time, but there is still some uncertainty about the simulated model. Other studies have shown that the model tends to overestimate PAR_{dif} when the fraction of diffuse PAR ($f\text{PAR}_{\text{dif}} < 0.3$ and $f\text{PAR}_{\text{dif}} > 0.9$), and underestimates PAR_{dif} values in the $f\text{PAR}_{\text{dif}}$ range of 0.3 and 0.9 (Oliphant & Stoy, 2018). Although the GEP model based on the diffuse beam partitioning model underestimated the observed GEP because of errors in the diffuse partitioning (Lee et al., 2018), the diffuse beam partitioning model is still more accurate than other GEP models (Zhou & Xin, 2019).

We concluded that the increase of diffuse radiation under cloudy conditions has a positive impact on the GEP of the desert steppe. Further detailed analysis of how ecosystem diffuse radiation and other environmental factors affect carbon exchange processes are needed. Additionally, more data should be collected to distinguish the effects of aerosol, cloudiness, and UV on photosynthesis in mid-high altitude areas.

CONCLUSION

Our results indicated that diffuse radiation could improve photosynthesis in the desert steppe ecosystem. The GEP peaked when the CI was around 0.5 with a maximum PAR_{dif} . The P_{max} was 23.7% higher under cloudy skies than under clear skies. When the sky was cloudy, the PAR_{dif} became a positive influence on the desert steppe GEP. In addition, lower VPD ($1 \leq 1 \text{ kPa}$), lower T_a ($T_a \leq 20 \text{ }^\circ\text{C}$), and non-stressed water conditions ($\text{REW} \geq 0.4$) were more conducive to ecosystem photosynthetic enhancement under cloudy skies than under clear skies due to the comprehensive effects of VPD and T_a on stomatal conductance. Our study showed that diffuse radiation did encourage a potential increase in the production of desert steppe ecosystems in Northwest China. Further changes in sky conditions and diffuse radiation should be considered when modeling the GEP of desert steppe ecosystems.

ACKNOWLEDGEMENTS

The authors thank the U.S.–China Carbon Consortium (USCCC) who supported this work via helpful discussions and idea exchanges.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This work was supported by the National Natural Science Foundation of China (NSFC; 31670710, 31670708), the Fundamental Research Funds for the Central Universities (Proj. No. 2015ZCQ-SB-02), and the Beijing Municipal Science & Technology Commission (Z161100001116084). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:

National Natural Science Foundation of China: NSFC; 31670710, 31670708.
Fundamental Research Funds for the Central Universities: Proj. No. 2015ZCQ-SB-02.
Beijing Municipal Science & Technology Commission: Z161100001116084.

Competing Interests

The authors declare there are no competing interests.

Author Contributions

- Cheng Li conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.
- Xin Jia conceived and designed the experiments, analyzed the data, authored or reviewed drafts of the paper, and approved the final draft.
- Jingyong Ma, Peng Liu, Ruizhi Yang, Yujie Bai and Muhammad Hayat performed the experiments, authored or reviewed drafts of the paper, and approved the final draft.
- Jinglan Liu conceived and designed the experiments, authored or reviewed drafts of the paper, financial support, and approved the final draft.
- Tianshan Zha conceived and designed the experiments, authored or reviewed drafts of the paper, and approved the final draft.

Data Availability

The following information was supplied regarding data availability:

The raw measurements are available in the [Supplementary Files](#).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.9043#supplemental-information>.

REFERENCES

- Alton PB. 2008.** Reduced carbon sequestration in terrestrial ecosystems under overcast skies compared to clear skies. *Agricultural and Forest Meteorology* **148**:1653–1653 DOI [10.1016/j.agrformet.2008.05.014](https://doi.org/10.1016/j.agrformet.2008.05.014).
- Alton PB, Ellis R, Los SO, North PR. 2007a.** Improved global simulations of gross primary product based on a separate and explicit treatment of diffuse and direct sunlight. *Journal of Geophysical Research Atmospheres* **112**:D07203 DOI [10.1029/2006jd008022](https://doi.org/10.1029/2006jd008022).
- Alton PB, North PR, Los SO. 2007b.** The impact of diffuse sunlight on canopy light-use efficiency, gross photosynthetic product and net ecosystem exchange in three forest biomes. *Global Change Biology* **13**:776–787 DOI [10.1111/j.1365-2486.2007.01316.x](https://doi.org/10.1111/j.1365-2486.2007.01316.x).
- Bai YF, Wang J, Zhang BC, Zhang ZH, Liang J. 2012.** Comparing the impact of cloudiness on carbon dioxide exchange in a grassland and a maize cropland in northwestern China. *Ecological Research* **27**:615–623 DOI [10.1007/s11284-012-0930.z](https://doi.org/10.1007/s11284-012-0930.z).

- Baldocchi DD. 2003.** Assessing the eddy covariance technique for evaluating carbon dioxide exchange rates of ecosystems: past, present and future. *Global Change Biology* 9:479–492 DOI [10.1046/j.1365-2486.2003.00629.x](https://doi.org/10.1046/j.1365-2486.2003.00629.x).
- Chen J, Shen M, Kato T. 2009.** Diurnal and seasonal variations in light-use efficiency in an alpine meadow ecosystem: causes and implications for remote sensing. *Journal of Plant Ecology* 2:173–185 DOI [10.1093/jpe/rtp020](https://doi.org/10.1093/jpe/rtp020).
- Cheng SJ, Bohrer G, Steiner AL, Hollinger DY, Suyker A, Phillips RP, Nadelhoffer KJ. 2015.** Variations in the influence of diffuse light on gross primary productivity in temperate ecosystems. *Agricultural and Forest Meteorology* 201:98–110 DOI [10.1016/j.agrformet.2014.11.002](https://doi.org/10.1016/j.agrformet.2014.11.002).
- Correia CM, Pereira JMM, Coutinho JF, Björn LO, Torres-Pereira JM. 2005.** Ultraviolet-B radiation and nitrogen affect the photosynthesis of maize: a Mediterranean field study. *European Journal of Agronomy* 22:337–347 DOI [10.1016/j.eja.2004.05.002](https://doi.org/10.1016/j.eja.2004.05.002).
- Dengel S, Grace J. 2010.** Carbon dioxide exchange and canopy conductance of two coniferous forests under various sky conditions. *Oecologia* 164:797–808 DOI [10.1007/s00442-010-1687-0](https://doi.org/10.1007/s00442-010-1687-0).
- Ekelund NGA. 2000.** Interactions between photosynthesis and ‘light-enhanced dark respiration’ (LEDR) in the flagellate *Euglena gracilis* after irradiation with ultraviolet radiation. *Journal of Photochemistry and Photobiology B: Biology* 55:63–69 DOI [10.1016/s1011-1344\(00\)00029-4](https://doi.org/10.1016/s1011-1344(00)00029-4).
- Falge E, Baldocchi D, Olson R, Anthoni P, Aubinet M, Bernhofer C, Burba G, Ceulemans R, Clement R, Dolman H, Granier A, Gross P, Grünwald T, Hollinger D, Jensen N-O, Katul G, Keronen P, Kowalski A, Lai CT, Law BE, Meyers T, Moncrieff J, Moors E, Munger JW, Pilegaard K, Rannik U, Rebmann C, Suyker A, Tenhunen J, Tu K, Verma S, Vesala T, Wilson K, Wofsy S. 2001.** A. Gap filling strategies for defensible annual sums of net ecosystem exchange. *Agricultural and Forest Meteorology* 107:43–69 DOI [10.1016/s0168-1923\(00\)00225-2](https://doi.org/10.1016/s0168-1923(00)00225-2).
- Farquhar GD, Roderick ML, Pinatubo II. 2003.** Diffuse light, and the carbon cycle. *Science* 299:1997–1998 DOI [10.1126/science.1080681](https://doi.org/10.1126/science.1080681).
- Farquhar GD, Sharkey TD. 1982.** Stomatal conductance and photosynthesis. *Annual Review of Plant Physiology* 33:317–345 DOI [10.1146/annurev.pp.33.060182.001533](https://doi.org/10.1146/annurev.pp.33.060182.001533).
- Feng W, Zhang Y, Wu B, Zha TS, Jia X, Qin SG, Shao CX, Liu JB, Lai ZR, Fa KY. 2013.** Influence of disturbance on soil respiration in biologically crusted soil during the dry season. *The Scientific World Journal* 2013: Article 408560 DOI [10.1155/2013/408560](https://doi.org/10.1155/2013/408560).
- Franklin LA, Forster RM. 1997.** Review the changing irradiance environment: consequences for marine macrophyte physiology, productivity and ecology. *European Journal of Phycology* 32:207–232 DOI [10.1080/09670269710001737149](https://doi.org/10.1080/09670269710001737149).
- Gao X, Gu F, Mei X, Hao W, Li H, Gong D, Li X. 2018.** Light and Water Use Efficiency as Influenced by Clouds and/or Aerosols in a Rainfed Spring Maize Cropland on the Loess Plateau. *Crop Science* 58:853–862 DOI [10.2135/cropsci2017.06.0341](https://doi.org/10.2135/cropsci2017.06.0341).
- Gong J, Jia X, Zha T, Wang B, Kellomäki S, Peltola H. 2016.** Modeling the effects of plant-interspace heterogeneity on water-energy balances in a semiarid ecosystem.

- Agricultural and Forest Meteorology* 221:189–206
DOI 10.1016/j.agrformet.2016.01.144.
- Goodrich JP, Campbell DI, Clearwater MJ, Rutledge S, Schipper LA. 2015.** High vapor pressure deficit constrains GPP and the light response of NEE at a Southern Hemisphere bog. *Agricultural and Forest Meteorology* 203:54–63
DOI 10.1016/j.agrformet.2015.01.001.
- Goudriaan J. 1977.** *Crop micrometeorology: a simulation study*. Wageningen: Pudoc.
- Granier A, Breda N, Biron P, Villette S. 1999.** A lumped water balance model to evaluate duration and intensity of drought constraints in forest stands. *Ecological Modelling* 116:269–283 DOI 10.1016/s0304-3800(98)00205-1.
- Gu L, Baldocchi D, Verma SB, Black TA, Vesala T, Falge EM, Dowty PR. 2002.** Advantages of diffuse radiation for terrestrial ecosystem productivity. *Journal of Geophysical Research: Atmospheres* 107:ACL 2–1–ACL 2–23 DOI 10.1029/2001jd001242.
- Gu L, Baldocchi DD, Wofsy SC, Munger JW, Michalsky JJ, Urbanski SP, Boden TA. 2003.** Response of a deciduous forest to the Mount Pinatubo eruption: enhanced photosynthesis. *Science* 299:2035–2038 DOI 10.1126/science.1078366.
- Gu L, Fuentes JD, Shugart HH, Staebler RM, Black TA. 1999.** Responses of net ecosystem exchanges of carbon dioxide to changes in cloudiness: results from two North American deciduous forests. *Journal of Geophysical Research: Atmospheres* 104:31421–31434 DOI 10.1029/1999jd901068.
- Guan DX, Wu JB, Zhao XS, Han SJ, Yu GR, Sun XM, Jin CJ. 2006.** CO₂ fluxes over an old, temperate mixed forest in northeastern China. *Agricultural and Forest Meteorology* 137:138–149 DOI 10.1016/j.agrformet.2006.02.003.
- He M, Ju W, Zhou Y, Chen J, He H, Wang S, Wang H, Guan D, Guan D, Yan J, Li Y, Hao Y, Zhao F. 2013.** Development of a two-leaf light use efficiency model for improving the calculation of terrestrial gross primary productivity. *Agricultural and Forest Meteorology* 173:28–39 DOI 10.1016/j.agrformet.2013.01.003.
- Hollinger DY, Kelliher FM, Byers JN, Hunt JE, McSeveny TM, Weir PL. 1994.** Carbon dioxide exchange between an undisturbed old-growth temperate forest and the atmosphere. *Ecology* 75:134–150 DOI 10.2307/1939390.
- Huang K, Wang S, Zhou L, Wang H, Zhang J, Yan J, Zhao L, Wang Y, Shi P. 2014.** Impacts of diffuse radiation on light use efficiency across terrestrial ecosystems based on eddy covariance observation in China. *PLOS ONE* 9(11):e110988
DOI 10.1371/journal.pone.0110988.
- Huxman TE, Turnipseed AA, Sparks JP, Harley PC, Monson RK. 2003.** Temperature as a control over ecosystem CO₂ fluxes in a high-elevation, subalpine forest. *Oecologia* 134:537–546 DOI 10.1007/s00442-002-1131-1.
- Jia X, Zha TS, Gong JN, Wu B, Zhang YQ, Qin SG, Chen GP, Feng W, Kellomäki S, Peltola H. 2016.** Energy partitioning over a semiarid shrubland in northern China. *Hydrological Processes* 30:972–985 DOI 10.1002/hyp.10685.
- Jia X, Zha TS, Wu B, Zhang YQ, Gong JN, Qin SG, Chen GP, Qian D, Kellomäki S, Peltola H. 2014.** Biophysical controls on net ecosystem CO₂ exchange

- over a semiarid shrubland in northwest China. *Biogeosciences* **11**:4679–4693 DOI [10.5194/bg-11-4679-2014](https://doi.org/10.5194/bg-11-4679-2014).
- Jing X, Huang J, Wang G, Higuchi K, Bi J, Sun Y, Yu H, Wang T. 2010.** The effects of clouds and aerosols on net ecosystem CO₂ exchange over semi-arid Loess Plateau of Northwest China. *Atmospheric Chemistry and Physics* **10**:8205–8218 DOI [10.5194/acp-10-8205-2010](https://doi.org/10.5194/acp-10-8205-2010).
- Kanniah KD, Beringer J, Hutley L. 2013.** Exploring the link between clouds, radiation, and canopy productivity of tropical savannas. *Agricultural and Forest Meteorology* **182**:304–313 DOI [10.1016/j.agrformet.2013.06.010](https://doi.org/10.1016/j.agrformet.2013.06.010).
- Keenan TF, Migliavacca M, Papale D, Baldocchi D, Reichstein M, Torn M, Wutzler T. 2019.** Widespread inhibition of daytime ecosystem respiration. *Nature Ecology & Evolution* **3**:407 DOI [10.1038/s41559-019-0809-2](https://doi.org/10.1038/s41559-019-0809-2).
- Kline RB. 2011.** *Principles and practice of structural equation modeling*. 3rd edition. New York/London: Guilford.
- Knobl A, Baldocchi DD. 2008.** Effects of diffuse radiation on canopy gas exchange processes in a forest ecosystem. *Journal of Geophysical Research: Biogeosciences* **113**:G02023 DOI [10.1029/2007jg000663](https://doi.org/10.1029/2007jg000663).
- Lee MS, Hollinger DY, Keenan TF, Ouimette AP, Ollinger SV, Richardson AD. 2018.** Model-based analysis of the impact of diffuse radiation on CO₂ exchange in a temperate deciduous forest. *Agricultural and Forest Meteorology* **249**:377–389 DOI [10.1016/j.agrformet.2017.11.016](https://doi.org/10.1016/j.agrformet.2017.11.016).
- Letts MG, Lafleur PM, Roulet NT. 2005.** On the relationship between cloudiness and net ecosystem carbon dioxide exchange in a peatland ecosystem. *Ecoscience* **12**:53–69 DOI [10.2980/i1195-6860-12-1-53.1](https://doi.org/10.2980/i1195-6860-12-1-53.1).
- Li FR, Peng SL, Chen BM, Hou YP. 2010.** A meta-analysis of the responses of woody and herbaceous plants to elevated ultraviolet-B radiation. *Acta Oecologica* **36**:1–9 DOI [10.1016/j.actao.2009.09.002](https://doi.org/10.1016/j.actao.2009.09.002).
- Lin X, Chen B, Chen J, Zhang H, Sun S, Xu G, Guo L, Ge M, Qu J, Li L, Kong Y. 2017.** Seasonal fluctuations of photosynthetic parameters for light use efficiency models and the impacts on gross primary production estimation. *Agricultural and Forest Meteorology* **236**:22–35 DOI [10.1016/j.agrformet.2016.12.019](https://doi.org/10.1016/j.agrformet.2016.12.019).
- Mercado LM, Bellouin N, Sitch S, Boucher O, Huntingford C, Wild M, Cox PM. 2009.** Impact of changes in diffuse radiation on the global land carbon sink. *Nature* **458**:1014 DOI [10.1038/nature07949](https://doi.org/10.1038/nature07949).
- Monteith JL. 1972.** Solar radiation and productivity in tropical ecosystems. *Journal of Applied Ecology* **9**:747–766 DOI [10.2307/2401901](https://doi.org/10.2307/2401901).
- Monteith JL. 1977.** Climate and the efficiency of crop production in Britain. *Philosophical Transactions of the Royal Society of London. B, Biological Sciences* **281**:277–294 DOI [10.1098/rstb.1977.0140](https://doi.org/10.1098/rstb.1977.0140).
- Niyogi D, Chang HI, Chen F, Gu L, Kumar A, Menon S, Pielke RA. 2007.** Potential impacts of aerosol–land–atmosphere interactions on the Indian monsoonal rainfall characteristics. *Natural Hazards* **42**:345–359 DOI [10.1007/s11069-006-9085-y](https://doi.org/10.1007/s11069-006-9085-y).

- Oliphant AJ, Dragoni D, Deng B, Grimmond CSB, Schmid HP, Scott SL. 2011.** The role of sky conditions on gross primary production in a mixed deciduous forest. *Agricultural and Forest Meteorology* **151**(7):781–791 DOI [10.1016/j.agrformet.2011.01.005](https://doi.org/10.1016/j.agrformet.2011.01.005).
- Oliphant AJ, Stoy PC. 2018.** An evaluation of semiempirical models for partitioning photosynthetically active radiation into diffuse and direct beam components. *Journal of Geophysical Research: Biogeosciences* **123**(3):889–901 DOI [10.1002/2017JG004370](https://doi.org/10.1002/2017JG004370).
- Oliveira PHF, Artaxo P, Pires C, Lucca SDe, Procópio A, Holben B, Schafer J, Cardoso LF, Wofsy SC, Rocha HR. 2007.** The effects of biomass burning aerosols and clouds on the CO₂ flux in Amazonia. *Tellus B: Chemical and Physical Meteorology* **59**:338–349 DOI [10.1111/j.1600-0889.2007.00270.x](https://doi.org/10.1111/j.1600-0889.2007.00270.x).
- Park SB, Knohl A, Lucas-Moffat AM, Migliavacca M, Gerbig C, Vesala T, Peltola O, Mammarella I, Kolle O, Lavrič JV, Prokushkin A, Heimann M. 2018.** Strong radiative effect induced by clouds and smoke on forest net ecosystem productivity in central Siberia. *Agricultural and Forest Meteorology* **250**:376–387 DOI [10.1016/j.agrformet.2017.09.009](https://doi.org/10.1016/j.agrformet.2017.09.009).
- Prior LD, Eamus D, Duff GA. 1997.** Seasonal and diurnal patterns of carbon assimilation stomatal conductance and leaf water potential in *Eucalyptus tetrodonta* saplings in a wet–dry savanna in northern Australia. *Australian Journal of Botany* **45**:241–258 DOI [10.1071/bt96017](https://doi.org/10.1071/bt96017).
- Ren XL, He HL, Zhang L, Zhou L, Yu GR, Fan JW. 2013.** Spatiotemporal variability analysis of diffuse radiation in China during 1981–2010. *Annales Geophysicae. Copernicus GmbH* **31**:277–289 DOI [10.5194/angeo-31-277-2013](https://doi.org/10.5194/angeo-31-277-2013).
- Rocha AV, Su HB, Vogel CS, Schmid HP, Curtis PS. 2004.** Photosynthetic and water use efficiency responses to diffuse radiation by an aspen dominated northern hardwood forest. *Forest Science* **50**:793–801 DOI [10.1093/forestscience/50.6.793](https://doi.org/10.1093/forestscience/50.6.793).
- Roderick ML, Farquhar GD, Berry SL, Noble IR. 2001.** On the direct effect of clouds and atmospheric particles on the productivity and structure of vegetation. *Oecologia* **129**:21–30 DOI [10.1007/s004420100760](https://doi.org/10.1007/s004420100760).
- Sangtarash MH, Qaderi MM, Chinnappa CC, Reid DM. 2009.** Differential responses of two *Stellaria longipes* ecotypes to ultraviolet-B radiation and drought stress. *Flora-Morphology, Distribution, Functional Ecology of Plants* **204**:593–603 DOI [10.1016/j.flora.2008.08.004](https://doi.org/10.1016/j.flora.2008.08.004).
- Shipley B. 2002.** *A user's guide to path analysis, structural equations and causal inference*. Cambridge: Cambridge University Press, 100135.
- Still CJ, Riley WJ, Biraud SC, Noone DC, Buening NH, Randerson JT, Torn MS, Welker J, White JWC, Vachon R, Farquhar GD, Berry JA. 2009.** Influence of clouds and diffuse radiation on ecosystem-atmosphere CO₂ and CO₁₈O exchanges. *Journal of Geophysical Research* **114**:G01018 DOI [10.1029/2007jg000675](https://doi.org/10.1029/2007jg000675).
- Tong X, Li J, Yu Q, Lin Z. 2014.** Biophysical controls on light response of net CO₂ exchange in a winter wheat field in the North China Plain. *PLOS ONE* **9**:e89469 DOI [10.1371/journal.pone.0089469](https://doi.org/10.1371/journal.pone.0089469).

- Tong X, Zhang J, Meng P, Li J, Zheng N. 2017. Light use efficiency of a warm-temperate mixed plantation in north China. *International Journal of Biometeorology* 61:1607–1615 DOI 10.1007/s00484-017-1339-8.
- Urban O, JANOŮŠ D, Acosta M, Czerný R, Marková I, Navrátil M, Pavelka M, Pokorný R, Šprtová M, Zhang R, ŠPUNDA V, Grace J, Marek MV. 2007. Eco-physiological controls over the net ecosystem exchange of mountain spruce stand. Comparison of the response in direct vs. diffuse solar radiation. *Global Change Biology* 13:157–168 DOI 10.1111/j.1365-2486.2006.01265.
- Urban O, Klem K, Ač A, Havránková K, Holíšová P, Navrátil M, Zitová M, Kozlová K, Pokorný R, Šprtová M, Tomášková I, Špunda V, Grace J. 2012. Impact of clear and cloudy sky conditions on the vertical distribution of photosynthetic CO₂ uptake within a spruce canopy. *Functional Ecology* 26:46–55 DOI 10.1111/j.1365-2435.2011.01934.x.
- Wang K, Dickinson RE, Liang S. 2008. Observational evidence on the effects of clouds and aerosols on net ecosystem exchange and evapotranspiration. *Geophysical Research Letters* 35:L10401 DOI 10.1029/2008gl034167.
- Wang B, Zha TS, Jia X, Wu B, Zhang YQ, Qin SG. 2014. Soil moisture modifies the response of soil respiration to temperature in a desert shrub ecosystem. *Biogeosciences* 11:259–268 DOI 10.5194/bg-11-259-2014.
- Webb EK, Pearman GI, Leuning R. 1980. Correction of flux measurements for density effects due to heat and water vapour transfer. *Quarterly Journal of the Royal Meteorological Society* 106:85–100 DOI 10.1002/qj.49710644707.
- Williams M, Rastetter EB, Van der Pol L, Shaver GR. 2014. Arctic canopy photosynthetic efficiency enhanced under diffuse light, linked to a reduction in the fraction of the canopy in deep shade. *New Phytologist* 202(4):1267–1276 DOI 10.1111/nph.12750.
- Xie J, Zha T, Jia X, Qian D, Wu B, Zhang Y, Bourque CPA, Chen J, Peltola H. 2015. Irregular precipitation events in control of seasonal variations in CO₂ exchange in a cold desert-shrub ecosystem in northwest China. *Journal of Arid Environments* 120:33–41 DOI 10.1016/j.jaridenv.2015.04.009.
- Xie J, Zha T, Zhou C, Jia X, Yu H, Yang B, Chen J, Zhang F, Wang B, Bourque CPA, Sun G, Ma H, Liu H, Peltola H. 2016. Seasonal variation in ecosystem water use efficiency in an urban-forest reserve affected by periodic drought. *Agricultural and Forest Meteorology* 221:142–151 DOI 10.1016/j.agrformet.2016.02.013.
- Xu H, Zhang Z, Chen J, Xiao J, Zhu M, Kang M, Cao W. 2018. Regulations of cloudiness on energy partitioning and water use strategy in a riparian poplar plantation. *Agricultural and Forest Meteorology* 262:135–146 DOI 10.1016/j.agrformet.2018.07.008.
- Zhang BC, Cao JJ, Bai YF, Yang SJ, Hu L, Ning ZG. 2011a. Effects of cloudiness on carbon dioxide exchange over an irrigated maize cropland in northwestern China. *Biogeosciences Discussions* 8:1669–1691 DOI 10.5194/bgd-8-1669-2011.
- Zhang WL, Chen SP, Chen J, Wei L, Han XG, Lin GH. 2007. Biophysical regulations of carbon fluxes of a steppe and a cultivated cropland in semiarid Inner Mongolia. *Agricultural and Forest Meteorology* 146:216–229 DOI 10.1016/j.agrformet.2007.06.002.

- Zhang M, Yu GR, Zhang LM, Sun XM, Wen XF, Han SJ, Yan JH. 2010.** Impact of cloudiness on net ecosystem exchange of carbon dioxide in different types of forest ecosystems in China. *Biogeosciences* 7:711–722 DOI [10.5194/bg-7-711-2010](https://doi.org/10.5194/bg-7-711-2010).
- Zhang M, Yu GR, Zhuang J, Gentry R, Fu YL, Sun XM, Zhang ML, Wen XF, Han SJ, Yan JH, Zhang YP, Wang YF, Li YN. 2011b.** Effects of cloudiness change on net ecosystem exchange, light use efficiency, and water use efficiency in typical ecosystems of China. *Agricultural and Forest Meteorology* 151:803–816 DOI [10.1016/j.agrformet.2011.01.011](https://doi.org/10.1016/j.agrformet.2011.01.011).
- Zhou X, Xin Q. 2019.** Improving satellite-based modelling of gross primary production in deciduous broadleaf forests by accounting for seasonality in light use efficiency. *International Journal of Remote Sensing* 40(3):931–955 DOI [10.1080/01431161.2018.1519285](https://doi.org/10.1080/01431161.2018.1519285).
- Zhou J, Zhang Z, Sun G, Fang X, Zha T, McNulty S, Chen J, Jin Y, Noormets A. 2013.** Response of ecosystem carbon fluxes to drought events in a poplar plantation in Northern China. *Forest Ecology and Management* 300:33–42 DOI [10.1016/j.foreco.2013.01.007](https://doi.org/10.1016/j.foreco.2013.01.007).
- Zhu Z, Sun X, Wen X, Zhou Y, Tian J, Yuan G. 2006.** Study on the processing method of nighttime CO₂ eddy covariance flux data in China FLUX. *Science in China* 49:36–46 DOI [10.1007/s11430-006-8036-5](https://doi.org/10.1007/s11430-006-8036-5).