

# CRISPR screening identifies M1AP as a new MYC regulator with a promoter-reporter system

Akiko Yamamoto<sup>Equal first author, 1</sup>, Morito Kurata<sup>Corresp., Equal first author, 1</sup>, Ichiroh Onishi<sup>1</sup>, Keisuke Sugita<sup>1</sup>, Miwa Matsumura<sup>1</sup>, Sachiko Ishibashi<sup>1</sup>, Masumi Ikeda<sup>1</sup>, Kouhei Yamamoto<sup>1</sup>, Masanobu Kitagawa<sup>1</sup>

<sup>1</sup> Department of Comprehensive Pathology, Graduate School of Medical and Dental Sciences, Tokyo Medical and Dental University, Tokyo, Japan

Corresponding Author: Morito Kurata  
Email address: kurata.pth2@tmd.ac.jp

**Background.** *MYC* is one of the proto-oncogenes contributing to tumorigenesis in many human cancers. Although the mechanism of *MYC* regulation is still not fully understood, learning about the comprehensive mechanism controlling the transcriptional activity of *MYC* will lead to therapeutic targets. The CRISPR/Cas9 library system is a simple and powerful screening technique. This study aims to identify new transcriptional upstream activators of *MYC* using the CRISPR activation library with new promoter-reporter systems.

**Methods and Results.** The *MYC* promoter-reporter system was developed with a photoconvertible fluorescent protein, Dendra2, and named “p*MYC*-promoter-Dendra2.” This *MYC* promoter-reporter system was designed to harbor a proximal *MYC* promoter at –3.1 kb. Both the CRISPR activation library and p*MYC*-promoter-Dendra2 were induced to HEK 293T cells, and Dendra2-positive cells, that are supposed that *MYC* should be upregulated, were collected individually by a cell sorter. Among the 169 cells collected, 12 clones were successfully established. Then, p*MYC*-promoter-Dendra2 was transfected again into these 12 clones, and 2 of 12 clones showed Dendra2 positivity. In this procedure, the cells with non-specific autofluorescence were correctly distinguished by utilizing the photoswitchable character of Dendra2. Using extracted guide DNA of these two Dendra2 positive clones, polymerase chain reaction (PCR) was performed to amplify the guide RNA (gRNA) containing region, which was introduced by the CRISPR activation library. Eventually, *PLEKHO2*, *MICU*, *MBTPS1*, and *M1AP* were identified, and these gRNAs were transfected individually into HEK 293T cells again using the CRISPR activation system. Only *M1AP* gRNA transfected cells showed Dendra2-positive fluorescence. Then, the overexpression vector for *M1AP* with a doxycycline-inducible vector confirmed that *M1AP* induced high *MYC* expression by real-time quantitative PCR and western blot. Furthermore, the dual-luciferase assay showed a significant increase of promoter activity, and *MYC* mRNA was higher in *M1AP*-overexpressing cells. *M1AP* is highly expressed in several cancers, though, a positive correlation between *M1AP* and *MYC* was observed only in human acute myeloid leukemia.

**Conclusion.** The present study confirmed that the experimental method using the CRISPR library technology functions effectively for the identification of molecules that activate endogenous *MYC*. This method will help elucidate the regulatory mechanism of *MYC* expression, as well as supporting further drug research against malignant tumors.

# CRISPR Screening Identifies M1AP as a New MYC Regulator with a Promoter-Reporter System

Akiko Yamamoto<sup>1, 2</sup>, Morito Kurata<sup>1, 2\*\*</sup>, Ichiroh Onishi<sup>1</sup>, Keisuke Sugita<sup>1</sup>, Miwa Matsumura<sup>1</sup>, Sachiko Ishibashi<sup>1</sup>, Masumi Ikeda<sup>1</sup>, Kohei Yamamoto<sup>1</sup>, Masanobu Kitagawa<sup>1</sup>

1. Department of Comprehensive Pathology, Graduate School of Medical and Dental Sciences, Tokyo Medical and Dental University, Tokyo, JAPAN

2. These authors contributed equally to this work.

**\*\*Corresponding Author:**

Morito Kurata,

1-5-45 Yushima, Bunkyo-ku, Tokyo 113-8510, JAPAN

E-mail address: kurata.pth2@tmd.ac.jp

# Abstract

**Background.** *MYC* is one of the proto-oncogenes contributing to tumorigenesis in many human cancers. Although the mechanism of *MYC* regulation is still not fully understood, learning about the comprehensive mechanism controlling the transcriptional activity of *MYC* will lead to therapeutic targets. The CRISPR/Cas9 library system is a simple and powerful screening technique. This study aims to identify new transcriptional upstream activators of *MYC* using the CRISPR activation library with new promoter-reporter systems.

**Methods and Results.** The *MYC* promoter-reporter system was developed with a photoconvertible fluorescent protein, Dendra2, and named “p*MYC*-promoter-Dendra2.” This *MYC* promoter-reporter system was designed to harbor a proximal *MYC* promoter at −3.1 kb. Both the CRISPR activation library and p*MYC*-promoter-Dendra2 were induced to HEK 293T cells, and Dendra2-positive cells, that are supposed that *MYC* should be upregulated, were collected individually by a cell sorter. Among the 169 cells collected, 12 clones were successfully established. Then, p*MYC*-promoter-Dendra2 was transfected again into these 12 clones, and 2 of 12 clones showed Dendra2 positivity. In this procedure, the cells with non-specific autofluorescence were correctly distinguished by utilizing the photoswitchable character of Dendra2. Using extracted guide DNA of these two Dendra2 positive clones, polymerase chain reaction (PCR) was performed to amplify the guide RNA (gRNA) containing region, which was introduced by the CRISPR activation library. Eventually, *PLEKHO2*, *MICU*, *MBTPS1*, and *MIAP* were identified, and these gRNAs were transfected individually into HEK 293T cells again using the CRISPR activation system. Only *MIAP* gRNA transfected cells showed Dendra2-positive fluorescence. Then, the overexpression vector for *MIAP* with a doxycycline-inducible vector confirmed that *MIAP* induced high *MYC* expression by real-time quantitative

PCR and western blot. Furthermore, the dual-luciferase assay showed a significant increase of promoter activity, and *MYC* mRNA was higher in *MIAP*-overexpressing cells.

*MIAP* is highly expressed in several cancers, though, a positive correlation between *MIAP* and *MYC* was observed only in human acute myeloid leukemia.

**Conclusion.** The present study confirmed that the experimental method using the CRISPR library technology functions effectively for the identification of molecules that activate endogenous *MYC*. This method will help elucidate the regulatory mechanism of *MYC* expression, as well as supporting further drug research against malignant tumors.

## Introduction

*MYC* is one of the most well-known proto-oncogenes, contributing to tumorigenesis in many human cancers. It can function as a transcriptional factor in both cancerous and non-cancerous cells, and it is involved in cell growth, proliferation, metabolism, and transformation (Dang, 2012). The Wnt- $\beta$ -catenin signaling pathway is one of the most well-described pathways involved in the regulation of *MYC*. Due to mutations in the *adenomatous polyposis coli* (*APC*) gene or abnormalities in *glycogen synthase kinase-3 $\beta$* , Wnt-signal is stimulated and the excessive accumulation of  $\beta$ -catenin occurs. Then, active  $\beta$ -catenin moves into the nucleus and binds to T cell factor (TCF), followed by inducing *MYC* expression (Liu et al., 2002) (Novak, Dedhar, 1999). Mitogen-activated protein kinase (MAPK), which is a serine/threonine kinase, and its subfamily extracellular signal-regulated kinase (ERK), regulate *MYC* transcriptional activity by inducing phosphorylation of *MYC* (MAPK/ERK signaling pathway). Furthermore, as non-coding RNA, microRNAs like *miR-34* (Kong et al., 2008) and *let-7* (Sampson et al., 2007) suppress *MYC* expression by binding to the 3'-untranslated region of *MYC* mRNA. As for long

noncoding RNAs, *PVT1* regulates the expression of *MYC* by inhibiting the degradation of *MYC* protein (Tseng et al., 2014). Thus, it has been reported that various regulatory mechanisms function in a complex fashion regarding *MYC* expression. Though, other molecules and elements like crosstalk between each pathway are not fully understood.

Concerning the high *MYC* expression rate of multiple malignant tumors, elucidating the mechanism controlling the transcriptional activity of *MYC* is a crucial issue because it will lead to therapeutic targets. Using the CRISPR/Cas9 library system, with its collection of more than 70,290 different guide RNAs (gRNAs), enables us to identify molecules that relate knockout or activation of endogenous genes. To date, the CRISPR library has been utilized for various studies. Shalem et al. (2014) identified genes whose loss is involved in resistance to vemurafenib by introducing the knockout library into a human melanoma cell line followed by treatment with *BRAF* inhibitor vemurafenib. In contrast, Konnermann et al. (2015) identified molecules that exhibit resistance to PLX-4720 by introducing an activation library. In this way, the CRISPR library has been used to identify the certain genes involved in drug sensitivity or resistance, toxin resistance, and hypoxia response (Konnermann et al., 2015; Gilbert et al., 2014; Kampmann, 2018; Kurata et al., 2018; Shalem et al., 2014; Wang et al., 2014; Koike-Yusa et al., 2014; Doench et al., 2016; Jain et al., 2016); however, a screening of molecules that act on the promoter region of a specific gene and control transcriptional activity has not yet been tested. In addition, autofluorescence is highly problematic for collecting fluorescent cells in the screening. Therefore, a photoconversion protein, Dendra2, was used in this system. Dendra2 is derived from the octocoral *Dendronephthya* sp. *T.* and can change color with UV-violet light (405 nm laser). This study aims to utilize the CRISPR activation library for the detection of new

transcriptional activators of *MYC*. For this purpose, we developed an original tool for evaluating *MYC* activity and establishing screening methods.

## Materials & Methods

### *MYC Promoter-Reporter System and Lentivirus Plasmids*

The Polymerase chain reaction (PCR) primers were designed across the proximal *MYC* promoter at -3.1 kb from the *MYC* transcriptional start site (TSS), and PCR was performed with KOD-FX (Toyobo). The primers were 5'-TTAGCTAGCGAGGGTTTCTTTGAGGGGC-3' (forward) and 5'-AACGGATCCCGGAGATTAGCGAGAGAGGA-3' (reverse).

A plasmid of Dendra2 (pDendra2) was purchased from Takara-Clontech. The cytomegalovirus immediate early promoter (PCMVIE) was removed with restriction enzymes (Afl III and Nhe I), and the multi-cloning site was replaced with Nhe I and BamH I. Amplified fragments of -3.1 kb of the *MYC* promoter region with Nhe I and BamH I restriction enzyme sites were integrated into pDendra2. The new reporter system with Dendra2 and the -3.1 kb *MYC* promoter region (p*MYC*-promoter-Dendra2) was used for further experiments. The double-stranded DNA fragments of *MIAP* with attB sites were purchased from gBlocks® (IDT) and generated into the pENTR221 vector using Gateway® BP clonase. Then they were transferred to the TripZ-TRE-DEST-IRES-GFP-EF1A-rtTA vector (Kurata *et al.*, 2016) using Gateway® LR clonase.

### *Cell Culture and SAM Library Screening*

Human embryonic kidney cells (HEK 293T) were maintained in Dulbecco's modified Eagle's medium (DMEM, Fuji Film) with 10% fetal bovine serum and penicillin/streptomycin with 5%

CO<sub>2</sub> at 37°C as previously used (Kurata *et al.*, 2016). To transduce the lentivirus, HEK 293T cells (5 x 10<sup>5</sup> cells/well) were seeded in a six-well cell culture plate one day before the transfection. The next day, 3 µg of lentivirus plasmid was transfected with 1 µg of pMDG and 2 µg of pCMV using Lipofectamine™ 3000 Reagent (Invitrogen) according to previous work (Kurata *et al.*, 2016). Twelve hours after transfection, the medium was changed to fresh DMEM. The virus supernatant was harvested at 48 h after transfection, then filtered with Millex™-HP 0.45 µm (Millipore). The target HEK 293T cells (5 x 10<sup>5</sup> cells/well) were seeded in a six-well plate one day prior to transduction and were transduced with this lentivirus supernatant with 5 µg/ml polybrene (Sigma). The used lentiviral plasmids were lentiMPHv2 (Addgene, #89308), lentiCAS-VP64\_blast (Addgene, #61425), and lentiSAMv2 (Addgene, #61597). At first, lentiMPHv2 was transduced into HEK 293T and treated with Hygromycin B for 2 weeks. Then, lentiCAS9-VP64 was induced in MPH-expressing HEK 293T cells and treated with Blasticidin for 2 weeks. The MPH-dCas9-VP64-expressing HEK 293T applied to further experiments. The control and *MYC* activation gRNA (Fig. S1C) were induced in MPH-dCas9-VP64-expressing HEK 293T cells. The CRISPR activation library, lentiSAMv2, was also transduced into MPH-dCas9-VP64-expressing HEK 293T cells and then treated with Zeocin (300 µg/ml, Invivogen) for 2 weeks.

### ***Collection of Dendra2-Positive Cells and Identification of gRNAs***

After 2 weeks of selection with Zeocin, p*MYC*-promoter-Dendra2 was transfected into SAM library lentivirus-transduced cells with Lipofectamine™ 3000 Reagent (Invitrogen). HEK 293T cells transfected with p*MYC*-promoter-Dendra2 were evaluated for Dendra2-positive cells by fluorescence microscope (Keyence). Cells were collected at 72 h and washed with phosphate-

buffered saline (PBS) and applied to the cell sorter. The Dendra2-positive cells were collected into a 96-well plate with MoFlo XDP (Beckman Coulter).

A total of 169 cells were collected, and 12 clones were successfully established. Collected clones were centrifuged and washed with PBS solution and pelletized. Lysis buffer solution with proteinase K (20 mg/ml) was added to cells and left for 30 min at 60°C. After treatment at 94°C for 10 min, the lysate was centrifuged for 15 min, and the supernatant was collected as genome DNA (gDNA) containing fractions. Using extracted gDNA of each collected clone, PCR was conducted with GoTaq® qPCR Master Mix (Promega) with the following primers: 5'-GAGGGCCTATTTCCCATGAT -3' (forward) and 5'-GAGGGCCTATTTCCCATGAT-3' (reverse). PCR product containing the gRNA region was employed for TA cloning with pGEM®-T Vector Systems (Promega) and gRNAs were sequenced by Sanger-sequencing.

# ***Preparation of CRISPR backbone plasmid with target gRNAs and Transcriptional activation for MYC using the CRISPR Activation System***

The oligonucleotides for each target gRNA sequence were designed using the list from the SAM library and integrated into the pENTR221/U6/stuffer/MS2 vector (Kurata et al., 2018). The MPH-dCas9-VP64-expressing HEK 293T cells (1 x 10<sup>5</sup> cells/ml) were seeded in a 24-well cell culture plate. The next day, 2.5 µg of individual gRNA vector and pMYC-promoter-Dendra2 were transfected with Lipofectamine™ 3000 Reagent. At 48 hours after the transfection, the MYC transcriptional activation was evaluated by Dendra2-positivity by Keyence.

# ***Evaluation of MYC expression***



A pTRIPZ-*MIAP-GFP* was transduced into the HEK 293T cells. These transduced cells were selected with puromycin (1 µg/ml, Invitrogen) for 2 weeks. Seventy-two hours after the induction of Doxycycline (DOX) (1 µg/ml), the collected cells were employed for RNA isolation.

RNA was isolated using an RNeasyR Mini Kit (Qiagen) according to the manufacturer's instructions. Complementary DNA (cDNA) was generated from RNA with ReverTra AceR qPCR RT Master Mix (Toyobo). *Beta-ACTIN* was used as an endogenous control. Using the ABI Prism 7900HT (Applied Biosystems), quantitative PCR (qPCR) analysis was performed to quantify the RNA level using SYBR Mix. The sequences for the PCR primers used in gene expression were as follows: *MYC*: 5'-CGACTCTGAGGAGGAACAAGAA-3' (forward) and 5'-CAGCAGAAGGTGATCCAGACT-3' (reverse), *β-ACTIN*: 5'-CACAGAGCCTCGCCTTTGCC-3' (forward) and 5'-CACAGAGCCTCGCCTTTGCC-3' (reverse). The mRNA level of the targeted gene was analyzed by comparison with the standard calibration curve. Western blot analysis was performed according to standard procedures (Kurata et al., 2016). The primary antibodies used were anti-c-Myc (Abcam ab32072) and anti-GAPDH (Santa Cruz sc47724), each at a dilution of 1:1000. The primary antibodies were detected with horseradish peroxidase (HRP)-conjugated secondary antibodies at 1:50000. Protein bands were visualized with Clarity™ Western ECL Substrate (Bio-Rad).

### ***Dual-Luciferase Reporter Assay***

The fragments of the *MYC* promoter was integrated into a pGL4 vector (Promega). A pT3.5-CAG-*MIAP* was generated with pENTR221-*MIAP* and pT3.5-CAG-DEST (Kurata et al., 2018) using Gateway® LR clonase. The luciferase reporter vectors, the pT3.5-CAG-*MIAP*

vector, and the pRL Renilla luciferase reporter vector were co-transfected in the HEK 293T cells. The samples were harvested 48 and 72 hours after the transfection. Each assay was biologically triplicated and repeated. Luciferase activities were measured using the dual-luciferase reporter assay system (Promega) and Lumat LB9507 (Perkin Elmer). RNA was harvested at the same time points and the expression of *MYC* mRNA was analyzed by qPCR.

### ***MIAP and MYC expressions in databases***

For the purpose of comparing gene expression between cancer and non-cancerous tissue, the GEPIA web server was employed (Tang et al., 2017) (<http://gepia2.cancer-pku.cn/>). Datasets for AML samples were used in BloodSpot database, that database was used to retrieve gene expression data from curated human and murine microarray and RNA-sequencing (RNA-Seq) datasets (Bagger et al., 2016). GSE42519, GSE13159, GSE15434, GSE61804, GSE14468, and The Cancer Genome Atlas (TCGA) datasets were used for analysis. In addition, following database were used; The Human Protein Atlas (<https://www.proteinatlas.org/>) and cBioPortal (<https://www.cbioportal.org>) using TCGA data (Cancer Genome Atlas Research Network et al., 2013; Gao et al., 2013; Cerami et al., 2012)

### ***Statistical analysis***

The data were statistically analyzed using the EZR version 1.36 software. A value of  $p < 0.05$  was considered statistically significant for all analyses.

## **Results**

### ***Development of the MYC Promoter-Reporter System for Library Screening***

Using PCR primers designed across the proximal *MYC* promoter at -3.1 kb (-3134 to -40) from the *MYC* TSS, PCR was performed. The *MYC* promoter region was integrated upstream of *Dendra2*. The pDendra2 harboring the 3.1 kb *MYC* promoter region was called “p*MYC*-promoter-Dendra2” for further experiments (Fig. 1SA). A plasmid p*MYC*-promoter-Dendra2 and gRNA of *MYC* and *HPRT* (as a negative control) were transfected in HEK 293T cells using the CRISPR activation system. We confirmed that Dendra2 showed positivity only in the cells with gRNA for *MYC* (Fig. 1SB, C). Therefore, p*MYC*-promoter-Dendra2 was utilized as a *MYC* promoter-reporter analysis tool in this study.

### ***MYC Transcriptional Activator Screening Using the CRISPR Activation Library***

To screen transcriptional activators working on the *MYC* promoter region, the CRISPR activation library was employed. The SAM library lentivirus was infected into MPH-dCas9-VP64-expressing HEK 293T cells ( $1 \times 10^7$  cells). After selection, p*MYC*-promoter-Dendra2 was transfected into these HEK 293T cells. The multiplicity of infection (MOI) ranged from 1.4 to 5.2.

At 72 h after transfection, cells were proceeded to Dendra2-positive cell collection using flow cytometry (positive ratio: 0.017%). Sorted single cells were collected into a 96-well plate and incubated for 2 weeks. Among the collected 169 cells, 12 clones were successfully established (Fig. 1).

### ***Second Screening to Identify Candidate Genes as MYC Activators***

For the second screening, p*MYC*-promoter-Dendra2 was again transfected into the 12 clones. Using fluorescence microscope, clones 3 and 5 were shown to be truly Dendra2 positive (Fig. 2A-L), and the photoswitchable attribute of Dendra2 was useful for distinguishing it from cells

with non-specific autofluorescence (Fig. 2S). Using extracted gDNA of clones 3 and 5, PCR was performed to amplify the gRNA containing region, which was introduced by the CRISPR activation library. Then, the PCR product was employed for TA cloning, followed by colony PCR. By this method, four candidate genes—*PLEKHO2*, *MICU*, *MBTPS1*, and *MIAP* were identified (Table 1).

Each candidate gRNA was transfected to the HEK 293T cells using the CRISPR activation system accompanied by pMYC-promoter-Dendra2. Because Dendra2 was positive only in the cells transfected with *MIAP* gRNA (Fig. 3A-D), *MIAP* was considered as the most promising gene that would increase *MYC* transcriptional activity. Then, double-stranded DNA fragments coding the full length of *MIAP* induced into Doxycycline-inducible lentiviral vector (pTRIPZ-*MIAP-GFP*, Fig. 4A). The pTRIPZ-*MIAP-GFP* was introduced into HEK 293T cells. Seventy-two hours after the induction *MIAP* with DOX (1 µg/ml), high *MYC* expression was observed in DOX-induced cells by RT qPCR (*t*-test,  $p < 0.05$ , Fig. 4B). Furthermore, in the western blots, an increase of *MYC* expression was also observed in DOX-induced cells (Fig. 4C).

Additionally, a dual-luciferase assay conducted with a reporter containing the *MYC* promoter region showed a significant increase of promoter activity in *MIAP*-overexpressing cells at both 48 and 72 hours (Fig. 4D and 3S). On the other hand, the *MYC* mRNA, which was isolated at the same time points, showed mildly higher in *MIAP*-overexpressing cells at 48 hours but not at 72 hours (Fig. 4E and 3S). Taken together, although *MIAP* was confirmed to activate the *MYC* promoter strongly, the *MYC* mRNA itself might be regulated also by other factors.

### ***Correlation Between MYC and MIAP in the Database***

According to the GEPIA web server (Tang Z *et al.*, 2017) (<http://gepia2.cancer-pku.cn/>), high *MIAP* expression is observed in cervical carcinoma, acute myeloid leukemia (AML), ovarian carcinoma, thyroid carcinoma, uterine endometrial carcinoma, and uterine carcinosarcoma when compared to normal tissue (Fig. 4S A). Among these carcinomas, a correlation analysis using cBioPortal revealed a mild positive correlation between *MYC* and *MIAP* only in AML (Fig. 5A, Fig. 4S B-F). Therefore, we focused on hematopoietic cells, and *MIAP* expression was analyzed using the BloodSpot database. As a result, *MIAP* was significantly higher in AML tissue compared to normal bone marrow tissue (Fig. 5B). Interestingly, *MIAP* was highly expressed in AML, especially with t (8;21) (Fig. 5C).

## Discussion

In this study, we created a new reporter vector that incorporates the *MYC* promoter region across the proximal -3.1 kb from the TSS and photoconversion fluorescent protein to evaluate the transcriptional activity of endogenous *MYC*. Ideally, a large region of the *MYC* promoter on Chr.8q24 would be applied to better understand the full *MYC* promoter mechanism. However, even short region on the *MYC* promoter region are still not fully understood. In designing this new vector, 3.1 kb of the promoter region was applied because of the limited length of effective DNA insertion into the vector. Future improved experiments on larger region will help to clarify the whole mechanism of *MYC* regulation via the promoter region.

By applying this tool to the CRISPR library, we screened molecules that increase the expression of endogenous *MYC*. According to previous reports, by introducing the CRISPR library into a malignant tumor cell line and carrying out selection with a drug, molecules that cause susceptibility or resistance to that drug can be identified (Konermann *et al.*, 2015; Gilbert *et al.*, 2014; Kampmann, 2018; Kurata *et al.*, 2018; Shalem *et al.*, 2014; Wang *et al.*, 2014;

*Koike-Yusa et al., 2014; Doench et al., 2016; Jain et al., 2016*). However, this is the first specific report of a molecule that activates transcription upstream of *MYC*.

*MYC* is an indispensable factor for cell growth and proliferation in normal tissues, but at the same time, it is one of the most well-known proto-oncogenes, with high amplification in various carcinomas. Moreover, *MYC* is widely involved in tumor development and growth (*Dang, 2012*). Elucidation of factors that regulate the transcriptional activity of *MYC* is crucial for establishing new therapeutic strategies for tumors. Although *MYC* expression is strictly regulated by mRNA and MYC protein, in 1998, *He et al.* reported that inactivation of *APC* located downstream of Wnt signaling pathway results in excessive accumulation of  $\beta$ -catenin and promotes endogenous *MYC* expression by acting on TCF present in the promoter region of *MYC*.

Regarding the research of the regulatory mechanism of *MYC*, *Vo et al.* (2018) transplanted CRISPR/dCas9-VP160 and gRNAs that activated *MYC* transcription in the mouse brain, resulting in the development of medulloblastoma. They concluded that this model is useful for future drug research by suppressing *MYC* expression.

Because this screening was adapted to about 70,290 kinds of gRNAs, initially, multiple molecules were expected to be identified; however, the screening eventually led to the identification of only one molecule, *MIAP*. The reason for this is that, on performing single-cell collection using flow cytometry, to avoid collecting false-positive cells, the cells with the highest fluorescence intensity were collected; as a result, the number of collected cells was extremely small. Furthermore, since the proliferation of the cells after the collection was poor, it is expected that collected cells included cells that were highly brightened due to cell senescence. Moreover, oncogene-induced cell senescence can be one explanation for why the survival rate of collected single cells was very low. Oncogene-induced senescence, which is now a well-known

phenomenon, causes cell growth arrest where the oncogene is excessive. This was first reported in *RAS* (Serrano *et al.*, 1997), and the same mechanism has been reported for *MYC*, where *MYC* oncogene overexpression results in inactivation of *MYC*, followed by the arrest of cell growth and proliferation (Wu *et al.*, 2007). Excessive *MYC* expression due to the CRISPR activation system may have induced cell senescence and caused a low survival rate of sorted cells. We would like to use a next-generation sequencing system and adjust the multiplicity of infection in future screenings.

By newly established experimental line in this study, *MIAP*, which was found as a transcription factor for *MYC*, was proved to function strongly as a *MYC* activator by promoter assay. On the other hand, *MYC* expression was mildly elevated in the *MIAP*-overexpressing cells collected at the same time as the luciferase assay. These results suggest that the *MYC* mRNA is complexly controlled by multiple factors and multiple pathways.

In conclusion, the present study confirmed that this experimental method applying the CRISPR library technology functioned effectively for the identification of the molecule that activates endogenous *MYC*. This method will help elucidate the regulatory mechanism of *MYC* expression, as well as facilitating further drug research against malignant tumors. However, more investigation is required to establish a more accurate and precise screening technique.

## Acknowledgments

The authors would like to thank for the member of stem cell laboratory Tokyo Medical and Dental University, for cell sorting and her technical assistance and advice.

## References

- Bagger FO, Sasivarevic D, Sohi SH, Laursen LG, Pundhir S, Sonderby CK, Winther O, Rapin N, Porse BT. 2016. BloodSpot: a database of gene expression profiles and transcriptional programs for healthy and malignant haematopoiesis. *Nucleic acids research*, 44:D917-24.
- Cancer Genome Atlas Research Network, Weinstein JN, Collisson EA, Mills GB, Shaw KR, Ozenberger BA, Ellrott K, Shmulevich I, Sander C, Stuart JM. 2013. The Cancer Genome Atlas Pan-Cancer analysis project. *Nature genetics*, 45:1113-1120.
- Cerami E, Gao J, Dogrusoz U, Gross BE, Sumer SO, Aksoy BA, Jacobsen A, Byrne CJ, Heuer ML, Larsson E, Antipin Y, Reva B, Goldberg AP, Sander C, Schultz N. 2012. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer discovery*, 2:401-404.
- Dang CV. 2012. MYC on the path to cancer. *Cell*, 149:22-35.
- Doench JG, Fusi N, Sullender M, Hegde M, Vaimberg EW, Donovan KF, Smith I, Tothova Z, Wilen C, Orchard R, Virgin HW, Listgarten J, Root DE. 2016. Optimized sgRNA design to maximize activity and minimize off-target effects of CRISPR-Cas9. *Nature biotechnology*, 34:184-191.
- Gao J, Aksoy BA, Dogrusoz U, Dresdner G, Gross B, Sumer SO, Sun Y, Jacobsen A, Sinha R, Larsson E, Cerami E, Sander C, Schultz N. 2013. Integrative analysis of complex cancer genomics and clinical profiles using the cBioPortal. *Science signaling*, 6:pl1.



347 Gilbert LA, Horlbeck MA, Adamson B, Villalta JE, Chen Y, Whitehead EH, Guimaraes C,  
 348 Panning B, Ploegh HL, Bassik MC, Qi LS, Kampmann M, Weissman JS. 2014. Genome-Scale  
 349 CRISPR-Mediated Control of Gene Repression and Activation. *Cell*, 159:647-661.

350 Jain IH, Zazzeron L, Goli R, Alexa K, Schatzman-Bone S, Dhillon H, Goldberger O, Peng J,  
 351 Shalem O, Sanjana NE, Zhang F, Goessling W, Zapol WM, Mootha VK. 2016. Hypoxia as a  
 352 therapy for mitochondrial disease. *Science (New York, N.Y.)*, 352:54-61.

353 Kampmann M. 2018. CRISPRi and CRISPRa Screens in Mammalian Cells for Precision Biology  
 354 and Medicine. *ACS chemical biology*, 13:406-416.

355 Koike-Yusa H, Li Y, Tan EP, Velasco-Herrera Mdel C, Yusa K. 2014. Genome-wide recessive  
 356 genetic screening in mammalian cells with a lentiviral CRISPR-guide RNA library. *Nature*  
 357 *biotechnology*, 32:267-273.

358 Konermann S, Brigham MD, Trevino AE, Joung J, Abudayyeh OO, Barcena C, Hsu PD, Habib  
 359 N, Gootenberg JS, Nishimasu H, Nureki O, Zhang F. 2015. Genome-scale transcriptional  
 360 activation by an engineered CRISPR-Cas9 complex. *Nature*, 517:583-588.

361 Kong YW, Cannell IG, de Moor CH, Hill K, Garside PG, Hamilton TL, Meijer HA, Dobbyn HC,  
 362 Stoneley M, Spriggs KA, Willis AE, Bushell M. 2008. The mechanism of micro-RNA-  
 363 mediated translation repression is determined by the promoter of the target gene. *Proceedings*  
 364 *of the National Academy of Sciences of the United States of America*, 105:8866-8871.

365 Kurata M, Rathe SK, Bailey NJ, Aumann NK, Jones JM, Veldhuijzen GW, Moriarity BS,  
 366 Largaespada DA. 2016. Using genome-wide CRISPR library screening with library resistant  
 367 DCK to find new sources of Ara-C drug resistance in AML. *Scientific reports*, 6:36199.

368 Kurata M, Wolf NK, Lahr WS, Weg MT, Kluesner MG, Lee S, Hui K, Shiraiwa M, Webber BR,  
369 Moriarity BS. 2018. Highly multiplexed genome engineering using CRISPR/Cas9 gRNA  
370 arrays. *PloS one*, 13:e0198714.

371 Kurata M, Yamamoto K, Moriarity BS, Kitagawa M, Largaespada DA. 2018. CRISPR/Cas9  
372 library screening for drug target discovery. *Journal of human genetics*, 63:179-186.

373 Liu C, Li Y, Semenov M, Han C, Baeg GH, Tan Y, Zhang Z, Lin X, He X. 2002. Control of  
374 beta-catenin phosphorylation/degradation by a dual-kinase mechanism. *Cell*, 108:837-847.

375 Novak A, Dedhar S. 1999. Signaling through beta-catenin and Lef/Tcf. *Cellular and molecular*  
376 *life sciences : CMLS*, 56:523-537.

377 Sampson VB, Rong NH, Han J, Yang Q, Aris V, Soteropoulos P, Petrelli NJ, Dunn SP, Krueger  
378 LJ. 2007. MicroRNA let-7a down-regulates MYC and reverts MYC-induced growth in Burkitt  
379 lymphoma cells. *Cancer research*, 67:9762-9770.

380 Serrano M, Lin AW, McCurrach ME, Beach D, Lowe SW. 1997. Oncogenic ras provokes  
381 premature cell senescence associated with accumulation of p53 and p16INK4a. *Cell*, 88:593-  
382 602.

383 Shalem O, Sanjana NE, Hartenian E, Shi X, Scott DA, Mikkelsen T, Heckl D, Ebert BL, Root  
384 DE, Doench JG, Zhang F. 2014. Genome-scale CRISPR-Cas9 knockout screening in human  
385 cells. *Science (New York, N.Y.)*, 343:84-87.

386 Tang Z, Li C, Kang B, Gao G, Li C, Zhang Z. 2017. GEPIA: a web server for cancer and normal  
387 gene expression profiling and interactive analyses. *Nucleic acids research*, 45:W98-W102.

388 Tseng YY, Moriarity BS, Gong W, Akiyama R, Tiwari A, Kawakami H, Ronning P, Reuland B,  
389 Guenther K, Beadnell TC, Essig J, Otto GM, O'Sullivan MG, Largaespada DA, Schwertfeger

KL, Marahrens Y, Kawakami Y, Bagchi A. 2014. PVT1 dependence in cancer with MYC copy-number increase. *Nature*, 512:82-86.

Vo BT, Kwon JA, Li C, Finkelstein D, Xu B, Orr BA, Sherr CJ, Roussel MF. 2018. Mouse medulloblastoma driven by CRISPR activation of cellular Myc. *Scientific reports*, 8:8733-018.

Wang T, Wei JJ, Sabatini DM, Lander ES. 2014. Genetic screens in human cells using the CRISPR-Cas9 system. *Science (New York, N.Y.)*, 343:80-84.

Wu CH, van Riggelen J, Yetil A, Fan AC, Bachireddy P, Felsher DW. 2007. Cellular senescence is an important mechanism of tumor regression upon c-Myc inactivation. *Proceedings of the National Academy of Sciences of the United States of America*, 104:13028-13033.

## Figure Legends

**Figure 1. A scheme for MYC transcriptional activator screening with the CRISPR activation library.**

**Figure 2. Dendra2-positive cells using fluorescence microscope.**

(A)-(L) correspond Clone 1-12. pMYC-promoter-Dendra2 was transfected into 12 clones that were successfully cloned after CRISPR activation screening. Clones 3 and 5 showed Dendra2 positivity.

**Figure 3. Validation experiments for candidate genes as MYC activators.**

(A) *PLEKHO2* (B) *MICU3* (C) *MBTPS1* (D) *MIAP*

With gRNAs of the four nominated genes, pMYC-promoter-Dendra2 was transfected into HEK 293T cells by applying the CRISPR activation system. Only *MIAP*-introduced cells were Dendra2 positive.

**Figure 4. *MYC* mRNA expression analyzed by real-time quantitative polymerase chain reaction.**

(A) Vector map of Dox-inducible *MIAP* expression vector. (B) High *MYC* expression was observed in DOX-induced cells ( $t$ -test,  $p < 0.05$ ). (C) Western blotting analysis showed increasing of *MYC* protein in DOX- induced cells. (D) Luciferase activity at 48 hours was significantly higher in *MIAP*-overexpressing cells ( $t$ -test,  $p < 0.05$ ). (E) *MYC* mRNA expression at 48 hours was mildly higher in *MIAP*-overexpressing cells ( $t$ -test,  $p < 0.05$ ).

**Figure 5. Correlation between *MYC* and *MIAP* in the database.**

(A) In cBioPortal using TCGA data, a correlation analysis between *MYC* and *MIAP* in AML revealed a mild positive correlation. (B) Using the Bloodspot database *MIAP* mRNA expression in acute myeloid leukemia (AML, N=989) was significantly higher than in normal tissue (N=397). (Mann–Whitney U test,  $p < 0.05$ ) (C) The BloodSpot database was used to retrieve gene expression data. *MIAP* expression was higher in each subtype of AML, especially with t (8;21). (Kruskal–Wallis test, \*:  $p < 0.01$ )

**Table 1. List of Candidate Genes for *MYC* Activation**

**Supplementary Figures**

**Fig. 1S. Construct of *MYC* promoter-reporter system.**

**Fig. 2S. Photoswitchable attribute of Dendra2.**

**Fig. 3S. Luciferase activity in *MIAP*-overexpressing cells and *MYC* mRNA expression at 72 hours. *MYC* promoters are strongly activated at 72 hours after induction of *MIAP* expression, but the mRNA of *MYC* was not significantly upregulated at 72 hours.**

**Fig. 4S. *MIAP* expression and correlation with *MYC* in databases.**

(A) Comparison of *MIAP* expression in normal and tumor tissue in GEPIA.

447 (B)-(F) Correlations between *MYC* and *MIAP* in cBioPortal.  
 448 (B) Cervical carcinoma. (C) Ovarian carcinoma. (D) Thyroid carcinoma. (E) Uterine endometrial  
 449 carcinoma. (F) Uterine carcinosarcoma.  
 450

# **Table 1**(on next page)


List of Candidate Genes for *MYC* Activation

**Table 1**

Sequence of gRNA	Gene name
GGGCGGGAAATGGGTGGGGA	<i>PLEKHO2</i>
TTGTGACACTACTCCAGCCT	<i>MICU1</i>
AGGATCCCCGAAAAGGAGCA	<i>MBTPS1</i>
TCTCAGGGTATCTAGGGACT	<i>MIAP</i>

# Figure 1

A scheme for *MYC* transcriptional activator screening with the CRISPR activation library.






# Figure 2

Dendra2-positive cells using fluorescence microscope.


(A)-(L) correspond Clone 1-12. p MYC- promoter-Dendra2 was transfected into 12 clones that were successfully cloned after CRISPR activation screening. Clones 3 and 5 showed Dendra2 positivity.



# Figure 3

Validation experiments for candidate genes as *MYC* activators

(A) *PLEKHO2* (B) *MICU3* (C) *MBTPS1* (D) *M1AP* With gRNAs of the four nominated genes, pMYC- promoter-Dendra2 was transfected into HEK 293T cells by applying the CRISPR activation system. Only *M1AP*-introduced cells were Dendra2 positive.




# Figure 4


*MYC* mRNA expression analyzed by real-time quantitative polymerase chain reaction.

(A) Vector map of Dox-inducible *M1AP* expression vector. (B) High *MYC* expression was observed in DOX-induced cells ( $t$ -test,  $p < 0.05$ ). (C) Western blotting analysis showed increasing of MYC protein in DOX- induced cells. (D) Luciferase activity at 48 hours was significantly higher in *M1AP*-overexpressing cells ( $t$ -test,  $p < 0.05$ ). (E) *MYC* mRNA expression at 48 hours was mildly higher in *M1AP*-overexpressing cells ( $t$ -test,  $p < 0.05$ ).


**A**




**B**




**C**



**D**



**E**



# Figure 5

Correlation between *MYC* and *M1AP* in the database.

(A) In cBioPortal using TCGA data, a correlation analysis between *MYC* and *M1AP* in AML revealed a mild positive correlation. (B) Using the Bloodspot database *M1AP* mRNA expression in acute myeloid leukemia (AML, N=989) was significantly higher than in normal tissue (N=397). (Mann-Whitney U test,  $p < 0.05$ ) (C) The BloodSpot database was used to retrieve gene expression data. *M1AP* expression was higher in each subtype of AML, especially with t (8;21). (Kruskal-Wallis test, \*:  $p < 0.01$ )

