

# On the importance of spatial scales on beta diversity of coral assemblages: a study from Venezuelan coral reefs

Emy Miyazawa<sup>Corresp., 1</sup>, Luis M Montilla<sup>1, 2</sup>, Esteban Alejandro Agudo-Adriani<sup>1, 3</sup>, Alfredo Ascanio<sup>1, 4</sup>, Gloria Mariño-Briceño<sup>1</sup>, Aldo Croquer<sup>Corresp. 1, 5</sup>

<sup>1</sup> Laboratorio de Ecología Experimental, Universidad Simón Bolívar, Caracas, Venezuela

<sup>2</sup> Integrative Marine Ecology Department, Stazione Zoologica Anton Dohrn, Naples, Italy

<sup>3</sup> Department of Biology, University of North Carolina at Chapel Hill, North Carolina, United States

<sup>4</sup> Department of Biology, Miami University of Ohio, Oxford, Ohio, United States

<sup>5</sup> Centro de Innovación Marina, The Nature Conservancy, Punta Cana, Dominican Republic

Corresponding Authors: Emy Miyazawa, Aldo Croquer

Email address: 12-11222@usb.ve, acroquer@usb.ve

Estimating variability across spatial scales has been a major issue in ecology because describing patterns in space is extremely valuable to propose specific hypotheses aim to unveil key processes behind these patterns. This paper aims to estimate the variability of the coral assemblage structure at different spatial scales in order to determine which scales explain the greatest variability on  $\beta$ -diversity. For this, a fully-nested design including a series of hierarchical-random factors encompassing three spatial scales: (1) regions, (2) localities and (3) reefs sites across the Venezuelan territory. The variability among spatial scales were tested with a permutation analysis of variance (Permanova) based on Bray-Curtis index. Dispersion in species presence/absence across scales (i.e.,  $\beta$ -diversity) was tested with a PermDisp analysis based on Jaccard's index. We found the highest variability in the coral assemblage structure between sites within locality (Pseudo- $F = 5.34$ ;  $p$ -value = 0.001, CV = 35.10%), We also found that longitude (Canonical corr = 0.867,  $p = 0.001$ ) is a better predictor of the coral assemblage structure in Venezuela, rather than latitude (Canonical corr = 0.552,  $p = 0.021$ ). Largest changes in  $\beta$ -diversity of corals occurred within sites ( $F = 2.764$ ,  $df_1 = 35$ ,  $df_2 = 107$ ,  $p = 0.045$ ) and within localities ( $F = 4.438$ ,  $df_1 = 6$ ,  $df_2 = 29$ ,  $p = 0.026$ ). Our results suggest that processes operating at spatial scales of hundreds of meters and hundreds of kilometers might both be critical to shape coral assemblage structure in Venezuela, whereas smaller scales (i.e., hundreds of meters) showed to be highly- important for the species turnover component of  $\beta$ -diversity. This result highlights the importance of creating scale-adapted management actions in Venezuela and likely across the Caribbean region.

# On the importance of spatial scales on beta diversity of coral assemblages: a study from Venezuelan coral reefs

Emy Miyazawa<sup>1</sup>, Luis M. Montilla<sup>1, 2</sup>, Esteban Alejandro Agudo-Adriani<sup>1, 3</sup>, Alfredo Ascanio<sup>1, 4</sup>, Gloria Mariño-Briceño<sup>1</sup>, and Aldo Cróquer<sup>1, 5</sup>

<sup>1</sup>Laboratorio de Ecología Experimental, Universidad Simón Bolívar

<sup>2</sup>Integrative Marine Ecology Department, Stazione Zoologica Anton Dohrn

<sup>3</sup>Department of Biology, University of North Carolina at Chapel Hill

<sup>4</sup>Department of Biology, Miami University of Ohio

<sup>5</sup>Centro de Innovación Marina, The Nature Conservancy

Corresponding author:

Emy Miyazawa, Aldo Cróquer<sup>1</sup>

Email address: 12-11222@usb.ve, acroquer@usb.ve

## ABSTRACT

Estimating variability across spatial scales has been a major issue in ecology because describing patterns in space is extremely valuable to propose specific hypotheses aim to unveil key processes behind these patterns. This paper aims to estimate the variability of the coral assemblage structure at different spatial scales in order to determine which scales explain the greatest variability on  $\beta$ -diversity. For this, a fully-nested design including a series of hierarchical-random factors encompassing three spatial scales: (1) regions, (2) localities and (3) reefs sites across the Venezuelan territory. The variability among spatial scales were tested with a permutation analysis of variance (Permanova) based on Bray-Curtis index. Dispersion in species presence/absence across scales (i.e.,  $\beta$ -diversity) was tested with a test of homogeneity of dispersions (PermDisp) based on Jaccard's index. We found the highest variability in the coral assemblage structure between sites within locality (Pseudo-F = 5.34; p-value = 0.001, CV = 35.10%). We also found that longitude (Canonical corr = 0.867, p = 0.001) is a better predictor of the coral assemblage structure in Venezuela, rather than latitude (Canonical corr = 0.552, p = 0.021). Largest changes in  $\beta$ -diversity of corals occurred within sites (F = 2.764, df1 = 35, df2 = 107, p = 0.045) and within localities (F = 4.438, df1 = 6, df2 = 29, p = 0.026). Our results suggest that processes operating at spatial scales of hundreds of meters and hundreds of kilometers might both be critical to shape coral assemblage structure in Venezuela, whereas smaller scales (i.e., hundreds of meters) showed to be highly- important for the species turnover component of  $\beta$ -diversity. This result highlights the importance of creating scale-adapted management actions in Venezuela and likely across the Caribbean region.

## INTRODUCTION

The importance of scales in ecology has been largely acknowledged for decades (Schneider, 2001; Mac Nally and Quinn, 1998; Wiens, 1989). MacArthur (1972) and Levin (1992) assays deeply influenced modern ecologists by promoting the view that ecological processes act at a variety of spatial and temporal scales, and they generate patterns that may differ from those at which processes act (Chave, 2013). Today it is known that ecological dynamics tends to be stochastic at small scales, but variability is conditional on the resolution of description (Chave, 2013; Levin, 1992). Furthermore, there has been an increased recognition that the problem of scale at which ecological processes act, should be considered as critical if it is wanted to produce general predictions about patterns in space and time (Chave, 2013). Thus, modern ecological thinking agrees that to understand a system (e.g. a community), it is important to study it at the appropriate scale (Chave, 2013).


It is clear that increasing consideration of scale is helping to address a key issue in ecology: the question of what influences the distribution and abundance of organisms (McGill, 2010). Species distributions

depend on four important processes: (1) climate, (2) species interactions, (3) habitat structure and (4) dispersal capabilities, each one operating with different strength at a range of spatial scales (McGill, 2010). Generally, the presence or absence of organisms within a community may depend on rare or large-scale (region-specific) dispersal and colonization events, while local abundance is more a function of frequent, fine-spatial scale processes such as biotic interactions and habitat heterogeneity, e.g. Ricklefs (1987). This implies that communities are structured by both abiotic and biotic factors nested along different spatial scales which often occur along environmental gradients (Johnson and Goedkoop, 2002; Whittaker and Heegaard, 2003). Concomitantly, the species richness of a community is also expected to be highly dependent on spatial scales evaluated (Barton et al., 2013; Field et al., 2009; Melchior et al., 2017; Whittaker et al., 2001).

Coral reefs are one of the most complex and diverse ecosystems of the planet. Reef species diversity has been estimated at 600,000 to more than 9 million species worldwide (Plaisance et al., 2011; Reaka-Kudla, 1997). The habitat and shelter for the majority of these species is largely provided by scleractinian corals (Alvarez-filip et al., 2009). There is compelling evidence indicating that ecological processes controlling the structure of coral assemblages (e.g. substrate availability, recruitment, competition, and herbivory) are strongly dependent on spatial scales (Pandolfi, 2002). In addition, oceanographic processes which partly define the environmental setting of a reef are also extremely variable within habitats, across sites, reef systems, and regions (Chollett et al., 2010; Eidens et al., 2015). Furthermore, biological and environmental factors may interact with each other to produce different patterns in species distribution across several spatial scales. In consequence, understanding the underlying factors controlling the coral species richness in a reef is not a simple task (Edmunds, 2013; Eidens et al., 2015; Karlson and Cornell, 1999) for it is a multi-scale problem (Komyakova et al., 2018).

Total species richness of a region, frequently named gamma diversity ( $\gamma$ ), can be partitioned in two components: (1)  $\alpha$ -diversity (i.e., the number of species by site), and (2)  $\beta$ -diversity (i.e., the variation in the species identities from site to site, *sensu* Whittaker (1960, 1972)). For decades, ecologists have debated ways to estimate and interpret  $\alpha$  and  $\beta$ -diversity; but in recent years, the study of  $\beta$ -diversity has gained a lot of interest for it is what actually makes assemblages of species more or less similar to one another at different places and times (Anderson et al., 2011; Vellend, 2010). Many different measures of  $\beta$ -diversity have been introduced, but there is no overall consensus about which ones are most appropriate for addressing particular ecological questions (Jurasinski et al., 2009; Tuomisto, 2010a,b). Anderson et al. (2011) distinguished two types of  $\beta$ -diversity: (a) turn-over and (b) variation. Turn-over refers to changes in community structure among sampling units distributed along well-defined environmental gradients, whereas variation portrays variability in species composition among sample units within a given spatial or temporal extent, or within a given category of a factor (such as a habitat type or experimental treatment). On the other hand, Baselga (2010) partitioned the total  $\beta$ -diversity into two components: (1) nestedness, i.e. when the species composition of sample units with low richness represent a subset of the species found in the richest sample units, and (2) species replacement, i.e. a turn-over of species (Gaston et al., 2007; Leibold et al., 2004; Svenning et al., 2011). Regardless the point of view, the study of each of these components is relevant to understand processes that control ecological communities and a range of ecosystem functions (Arias-González et al., 2008; Harborne et al., 2006).

While spatial patterns of  $\gamma$  and  $\alpha$ -diversity of coral assemblages have been studied extensively; only few studies have focused on measuring  $\beta$ -diversity, e.g. Arias-González et al. (2008); Connell et al. (2004); Harborne et al. (2006). This is the case of Venezuela, where most of the papers published to date have only been focused on site descriptions based on species composition and abundance, whereas the influence of spatial variation across different scales on coral assemblages remains poorly explored. The Venezuelan coast is highly heterogeneous with clear longitudinal environmental gradients (Chollett et al., 2010; Miloslavich et al., 2003) which are deeply influenced by up-welling regimes that play an important role for the distribution of marine biodiversity (Miloslavich et al., 2010). In fact, algal communities in rocky shores (Cruz-Motta, 2007) and sessile organisms associated to mangrove roots have been found to vary at different spatial scales along the Venezuelan coast (Guerra-Castro et al., 2011). Thus, it should not be surprising to find coral assemblages to be extremely variable across spatial scales in Venezuela. We expected that greater changes in community structure and  $\beta$ -diversity of coral assemblages will occur at scales of thousand of kilometers (i.e., between the eastern and western regions) and within sites (i.e., hundreds of meters). This is because of existing contrasting environmental settings driven by upwelling spots that have been described along the Venezuelan coast line (Chollett et al., 2010; Miloslavich et al.,



**Figure 1.** Map of the Venezuelan coast with the seven locations used in this study. Western region, represented in blue, include: MOR= Morrocoy National Park and OCU= Ocumare de la Costa. Central region, represented in red, include: ROQ= Archipiélago Los Roques National Park and CHI= Chichiriviche de la Costa. Eastern region, represented in green, included MOC= Mochima National Park, CUB= Cubagua and FRA= Los Frailes.

2003). The goal of this study was two-fold: (1) quantify spatial variation of coral assemblages from hundreds meters to hundreds of kilometers, and (2) to determine if there are patterns of  $\beta$ -diversity across these scales.

## METHODS

### Study area

We conducted a multi-scale sampling design comprising coastal areas as well as continental and oceanic islands (Figure 1). Specifically, seven locations were sampled along the Venezuelan territory encompassing three contrasting regions (Figure 1). The western region, included two localities: (1) Morrocoy National Park (MNP) and (2) Ocumare de la Costa. The former is a continental reef system formed by a group of keys and lagoons surrounded by fringing and patch reefs located nearby mangroves and seagrass beds (Weil, 2003); whereas the latter is a small bay protected by reef barriers with seagrass and mangroves dominating the inner and shallower habitats. Likewise, the central region, entailed two localities: (3) Los Roques National Park (LRNP) which is an oceanic archipelago with a central lagoon, characterized by extensive reef banks/patches and two large coralline barriers located south and east of the archipelago (Weil, 2003); and (4) Chichirivivhe de la Costa, a location of rocky reefs with steep slopes and scattered coral assemblages. Finally, in the eastern region three localities were included: (5) Mochima National Park (MoNP), (6) Los Frailes and (7) Cubagua. In Mochima, seagrass beds and mangroves border a rocky coastline with steep slopes and fringing reef communities. Los Frailes and Cubagua are islands lying at the continental shelf and dominated by small patch reefs with scattered coral assemblages bordering their coastlines. The whole eastern coast of Venezuela and its continental islands are subjected to seasonal upwelling due to its connection with the Cariaco trench (Weil, 2003). The selection of these locations aimed to cover the vast majority of reef habitats described for Venezuela (Weil, 2003). Permit for taking pictures at marine protected areas was given by Ministerio del Poder Popular para el Ecosocialismo y Aguas - Dirección General de Diversidad Biológica under the office n° 0033.

### Experimental design

A fully-nested design encompassing three hierarchical-random factors (i.e., site, locality and region) was used to determine spatial variation on coral assemblage structure (i.e., absolute cover of coral species)

and  $\beta$ -diversity from hundreds of meters (sites) to hundreds of kilometers (region). The factor region encompassed three levels (West, Center, and East); nested within region there were two/three localities, four to seven reef sites within each locality and four 30m-long transects within each site, understood as the operational unit.

### Benthic surveys

At each reef site, benthic surveys were conducted during 2017 and 2018, following the guidelines outlined by the Global Coral Reef Monitoring Network-Caribbean (GCRMN) (GCRMN, 2014) with slight modifications. In order to increase the number of sampled sites, we surveyed four instead of five 30m-long transects parallel to the shoreline following the bottom contour between 8-10 m depth. Transects were set randomly, with the first transect being always layout at the first spot of diving. From that point, each transect was moved up or down from the first transect. Distance among transects varied from 5 to 6 m, so each operational unit was inter-spaced across the sampled reef habitat. For each transect, 15 80x90-cm photos were taken every other meter to determine the benthic community structure (N = 60 photos per site). A reference frame was used in the field to calibrate each photograph in the laboratory for further analysis of benthic cover.

### Analysis of photo-quadrats

The photo quadrat analysis was performed using PhotoQuad (Trygonis and Sini, 2012). For this, every coral was identified to species level and the percentage cover was estimated from 25 points randomly set in an area of approximately 7,200  $cm^2$ . From the analysis of photo quadrats, we obtained two matrices: (1) absolute cover of coral species and (2) coral species presence/absence. Data cleaning and quality control were performed using R (R Core Team, 2019). Thus, coral cover estimates were done from a randomly-selected sample composed of 375 points per transect (15 photos x 25 points = 375).

### Data analysis

A  $\text{Log}(x + 1)$  transformation was applied to the data and the Bray-Curtis index was used to build the similarity matrices. Hypothesis test were performed with a permutation-based analysis of variance (PERMANOVA). A test of homogeneity of dispersions (PermDisp) was performed before running the PERMANOVA to verify dispersion of data across different spatial scales. For the majority of the spatial scales the PermDisp confirmed homogeneity in dispersion except for sites (Supp. 2). However the PERMANOVA test has been shown to be robust to identify dispersion versus location effects for balanced designs like the one presented here (Anderson and Walsh, 2013).

With the PERMANOVA, we tested the spatial variability of coral assemblage structure (i.e., absolute abundance of each species) from meters to hundreds of kilometers. The variance components were estimated depending on each source of variation in the analysis following the procedures outlined by Clarke and Gorley (2006); Anderson et al. (2008). Spatial patterns (if any) were visualized by using an Analysis of Principal Coordinates (PCO) aimed to represent the position of centroids of each site in a Bray-Curtis space. Also, a Similarity Percentage Analysis (SIMPER) was carried out at each spatial scale to identify and estimate the species that contribute the most to the patterns observed. SIMPER results were represented as heat maps to show the relative contribution of each coral specie to the average Bray-Curtis similarity. Additionally, a Canonical Analysis of Principal Coordinates (CAP) based on Bray-Curtis similarity index was carried out. We used the scale that most contributed to the variation in the dissimilarity of coral species to determine the correlation between the coverage of scleractinian corals and their latitudinal and longitudinal position. For coral cover we used the same variables as in the PERMANOVA, and the latitude and longitude were understood as proxies of distance respect to the coast and position along the coast, respectively. All analyses were carried out using Primer 6 - Permanova + (Clarke and Gorley, 2006; Anderson et al., 2008).

Differences on  $\beta$ -diversity were tested across different spatial scales using a test of homogeneity of dispersions (PermDisp) (Anderson, 2006) based on presence/absence Jaccard index. To maintain the nested design we used transects to test dispersion across sites, the composition of sites to test dispersion across localities, and the composition of localities to test dispersion across regions. The Jaccard index was then split into the components of nestedness and turn-over using the *BetaPart* package in R (Baselga and Orme, 2012).

Due to the life history strategies of scleractinian corals (i.e. large colonies of a single species may often occupy an entire quadrat), rare species in the assemblages might have been underestimated (Chao

et al., 2014). To verify this, we estimated sampling coverage (i.e. the relationship of interpolation and extrapolation of the effective number of species in the community) (Hsieh et al., 2016; Chao et al., 2014). This methodological approach allows statistical inferences without the bias of underestimated rare species in assemblies and/or assigning them the same weight as abundant species (Chao et al., 2014). The sampling coverage was calculated using the *iNEXT* package in the R software (Hsieh et al., 2016). We found that 80% of the sites had a sample coverage greater than 0.7 (Table 1, Supp. 3), which is considered a proper sampling effort to describe species assemblages Hsieh et al. (2016).

## RESULTS

### Patterns of coral assemblage structure

The results show that species composition and abundance of corals in Venezuela varied across different spatial scales. The greatest variability was found at the scale of sites (Pseudo-F = 5.34, p-value = 0.001) which accounted for 35.1% of the total variance (table 2). The scale of hundreds of kilometers was the second most important source of variation in the analysis (Pseudo-F = 3.35, p-value = 0.01, CV = 21.35% table 2). This result indicates that coral assemblages in Venezuela only vary by 21.35% at the scale of region. Also, we found statistical significance at the scale of locations within regions, explaining 11.42% of the total variance (table 2). Thus, our results indicate that coral assemblages are much variable at small to medium scales (i.e., from hundreds of meters to tens of kilometers) rather than hundreds of kilometers (i.e., regions) alone.

With over 70% of the total variance explained by the first three PCO axes, localities within regions ordinate according to changes in cover of coral species (Figure 2). Our results show that the eastern and central regions are much more similar to each other concerning the western region. The SIMPER analysis showed clear patterns defining regions and locations across Venezuela, however, sites showed variable species composition and cover (Figure 3). Overall, Western region was largely composed of *Orbicella faveolata* reefs, whereas the eastern reefs were dominated by *Pseudodiploria strigosa* (Figure 3). On the other hand, across the Central regions which included Oceanic and Coastal Reefs, mixed coral communities were found. These species accounted for more than 75% of dissimilarities across localities, sites and regions (Supp. 4-6).

Longitude (correlation = 0.867, p = 0.001) was highly correlated with observed spatial patterns in contradiction to latitude (correlation = 0.552, p = 0.021). This result indicates that the relative position of each site along the Venezuelan coast (i.e., longitudinal variation), is an important factor to determine the features of coral assemblages in Venezuela, instead of the proximity to the coast (i.e., latitudinal variation).

### Patterns of beta diversity

When assessing  $\beta$ -diversity, we found the highest variation in species presence/absence occurring between transect of the same site (F = 2.764, p = 0.045) and between sites of the same locality (F = 4.438, p = 0.026). On the other hand, at larger scales, we found no significant dispersion in species composition between localities of the same region (table 3, Supp. 7-9). In addition, site and locality were the spatial scales with the largest Jaccard dissimilarity, with turn-over component as the main contributor. Furthermore, at larger scales, Jaccard dissimilarity decreased and the contribution of turn-over and nestedness became even (Figure 4). Thus, our result clearly shows that in Venezuela it is more likely to find changes in coral species composition at small to medium scales (i.e., hundreds of meters to tens of kilometers) than at larger scales (i.e., hundreds of kilometers). Finally, the results indicate that coral species found between the western, central and eastern region of Venezuela can result either from species replacement or from species loss, which is interpreted as a subset of a total pool of species.

## DISCUSSION


While coral assemblages have been extensively studied in Venezuela, this is the first multi-scale assessment to show the importance of spatial scales in determining the structure of these communities. Overall, we found that coral assemblages in Venezuela are variable from hundreds of meters and hundreds of kilometers. Additionally, the largest changes in the composition of coral species occurred at a small scale with a clear predominance of species turn-overs. Also, longitude and latitude are a good predictors of

**Table 1.** Sampling coverage for each reef site.

Locality	Site	Replicates	Incidence	Richness	Sampling coverage
Morrocoy	Boca Seca	4	12	6	0.900
	Bajo Caimán	4	9	4	0.952
	Medio	4	13	7	0.736
	Mero	4	16	10	0.796
	Norte	4	20	7	0.979
	Sombrero	4	24	11	0.836
	Sur	4	18	8	0.885
Ocumare	Ciénaga este	4	15	8	0.706
	Ciénaga interna	4	24	11	0.779
	Ciénaga oeste	4	25	10	0.917
	Guabinitas	4	16	7	0.925
Los Roques	Cayo Agua	4	25	11	0.893
	Boca de Cote	4	33	12	0.937
	Dos Mosquises sur	4	21	10	0.790
	Madriski	4	17	7	0.912
	Pelona de Rabusqui	4	14	6	0.839
	Salinas	4	23	10	0.851
	La Venada	4	7	4	0.679
Chichiriviche de la costa	La Pared	4	21	9	0.901
	Media Legua	4	15	9	0.640
	Punta de Media Legua	4	17	9	0.741
	Petaquire	3	11	7	0.636
	Playa Tiburón	4	20	7	0.979
	Punta Mono	4	19	8	0.828
Mochima	San Agustín	4	1	1	1.000
	Blanca	4	23	9	0.893
	Carabela	4	2	2	0.400
	Punta Cruz	4	14	6	0.893
	Gabarra	4	11	5	0.864
	Garrapata	4	20	7	0.940
Cubagua	Charagato	4	6	4	0.625
	Punta Conejo	4	6	4	0.625
	La Muerta	4	6	4	0.625
Los Frailes	Cominoto	4	6	2	1.000
	La Pecha	4	1	1	1.000
	Puerto Real	4	8	2	1.000

**Table 2.** Three-way permutation-based analysis of variance (PERMANOVA) based on Bray-Curtis Similarity to test differences in coral assemblage structure.


Source	df	SS	MS	Pseudo-F	P(perm)	Unique perms	%CV
Region	2	7.11x10 <sup>4</sup>	35529	3.35	0.01	998	21.35
Locality	4	4.09x10 <sup>4</sup>	10219	2.28	0.002	998	11.42
Site	29	1.30x10 <sup>5</sup>	4483	5.34	0.001	994	35.10
Residuals	107	8.99x10 <sup>4</sup>	840.17				32.14
Total	142	3.36x10 <sup>5</sup>					




**Figure 2.** Principal coordinates ordination plot (PCO) showing the variation of coral assemblages at the site level nested into locality. The distance between samples are interpreted as % of similarity. Panels A, B, and C represent the different combined pairs of the first three principal coordinates axes.

**Table 3.** Test of homogeneity of dispersions (PermDisp) to compare the distances from each factor to centroids as a test for similarity in  $\beta$  diversity among factors.

	Df	F	N.Perm	P(perm)
Site	35	2.764	999	0.045
	107			
Locality	6	4.438	999	0.026
	29			
Region	2	8.216	999	0.283
	4			



**Figure 3.** Heat map showing the species contribution to similarity (Cut off for low contributions = 75%). Panels A, B, and C represent the three scales considered in the study.



**Figure 4.** Total dissimilarity as Jaccard index and their respective turn-over and nestedness components for each spatial scale.

231 coral assemblage structure (i.e., species composition and abundance) further indicating that large-scale  
232 processes are also important to determine the structure of these communities.

### 233 **Patterns of coral assemblage structure**

234 Previous studies have acknowledged the importance of spatial scales on coral assemblages in the Caribbean  
235 and in Venezuela (Weil, 2003). Particularly, the effect of upwelling and other related oceanographic  
236 processes has been pinpointed as strong factors that shape coral assemblage structure along the Venezuelan  
237 coast where at least 12 upwelling points have been targeted (Miloslavich et al., 2003; Chollett et al., 2012).  
238 Our study shows that coral assemblages in Venezuela are much variable within and between localities than  
239 we originally expected. We found two fold higher variability at small to medium scales when compared to  
240 regions. However, coral assemblages between the western, central and eastern regions differed by 21.35%,  
241 further indicating that differences at scale of region cannot be neglected. Moreover, about 32% of the  
242 total variance in coral species composition and abundance was associated to the residual which indicates  
243 that other variables like levels of anthropogenic disturbance, oceanic influence or other intermediate  
244 scales between those taken into consideration might also be relevant to determine the structure of these  
245 assemblages.

246 In Venezuela, encrusting communities associated with mangrove roots have been studied following  
247 a spatial hierarchical design (Guerra-castro et al., 2016). Similar to our results, higher variation for  
248 this assemblages were found at the smallest and biggest scales (Guerra-castro et al., 2016). Moreover,  
249 algal assemblages associated rocky platforms have previously been found to be highly variable at tens of  
250 kilometers and not between localities or regions, further illustrating the importance of local processes in  
251 providing structure to different assemblages of sessile organisms in the country (Cruz-Motta, 2007).

252 Only a few numbers of studies encompassing multiple hierarchical spatial scales have been conducted  
253 in the Caribbean. For example, in a multi-scale study Williams et al. (2015) surveyed a series of reef sites  
254 across locations and different bio-regions in the Caribbean for three major coral taxonomic groups: corals,  
255 sponges and octocorals. They concluded these faunas exhibited considerable biogeographical variability

at broad spatial scales (hundreds of kilometers). However similar to our study, they found a higher degree of variability within sites highlighting the relevance of local ecological drivers (e.g. rugosity and wave exposure) in structuring coral assemblages. Other studies have also taken into account the importance of spatial scales for coral assemblages, but must of then have focused on total live coral cover (Murdoch and Aronson, 1999) and total abundance of colonies (Hughes et al., 1999).

It is widely acknowledged that behind patterns are ecological processes that shape communities (Barton et al., 2013). In coral reef ecosystems, processes such as predation (e.g. herbivory) and competition have profound impacts on species abundance and composition at scales of a few meters. For example, very high densities or the absence of *Diadema antillarum* can determine the composition of corals in patches of few  $m^2$  (Sammarco, 1980). Also, reef fish can preferentially prey on certain coral species, thus decreasing their abundance or making them less competitive than other neighboring corals. In addition, the presence of vermetids could potentially modify the survival rates of coral species (Lenihan et al., 2011). Furthermore, factors such as structural complexity (e.g. rugosity and micro scale habitat heterogeneity) may influence competition and survival of colonies depending on their sizes (Almany, 2004; Zilberberg and Edmunds, 2001). Coral reefs in Venezuela are known to be highly variable within and between sites but the processes responsible for these patterns have not been firmly established. However, spatial variation of coral assemblages in Los Roques has been associated to changes in reef slopes (Cróquer and Villamizar, 1998).

Various anthropogenic stressors can alter biological interactions thereby affecting the structure of coral assemblages. For example, over fishing often leads to the dominance of macroalgae which escape to herbivory control (Hughes, 1994). The selective extraction of species of carnivorous fish can lead to an increment in the abundance of echinoderms, which may also steer to increasing erosion, loss of topographic complexity and live coral cover (Roberts, 1995; McClanahan and Muthiga, 1988). Also, high intensity of recreational activities represents an important disturbance to marine communities (Milazzo et al., 2002), for example, coral cover and the proportion of massive corals is being found to be lower in places with high recreational diving intensity (Tratalos and Austin, 2001). Thus, spatial variability recorded within sites of Morrocoy and Los Roques National Parks could be explained by their differences in touristic use for not all sites within these MPAs are exposed to the same human pressures.

In addition, experimental evidence shows that some coral species differ in resistance to environmental stressors such as sedimentation (McClanahan and Obura, 1997; Rogers, 1983) which could explain the patterns observed within sites and between locations showing higher sedimentary regimes and river presence (e.g. coastal versus oceanic reef sites) (Dikou and van Woesik, 2006). Furthermore, local oceanographic events can generate mortality which leads to changes in the structure of coral assemblages. Differences between Playa Caimán, Cayo Norte and Sombrero (Morrocoy National Park) represents a example of how abnormal oceanographic conditions can alter benthic communities by killing dominant species in specific sites while promoting stable alternative states which hampers recovery (Bastidas et al., 2006). Our results seem to support that each site/location in Venezuela possess different communities because they may have been affected/unaffected by different stressors and/or mortality events in different times. Thus, high spatial variability on coral assemblages in Venezuela could be related to the differences in the disturbance regime and local history as noticed in other studies (Pisapia et al., 2019).

Coral bleaching mortality events may be patchy (West and Salm, 2003) and could potentially affect coral assemblages at different spatial scales (e.g. within sites, localities and regions). For example, in 2010 an increase in seawater temperature in Los Roques National Park affected 72% of the colonies at the study sites, showing bleaching and prevalence of diseases such as black band and white plague. Extensive mortality caused changes in the community structure a year later (Bastidas et al., 2012). Other bleaching events recorded in Venezuela since 1998 primarily affected reefs in several oceanic islands and the western and central coast of Venezuela, but these events did not produce extensive mortality events like the one reported in 2010 (Rodríguez et al., 2010; Bastidas et al., 2012; Mónaco et al., 2012).

It is likely that the eastern coast of Venezuela remained less affected by bleaching events because of seasonal upwelling and unique environmental conditions. In fact, Nakamura and van Woesik (2001) found differences in coral mortality rates during bleaching events, according to local environmental settings (e.g. light intensity, penetration, temperature and currents). However, Chollett et al. (2010) argued that upwelling does not necessarily guarantee a refuge for corals. Thus, in Venezuela differences between geographical regions could be strongly influenced by factors such as nutrient input and temperature decrease associated with the upwelling season (Birkeland, 1988; Weil, 2003). In fact, it has been

suggested that during these periods the assemblages of macroalgae become more dominant (Diaz-Pulido and Garzón-Ferreira, 2002) which could modify the coral assemblage structure through competitive processes.

The Venezuelan coast is characterized by an upwelling period that occurs between January and June, in particular, the eastern region of the country is characterized by a large area of upwelling (Castellanos et al., 2002). However, studies such as those of Jiménez and Cortés (2003) and Rodríguez et al. (2009) did not find an effect of the upwelling on factors such as the reproductive behavior of spawn corals and the growth rates of colonies. On the other hand, the dynamic of black band disease, one of the most important factors producing rapid coral mortality in Cubagua has been shown to be deeply influenced by upwelling events (Rodríguez and Croquer, 2008). Thus, our results indicate that upwelling alone is not sufficient to explain the extremely variable nature of coral assemblage.

## Patterns of beta diversity

We found that differences in coral species composition occurred at spatial scales of hundreds of meters to tens of kilometers. Although it is known that  $\beta$ -diversity depends on the spatial scale at which it is measured, opposite to our results, in most studies  $\beta$ -diversity is assumed to be homogeneous at small spatial scales (Hewitt et al., 2005; Whittaker, 1975). Changes in species composition at scales of tens of meters often occurs in highly-heterogeneous habitats sampled with enough resolution to detect these changes (Hewitt et al., 2005). In our study, we found more likely to have different species composition within transects of a single site than between localities belonging to different regions. Thus, our results clearly support that coral habitats in Venezuela are extremely variable at local scales, suggesting significant environmental heterogeneity within reef habitats, with coral species probably forming mosaics or patches within a single habitat. However, it is not clear what are the conditions favoring this heterogeneity within the reef sites. This variation at the smallest scale could mean that Venezuelan coral assemblages are in good condition, as well as there are patches of mortality, with more or fewer species, reflected in a high turnover rate.

At larger scales (i.e., between the eastern, central and western Venezuelan coast) we found quite similar and homogeneous coral species composition, which may be partly due to the reduced pool of species that exist in the Caribbean when compared with the Indo Pacific region. In regions with larger species pools such as the Indo Pacific,  $\beta$ -diversity tends to be higher at larger spatial scales because species represent a subset of a total species pool, e.g. Zvuloni et al. (2010). In Venezuela, coral reefs located at the oceanic sites and the western coast are dominated by *Orbicellas*, whereas in the eastern coast and the majority of sites located at the central coast *Pseudodiplorias* and *Colpophyllias* become more important.

Our results indicate that at scales of tens of kilometers species nestedness (loss) becomes as important as species turnover (replacement). These two components arise from different ecological phenomena (Baselga, 2010). Species nestedness occurs when the biotas of sites with smaller numbers of species are subsets of the biotas at richer sites, reflecting a non-random process of species loss as a consequence of any factor that promotes the orderly disaggregation of assemblages (Baselga, 2010). On the other hand, species turn-over (replacement) occurs as a consequence of environmental sorting and spatial and historical constraints (i.e., stochastic process) (Qian et al., 2004). For example, processes such as settlement selectivity of coral larvae could explain species turn overs at tens of meters and kilometers. Coral larvae are known to select certain characteristics in the habitat to settle down, e.g. presence of certain species of coralline algae (Ritson-Williams et al., 2010) or sounds of the reef (Vermeij et al., 2010). Our results therefore indicate that coral assemblage structure in Venezuela is probably regulated by a series of interconnected processes acting alone and/or in combination at various spatial scales. This result highlights the importance of creating scale-adapted management actions in Venezuela since the smallest scales reflect the greatest variability. However, very small MPAs are often ineffective in achieving their conservation goals, so they must necessarily be chained into a large-scale strategy (Agardy et al., 2011).

In summary, coral assemblage structure in Venezuela is highly variable at different spatial scales but within locality variability seem to be very important. The processes that could underlie these patterns are diverse and complex and little experimental efforts to untangle the specific contribution of each factor have been conducted. Longitude is a good predictor of coral assemblages in Venezuela. Upwelling-related processes could be targeted as potential candidates to explain longitudinal variation of coral assemblages,

whereas oceanographic/coastal processes could explain latitudinal variability. Regarding  $\beta$ -diversity, coral assemblages are fairly homogeneous across the Venezuelan coast, while increasing spatial resolution shows greater heterogeneity, with smaller scales revealing a greater change in species composition. In addition, the replacement of species is a relevant phenomenon to explain these diversity patterns. Results from this study highlights the importance of taking into account local variability during the design and implementation of specific conservation efforts.

## ACKNOWLEDGMENTS

This research was funded by the Waitt Foundation through the Rapid Ocean Conservation (ROC) Grant Program. We thank the support provided by Fundación para la Defensa de la Naturaleza (FUDENA), Instituto de Estudios Avanzados (IDEA) and the Escuela de Ciencias Aplicadas del Mar (ECAM), as well as José Cappelletto, Zlatka Rebolledo and Alejandra Verde who helped with the field work. We also thank Rita Peachey and all the 39th AMLC meeting committee for their help to attend the conference.

## REFERENCES

- Agardy, T., Di Sciara, G. N., and Christie, P. (2011). Mind the gap: addressing the shortcomings of marine protected areas through large scale marine spatial planning. *Marine Policy*, 35(2):226–232.
- Almany, G. R. (2004). Does increased habitat complexity reduce predation and competition in coral reef fish assemblages? *Oikos*, 106(2):275–284.
- Alvarez-filip, L., Dulvy, N. K., Gill, J. A., Watkinson, A. R., and Cote, I. M. (2009). Flattening of Caribbean coral reefs: region-wide declines in architectural complexity. *Proceedings of the Royal Society B: Biological Sciences*, 276(1):3019–3025.
- Anderson, M. J. (2006). Distance-based tests for homogeneity of multivariate dispersions. *Biometrics*, 62(1):245–253.
- Anderson, M. J., Crist, T. O., Chase, J. M., Vallend, M., Inouye, B. D., Freestone, A. L., Sanders, N. J., Cornell, H. V., Comita, L. S., Davies, K. F., Harrison, S. P., Kraft, N. J., Stegen, J. C., and Swenson, N. G. (2011). Navigating the multiple meanings of beta-diversity: a roadmap for the practicing ecologist. *Ecology Letters*, 14:19–28.
- Anderson, M. J., Gorley, R. N., and Clarke, K. R. (2008). *PERMANOVA+ for PRIMER: Guide to Software and Statistical Methods*. PRIMER-E, Plymouth.
- Anderson, M. J. and Walsh, D. C. (2013). Permanova, anosim, and the mantel test in the face of heterogeneous dispersions: what null hypothesis are you testing? *Ecological monographs*, 83(4):557–574.
- Arias-González, J. E., Legendre, P., and Rodríguez-zaragoza, F. A. (2008). Journal of Experimental Marine Biology and Ecology Scaling up beta diversity on Caribbean coral reefs. *Journal of Experimental Marine Biology and Ecology*, 366(1-2):28–36.
- Barton, P. S., Cunningham, S. A., Manning, A. D., Gibb, H., Lindenmayer, D. B., and Didham, R. K. (2013). The spatial scaling of beta diversity. *Global Ecology and Biogeography*, 22(6):639–647.
- Baselga, A. (2010). Partitioning the turnover and nestedness components of beta diversity. *Global Ecology and Biogeography*, 19(1):134–143.
- Baselga, A. and Orme, C. D. L. (2012). Betapart: An R package for the study of beta diversity. *Methods in Ecology and Evolution*, 3(5):808–812.
- Bastidas, C., Bone, D., Croquer, A., Debrot, D., Garcia, E., Humanes, A., Ramos, R., and Rodríguez, S. (2012). Massive hard coral loss after a severe bleaching event in 2010 at los roques, venezuela. *Revista de Biología Tropical*, 60:29–37.
- Bastidas, C., Croquer, A., and Bone, D. (2006). Shift of dominant species after a mass mortality on a Caribbean reef. *Proceedings of 10th International Coral Reef Symposium, Okinawa, Japan.*, pages 989–993.
- Birkeland, C. (1988). Geographic Comparisons of Coral-Reef Community Processes. *Proceedings of the 6th International Coral Reef Symposium, Australia*, 1:211–220.
- Castellanos, P., Varela, R., and Muller-Karger, F. (2002). Descripción de las áreas de surgencia al sur del mar caribe examinadas con el sensor infrarrojo avhrr. *Memoria de la Fundación La Salle de Ciencias Naturales*, 154:55–76.
- Chao, A., Gotelli, N. J., Hsieh, T., Sander, E. L., Ma, K., Colwell, R. K., and Ellison, A. M. (2014).

- 417 Rarefaction and extrapolation with hill numbers: a framework for sampling and estimation in species  
418 diversity studies. *Ecological monographs*, 84(1):45–67.
- 419 Chave, J. (2013). The problem of pattern and scale in ecology: What have we learned in 20 years?  
420 *Ecology Letters*, 16(SUPPL.1):4–16.
- 421 Chollett, I., Mumby, P. J., and Cortés, J. (2010). Upwelling areas do not guarantee refuge for coral reefs  
422 in a warming ocean. *Marine Ecology Progress Series*, 416(1):47–56.
- 423 Chollett, I., Mumby, P. J., Müller-Karger, F. E., and Hu, C. (2012). Physical environments of the caribbean  
424 sea. *Limnology and Oceanography*, 57(4):1233–1244.
- 425 Clarke, K. R. and Gorley, R. N. (2006). *PRIMER v6: User Manual/Tutorial*. PRIMER-E, Plymouth.
- 426 Connell, J. H., Hughes, T. P., Wallace, C. C., Tanner, J. E., Harms, K. E., and Kerr, A. M. (2004). A  
427 long-term study of competition and diversity of corals. *Ecological Monographs*, 74(2):179–210.
- 428 Cróquer, A. and Villamizar, E. (1998). Las variaciones de la pendiente topográfica, un factor a considerar  
429 en la evaluación de la estructura de una comunidad arrecifal. *Revista de biología tropical*, pages 29–40.
- 430 Cruz-Motta, J. J. (2007). Análisis espacial de las comunidades tropicales intermareales asociadas a los  
431 litorales rocosos de Venezuela. *Ciencias marinas*, 33(2):133–148.
- 432 Diaz-Pulido, G. and Garzón-Ferreira, J. (2002). Seasonality in Algal Assemblages on Upwelling-  
433 influenced Coral Reefs in the Colombian Caribbean. *Botanica Marina*, 45(3):284–292.
- 434 Dikou, A. and van Woesik, R. (2006). Survival under chronic stress from sediment load: Spatial  
435 patterns of hard coral communities in the southern islands of Singapore. *Marine Pollution Bulletin*,  
436 52(11):1340–1354.
- 437 Edmunds, P. J. (2013). Decadal-scale changes in the community structure of coral reefs of St . John , US  
438 Virgin Islands. *Marine Ecology Progress Series*, 489:107–123.
- 439 Eidens, C., Hauffe, T., Bayraktarov, E., Wild, C., and Wilke, T. (2015). Multi-scale processes drive  
440 benthic community structure in upwelling-affected coral reefs. *Frontiers in Marine Science*, 2(2):1–11.
- 441 Field, R., Hawkins, B. A., Cornell, H. V., Currie, D. J., Diniz-Filho, J. A. F., Guégan, J. F., Kaufman,  
442 D. M., Kerr, J. T., Mittelbach, G. G., Oberdorff, T., O'Brien, E. M., and Turner, J. R. (2009). Spatial  
443 species-richness gradients across scales: A meta-analysis. *Journal of Biogeography*, 36(1):132–147.
- 444 Gaston, K. J., Davies, R. G., Orme, C. D. L., Olson, V. A., Thomas, G. H., Ding, T.-s., Rasmussen, P. C.,  
445 Lennon, J. J., Bennett, P. M., Owens, I. P. F., and Blackburn, T. M. (2007). Spatial turnover in the  
446 global avifauna. *Proceedings of the Royal Society B: Biological Sciences*, 274(1):1567–1574.
- 447 GCRMN (2014). Report of the Workshop of GCRMN for the Wider Caribbean: Review, improve and  
448 revitalize the network and the nodes for more effective coral reef monitoring and data management  
449 Curaçao. Technical report, UNEP-CEP / SPAW-RAC /Ministry of Economic Affairs of the Netherlands,  
450 Curaçao.
- 451 Guerra-Castro, E., Cruz-Motta, J. J., and Conde, J. E. (2011). Cuantificación de la diversidad de especies  
452 incrustantes asociadas a las raíces de *Rhizophora mangle* L. en el Parque Nacional Laguna de la  
453 Restinga. *Interciencia*, 36(12):923–930.
- 454 Guerra-castro, E. J., Conde, J. E., and Cruz-motta, J. J. (2016). Scales of spatial variation in tropical  
455 benthic assemblages and their ecological relevance: epibionts on Caribbean mangrove roots as a model  
456 system. *Marine Ecology Progress Series*, 548:97–110.
- 457 Harborne, A. R., Mumby, P. J., Zychaluk, K., Hedley, J. D., and Blackwell, P. G. (2006). Modeling the  
458 beta diversity of coral reefs. *Ecological Society of America*, 87(11):2871–2881.
- 459 Hewitt, J. E., Thrush, S. F., Halliday, J., and Duffy, C. (2005). The importace of small-scale habitat  
460 structure for maintaining Beta Diversity. *Ecology*, 86(6):1619–1626.
- 461 Hsieh, T., Ma, K., and Chao, A. (2016). inext: an r package for rarefaction and extrapolation of species  
462 diversity (h ill numbers). *Methods in Ecology and Evolution*, 7(12):1451–1456.
- 463 Hughes, T. P. (1994). Catastrophes, phase shifts, and large-scale degradation of a Caribbean coral reef.  
464 *Science*, 265(5178):1547–1551.
- 465 Hughes, T. P., Baird, A. H., Dinsdale, E. A., Moltschaniwskyj, N. A., Pratchett, M. S., Tanner, J. E., and  
466 Willis, B. L. (1999). Patterns of recruitment and abundance of corals along the Great Barrier Reef.  
467 *Nature*, 397(6714):59–63.
- 468 Jiménez, C. and Cortés, J. (2003). Growth of seven species of scleractinian corals in an upwelling  
469 environment of the eastern pacific (golfo de papagayo, costa rica). *Bulletin of Marine Science*,  
470 72(1):187–198.
- 471 Johnson, R. K. and Goedkoop, W. (2002). Littoral macroinvertebrate communities: Spatial scale and

- ecological relationships. *Freshwater Biology*, 47(1):1840–1854.
- Jurasinski, G., Retzer, V., and Beierkuhnlein, C. (2009). Inventory , differentiation , and proportional diversity : a consistent terminology for quantifying species diversity. *Oecologia*, 159(1):15–26.
- Karlson, R. H. and Cornell, H. V. (1999). Integration of Local and Regional Perspectives on the Species Richness of Coral Assemblages. *American Zoologist*, 39:104–112.
- Komyakova, V., Jones, G. P., and Munday, P. L. (2018). Strong effects of coral species on the diversity and structure of reef fish communities : A multi-scale analysis. *PLoS ONE*, 13(8):1–20.
- Leibold, M. A., Holyoak, M., Mouquet, N., Amarasekare, P., Chase, J. M., Hoopes, M. F., Holt, R. D., Shurin, J. B., Law, R., Tilman, D., Loreau, M., and Gonzalez, A. (2004). The metacommunity concept : a framework for multi-scale community ecology. *Ecology Letters*, 7:601–613.
- Lenihan, H. S., Holbrook, S. J., Schmitt, R. J., and Brooks, A. J. (2011). Influence of corallivory, competition, and habitat structure on coral community shifts. *Ecology*, 92(10):1959–1971.
- Levin, S. A. (1992). The Problem of Pattern and Scale in Ecology. *Source: Ecology Ecology*, 73(736):1943–1967.
- Mac Nally, R. and Quinn, G. (1998). Symposium introduction: The importance of scale in ecology. *Australian Journal of Ecology*, 23(1):1–7.
- MacArthur, R. (1972). *Geographical Ecology*. Princeton University Press, Princeton, USA.
- McClanahan, T. and Obura, D. (1997). Sedimentation effects on shallow coral communities in Kenya. *Journal of Experimental Marine Biology and Ecology*, 209(1-2):103–122.
- Mcclanahan, T. R. and Muthiga, N. A. (1988). Changes in Kenyan coral reef community structure and function due to exploitation. *Hydrobiologia*, 166:269–276.
- McGill, B. J. (2010). Matters of scale. *Science*, 328(1):575–576.
- Melchior, L. G., Rossa-Feres, D. d. C., and da Silva, F. R. (2017). Evaluating multiple spatial scales to understand the distribution of anuran beta diversity in the Brazilian Atlantic Forest. *Ecology and Evolution*, 7(1):2403–2413.
- Milazzo, M., Chemello, R., Badalamenti, F., Camarda, R., and Riggio, S. (2002). The Impact of Human Recreational Activities in Marine Protected Areas : What Lessons Should Be Learnt in the Mediterranean Sea? *Marine Ecology*, 23:280–290.
- Miloslavich, P., Díaz, J. M., Klein, E., Alvarado, J. J., Díaz, C., Gobin, J., Escobar-Briones, E., Cruz-motta, J. J., Weil, E., Cortés, J., Bastidas, C., Robertson, R., Zapata, F., Martín, A., Castillo, J., Kazandjian, A., and Ortiz, M. (2010). Marine Biodiversity in the Caribbean : Regional Estimates and Distribution Patterns. *PLoS ONE*, 5(8):e11916.
- Miloslavich, P., Klein, E., Yarena, E., and Martin, A. (2003). Marine biodiversity in Venezuela: Status and perspectives. *Gayana (Concepción)*, 67(2):275–301.
- Mónaco, C. d., Haiek, G., Narciso, S., and Galindo, M. (2012). Massive bleaching of coral reefs induced by the 2010 enso, puerto cabello, venezuela. *Revista de biologia tropical*, 60(2):527–538.
- Murdoch, T. J. and Aronson, R. B. (1999). Scale-dependent spatial variability of coral assemblages along the Florida Reef Tract. *Coral Reefs*, 18(4):341–351.
- Nakamura, T. and van Woesik, R. (2001). Water-flow rates and passive diffusion partially explain differential survival of corals during the 1998 bleaching event. *Marine Ecology Progress Series*, 212(2):301–304.
- Pandolfi, J. M. (2002). Coral community dynamics at multiple scales. *Coral Reefs*, 21(1):13–23.
- Pisapia, C., Burn, D., and Pratchett, M. S. (2019). Changes in the population and community structure of corals during recent disturbances (February 2016-October 2017) on Maldivian coral reefs. *Scientific Reports*, 9(1):8402.
- Plaisance, L., Caley, M. J., Brainard, R. E., and Knowlton, N. (2011). The Diversity of Coral Reefs: What Are We Missing? *PLoS ONE*, 6(10):e25026.
- Qian, H., Ricklefs, R. E., and White, P. S. (2004). Beta diversity of angiosperms in temperate floras of eastern Asia and eastern North America. *Ecology Letters*, 8(1):15–22.
- R Core Team (2019). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria.
- Reaka-Kudla, M. L. (1997). The global biodiversity of coral reefs: a comparison with rain forests. In *Biodiversity II: Understanding and protecting our biological resources*, page 551.
- Ricklefs, R. E. (1987). Community diversity: relative roles of local and regional processes. *Science*, 235(4785):167–171.

- 527 Ritson-Williams, R., Paul, V. J., Arnold, S. N., and Steneck, R. S. (2010). Larval settlement preferences  
528 and post-settlement survival of the threatened Caribbean corals *Acropora palmata* and *A. cervicornis*.  
529 *Coral Reefs*, 29(1):71–81.
- 530 Roberts, C. M. (1995). Effects of fishing on the ecosystem structure of coral reefs. *Conservation Biology*,  
531 9(5):988–995.
- 532 Rodríguez, S., Alvizu, A., Tagliafico, A., and Bastidas, C. (2009). Low natural repopulation of marginal  
533 coral communities under the influence of upwelling. *Hydrobiologia*, 624(1):1–11.
- 534 Rodríguez, S. and Croquer, A. (2008). Dynamics of black band disease in a *diploria strigosa* population  
535 subjected to annual upwelling on the northeastern coast of venezuela. *Coral Reefs*, 27(2):381–388.
- 536 Rodríguez, S., Cróquer, A., Bone, D., and Bastidas, C. (2010). Severity of the 1998 and 2005 bleaching  
537 events in venezuela, southern caribbean. *Revista de biología tropical*, 58:189–196.
- 538 Rogers, C. S. (1983). Sublethal and lethal effects of sediments applied to common Caribbean Reef corals  
539 in the field. *Marine Pollution Bulletin*, 14(10):378–382.
- 540 Sammarco, P. W. (1980). *Diadema* and its relationship to coral spat mortality: Grazing, competition, and  
541 biological disturbance. *Journal of Experimental Marine Biology and Ecology*, 45(2):245–272.
- 542 Schneider, D. C. (2001). The Rise of the Concept of Scale in Ecology. *BioScience*, 51(7):545–553.
- 543 Svenning, J.-c., Fløjgaard, C., and Baselga, A. (2011). Climate, history and neutrality as drivers of  
544 mammal beta diversity in Europe: insights from multiscale deconstruction. *Journal of Animal Ecology*,  
545 80:393–402.
- 546 Tratalos, J. A. and Austin, T. J. (2001). Impacts of recreational SCUBA diving on coral communities of  
547 the Caribbean island of Grand Cayman. *Biological Conservation*, 102(1):67–75.
- 548 Trygonis, V. and Sini, M. (2012). PhotoQuad: A dedicated seabed image processing software, and a  
549 comparative error analysis of four photoquadrat methods. *Journal of Experimental Marine Biology and*  
550 *Ecology*, 424-425:99–108.
- 551 Tuomisto, H. (2010a). A diversity of beta diversities: Straightening up a concept gone awry. Part 1.  
552 Defining beta diversity as a function of alpha and gamma diversity. *Ecography*, 33(1):2–22.
- 553 Tuomisto, H. (2010b). A diversity of beta diversities: Straightening up a concept gone awry. Part 2.  
554 Quantifying beta diversity and related phenomena. *Ecography*, 33(1):23–45.
- 555 Vellend, M. (2010). Conceptual synthesis in community ecology. *The Quarterly Review of Biology*,  
556 85(2):183–206.
- 557 Vermeij, M. J. A., Marhaver, K. L., Huijbers, C. M., Nagelkerken, I., and Simpson, S. D. (2010). Coral  
558 Larvae Move toward Reef Sounds. *PLoS ONE*, 5(5):e10660.
- 559 Weil, E. (2003). The corals and coral reefs of Venezuela. In *Latin American Coral Reefs*, pages 303–330.  
560 Elsevier.
- 561 West, J. M. and Salm, R. V. (2003). Resistance and Resilience to Coral Bleaching: Implications for Coral  
562 Reef Conservation and Management. *Conservation Biology*, 17(4):956–967.
- 563 Whittaker, R. H. (1960). Vegetation of the Siskiyou Mountains, Oregon and California. *Ecological*  
564 *Monographs*, 30:279–338.
- 565 Whittaker, R. H. (1972). Evolution and measurement of species diversity. *Taxon*, 21:213–251.
- 566 Whittaker, R. H. (1975). *Communities and Ecosystems*. MacMillan, New York, 2nd edition.
- 567 Whittaker, R. J. and Heegaard, E. (2003). What is the observed relationship between species richness and  
568 productivity? *Ecology*, 84(12):3384–3390.
- 569 Whittaker, R. J., Willis, K. J., and Field, R. (2001). Scale and species richness: towards a general,  
570 hierarchical theory of species diversity. *Journal of Biogeography*, 28:453–470.
- 571 Wiens, J. A. (1989). Spatial scaling in ecology. *Functional Ecology*, 3(4):385–397.
- 572 Williams, S. M., Chollett, I., Roff, G., Cortés, J., Dryden, C. S., and Mumby, P. J. (2015). Hierarchical  
573 spatial patterns in caribbean reef benthic assemblages. *Journal of Biogeography*, 42(7):1327–1335.
- 574 Zilberberg, C. and Edmunds, P. (2001). Competition among small colonies of *Agaricia*: the importance of  
575 size asymmetry in determining competitive outcome. *Marine Ecology Progress Series*, 221:125–133.
- 576 Zvuloni, A., van Woesik, R., and Loya, Y. (2010). Diversity Partitioning of Stony Corals Across Multiple  
577 Spatial Scales Around Zanzibar Island, Tanzania. *PLoS ONE*, 5(3):e9941.