

Fear and stressing in predator-prey ecology: considering the twin stressors of predators and people on mammals

Loren L Fardell^{Corresp., 1}, Chris R Pavey², Chris R Dickman¹

¹ School of Life and Environmental Sciences, University of Sydney, Sydney, NSW, Australia

² Land and Water, CSIRO, Winnellie, Northern Territory, Australia

Corresponding Author: Loren L Fardell
Email address: lfard3796@uni.sydney.edu.au

Predators induce stress in prey and can have beneficial effects in ecosystems, but can also have negative effects on biodiversity if they are overabundant or have been introduced. The growth of human populations is, at the same time, causing degradation of natural habitats and increasing interaction rates of humans with wildlife, such that conservation management routinely considers the effects of human disturbance as tantamount to or surpassing those of predators. The need to simultaneously manage both of these threats is particularly acute in urban areas that are, increasingly, being recognized as global hotspots of wildlife activity. Pressures from altered predator-prey interactions and human activity may each initiate fear responses in prey species above those that are triggered by natural stressors in ecosystems. If fear responses are experienced by prey at elevated levels, on top of responses to multiple environmental stressors, chronic stress impacts may occur. Despite common knowledge of the negative effects of stress, however, it is rare that stress management is considered in conservation, except in intensive *ex situ* situations such as in captive breeding facilities or zoos. We propose that mitigation of stress impacts on wildlife is crucial for preserving biodiversity, especially as the value of habitats within urban areas increases. As such, we highlight the need for future studies to consider fear and stress in predator-prey ecology to preserve both biodiversity and ecosystem functioning, especially in areas where human disturbance occurs. We suggest, in particular, that non-invasive *in situ* investigations of endocrinology and ethology be partnered in conservation planning with surveys of habitat resources to incorporate and reduce the effects of fear and stress on wildlife.

Fear and stressing in predator-prey ecology: considering the twin stressors of predators and people on mammals

Loren L. Fardell¹, Chris R. Pavey², Chris R. Dickman¹

¹School of Life and Environmental Sciences, University of Sydney, NSW 2006, Australia.

²CSIRO, Land and Water, PMB 44 Winnellie, 0822, Northern Territory, Australia.

Corresponding author:

Loren L. Fardell¹

University of Sydney, Heydon-Laurence Building (A08), Room 325, Camperdown, NSW 2006, Australia.

Email address:loren.fardell@gmail.com

23 Abstract

24 Predators induce stress in prey and can have beneficial effects in ecosystems, but can also have
 25 negative effects on biodiversity if they are overabundant or have been introduced. The growth of
 26 human populations is, at the same time, causing degradation of natural habitats and increasing
 27 interaction rates of humans with wildlife, such that conservation management routinely considers
 28 the effects of human disturbance as tantamount to or surpassing those of predators. The need to
 29 simultaneously manage both of these threats is particularly acute in urban areas that are,
 30 increasingly, being recognized as global hotspots of wildlife activity. Pressures from altered
 31 predator-prey interactions and human activity may each initiate fear responses in prey species
 32 above those that are triggered by natural stressors in ecosystems. If fear responses are
 33 experienced by prey at elevated levels, on top of responses to multiple environmental stressors,
 34 chronic stress impacts may occur. Despite common knowledge of the negative effects of stress,
 35 however, it is rare that stress management is considered in conservation, except in intensive *ex*
 36 *situ* situations such as in captive breeding facilities or zoos. We propose that mitigation of stress
 37 impacts on wildlife is crucial for preserving biodiversity, especially as the value of habitats
 38 within urban areas increases. As such, we highlight the need for future studies to consider fear
 39 and stress in predator-prey ecology to preserve both biodiversity and ecosystem functioning,
 40 especially in areas where human disturbance occurs. We suggest, in particular, that non-invasive
 41 *in situ* investigations of endocrinology and ethology be partnered in conservation planning with
 42 surveys of habitat resources to incorporate and reduce the effects of fear and stress on wildlife.

43

44 Introduction

Predators in many systems positively influence the local distribution and abundance of their prey (e.g. Polis *et al.*, 1998; Ayal, 2007; Estes *et al.*, 2011; Weissburg, Smee & Ferner, 2014), and indirectly but positively influence the functioning of whole ecosystems via trophic cascades (Prugh *et al.*, 2009; Ritchie & Johnson, 2009; Estes *et al.*, 2011; Ripple *et al.*, 2014). Predators often elicit fear responses in prey that affect prey behaviour, energy budgets and the way they interact with the environment (Brown & Kotler, 2004; Clinchy *et al.*, 2004; Romero, 2004). For example, the fear response of an individual of a prey species to a predator may be to change the locations or times in which they forage, which can have positive impacts on the structure of the environment (Laundré, Hernández & Ripple, 2010). Such fear responses can in turn influence ecosystem function (Schmitz, Krivan & Ovadia, 2004, Schmitz *et al.* 1997, Hawlena *et al.*, 2012). These effects arise due to the ‘*landscape of fear*’ that prey individuals perceive – that is, the spatial distribution of perceived predation risk that influences prey movement and behaviour as prey individuals attempt to mitigate risk and obtain essential resources (Laundré, Hernández & Altendorf, 2001; Laundré, Hernández & Ripple, 2010).

Recent studies suggest that the effects of predators are being altered, and often amplified, by human activities. For example, the demise of top predators in many parts of the world has ‘released’ populations of smaller predators, or mesopredators that have since become overabundant and now exert strong pressure on smaller species (Prugh *et al.*, 2009). Predators that have been introduced to new areas also exert much stronger pressure on prey populations than do native predators (Salo *et al.*, 2007). Climate change and associated shifts in primary productivity can also have bottom-up impacts on predator-prey interactions (Laws, 2017). Such novel alterations to predator-prey dynamics can depress prey populations and disrupt community

structure, and as such require novel approaches to recognize and manage their effects (Carthey & Blumstein, 2018; Fleming & Bateman, 2018; Guiden *et al.*, 2019).

Altered predator-prey interactions may contribute to declines in biodiversity, but a primary cause of biodiversity loss is the rapid global increase in human populations, driving agricultural, industrial, and urban expansions that change or destroy natural habitats (Madsen *et al.*, 2010). Such expansions are resulting in novel human – wildlife interactions and secondary impacts that arise from human activities in close proximity to natural environments, and thus increasingly are becoming an additional stressor that influences ecosystem function. Pressures exerted both directly and indirectly by human activities have been likened to the pressures exerted by the presence of a top predator on prey (e.g. Frid & Dill, 2002; Rehnus, Wehrle & Palme, 2014; Patten & Burger, 2018). Natural predation pressures coupled with human-imposed predation-like pressures and / or additional exogenous stressors, such as pollution, arising from anthropogenic activities are likely to negatively affect prey species by increasing their levels or frequency of stress. Current research is developing a more nuanced understanding of the effects of human activity and predator presence on the stress of prey (e.g. Arlettaz *et al.*, 2015; Jaatinen, Seltmann & Öst, 2014, respectively). However, few studies in terrestrial ecosystems have considered both of these pressures simultaneously or have linked behavioural responses with endocrinological evidence of stress. In one instructive recent study, human activity was shown to be positively correlated with physiological stress in ungulates, whilst in areas occupied by large predators stress was found to be lower and less variable (Zbyryt *et al.*, 2017).

Conservation management and associated scientific research is often viewed from the perspective of a single discipline. However, ecological changes arising from anthropogenic disturbances, including the alteration of predator-prey encounter rates, creation of novel species

interactions (including between people and wildlife), introduction of novel species, and the alteration of natural habitats, call for multidisciplinary solutions. This is particularly pertinent as urban edge habitats are becoming increasingly valuable global hotspots of wildlife activity (Ives *et al.*, 2016), and human influences are spreading further into natural habitats (Otto, 2018). There is growing evidence that multiple introduced stressors (e.g. altered predator-prey interactions and disturbance from humans) can have compounding impacts on wildlife, especially when interacting with natural stressors in ecosystems (Geary *et al.*, 2019). For these reasons, it is timely to review relevant information from the broad knowledge bases of conservation physiology, ethology, and ecology to communicate the urgent need for wildlife managers and urban planners to routinely consider how fear and stress effects from multiple sources, particularly novel predator interactions and human activities, can affect prey species and ecosystem functioning in increasingly altered systems. Our focus in this review is on mammals because of the wealth of information on this group, but we note relevant studies from other groups where appropriate. We begin our review by describing physiological responses of mammals to fear and stress, then we consider the behavioural responses of mammals to these factors, and argue that human activity should be considered as part of the ecosystem so that overall stress impacts can be managed accordingly. We then demonstrate how fear and stress can influence habitat use, how vegetation and microhabitat management potentially may be used to alleviate stress, and how to monitor for fear and stress to then create management change. Finally, we use Australia as an example to show the benefits of considering cumulative fear and stress impacts to mitigate the effects of introduced eutherian predators, increased human activity, and the possible interactive effects of the two.

113 **Review method**

114 This is not a systematic review, but rather a novel synthesis of existing knowledge. It seeks to
 115 extend ideas on the physiological impacts of fear and stress, behavioural ecology, predator-prey
 116 dynamics, and conservation management to explore the influence of humans and altered
 117 predator-prey interactions in the urban environment. To ensure that all key topic areas were
 118 thoroughly covered we conducted literature searches via: Google Scholar[®], Web of Science[®],
 119 JSTOR[®], and Wiley Interscience Online Library[®]. Manuscripts were mined for information
 120 relevant to wildlife fear and stress in terms of ecological management under anthropogenic
 121 pressures, including introduced predators and altered predator-prey interactions. The literature
 122 searches were undertaken using the following key terms as part of their title, keywords, and / or
 123 within the abstract: “acute stress”, “chronic stress”, “homeostasis”, “cumulative stress”,
 124 “multiple stressors”, “multiple threats”, “allostatic load”, “allostatic overload”, “acclimitisation”,
 125 “glucocorticoid response”, “hypothalamic pituitary adrenal axis”, “fear arousal”, “fear
 126 evolution”, “fear predation”, “amygdala” “fear”, “Pavlovian fear conditioning”, “glucocorticoid
 127 assay”, “f(a)ecal glucocorticoid”, “non-invasive glucocorticoid assay”, “reactive scope model”,
 128 “landscape(s) of fear”, “risk allocation hypothesis”, “olfaction” + “fear” + “mammal”, “post-
 129 traumatic stress disorder”, “predator” + “odo(u)r” + “fear”, “predator cue(s)”, “predation stress
 130 hypothesis”, “predator-sensitive food hypothesis”, “human activity” + “stress” + “wildlife”,
 131 “human disturbance”, “interactive stress”, “multiple stress(ors)”, “additive stress impacts”,
 132 “synergistic stress impacts”, “antagonistic stress impacts”, “human” + “wildlife” + “resource
 133 subsidies”, “predator trophic cascade”, “wildlife urban adaptation”, “urban ecology” +
 134 “wildlife”, “Australian mammal extinction”, “critical weight range mammal”, “diet” + “change”
 135 + “Australian” + “predator”, “red fox” + “Australia”, “domestic cat” + “Australia”, “human

activity” + “wildlife” + “Australia”, “Australia” + “biodiversity” + “conservation policy” + “urban hotspot”, “Australian environment protection and biodiversity conservation act” + “cumulative stress”, “Australia” + “conservation” + “glucocorticoids”, “introduced predator control”, “habitat structural complexity”, “habitat structural diversity”, “vegetation diversity”, “vegetation heterogeneity”, “habitat heterogeneity hypothesis”, “cumulative effects assessments”. Supplementary articles supporting the review narrative, suggested by reviewers on earlier versions of the manuscript, have also been included.

Mammalian physiological responses to fear and stress

Physiological stress can be broadly defined as a change in the physiological well-being of an individual following exposure to an aversive extrinsic stimulus - frequently referred to as a stressor (Seyle, 1936). The physiological stress response that follows aims to restore internal homeostasis (Cannon, 1932). An individual animal experiences stress in response to conditions that threaten its survival or compromise its ability to maintain homeostasis. Examples include acute or chronic encounters with predators, inclement weather, significant natural disturbances including fire and flood, reduced oxygen availability, and depleted food resources (Lima, 1998; King & Bradshaw, 2010; Malcolm *et al.*, 2014; Santos *et al.*, 2014; Crocker, Khudyakov & Champagne, 2016).

In mammals, activation of the hypothalamic-pituitary-adrenal (HPA) axis is a common stress response, but this may vary depending on the nature of the stressor (Mason, 1971); it is important to consider this when assessing stress impacts. Changes in abiotic conditions, such as in food availability, or the introduction of toxins or diseases can be stressful, but such changes do

not arouse fear. However, a stress response involving peripheral autonomic and neuroendocrine changes (Yates, Russell & Maran, 1971; Sapolsky, Romero & Munck, 2000; McEwen & Wingfield, 2003), can be initiated by fear arousal (LeDoux, 2003; LaBar & LeDoux, 2011). Animals with sophisticated nervous systems, such as mammals, exhibit a central motive state between threat stimulus and response that is driven by the amygdala (Pitkanen, 2000) and can be identified as ‘fear’ (Mineka, 1979; Öhman, 2000). Fear has been defined as a psychological state that triggers physiological responses to avoid or escape from a stressor (Epstein, 1972; LeDoux, 1996). The development of successful defense mechanisms to fear-inducing stressors has clear survival benefits for animals, and thus fear can be seen as a driver of evolutionary adaptations (Tooby & Cosmides, 1990). Common strategies of escape and avoidance allow animals to cope with recurrent stressors, such as the fear-inducing threat of predation (Lima & Dill, 1990).

The sections of the mammalian amygdala associated with fear behaviour serve as an interface between sensory input and information transport and processing, endocrine response, and motor output (Davis & Whalen, 2001). These interactions are associated with learning and memory via the involvement of the lateral and basal nuclei, as demonstrated on captive rodents using neurotoxic lesions on the basal and lateral nuclei of the amygdala (e.g. Wallace & Rosen, 2001), and are evident in Pavlovian fear conditioning paradigms (Davis, 1992; Maren, 2001). Activation of hormones, specifically glucocorticoids and norepinephrine in stress responses initiated by fear arousal provides feedback to the brain that influences emotion control and cognition, which contributes to fear conditioning (Rodrigues, LeDoux & Sapolsky, 2009). Fear responses may, therefore, be both conditioned as aversive learnt behaviours (e.g. Fanselow & Poulos, 2005), and unconditioned as innate freezing responses (e.g. Schulkin, Thompson & Rosen, 2003).

Given that fear motivates a stress response – initiating the freeze, and fight or flight actions – quantifying glucocorticoid outputs from the autonomic nervous and HPA systems should yield a measurable indication of fear from predation as a stressor. Minimally invasive techniques are available that assay glucocorticoid levels in faeces, fur, or feathers (Sheriff *et al.*, 2011; Cook, 2012; Palme, 2019). Using minimally invasive methods to measure glucocorticoid levels in the wild does not necessarily yield an isolated 'output' of a fear or predator/human-induced stress response (Rosen & Schulkin, 2004). This situation results from glucocorticoids not being a molecule of fear *per se* but having a fundamental role in maintaining energy balance (Rosen & Schulkin, 2004). As such, glucocorticoids may change not just because of fear but also in response to other exogenous stressors, like food shortage (McEwen & Wingfield, 2003). Nevertheless, such methods could usefully compare longer-term stress responses among habitats or along disturbance gradients with variable exposure to predators, by following experimental designs that are considerate of additional stressors that may be present (see: MacDougall-Shackleton *et al.*, 2019).

If two or more stressors are present, the resultant combined stress may present severe challenges to an individual's physiological systems (Johnstone, Lill & Reina, 2012; Brearley *et al.*, 2013; Malcolm *et al.*, 2014; Arlettaz *et al.*, 2015; Geary *et al.*, 2019; Legge *et al.*, 2019). Interactive fear and stress effects are context-dependent (Belarde & Railsback, 2016). They can be additive and combine the multiple impacts, synergistic whereby the presence of one threat amplifies another (Doherty *et al.*, 2015), or antagonistic whereby one threat cancels the effects of the other. A reduction in the activity of mesopredators in the presence of humans provides an example of antagonism (Clinchy *et al.*, 2016). Additive, synergistic, and antagonistic reactions have each been observed in mammalian prey in natural systems in response to exposure to

multiple stressors (Crain *et al.*, 2008; Côté, Darling & Brown, 2016; Gunderson *et al.*, 2016; Jackson *et al.*, 2016; Geary *et al.*, 2019; Legge *et al.*, 2019).

Exposure to a stressor(s) that is prolonged, constant, or recurring can have chronic impacts, as recovery from a stressor cannot occur whilst the threat remains (Sapolsky, Romero & Munck, 2000). Acute stress occurs as the initial response to a threat to sustain fitness in the short term; it subsides once the responding action—be it freezing, fighting, or fleeing—diminishes the threat (Wingfield & Kitaysky, 2002). Activation of the HPA axis in an acute stress response has rapid effects that increase immune system function, energise muscles via enhanced cardiovascular tone, and heighten cognition, including memory (Sapolsky, Romero & Munck, 2000). These responses occur via increased cerebral perfusion rates and use of glucose, all of which come at the cost of decreased appetite and reproductive behaviours (Sapolsky, Romero & Munck, 2000). Effectively, the acute stress response suspends non-essential behaviours in favour of altered behaviours that minimize the threat (Wingfield & Kitaysky, 2002).

Continued exposure to a stressor, or stressors, creates a state of chronic stress, which is classically described as allostatic overload (Dantzer *et al.*, 2014). Allostatic load describes the body's ability to maintain homeostasis in response to a stressor (Sterling & Eyer, 1988; McEwen & Stellar, 1993; McEwen & Wingfield, 2003). Allostatic overload, by extension, refers to the inability to maintain homeostasis and thus an organism's increased susceptibility to external stressors (Sterling & Eyer, 1988; McEwen & Stellar, 1993; McEwen & Wingfield, 2003). Chronic stress reduces an organism's resilience to future stressors by inducing extended behavioural changes in feeding, fighting, and mating (Mineur *et al.*, 2006). Chronic stress also affects an organism's physiological state by suppressing or impairing the reproductive system and decreasing physiological resistance to pathogens and toxins through the suppression of

immune function (Dhabhar & McEwen, 1999; McEwen & Wingfield, 2003; Romero, 2004; Travers *et al.*, 2010; Feng *et al.*, 2012). Repeated exposure to a stressor can have population and community level consequences as homeostasis is compromised, and it increases susceptibility to additional stressors and can have flow-on effects. For example, chronic risk of predation, which facilitates changes in nutrient demand for prey species and thus changes in the elemental composition of their excreta, can alter nutrient cycling in the community and ecosystem (Hawlena & Schmitz, 2010).

Cases of acclimatisation to chronic or repeated acute stressors have been observed, although the process often results in enhanced activation of the HPA axis to novel stressors, and thus may not be beneficial to fitness (Romero, 2004). Instead of acclimatising to a chronic or repeated stressor, glucocorticoid levels can remain the same, or become chronically elevated, or the HPA axis can shut down completely and render an animal vulnerable to future threats (Romero, 2004). Physiological impairment of the neurological, cardiovascular and musculoskeletal systems may also result from chronic stress: neurons of the brain can atrophy and impair memory, or grow and enhance fearfulness with extensive releases of adrenaline and cortisol (Roozendaal, 2000; McEwen, 2004); atherosclerotic plaques also may form and impede blood flow from repeated elevation of blood pressure (Manuck *et al.*, 1988), and skeletal muscle can suffer severe protein loss (Wingfield & Kitaysky, 2002).

Although chronic and acute stress responses have been well defined for mammals in laboratory studies, results from *in situ* studies show that reactions vary with the stressor type that animals are exposed to, as well as habituation potential, food availability, social interactions, and density (Dickens & Romero, 2013). *In situ* mammalian endocrine responses to a consistent stressor, such as human activity, have not been studied sufficiently to be able to determine if

there is a particular pattern of response. Avian endocrine responses to such stressors have been more extensively studied and they reveal that urban habitats, perceived as a consistent stressor, can shape endocrine responses in birds. However, a consistent stress response pattern has yet to be observed in birds (Bonier, 2012). Inconsistent behavioural responses of mammals and other animals to human activities that are often observed can perhaps be linked to modulating factors such as the level of human activity, the species and condition of the animal being observed, and the spatio-temporal context (Tablado & Jenni, 2017).

Mammalian behavioural responses to fear and stress: the landscape of fear concept

The landscape of fear concept (Laundré, Hernández & Altendorf, 2001) postulates that prey are aware of microhabitat patches associated with high and low predation risk, where predators are either active and ubiquitous, or scarce (Laundré, Hernández & Altendorf, 2001; Shrader *et al.*, 2008; van der Merwe & Brown, 2008; Laundré, Hernández & Ripple, 2010). Theoretically, landscape of fear effects should increase with landscape heterogeneity, since the differences between high and low risk sites become more pronounced and prey can more easily avoid high-risk sites (Bleicher, 2017; Gaynor *et al.*, 2019). However, this relationship can depend on the species and condition of the habitat and, for some species, simple habitats can be the safest (e.g. Hammerschlag, *et al.*, 2015; Schmidt & Kuijper, 2015; Atuo & O'Connell, 2017). Landscape of fear effects will also be more pronounced in systems where interactions between predators and prey are less frequent (Schmitz, 2008), as is evident in findings from Pavlovian fear conditioning, and the '*risk allocation hypothesis*' – which states that animals exposed to constant high predation risk will increase their foraging risks over time (Lima & Bednekoff, 1999; Van Buskirk *et al.*, 2002). Fear arousal in a landscape of fear results in two predictable outcomes:

either avoidance of high risk areas, or modulation of behaviour (e.g. increased vigilance) to reduce predation risk when foraging in such areas (Gaynor *et al.*, 2019). These outcomes indicate the potential positive benefits to conservation management of considering fear arousal and stress levels, and the direct and indirect cues that may trigger these effects (Atkins *et al.*, 2017).

The likelihood of survival of a prey species is improved by having a well-developed perception of high predation risk, as it allows adaptive behavioural responses to be established (Bókonyi *et al.*, 2009). Habitat shifts, temporal shift, grouping, vigilance, and freeze, fly, and fight responses are the most studied such responses to the non-consumptive effects of predators (Say-Sallaz *et al.*, 2019). These often develop due to increased interaction rates between predators and prey. However, chronic stress impacts that are sufficient to affect reproduction and long-term survival can be experienced by prey species perceiving recurring predation risks (Thomson *et al.*, 2010; Clinchy *et al.*, 2011). Chronic perceived predation risk may also result in altered foraging activity driven by fear. This can affect where and what prey eat (Schmitz, Krivan & Ovadia, 2004), causing them to move from risky to sheltered microhabitats (Trussell, Ewanchuk & Matassa, 2006), in turn altering the distribution and availability of resources. Fear-based adaptive behavioural responses such as these underlie the landscape of fear concept.

For mammals, fear arousal can be triggered both by a predator's presence or by a predator cue, such as an associated scent. Olfaction can be a key driver of fear arousal (Soso *et al.*, 2014; Banks, Daly & Bytheway, 2016; Jones *et al.*, 2016; Parsons *et al.*, 2017). The mechanics of this are best understood in mammals: odours are detected by the accessory olfactory bulb that transmits information directly to the amygdala and hypothalamus, where fight or flight responses are developed (Fogaca *et al.*, 2012; Canteras, Pavesi & Carobrez, 2015). Laboratory studies exploring the effects of post-traumatic stress disorder have exposed small

mammals to predators or their cues to induce stress, and in doing so revealed that exposure to predator cues alone can affect the neural circuitry associated with fear (Rosen & Sculkin, 1998, 2004).

Subtle cues such as predator odours may precede threats and allow for a mammalian prey animal's fear state to be conditioned to a cue that occurs before, or in correlation with, a previously encountered predation threat (Rescorla & Solomon, 1967; Rosen, 2004). Predation risk may, therefore, be perceived by prey species eavesdropping on predator scent marks, such as urine, faeces, or fur in the environment (Banks, Daly & Bytheway, 2016; Jones *et al.*, 2016). Such odours have been observed experimentally to induce fear-like responses of freezing (Wallace & Rosen, 2000), vigilance (Nersesian, Banks & McArthur, 2012), fleeing (Anson & Dickman, 2013), and avoidance (Hayes, Nahrung & Wilson, 2006), across a wide range of species in both field and laboratory experiments (Apfelbach *et al.*, 2005, 2015). Consequently, landscape of fear topography, where predators indirectly influence prey behaviour across a range of microhabitats, can arise from the influence of predator olfactory cues on mammalian prey foraging behaviours as much as it can from the direct threat of predation (e.g. Brown & Kotler, 2004; Parsons & Blumstein, 2010; Cremona, Crowther & Webb, 2014; Mella, Banks & McArthur, 2014; Hoffman, Sitvarin & Rypstra, 2016). It is worth noting, however, that whilst predator olfactory cues can elicit a fear or stress response, they do not always do so (Apfelbach *et al.*, 2005). This may be because predators often leave the site after depositing a cue, and the cue intensity diminishes. Predator cues elicit responses that are modulated according to the differential intensity of the perceived threat, prior experience, or pending further information gathering, and thus can be perceived as a low risk by prey (Bedoya-Pérez *et al.*, 2019).

Fear arousal due to predator presence or cues can deplete a prey individual's energy budget, resulting in poor reproduction and health either via energy exhaustion from stress (i.e. the predation stress hypothesis: Boonstra *et al.*, 1998; Clinchy *et al.*, 2004; Romero, 2004; Støen *et al.*, 2015), or reduced nutrition from foraging compromises (Brown & Kotler, 2004; Clinchy *et al.*, 2016). As food resources become limited, additionally, prey may take greater risks to meet nutritional needs (i.e. the predation-sensitive food hypothesis: Sinclair & Arcese, 1995), but at a possible cost of additional stress. In areas of high human activity, this may result in wildlife increasing the stressors they are exposed to as they seek human-based food sources, such as waste or pet food, and in doing so increase their exposure to domestic pets, altered light, increased sound, and vehicles (Navara & Nelson, 2007; Morgan *et al.*, 2009; Riley *et al.*, 2014; Shannon *et al.*, 2016; Doherty *et al.*, 2017). For managers mitigating fear arousal, mapping landscapes of fear (e.g. van der Merwe & Brown, 2008; Kauffman, Brodie & Jules, 2010; Iribarren & Kotler, 2012, Smith *et al.*, 2019) to identify, protect and extend safe foraging areas, could assist in the conservation of wildlife subject to human activity or multiple stressors such as human activity and predators.

Human activity as a fear-inducing stressor

Human activity can influence wildlife in a wide variety of ways (e.g. Albert & Bowyer, 1991; Bowyer *et al.*, 1999). Pressure from increased proximity to human activity or development that encroaches upon animal home ranges, result in wildlife becoming either: 1) 'urban avoiders' that move away from human activity; 2) 'urban adapters' that make some use of anthropogenic resources but still rely largely upon those found naturally; or 3) 'urban exploiters' that are synanthropic and make full use of anthropogenic resources (McKinney, 2006). The effects of

human activity can therefore differ between species and trophic levels, and even within the same species, depending on personality. Furthermore, the negative effects of a disturbance on one species may result in flow-on responses of positive consequences for its prey, predators, or competing species (e.g. Gill, Sutherland & Watkinson, 1996; Crooks & Soulé, 1999; Leighton, Horrocks & Kramer, 2010). These effects, however, depend strongly on the availability of necessary resources within the human occupied / disturbed areas, and will often differ between systems.

It has been postulated that humans may impose widespread effects on ecosystem function if they induce greater fear responses in ubiquitous small predators than in top predators (Clinchy *et al.*, 2016), or by increasing physiological stress in large ungulates through antagonistic behaviours (Vijayakrishnan *et al.*, 2018). Indeed, human activity can be comparable to the above-mentioned impacts of predation, creating landscapes of fear (Hofer & East, 1998; Frid & Dill, 2002; Rehnus, Wehrle & Palme, 2014; Patten & Burger, 2018), and disrupting foraging (Ciuti *et al.*, 2012; Clinchy *et al.*, 2016). Humans can also act as “super-predators” that disproportionately kill carnivores and drive trophic cascades (Darimont *et al.*, 2015). Trophic cascades can also arise when human activity has non-consumptive effects on the ecological roles of large predators (Smith *et al.*, 2017). For example, large predators may increase their hunting efforts in urban areas: if their own fear response to human activity results in less time spent consuming a kill, this in turn may increase their kill rate of prey in urban areas to meet their energy needs (Smith *et al.*, 2017). When large predators avoid human activity, and / or meso-predators reduce their foraging around human activity, prey species may respond by increasing their own foraging activity (Suraci *et al.*, 2019), effectively taking refuge behind the “human

shield” that is reducing predation pressure for them (Berger, 2007; Leighton, Horrocks & Kramer, 2010; Kuijper *et al.*, 2015).

Human activity can affect the trophic structure of communities not only via top-down pressure, but also by exerting bottom-up pressure through the constant provision of alternative resources (Fischer *et al.*, 2012), such as subsidies of shelter, water and food with year-round primary production. These resources can outweigh the stresses of human disturbance and influence population and community behaviours (Parris, 2016), resulting in urban colonisation by wildlife (Shochat, Lerman & Fernández-Juricic, 2010; Jokimäki, *et al.*, 2011). As such, human activity does not always result in fear responses, particularly for prey species that benefit from the human shield and access to food, water or shelter subsidies (Lyons *et al.*, 2017). It is possible that, given these conditions, and provided that resource subsidies are nutritionally rich and accessing them does not increase pathogen transmission risk (Murray *et al.*, 2016), prey species like small mammals would experience fitness benefits in urban areas. Resource subsidies can also be exploited by predators, resulting in increased predator activity in urban areas but also in altered foraging behaviours that either decrease or increase predation on small prey (Iossa *et al.*, 2010; Bateman & Fleming, 2012; Fischer *et al.*, 2012; Newsome *et al.*, 2014, 2015).

Despite such positive effects of human activity on wildlife, negative impacts are more pervasive for many species. The negative effects of high levels of disturbance, for example from roads and vehicles, or of active interference from recreational activities conducted in designated conservation areas, can lead to physiological stress or displacement, countervailing any positive effects of humans on wildlife (Kloppers, St Clair & Hurd, 2005; Banks & Bryant, 2007; Berger, 2007; Ciuti *et al.*, 2012; Rehnus, Wehrle & Palme, 2014; Arlettaz *et al.*, 2015; Patten & Burger, 2018; Vijayakrishnan *et al.*, 2018).

A common response in mammals to increased exposure to human activity is a temporal shift in activity patterns, from diurnal to crepuscular or nocturnal, to avoid interaction with humans (McClennen *et al.*, 2001; Tigas, Van Vuren & Sauvajot, 2002; Riley *et al.*, 2003; Ditchkoff, Saalfeld & Gibson, 2006; Gaynor *et al.*, 2018); the same or reverse may occur too, to reduce interactions with predators (Brown, 2000; Laundré, Hernández & Altendorf, 2001; Kohl *et al.*, 2018; Smith *et al.*, 2019). Temporal shifts can have strongly negative effects if they limit a forager's ability to locate and capture prey (Ditchkoff, Saalfeld & Gibson, 2006). Regardless of the response, it is evident that human activity can have profound indirect effects on community interactions through altering individual behaviour, particularly foraging, through either fear arousal or a stress response (e.g. Frid & Dill, 2002; Werner & Peacor, 2003).

The ability of wildlife to cope with urban environments can occur through either plastic or evolved shifts in behaviour, foraging, food preferences, or predator avoidance, and in shifts in the timing of breeding (Ditchkoff, Saalfeld & Gibson, 2006; Møller, 2009; Shochat, Lerman & Fernández-Juricic, 2010; Rodriguez, Hausberger & Clergeau, 2010; Alberti, 2015; McDonnell & Hahs, 2015; Otto, 2018; Santini *et al.*, 2019). It is most common for urban-occupying species to plastically adjust to human imposed stressors (Donihue & Lambert, 2015), with some ultimately evolving tolerance to urban environments. Human activity can therefore impose pressures that are strong enough to influence changes in behaviour and in physiology — endocrinology in particular (Bonier, 2012; Snell-Rood & Wick, 2013; Otto, 2018). Bolder individuals are most likely to exploit novel opportunities and have reduced stress responses to human activity (Atwell *et al.*, 2012). Animals that routinely encounter human activity, typically provide parental care, and are capable of reproducing multiple times across their life span may produce offspring that are acclimated to human activity and accordingly show lower fear/stress responses (Schell *et al.*,

2018). Habituation of groups of animals to human activity is being increasingly observed, often as a result of bold individuals spending significant amounts of time around human-disturbed areas and displaying reduced fear responses (Stillfried *et al.*, 2017). This counters the idea that human activity always imposes landscapes of fear, but may reflect the combination of reduced resource availability and bold personalities that allow for urban habituation (Lowry, Lill & Wong, 2012). In some instances this may be problematic for humans if increased interactions with habituated animals increase vehicle accidents and / or transfer of zoonotic diseases; this highlights the potential need to manage some mammalian prey species in urban areas (Honda *et al.*, 2018).

Fear and acute/chronic stress may be constant hurdles faced by wildlife, but adding the effects of anthropogenic disturbance could result in the elevation of acute stress responses to prolonged and widespread chronic stress, and upscale the possible impacts from individuals to populations (Rehnus, Wehrle & Palme, 2014). As we have outlined, human activity can both indirectly and directly create landscapes of fear, influencing and amplifying existing landscapes of fear through interactive effects. Predator presence and human activity can also interact to alter ecosystem structure and modify the predation risk perceived by prey species, with either positive or negative impacts dependent upon the circumstances. The combination of stress resulting from the pressures of altered predator-prey interactions and human activities is yet to be investigated extensively. Understanding the cumulative effects that both may impose is critical for effective conservation of wildlife in an increasingly human-influenced world.

Alleviating fear and stress for wildlife conservation

431 As we have illustrated, fear can produce stress and simultaneous multiple stressors can have
 432 chronic effects that negatively influence both predator and prey species populations (Allan *et al.*,
 433 2013). To synthesize these impacts, we note that in modern urban situations both habitat
 434 stressors and introduced stressors now collide (Figure 1). Habitat stressors result from naturally
 435 occurring environmental factors such as native predator presence (especially for prey), social and
 436 reproductive pressures, limited access to refuges / nests resource, food and water, disease and
 437 parasite prevalence, and produce predictive homeostasis responses by prey species (Figure 1).
 438 Introduced stressors, by contrast, are additional stressors that are imposed by human actions that
 439 result in reactive homeostasis responses. These introduced anthropogenic stressors include
 440 structural developments that change landscapes, human recreational activities in natural habitats
 441 (including use of vehicles), introduced pollutants and toxins that alter landscapes and resource
 442 quality, and introduced predators and altered predator-prey interactions (Figure 1). The
 443 combination of impacts from both naturally occurring and introduced anthropogenic stressors
 444 has the potential to cause a cumulative stress response that can result in homeostatic overload or
 445 failure, as defined by Romero, Dickens and Cyr (2009), and may result in population collapses
 446 due to increased susceptibility of individuals to the additional stressors (as we have demonstrated
 447 above, and Figure 1). Cumulative stress may be additive or synergistic, and is likely to be
 448 particularly detrimental for prey species. For example, if prey species face multiple stressors,
 449 they may take greater foraging risks, or be less able to allocate energy to vigilance or flight
 450 behaviours, and thus become more susceptible to predation or additional stressors. In the case of
 451 population-impacting stressors, the local population may be impacted to a high degree such that
 452 it becomes threatened (e.g. Sweitzer, 1996; Doherty *et al.*, 2015). As such, conservation action in

areas where simultaneous introduced stressors occur in addition to natural ecosystem pressures, like in urban and peri-urban habitats, may urgently needed.

The most direct way to alleviate stress in wildlife is to remove or reduce the key stressors. This may be difficult if the stressors are anthropogenic, as the needs of the expanding human population often usurp those of wildlife; alternative solutions then need to be considered. A suite of activities related to managing vegetation structure, density, and/or heterogeneity may well provide such an alternative solution. These activities result in increased availability of potential refuge sites and may be readily achieved through ongoing habitat engineering that, for example, supports the growth of structurally complex plants or adds logs / rock piles to reduce areas of open space. In order to mitigate introduced fear and stress effects, it is important that future studies investigate whether vegetation management and other habitat components can alleviate some of the pressures associated with multiple introduced stressors for target species.

Managing habitat to alleviate wildlife fear and stress

The impact of predators (or human activity) can be mitigated by changing the configuration of risky areas within a habitat (Hopcraft, Sinclair & Packer, 2005; Lone *et al.*, 2014). Food availability often drives habitat selection (e.g. Sherman, 1984; Johnson & Sherry, 2001), but predation risk and human activity affect a suite of correlated factors such as movement decisions (Turcotte & Desrochers, 2003), foraging patterns (Gil, Zill & Ponciano, 2017), social organisation (Rodríguez, Andrén & Jansson, 2001), and reproductive success (Zanette *et al.*, 2011). Supplementation of essential resources including refuges, nests or roosts, food, and water to wildlife is a growing conservation method due to its ease of implementation, immediate

results, and favourable portrayal in the media (Sergio *et al.*, 2018). One example of positive results includes increased parasite resistance in nestling birds after supplementation with high-quality food during the stressful young-rearing stage (Knutie, 2020). Another example is the reduction in impact from introduced predators on small desert mammals following the addition of artificial refuge structures (Bleicher & Dickman, 2020). Despite these successes, negative impacts can also ensue following supplementation. Built-up refuges may be primarily investigated by invasive species, such as the black rat (*Rattus rattus*), which could potentially deter commensal native species and support the population growth of invasive species (Price & Banks, 2018). Artificial nests or roosts can support increased predation rates owing to the structure supporting a sit-and-wait predator at the exit (e.g. McComb *et al.*, 2019). Food and water subsidies can increase pathogen transmission, reduce movement and migratory behaviours, increase predator-prey interaction rates, and competitor aggression interactions (Murray *et al.*, 2016). However, owing to the benefits of supplementation, especially during times of stress, it may be used as a tool to mitigate the negative impacts of stress on wildlife if conducted in a targeted, monitored and well-considered manner (e.g. Freeman *et al.*, 2020).

Considering the existing knowledge base on food and water supplementation, meta-analyses have suggested that the associated disease risk may be managed by focusing on specific natural food sources and pathogen groups subject to the target ecosystem and by increasing the spatial extent of feeding stations across potential linked habitats, which also limits microparasite host aggregations (see: Becker, Streicker, & Altizier, 2015; Becker *et al.*, 2018). In considering the specific needs of target animals and ecosystems, chainsaw-carved cavities have more favourable thermodynamic properties for small mammals and birds than do artificial nest boxes (Griffiths *et al.*, 2018). As they mimic natural hollows they are also less likely to increase rates

of predation. Supplementation projects are therefore moving forward using this growing knowledge base to adaptively tailor methods to specific conditions, and to orchestrate pre- and post-monitoring that ensures successful stress mitigation (Civitello *et al.*, 2018).

Survivorship of prey, in the face of predation stressors, is often positively correlated with increased structural complexity of the habitat (Hopcraft, Sinclair & Packer, 2005; Lone *et al.*, 2014; Leahy *et al.*, 2016). This has been demonstrated in many studies to be a consequence of the increased opportunities for prey to escape and hide, thereby mediating predator-prey dynamics by reducing encounter rates, as only a proportion of the total prey population remains available to predators (e.g. Holt, 1984; Kotler & Brown, 1988; Bianchi, Schellhorn & Van Der Werf, 2009; Rieucau, Vickery & Doucet, 2009; Klecka & Boukal, 2014; Laundré *et al.*, 2014). The extensive research on the benefits to prey of habitat structural complexity builds on the ‘*habitat heterogeneity hypothesis*,’ which postulates that structurally complex habitats support increased species diversity by offering a wide range of niches and diverse ways of exploiting resources (Bazzaz, 1975). Habitat complexity, by extension, has been generalized to be a primary driver of biodiversity (Pianka, 2011). The landscape of fear concept accepts this principle too, in that a wide range of microhabitat types offers multiple foraging and shelter conditions with varied predation risks for species. For example, northern quolls (*Dasyurus hallucatus*) of the semi-arid Pilbara region in Western Australia utilise complex rocky habitats in preference to open grasslands where the threat of predation from feral cats (*Felis catus*) is greater (Hernandez-Santin, Goldizen & Fisher, 2016).

Animals in habitats with high predation pressure may display foraging preferences for microhabitats or times that they perceive to be safe (Brown, 2000; Laundré, Hernández & Altendorf, 2001; Laundré, Hernández & Ripple, 2010). Some small mammals seek structurally

complex vegetation owing to the reduced risk of predation and increased reward of foraging they find there (Lima & Dill, 1990; Andruskiw *et al.*, 2008). Others, such as the Australian hopping mouse (*Notomys alexis*) that exhibit bursts of speed in open habitat, are less adept at moving through complex vegetation (Spencer, Crowther & Dickman, 2014). No matter the species, in landscapes of fear a prey species' use of the topography, refuges, and escape substrates can indicate its perceived risk of predation (Brown & Kotler, 2004; van der Merwe & Brown, 2008; Shrader *et al.*, 2008), and the associated fear and stress that may arise or be alleviated due to habitat structure. The same principle should be relevant to reducing stress in proximity to human activity, considering that prey responses to human activity and predators are similar.

Human activity and disturbance can quickly reduce habitat complexity (Western, 2001). However, enhancing habitat complexity and heterogeneity is being incorporated increasingly into restoration and management efforts, with some success (Brown, 2003; Bernhardt & Palmer, 2007; Palmer, Menninger & Bernhardt, 2010). To balance the needs of people and biodiversity, local planning procedures are at times now incorporating green spaces and greening initiatives into urban areas. As habitat complexity and diversity are of particular importance in supporting biodiversity and population sustainability, it is important that habitat structure underpins the engineering of wildlife habitats in urban and urban-adjacent areas (Threlfall *et al.*, 2016). The effectiveness of any management regime depends on recognising the direct and indirect impacts that occur across sustainable ecosystems. Consequently, habitat complexity as a management objective requires that urban landscapes are approached on a case by case basis, with full assessment before the habitat is ecologically engineered (Tews *et al.*, 2004).

Management tools to observe and alleviate fear and stress for wildlife conservation

Cumulative stress maps of human activity and stressors that occur naturally in ecosystems have allowed returns on restoration investments to be maximized by indicating key areas that will or do need intervention to mitigate the effects of severe stress on wildlife, as opposed to unsuccessful piecemeal management that focuses on one or two stressors broadly (Allan *et al.*, 2013). Physiological and behavioural stress in wildlife may be cost-effectively and straightforwardly observed by comparing the outcomes of simultaneous measurements taken across a disturbance gradient where multiple stressors are both abundant and low, using several well-established methods. Specifically, assays of faecal/urine/fur/feather glucocorticoid metabolites can provide insight into the level of physiological stress an animal is experiencing under comparative circumstances (see Cook, 2012; Cooke *et al.*, 2013; Sheriff *et al.*, 2011; Palme, 2019). Pairing such results for each location with those of giving-up density surveys (Brown, 1988) that can also be filmed by infrared motion sensor cameras to observe foraging behavioural responses (e.g. Leo, Reading & Letnic, 2015), yields an understanding of both physiological stress responses and foraging responses. Comparing these results may further indicate if behavioural responses are changing to moderate physiological stress impacts (e.g. Carlstead, Brown, & Seidensticker, 1993; Carlstead, Brown, & Strawn, 1993). Simultaneous measurements of habitat quality that include vegetation cover, refuges, distance to food / water source, and distance to disturbances (e.g. human activity or predator sighting based on spotlighting or in-field camera traps) will build a dataset for spatial correlation analyses that can geographically map the landscape of fear. Such measurements should also identify what factors influence the landscape, such as by use of behavioural data (e.g. Willems & Hill, 2009) and / or giving-up density data (e.g. van der Merwe & Brown, 2008).

Spatial mapping of the physiological stress response, based on the collection locations of samples, could be overlaid and compared to these results also. Such spatially mapped results may clarify important areas that are in need of protection or the habitat types that could be extended to mitigate stress impacts. It is worth noting that several authors have emphasized that these methods need to be used correctly to ensure that stress and habitat factors are linked (e.g. McMahon *et al.*, 2018; Bleicher, 2017; Bedoya-Perez *et al.*, 2013; MacDougall-Shackleton *et al.*, 2019). For example, collection of fresh faecal samples of a target species along a gradient of high to low human activity would allow glucocorticoid metabolites to be assayed, providing insight into potential stress induced by human activity along the gradient (Rehnus, Wehrle & Palme, 2014). A similar approach could be used to assay stress perceived by prey species in areas of high and low predator activity, and sex hormones also could be assayed to explore sex-related effects of stress. By allowing pathways of fear and stress to be mapped, including sites where stressors from predators and human activity occur simultaneously, such methods should allow managers to reliably identify where best to intervene to preserve at-risk populations and maintain community stability. We consider potential interventions in more detail below.

If cumulative stressors occur and are suspected to have detrimental effects on target wildlife populations or communities, we suggest that a potentially powerful mitigation approach could be developed based on Cumulative Effects Assessments (CEA). Devised in the 1990s amid growing concerns that Environmental Impact Assessments (EIA) did not consider all the effects of urban and peri-urban developments (Smit & Spaling, 1995), CEAs have been advocated as an effective tool for use by on-ground practitioners (Duinker *et al.*, 2012). Numerous countries have mandated that CEAs be incorporated into EIAs, namely: the United States of America, Canada, Europe, New Zealand, and Australia each advocate some level of CEAs (Therivel & Ross,

2007). Despite this legal requirement, and the CEA concept being widely known in scientific literature, it is rarely applied in practice (Ma, Becker & Kilgore, 2009; Foley *et al.*, 2017).

We propose further that the principles of the ‘*reactive scope model*’ be used to develop CEAs, to identify where cumulative stressors occur, and thus better inform conservation management initiatives in areas where wildlife is subject to homeostasis overload or failure (Figure 2). The results from the above mentioned methods and / or other applicable established methods may be used to answer our suggested CEA questions. The reactive scope model (Romero, Dickens & Cyr, 2009) provides useful insight into the range of physiological mediators available in response to a stressor. It maps the homeostasis range of a given species in four stages: 1) predictive – change occurs in response to routine environmental change, such as seasons or day to night; 2) reactive – change occurs in response to an unpredictable change, allowing survival via classic stress responses; 3) overload – consistent changes occur in response to a stressor, and chronic stress impacts start to occur; and 4) failure – the species shows inability to sustain homeostasis, and is very susceptible to additional stressors and death (Romero, Dickens & Cyr, 2009). If our proposed CEA approach (Figure 2), based on the reactive scope model, were applied to small mammals in urban, urban adjacent, and peri-urban ecosystems, for example, then areas of conservation concern could be identified where the additive or synergistic impacts of human disturbance and introduced predators combine with stressors that occur naturally. Appropriate management, such as increasing habitat complexity or reducing human activity in conservation areas (e.g. Bleicher & Dickman 2020), could then be implemented to alleviate these stressors.

Fear and stressing in small mammal ecology in Australia; the need to consider the twin stressors of introduced predators and people in wildlife management

Small mammals are often somewhat resilient to threatening processes owing to their high population growth rates (Cardillo *et al.*, 2005, 2006). However, in Australia small mammal populations are declining quickly, and due to causes dissimilar to those driving global declines (Woinarski *et al.* 2015). Declines in Australia have been attributed to a wide range of habitat stressors: habitat loss, altered fire regimes, disease, increasing temperatures, decreasing water availability, depleted soil quality, and salinity (Woinarski *et al.*, 2015). However, mammals within a 'critical weight range' (CWR: 35 to 5550 g) are particularly vulnerable (Chisholm & Taylor, 2007; Woinarski, Burbidge & Harrison, 2015) owing to the effects of predation by two introduced carnivores, the red fox (*Vulpes vulpes*) and domestic house cat (*Felis catus*) that arrived in Australia over 150 years ago (Johnson, 2006). If predators are introduced, their impacts on prey are likely to be exacerbated owing to prey naïveté (Doherty *et al.*, 2016). Such impacts have been particularly acute on wildlife in Australia, where eutherian carnivores are recent arrivals (Salo *et al.*, 2007). However, the question of Australian native animal stress responses and the extent of their naïveté to these introduced eutherian predators is debatable (see: Banks & Dickman, 2007; Carthey & Banks, 2016). Although CWR mammals are of high conservation concern, predation from introduced predators poses a threat also to all native predators by reducing their food resources, which in turn may increase predation and predation-associated stress on alternative food sources such as smaller (< 35 g) or larger (> 5550 g) mammal species. For example, a study examining the diet of a nocturnal, avian predator, the sooty owl (*Tyto tenebricosa*) before and after red fox introduction, revealed a dietary shift post introduction, with owls consuming more arboreal than terrestrial prey species after fox arrival

(Bilney, Cooke & White, 2006). This shift to consuming arboreal prey increased dietary overlap with the sympatric powerful owl (*Ninox strenua*), providing disproportionate predation pressure on prey in the ecosystems of East Gippsland, Victoria.

Predation by red foxes and cats is prevalent not only in natural habitats but also in agricultural and urban habitats (Dickman, 1996; Morgan *et al.*, 2009; Bino *et al.*, 2010). The paths and roads that fragment urban and agricultural habitats are used frequently by these predators, which exacerbates their predation pressure on prey species by combining with impacts imposed by human activities (Latham *et al.*, 2011, Červinka *et al.*, 2013). Consequently, this can result in increased abundances of red foxes and cats in human-modified landscapes that border or include natural habitats (Towerton *et al.*, 2011; Graham, Maron & McAlpine, 2012). Urban and agricultural habitats present many threats to wildlife, but they may also offer food and shelter opportunities (Pickett *et al.*, 2001; Gaston *et al.*, 2005; Hobbs, Higgs & Harris, 2009). As reported by Ives *et al.* (2016), 46 percent of threatened Australian animals occur in or near Australian cities. Thus, the fate of many species could depend on accommodating their needs in urban and agricultural habitats (Ives *et al.*, 2016). A recent assessment of data collected at the Wildlife Rehabilitation Centre in Queensland Zoo, Australia, revealed that pet cat or dog attack, car strike, and entanglement in human-placed objects represented 56.4% of the causes of submission of injured wildlife; mortality rates associated with these traumas were also high, with 61.3% of admitted animals dying from their injuries (Taylor-Brown *et al.*, 2019). These threats may contribute to landscapes of fear, through fear arousal, altered foraging behaviour, post-traumatic stress reactions, and cumulative stress exposures resulting in chronic stress responses.

Despite growing knowledge of anthropogenic impacts on threatened species, Australian conservation planning continues to exclude urban and agricultural habitats from consideration

(Dales, 2011). There is recognition of the scope of this issue in Australia (e.g. Hill *et al.*, 2007; Carthey & Banks, 2012; Threlfall, Law & Banks, 2012; McCauley *et al.*, 2013; Banks & Smith, 2015; Ives *et al.*, 2016), but few studies have explored the interactive effects of introduced predators and human activity on the survival of prey species. Despite the development of glucocorticoid analysis techniques to determine stress in animals dating to the 1960s (Jones *et al.*, 1964), only six of the 60 extant small mammals of conservation concern in Australia have been subject to studies seeking to better understand their glucocorticoid response to stressors (Hing *et al.*, 2014), such as predation by introduced species or human activity.

There is a need for alternative management regimes to mitigate the additional stressors of human-imposed impacts on wildlife, and the situation we have outlined in Australia underlines the urgency of this need. As there are difficulties with the current control of introduced predators in Australia (Glen & Dickman, 2005; Rayner *et al.*, 2007; Bergstrom *et al.*, 2009; Norton, 2009; Warburton & Norton, 2009; Carroll, 2011; Newsome *et al.*, 2017), there is also an urgent need for alternative management regimes to mitigate the additional stressors of introduced predator-imposed impacts. As we have argued throughout this review, it is logical that biodiversity conservation managers consider stress from the combined impacts of introduced predators, altered predator interactions, human activity, and stressors that occur naturally in ecosystems. Using the example methods to conduct our proposed CEA, based on the reactive scope model, to draft appropriate management plans, such as increasing habitat complexity or reducing human interaction in areas, would be a progressive response towards preserving many species of Australian small mammals and their constituent communities.

Conclusion

Considering ongoing global urbanization and the acknowledged importance of urban areas to biodiversity conservation, there is a great need for increased focus on the management of urban biodiversity. Management decisions require information about fear and stress impacts on wildlife, including impacts from both human activities and predators, especially if they are novel or introduced. Understanding the impacts of human activities is a research priority for modern science. There are many gaps in our current understanding of fear and stress impacts on wildlife, and of associated impacts of altered predation pressures and the persistence of target populations. To sustain biodiversity in urban and urban-adjacent green space habitats, reserves and conservation areas, it is vital that we establish better understanding and management of the multiple stressors that operate in these systems.

References

- Albert DM, Bowyer RT. 1991.** Factors related to grizzly bear: human interactions in Denali National Park. *Wildlife Society Bulletin* (1973-2006), **19**(3): 339-349.
- Alberti M. 2015.** Eco-evolutionary dynamics in an urbanizing planet. *Trends in Ecology & Evolution*, **30**(2): 114-126.
- Allan JD, McIntyre PB, Smith SD, Halpern BS, Boyer GL, Buchsbaum A, Burton Jr. GA, Campbell LM, Chadderton WL, Ciborowski JJH, Doran PJ, Eder T, Infante DM, Johnson LB, Joseph CA, Marino AL, Prusevich A, Read JG, Rose JB, Rutherford ES, Sowa SP, Steinman AD. 2013.** Joint analysis of stressors and ecosystem services to enhance restoration effectiveness. *Proceedings of the National Academy of Sciences USA*, **110**(1): 372-377.
- Andruskiw M, Fryxell JM, Thompson ID, Baker JA. 2008.** Habitat-mediated variation in predation risk by the American marten. *Ecology*, **89**(8): 2273-2280.
- Anson JR, Dickman CR. 2013.** Behavioral responses of native prey to disparate predators: naivité and predator recognition. *Oecologia*, **17**: 367–377. DOI: 10.1007/s00442-012-2424-7.
- Apfelbach R, Blanchard CD, Blanchard RJ, Hayes RA, McGregor IS. 2005.** The effects of predator odors in mammalian prey species: a review of field and laboratory studies. *Neuroscience & Biobehavioral Reviews*, **29**(8): 1123-1144.

- 708 **Apfelbach, R., Parsons MH, Soini HA, Novotny MV. 2015.** Are single odorous components of
709 a predator sufficient to elicit defensive behaviors in prey species?. *Frontiers in Neuroscience*, **9**:
710 263.
- 711 **Arlettaz R, Nusslé S, Baltic M, Vogel P, Palme R, Jenni-Eiermann S, Pathey P, Genoud M.**
712 **2015.** Disturbance of wildlife by outdoor winter recreation: allostatic stress response and altered
713 activity–energy budgets. *Ecological Applications*, **25**(5): 1197-1212.
- 714 **Atkins A, Redpath SM, Little RM, Amar A. 2017.** Experimentally manipulating the landscape
715 of fear to manage problem animals. *The Journal of Wildlife Management*, **81**(4): 610-616.
- 716 **Atwell JW, Cardoso GC, Whittaker DJ, Campbell-Nelson S, Robertson KW, Ketterson**
717 **ED. 2012.** Boldness behavior and stress physiology in a novel urban environment suggest rapid
718 correlated evolutionary adaptation. *Behavioral Ecology*, **23**(5): 960-969.
- 719 **Atuo FA, O'Connell TJ. 2017.** The landscape of fear as an emergent property of heterogeneity:
720 Contrasting patterns of predation risk in grassland ecosystems. *Ecology and Evolution*, **7**(13):
721 4782-4793.
- 722 **Ayal Y. 2007.** Trophic structure and the role of predation in shaping hot desert communities.
723 *Journal of Arid Environments*, **68**(2): 171-187.
- 724 **Banks PB, Bryant JV. 2007.** Four-legged friend or foe? Dog walking displaces native birds
725 from natural areas. *Biology Letters*, **3**: 611-613.
- 726 **Banks PB, Dickman CR. 2007.** Alien predation and the effects of multiple levels of prey
727 naiveté. *Trends in Ecology and Evolution*, **22**(5): 229-230.
- 728 **Banks PB, Smith HM. 2015.** The ecological impacts of commensal species: black rats, *Rattus*
729 *rattus*, at the urban-bushland interface. *Wildlife Research*, **42**(2): 86-97.
- 730 **Banks PB, Daly A, Bytheway JP. 2016.** Predator odours attract other predators, creating an
731 olfactory web of information. *Biology Letters*, **12**(5): 20151053.
- 732 **Bateman PW, Fleming PA. 2012.** Big city life: carnivores in urban environments. *Journal of*
733 *Zoology*, **287**(1): 1-23.
- 734 **Bazzaz FA. 1975.** Plant species diversity in old-field successional ecosystems in southern
735 Illinois. *Ecology*, **56**(2): 485-488.
- 736 **Becker DJ, Streicker DG, Altizer S. 2015.** Linking anthropogenic resources to wildlife–
737 pathogen dynamics: a review and meta-analysis. *Ecology Letters*, **18**(5): 483-495.
- 738 **Becker DJ, Snedden CE, Altizer S, Hall RJ. 2018.** Host dispersal responses to resource
739 supplementation determine pathogen spread in wildlife metapopulations. *The American*
740 *Naturalist*, **192**(4): 503-517.

- 741 **Bedoya-Perez MA, Carthey AJR, Mella VSA, McArthur C, Banks PB. 2013.** A practical
742 guide to avoid giving up on giving-up densities, *Behavioural Ecology and Sociobiology*, **67**(10):
743 1541-1553.
- 744 **Bedoya-Pérez MA, Smith KL, Kevin RC, Luo JL, Crowther MS, McGregor IS. 2019.**
745 Parameters that affect fear responses in rodents and how to use them for management. *Frontiers*
746 *in Ecology and Evolution*, **7**: 136. DOI: 10.3389/fevo.2019.00136.
- 747 **Belarde TA, Railsback SF. 2016.** New predictions from old theory: emergent effects of
748 multiple stressors in a model of piscivorous fish. *Ecological Modelling*, **326**: 54-62.
- 749 **Bernhardt ES, Palmer MA. 2007.** Restoring streams in an urbanizing world. *Freshwater*
750 *Biology*, **52**(4): 738-751.
- 751 **Berger J. 2007.** Fear, human shields and the redistribution of prey and predators in protected
752 areas. *Biology Letters*, **3**(6): 620-623.
- 753 **Bergstrom DM, Lucieer A, Kiefer K, Wasley J, Belbin L, Pedersen TK, Chown SL. 2009.**
754 Management implications of the Macquarie Island trophic cascade revisited: a reply to Dowding
755 et al. (2009). *Journal of Applied Ecology*, **46**(5): 1133-1136.
- 756 **Bianchi FJJ, Schellhorn NA, Van Der Werf W. 2009.** Foraging behaviour of predators in
757 heterogeneous landscapes: the role of perceptual ability and diet breadth. *Oikos*, **118**(9): 1363-
758 1372.
- 759 **Bilney RJ, Cooke R, White J. 2006.** Change in the diet of sooty owls (*Tyto tenebricosa*) since
760 European settlement: from terrestrial to arboreal prey and increased overlap with powerful
761 owls. *Wildlife Research*, **33**(1): 17-24.
- 762 **Bino G, Dolev A, Yosha D, Guter A, King R, Saltz D, Kark S. 2010.** Abrupt spatial and
763 numerical responses of overabundant foxes to reduction in anthropogenic resources. *Journal of*
764 *Applied Ecology*, **47**(6): 1262-1271.
- 765 **Bleicher SS. 2017.** The landscape of fear conceptual framework: definition and review of
766 current applications and misuses. *PeerJ*, **5**: e3772.
- 767 **Bleicher SS, Dickman CR. 2020.** On the landscape of fear: shelters affect foraging by dunnarts
768 (Marsupialia, *Sminthopsis* spp.) in a sandridge desert environment. *Journal of Mammalogy*,
769 **101**(1): 281-290.
- 770 **Bonier F. 2012.** Hormones in the city: endocrine ecology of urban birds. *Hormones and*
771 *Behaviour*, **61**(5): 763-772.
- 772 **Boonstra R, Hik D, Singleton GR, Tinnikov A. 1998.** The impact of predator-induced stress on
773 the snowshoe hare cycle. *Ecological Monographs*, **68**(3): 371-394.

- 774 **Bókony V, Lendvai AZ, Liker A, Angelier F, Wingfield JC, Chastel O. 2009.** Stress response
775 and the value of reproduction: are birds prudent parents? *The American Naturalist*, **173**(5): 589-
776 598.
- 777 **Bonier F. 2012.** Hormones in the city: endocrine ecology of urban birds. *Hormones and*
778 *Behavior*, **61**(5): 763-772.
- 779 **Bowyer RT, Van Ballenberghe V, Kie JG, Maier JA. 1999.** Birth-site selection by Alaskan
780 moose: maternal strategies for coping with a risky environment. *Journal of Mammalogy*, **80**(4):
781 1070-1083.
- 782 **Brearley G, Rhodes J, Bradley A, Baxter G, Seabrook L, Lunney D, ... McAlpine C. 2013.**
783 Wildlife disease prevalence in human-modified landscapes. *Biological Reviews*, **88**(2): 427-442.
- 784 **Brown JS. 1988.** Patch use as an indicator of habitat preference, predation risk, and competition.
785 *Behavioral Ecology and Sociobiology*, **22**(1): 37-47.
- 786 **Brown JS. 2000.** Foraging ecology of animals in response to heterogeneous environments. *The*
787 *ecological consequences of environmental heterogeneity*. Blackwell, Oxford, UK. 181-214.
- 788 **Brown BL. 2003.** Spatial heterogeneity reduces temporal variability in stream insect
789 communities. *Ecology Letters*, **6**(4): 316-325.
- 790 **Brown JS, Kotler BP. 2004.** Hazardous duty pay and the foraging cost of predation. *Ecology*
791 *Letters*, **7**(10): 999-1014.
- 792 **Cannon WB. 1932.** *The wisdom of the body*, W. W. Norton and Company, New York, 177-201.
- 793 **Canteras NS, Pavesi E, Carobrez AP. 2015.** Olfactory instruction for fear: neural system
794 analysis. *Frontiers in Neuroscience*, **9**: 276 (<https://doi.org/10.3389/fnins.2015.00276>).
- 795 **Cardillo M, Mace GM, Jones KE, Bielby J, Bininda-Emonds OR, Sechrest W, Orme CDL,**
796 **Purvis A. 2005.** Multiple causes of high extinction risk in large mammal
797 species. *Science*, **309**(5738): 1239-1241.
- 798 **Cardillo M, Mace GM, Gittleman, JL, Purvis A. 2006.** Latent extinction risk and the future
799 battlegrounds of mammal conservation. *Proceedings of the National Academy of Sciences of the*
800 *United States of America*, **103**(11): 4157-4161.
- 801 **Carlstead K, Brown JL, Seidensticker J. 1993.** Behavioral and adrenocortical responses to
802 environmental changes in leopard cats (*Felis bengalensis*). *Zoo Biology*, **12**(4): 321-331.
- 803 **Carlstead K, Brown JL, Strawn W. 1993.** Behavioral and physiological correlates of stress in
804 laboratory cats. *Applied Animal Behaviour Science*, **38**(2): 143-158.
- 805 **Carroll SP. 2011.** Conciliation biology: the eco-evolutionary management of permanently
806 invaded biotic systems. *Evolutionary Applications*, **4**(2): 184-199.

- 807 **Carthey AJ, Banks PB. 2012.** When does an alien become a native species? A vulnerable native
808 mammal recognises and responds to its long-term alien predator. *PLoS ONE*, **7**(2): e31804.
- 809 **Carthey AJR, Banks PB. 2016.** Naiveté is not forever: responses of a vulnerable native rodent
810 to its long term alien predators, *Oikos*, **125**(7): 918-926.
- 811 **Carthey JR, Blumstein DT. 2018.** Predicting predator recognition in a changing world. *Trends*
812 *in Ecology and Evolution*, **33**(2): 106-115.
- 813 **Červinka J, Šálek M, Padyšáková E, Šmilauer P. 2013.** The effects of local and landscape-
814 scale habitat characteristics and prey availability on corridor use by carnivores: a comparison of
815 two contrasting farmlands. *Journal for Nature Conservation*, **21**(2): 105-113.
- 816 **Chisholm R, Taylor R. 2007.** Null-hypothesis significance testing and the Critical Weight
817 Range for Australian mammals. *Conservation Biology*, **21**(6): 1641-1645.
- 818 **Ciuti S, Northrup JM, Muhly TB, Simi S, Musiani M, Pitt JA, Boyce MS. 2012.** Effects of
819 humans on behaviour of wildlife exceed those of natural predators in a landscape of fear, *PLoS*
820 *ONE*, **7**(11): e50611. <https://doi.org/10.1371/journal.pone.0050611>.
- 821 **Civitello DJ, Allman BE, Morozumi C, Rohr JR. 2018.** Assessing the direct and indirect
822 effects of food provisioning and nutrient enrichment on wildlife infectious disease
823 dynamics. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **373**(1745):
824 p.20170101.
- 825 **Clinchy M, Zanette L, Boonstra R, Wingfield JC, Smith JN. 2004.** Balancing food and
826 predator pressure induces chronic stress in songbirds. *Proceedings of the Royal Society of*
827 *London B: Biological Sciences*, **271**(1556): 2473-2479.
- 828 **Clinchy M, Zanette L, Charlier TD, Newman AE, Schmidt KL, Boonstra R, Soma KK.**
829 **2011.** Multiple measures elucidate glucocorticoid responses to environmental variation in
830 predation threat. *Oecologia*, **166**(3): 607-614.
- 831 **Clinchy M, Zanette LY, Roberts D, Suraci JP, Buesching CD, Newman C, Macdonald DW.**
832 **2016.** Fear of the human “super predator” far exceeds the fear of large carnivores in a model
833 mesocarnivore. *Behavioral Ecology*, **27**(6): 1826-1832.
- 834 **Cook NJ. 2012.** Minimally invasive sampling media and the measurement of corticosteroids as
835 biomarkers of stress in animals. *Canadian Journal of Animal Science*, **92**(3): 227-259.
- 836 **Cooke SJ, Sack L, Franklin CE, Farrell AP, Beardall J, Wikelski M, Chown SL. 2013.**
837 **What is conservation physiology? Perspectives on an increasingly integrated and essential**
838 **science.** *Conservation Physiology*, **1**(1): cot001.
- 839 **Côté IM, Darling ES, Brown CJ. 2016.** Interactions among ecosystem stressors and their
840 importance in conservation. *Proceedings of the Royal Society B: Biological Sciences*, **283**(1824):
841 20152592.

- 842 **Crain CM, Kroeker K, Halpern BS. 2008.** Interactive and cumulative effects of multiple
843 human stressors in marine systems. *Ecology Letters*, **11**(12): 1304-1315.
- 844 **Cremona T, Crowther MS, Webb JK. 2014.** Variation of prey responses to cues from a
845 mesopredator and an apex predator. *Austral Ecology*, **39**(7): 749-754.
- 846 **Crocker DE, Khudyakov JI, Champagne CD. 2016.** Oxidative stress in northern elephant
847 seals: Integration of omics approaches with ecological and experimental studies. *Comparative*
848 *Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, **200**: 94-103.
- 849 **Crooks KR, Soulé ME. 1999.** Mesopredator release and avifaunal extinctions in a fragmented
850 system. *Nature*, **400**(6744): 563.
- 851 **Dales JT. 2011.** Death by a thousand cuts: incorporating cumulative effects in Australia's
852 Environment Protection and Biodiversity Conservation Act. *Pacific Rim Law & Policy Journal*,
853 **20**: 149 - 178.
- 854 **Dantzer B, Fletcher QE, Boonstra R, Sheriff MJ. 2014.** Measures of physiological stress: a
855 transparent or opaque window into the status, management and conservation of species?,
856 *Conservation Physiology*, **2**(1): cou023.
- 857 **Darimont CT, Fox CH, Bryan HM, Reimchen TE. 2015.** The unique ecology of human
858 predators. *Science*, **349**(6250): 858-860.
- 859 **Davis M. 1992.** The role of the amygdala in fear and anxiety. *Annual Review of*
860 *Neuroscience*, **15**(1): 353-375.
- 861 **Davis M, Whalen PJ. 2001.** The amygdala: vigilance and emotion. *Molecular Psychiatry*, **6**(1):
862 13-34.
- 863 **Dhabhar FS, McEwen BS. 1999.** Enhancing versus suppressive effects of stress hormones on
864 skin immune function. *Proceedings of the National Academy of Sciences USA*, **96**(3): 1059-
865 1064.
- 866 **Dickens MJ, Romero ML. 2013.** A consensus endocrine profile for chronically stressed wild
867 animals does not exist. *General and Comparative Endocrinology*, **191**: 177-189.
- 868 **Dickman CR. 1996.** Impact of exotic generalist predators on the native fauna of Australia.
869 *Wildlife Biology*, **2**(3): 185-195.
- 870 **Ditchkoff SS, Saalfeld ST, Gibson CJ. 2006.** Animal behavior in urban ecosystems:
871 modifications due to human-induced stress. *Urban Ecosystems*, **9**(1): 5-12.
- 872 **Doherty TS, Dickman CR, Nimmo DG, Ritchie EG. 2015.** Multiple threats, or multiplying the
873 threats? Interactions between invasive predators and other ecological disturbances. *Biological*
874 *Conservation*, **190**: 60-68.

875 **Doherty TS, Glen AS, Nimmo DG, Ritchie EG, Dickman CR. 2016.** Invasive predators and
876 global biodiversity loss. *Proceedings of the National Academy of Sciences USA*, **113**(40): 11261-
877 11265.

878 **Doherty TS, Dickman CR, Glen AS, Newsome TM, Nimmo DG, Ritchie EG, Vanak AT,**
879 **Wirsing AJ. 2017.** The global impacts of domestic dogs on threatened vertebrates. *Biological*
880 *Conservation*, **210**: 56-59.

881 **Donihue CM, Lambert MR. 2015.** Adaptive evolution in urban ecosystems. *Ambio*, **44**(3): 194-
882 203.

883 **Duinker PN, Burbidge EL, Boardley SR, Greig LA. 2012.** Scientific dimensions of
884 cumulative effects assessment: toward improvements in guidance for practice. *Environmental*
885 *Reviews*, **21**(1): 40-52.

886 **Estes JA, Terborgh J, Brashares JS, Power ME, Berger J, Bond WJ, Carpenter SR,**
887 **Essington TE, Holt RD, Jackson JBC, Marquis RJ, Oksanen L, Oksanen T, Paine RT,**
888 **Pikitch EK, Ripple WJ, Sandin SA, Scheffer M, Schoener TW, Shurin JB, Sinclair ARE,**
889 **Soulé ME, Virtanen R, Wardle DA. 2011.** Trophic downgrading of planet Earth. *Science*,
890 **333**(6040): 301-306.

891 **Epstein S. 1972.** The nature of anxiety with emphasis upon its relationship to
892 expectancy. *Anxiety; Current Trends in Theory and Research*, 291-337.

893 **Fanselow MS, Poulos AM. 2005.** The neuroscience of mammalian associative learning. *Annual*
894 *Review of Psychology*, **56**: 207-234.

895 **Feng SF, Shi TY, Wang WN, Chen YC, Tan QR. 2012.** Long-lasting effects of chronic rTMS
896 to treat chronic rodent model of depression. *Behavioural Brain Research*, **232**(1): 245-251.

897 **Fischer JD, Cleeton SH, Lyons TP, Miller JR. 2012.** Urbanization and the predation paradox:
898 the role of trophic dynamics in structuring vertebrate communities. *Bioscience*, **62**(9): 809-818.

899 **Fleming PA, Bateman PW. 2018.** Novel predation opportunities in anthropogenic landscapes.
900 *Animal Behaviour*, **138**: 145-155.

901 **Fogaca MV, Lisboa SF, Aguiar DC, Moreira FA, Gomes FV, Casarotto PC, Guimaraes FS.**
902 **2012.** Fine-tuning of defensive behaviors in the dorsal periaqueductal gray by atypical
903 neurotransmitters. *Brazilian Journal of Medical and Biological Research*, **45**: 357-365.

904 **Foley MM, Mease LA, Martone RG, Prahlér EE, Morrison TH, Murray CC, Wojcik D.**
905 **2017.** The challenges and opportunities in cumulative effects assessment. *Environmental Impact*
906 *Assessment Review*, **62**: 122-134.

907 **Freeman NE, Norris DR, Sutton AO, Newman AE. 2020.** Raising young with limited
908 resources: supplementation improves body condition and advances fledging of Canada
909 jays. *Ecology*, **101**(1): p.e02909.

- 910 **Frid A, Dill L. 2002.** Human-caused disturbance stimuli as a form of predation
911 risk. *Conservation Ecology*, **6**(1):11.
- 912 **Gaston KJ, Warren PH, Thompson K, Smith RM. 2005.** Urban domestic gardens (IV): the
913 extent of the resource and its associated features. *Biodiversity and Conservation*, **14**(14): 3327-
914 3349.
- 915 **Gaynor KM, Hojnowski CE, Carter NH, Brashares JS. 2018.** The influence of human
916 disturbance on wildlife nocturnality. *Science*, **360**:1232–1235. DOI: 10.1126/science.aar7121.
- 917 **Gaynor KM, Brown JS, Middleton AD, Power ME, Brashares, JS. 2019.** Landscapes of fear:
918 spatial patterns of risk perception and response. *Trends in Ecology & Evolution*, **34**(4): 355-368.
- 919 **Geary WL, Nimmo DG, Doherty TS, Ritchie EG, Tulloch AI. 2019.** Threat webs: reframing
920 the co-occurrence and interactions of threats to biodiversity. *Journal of Applied Ecology*, **56**:
921 1992-1997.
- 922 **Gil MA, Zill J, Ponciano JM. 2017.** Context-dependent landscape of fear: algal density elicits
923 risky herbivory in a coral reef. *Ecology*, **98**(2): 534-544.
- 924 **Gill JA, Sutherland WJ, Watkinson AR. 1996.** A method to quantify the effects of human
925 disturbance on animal populations. *Journal of Applied Ecology*, **33**(4): 786-792.
- 926 **Glen AS, Dickman CR. 2005.** Complex interactions among mammalian carnivores in Australia,
927 and their implications for wildlife management. *Biological Reviews*, **80**(3): 387-401.
- 928 **Graham CA, Maron M, McAlpine CA. 2013.** Influence of landscape structure on invasive
929 predators: feral cats and red foxes in the brigalow landscapes, Queensland, Australia. *Wildlife*
930 *Research*, **39**(8): 661-676.
- 931 **Griffiths SR, Lentini PE, Semmens K, Watson SJ, Lumsden LF, Robert KA. 2018.**
932 Chainsaw-carved cavities better mimic the thermal properties of natural tree hollows than nest
933 boxes and log hollows. *Forests*, **9**(5): 235.
- 934 **Guiden PW, Bartel SL, Byer NW, Shipley AA, Orrock JL. 2019.** Predator-prey interactions
935 in the Anthropocene: reconciling multiple aspects of novelty. *Trends in Ecology and Evolution*,
936 **34**(7):616-627.
- 937 **Gunderson AR, Armstrong EJ, Stillman JH. 2016.** Multiple stressors in a changing world: the
938 need for an improved perspective on physiological responses to the dynamic marine
939 environment. *Annual Review of Marine Science*, **8**: 357-378.
- 940 **Hammerschlag N, Broderick AC, Coker JW, Coyne MS, Dodd M, Frick MG, Godfrey MH,**
941 **Godley BJ, Griffin DB, Hartog K, Murphy SR. 2015.** Evaluating the landscape of fear
942 between apex predatory sharks and mobile sea turtles across a large dynamic seascape. *Ecology*,
943 **96**(8): 2117-2126.

- 944 **Hawlena D, Schmitz OJ. 2010.** Physiological stress as a fundamental mechanism linking
945 predation to ecosystem functioning. *The American Naturalist*, **176**:537–556. DOI:
946 10.1086/656495.
- 947 **Hawlena D, Strickland MS, Bradford MA, Schmitz OJ. 2012.** Fear of predation slows litter
948 decomposition. *Science*, **336**:1434–1438. DOI 10.1126/science.1220097.
- 949 **Hayes RA, Nahrung HF, Wilson JC. 2006.** The response of native Australian rodents to
950 predator odours varies seasonally: a by-product of life history variation? *Animal Behaviour*,
951 **71**(6): 1307–1314.
- 952 **Hill NJ, Carbery KA, Deane EM. 2007.** Human-possum conflict in urban Sydney, Australia:
953 public perceptions and implications for species management. *Human Dimensions of Wildlife*,
954 **12**(2): 101-113.
- 955 **Hing S, Narayan E, Thompson RA, Godfrey S. 2014.** A review of factors influencing the
956 stress response in Australian marsupials. *Conservation Physiology*, **2**(1): cou027.
- 957 **Hobbie SE, Finlay JC, Janke BD, Nidzgorski DA, Millet DB, Baker LA. 2017.** Contrasting
958 nitrogen and phosphorus budgets in urban watersheds and implications for managing urban water
959 pollution. *Proceedings of the National Academy of Sciences USA*, **114**(16): 4177-4182.
- 960 **Hobbs RJ, Higgs E, Harris J A. 2009.** Novel ecosystems: implications for conservation and
961 restoration. *Trends in Ecology and Evolution*, **24**(11): 599-605.
- 962 **Hofer H, and East ML. 1998.** Biological conservation and stress. *Advances in the Study of*
963 *Behavior*, **27**: 405-525.
- 964 **Hoffman CR, Sitvarin MI, and Rypstra AL. 2016.** Information from familiar and related
965 conspecifics affects foraging in a solitary wolf spider. *Oecologia*, **181**(2): 359-367.
- 966 **Holt RD. 1984.** Spatial heterogeneity, indirect interactions, and the coexistence of prey
967 species. *The American Naturalist*, **124**(3): 377-406.
- 968 **Honda T, Iijima H, Tsuboi J, Uchida K. 2018.** A review of urban wildlife management from
969 the animal personality perspective: The case of urban deer. *Science of the Total Environment*,
970 **644**: 576-582.
- 971 **Hopcraft JGC, Sinclair ARE, Packer C. 2005.** Planning for success: Serengeti lions seek prey
972 accessibility rather than abundance. *Journal of Animal Ecology*, **74**(3): 559-566.
- 973 **Iossa G, Soulsbury CD, Baker PJ, Harris S. 2010.** A taxonomic analysis of urban carnivore
974 ecology. In: Gehrt SD, Riley SPD, Cypher BL (eds) *Urban Carnivores: Ecology, Conflict, and*
975 *Conservation*: 173–180. Johns Hopkins University Press, Baltimore, Maryland, USA.
- 976 **Iribarren C, Kotler BP. 2012.** Foraging patterns of habitat use reveal landscape of fear of
977 Nubian ibex *Capra nubiana*. *Wildlife Biology*, **18**(2): 194-202.

- 978 **Ives CD, Lentini PE, Threlfall CG, Ikin K, Shanahan DF, Garrard GE, Bekessy SA, Fuller**
979 **RA, Mumaw L, Rayner L, Rowe R, Valentine LE, Kendal D. 2016.** Cities are hotspots for
980 threatened species. *Global Ecology and Biogeography*, **25**(1): 117-126.
- 981 **Jaatinen K, Seltmann MW, Öst M. 2014.** Context-dependent stress responses and their
982 connections to fitness in a landscape of fear. *Journal of Zoology*, **294**(3): 147-153.
- 983 **Jackson MC, Loewen CJG, Vinebrooke RD, Chimimba CT. 2016.** Net effects of multiple
984 stressors in freshwater ecosystems: a meta-analysis. *Global Change Biology*, **22**: 180–189.
- 985 **Johnson MD, Sherry TW. 2001.** Effects of food availability on the distribution of migratory
986 warblers among habitats in Jamaica. *Journal of Animal Ecology*, **70**(4): 546-560.
- 987 **Johnson C. 2006.** *Australia's mammal extinctions: a 50,000-year history*. Cambridge University
988 Press.
- 989 **Johnstone CP, Lill A, Reina RD. 2012.** Does habitat fragmentation cause stress in the agile
990 antechinus? A haematological approach. *Journal of Comparative Physiology B*, **182**(1): 139-155.
- 991 **Jokimäki J, Kaisanlahti-Jokimäki ML, Suhonen J, Clergeau P, Pautasso M, Fernández-**
992 **Juricic E. 2011.** Merging wildlife community ecology with animal behavioral ecology for a
993 better urban landscape planning. *Landscape and Urban Planning*, **100**(4): 383-385.
- 994 **Jones IC, Vinson GP, Jarrett IG, Sharman GB. 1964.** Steroid components in the adrenal
995 venous blood of *Trichosurus vulpecula* (Kerr). *Journal of Endocrinology*, **30**(1): 149-150.
- 996 **Jones ME, Apfelbach R, Banks PB, Cameron EZ, Dickman CR, Frank A, McLean S,**
997 **McGregor IS, Müller-Schwarze D, Parsons MH, Sparrow EA, Blumstein DT. 2016.** A nose
998 for death: integrating trophic and informational networks for conservation and management.
999 *Frontiers in Ecology and Evolution*, **4**: 124. DOI: 10.3389/fevo.2016.00124.
- 1000 **Kauffman MJ, Brodie JF, Jules ES. 2010.** Are wolves saving Yellowstone's aspen? A
1001 landscape-level test of a behaviorally mediated trophic cascade. *Ecology*, **91**(9): 2742-2755.
- 1002 **King JM, Bradshaw SD. 2010.** Stress in an island kangaroo? The Barrow Island euro,
1003 *Macropus robustus isabellinus*. *General and Comparative Endocrinology*, **167**(1): 60-67.
- 1004 **Klecka J, Boukal DS. 2014.** The effect of habitat structure on prey mortality depends on
1005 predator and prey microhabitat use. *Oecologia*, **176**(1): 183-191.
- 1006 **Kloppers EL, St. Clair CC, Hurd TE. 2005.** Predator-resembling aversive conditioning for
1007 managing habituated wildlife. *Ecology and Society*, **10**(1): 31.
- 1008 **Knutie SA. 2020.** Food supplementation affects gut microbiota and immunological resistance to
1009 parasites in a wild bird species. *bioRxiv*, DOI: 10.1101/1365-2664.13567.

- 1010 **Kohl MT, Stahler DR, Metz MC, Forester JD, Kauffman MJ, Varley N, White PJ, Smith**
1011 **DW, MacNulty DR. 2018.** Diel predator activity drives a dynamic landscape of fear. *Ecological*
1012 *Monographs*, **88**:638–652. DOI: 10.1002/ecm.1313.
- 1013 **Kotler BP, Brown JS. 1988.** Environmental heterogeneity and the coexistence of desert
1014 rodents. *Annual Review of Ecology and Systematics*, **19**(1): 281-307.
- 1015 **Kuijper DP, Bubnicki JW, Churski M, Mols B, Van Hooft P. 2015.** Context dependence of
1016 risk effects: wolves and tree logs create patches of fear in an old-growth forest. *Behavioral*
1017 *Ecology*, **26**(6): 1558-1568.
- 1018 **Labar KS, Ledoux JE. 2011.** Coping with danger: the neural basis of defensive behavior and
1019 fearful feelings. *Comprehensive Physiology*, **8**: 139 – 154.
- 1020 **Latham ADM, Latham MC, Boyce MS, Boutin S. 2011.** Movement responses by wolves to
1021 industrial linear features and their effect on woodland caribou in northeastern Alberta.
1022 *Ecological Applications*, **21**(8): 2854-2865.
- 1023 **Laundré JW, Hernández L, Altendorf KB. 2001.** Wolves, elk, and bison: reestablishing the"
1024 landscape of fear" in Yellowstone National Park, USA. *Canadian Journal of Zoology*, **79**(8):
1025 1401-1409.
- 1026 **Laundré JW, Hernández L, Ripple WJ. 2010.** The landscape of fear: ecological implications
1027 of being afraid. *Open Ecology Journal*, **3**:1–7.
- 1028 **Laundré JW, Hernández L, Medina PL, Campanella A, López-Portillo J, González-**
1029 **Romero A, Grajales-Tam KM, Burke AM, Gronemeyer P, Browning DM. 2014.** The
1030 landscape of fear: the missing link to understand top-down and bottom-up controls of prey
1031 abundance? *Ecology*, **95**(5): 1141-1152.
- 1032 **Laws AN. 2017.** Climate change effects on predator-prey interactions. *Current Opinion in Insect*
1033 *Science*, **23**: 28-34.
- 1034 **Leahy L, Legge SM, Tuft K, McGregor HW, Barmuta LA, Jones ME, Johnson CN. 2016.**
1035 Amplified predation after fire suppresses rodent populations in Australia's tropical
1036 savannas. *Wildlife Research*, **42**(8): 705-716.
- 1037 **LeDoux JE. 1996.** *The Emotional Brain*, Simon and Schuster, New York.
- 1038 **LeDoux JE. 2003.** The emotional brain, fear, and the amygdala, *Cell and Molecular*
1039 *Neurobiology*, **23**: 727-738.
- 1040 **Legge S, Smith JG, James A, Tuft KD, Webb T, Woinarski JC. 2019.** Interactions among
1041 threats affect conservation management outcomes: Livestock grazing removes the benefits of fire
1042 management for small mammals in Australian tropical savannas. *Conservation Science and*
1043 *Practice*, e52.

- 1044 **Leighton PA, Horrocks JA, Kramer DL. 2010.** Conservation and the scarecrow effect: Can
1045 human activity benefit threatened species by displacing predators? *Biological Conservation*,
1046 **143**(9): 2156-2163.
- 1047 **Leo V, Reading RP, Letnic M. 2015.** Interference competition: odours of an apex predator and
1048 conspecifics influence resource acquisition by red foxes. *Oecologia*, **179**(4): 1033-1040.
- 1049 **Lima SL, Dill LM. 1990.** Behavioral decisions made under the risk of predation: a review and
1050 prospectus. *Canadian Journal of Zoology*, **68**(4): 619-640.
- 1051 **Lima SL. 1998.** Stress and decision making under the risk of predation: recent developments
1052 from behavioral, reproductive, and ecological perspectives. *Advances in the Study of*
1053 *Behavior*, **27**: 215-290.
- 1054 **Lima SL, Bednekoff PA. 1999.** Temporal variation in danger drives antipredator behavior: the
1055 predation risk allocation hypothesis. *The American Naturalist*, **153**(6): 649-659.
- 1056 **Lone K, Loe LE, Gobakken T, Linnell JD, Odden J, Remmen J, Mysterud A. 2014.** Living
1057 and dying in a multi-predator landscape of fear: roe deer are squeezed by contrasting pattern of
1058 predation risk imposed by lynx and humans. *Oikos*, **123**(6): 641-651.
- 1059 **Lowry H, Lill A, Wong BBM. 2012.** Behavioural responses of wildlife to urban environments,
1060 *Biological Reviews*, **88**: 537-549.
- 1061 **Lyons J, Mastro Monaco G, Edwards DB, Schulte-Hostedde AI. 2017.** Fat and happy in the
1062 city: Eastern chipmunks in urban environments. *Behavioural Ecology*, **28**(6): 1464-1471.
- 1063 **Ma Z, Becker DR, Kilgore MA. 2009.** Assessing cumulative impacts within state
1064 environmental review frameworks in the United States. *Environmental Impact Assessment*
1065 *Review*, **29**(6): 390-398.
- 1066 **MacDougall-Shackleton SA, Bonier F, Romero LM, Moore IT. 2019.** Glucocorticoids and
1067 “stress” are not synonymous. *Integrative Organismal Biology*, **1**(1): obz017, [https://doi-](https://doi-org.ezproxy1.library.usyd.edu.au/10.1093/iob/obz017)
1068 [org.ezproxy1.library.usyd.edu.au/10.1093/iob/obz017](https://doi-org.ezproxy1.library.usyd.edu.au/10.1093/iob/obz017)
- 1069 **Madsen B, Carroll N, Moore Brands K. 2010.** *State of Biodiversity Markets Report: Offset and*
1070 *Compensation Programs Worldwide*, available at:
1071 <http://www.ecosystemmarketplace.com/documents/acrobat/sbdlmr.pdf>
- 1072 **Malcolm KD, McShea WJ, Garshelis DL, Luo SJ, Van Deelen TR, Liu F, Li S, Miao L,**
1073 **Wang D, Brown JL. 2014.** Increased stress in Asiatic black bears relates to food limitation, crop
1074 raiding, and foraging beyond nature reserve boundaries in China. *Global Ecology and*
1075 *Conservation*, **2**: 267-276.
- 1076 **Manuck SB, Kaplan JR, Adams MR, Clarkson TB. 1988.** Studies of psychosocial influences
1077 on coronary artery atherogenesis in cynomolgus monkeys. *Health Psychology*, **7**(2): 113.

- 1078 **Maren S. 2001.** Neurobiology of Pavlovian fear conditioning. *Annual Review of Neuroscience*,
1079 **24**(1): 897-931.
- 1080 **Mason JW. 1971.** A re-evaluation of the concept of ‘non-specificity’ in stress theory. *Journal of*
1081 *Psychiatric Research*, **8**(3-4): 323-333.
- 1082 **McCauley LA, Jenkins DG, Quintana-Ascencio PF. 2013.** Isolated wetland loss and
1083 degradation over two decades in an increasingly urbanised landscape. *Wetlands*, **33**(1): 117-127.
- 1084 **McClennen NA, Wigglesworth RR, Anderson SH, Wachob DG. 2001.** The effect of suburban
1085 and agricultural development on the activity patterns of coyotes (*Canis latrans*). *The American*
1086 *Midland Naturalist*, **146**(1): 27-37.
- 1087 **McComb LB, Lentini PE, Harley DK, Lumsden LF, Antrobus JS, Eyre AC, Briscoe NJ.**
1088 **2019.** Feral cat predation on Leadbeater’s possum (*Gymnobelideus leadbeateri*) and observations
1089 of arboreal hunting at nest boxes. *Australian Mammalogy*, **41**(2): 262-265.
- 1090 **McDonnell MJ, Hahs AK. 2015.** Adaptation and adaptedness of organisms to urban
1091 environments. *Annual Review of Ecology, Evolution, and Systematics*, **46**: 261-280.
- 1092 **McEwen BS, Stellar E. 1993.** Stress and the individual: mechanisms leading to
1093 disease. *Archives of Internal Medicine*, **153**(18): 2093-2101.
- 1094 **McEwen BS, Wingfield JC. 2003.** The concept of allostasis in biology and
1095 biomedicine. *Hormones and Behavior*, **43**(1): 2-15.
- 1096 **McEwen BS, 2004.** Protection and damage from acute and chronic stress: allostasis and
1097 allostatic overload and relevance to the pathophysiology of psychiatric disorders. *Annals of the*
1098 *New York Academy of Sciences*, **1032**(1): 1-7.
- 1099 **McKinney ML. 2006.** Urbanization as a major cause of biotic homogenization. *Biological*
1100 *Conservation*, **127**(3): 247-260.
- 1101 **McMahon JD, Lashley MA, Brooks CP, Barton BT. 2018.** Covariance between predation risk
1102 and nutritional preferences confounds interpretations of giving-up density
1103 experiments. *Ecology*, **99**(7), 1517-1522.
- 1104 **Mella VSA, Banks PB, McArthur C. 2014.** Negotiating multiple cues of predation risk in a
1105 landscape of fear: what scares free-ranging brushtail possums? *Journal of Zoology*, **294**(1): 22-
1106 30.
- 1107 **Mineka S. 1979.** The role of fear in theories of avoidance learning, flooding, and extinction.
1108 *Psychological Bulletin*, **86**(5): 985.
- 1109 **Mineur YS, Belzung C, Crusio WE. 2006.** Effects of unpredictable chronic mild stress on
1110 anxiety and depression-like behavior in mice. *Behavioural Brain Research*, **175**(1): 43-50.

- 1111 **Møller AP. 2009.** Successful city dwellers: a comparative study of the ecological characteristics
1112 of urban birds in the Western Palearctic. *Oecologia*, **159**(4): 849-858.
- 1113 **Morgan SA, Hansen CM, Ross JG, Hickling GJ, Ogilvie SC, Paterson AM. 2009.** Urban cat
1114 (*Felis catus*) movement and predation activity associated with a wetland reserve in New Zealand.
1115 *Wildlife Research*, **36**(7): 574-580.
- 1116 **Murray MH, Becker DJ, Hall RJ, Hernandez SM. 2016.** Wildlife health and supplemental
1117 feeding: A review and management recommendations. *Biological Conservation*, **204**: 163-174.
- 1118 **Navara KJ, Nelson RJ. 2007.** The dark side of light at night: physiological, epidemiological,
1119 and ecological consequences. *Journal of Pineal Research*, **43**(3): 215-224.
- 1120 **Newsome TM, Ballard GA, Fleming PJ, van de Ven R, Story GL, Dickman CR. 2014.**
1121 Human-resource subsidies alter the dietary preferences of a mammalian top
1122 predator. *Oecologia*, **175**(1): 139-150.
- 1123 **Newsome TM, Dellinger JA, Pavey CR, Ripple WJ, Shores CR, Wirsing AJ, Dickman CR.**
1124 **2015.** The ecological effects of providing resource subsidies to predators. *Global Ecology and*
1125 *Biogeography*, **24**(1): 1-11.
- 1126 **Newsome T, van Eeden L, Lazenby B, Dickman C. 2017.** Does culling work? *Australasian*
1127 *Science*, **38**(1): 28-30.
- 1128 **Nersesian CL, Banks PB, McArthur C. 2012.** Behavioural responses to indirect and direct
1129 predator cues by a mammalian herbivore, the common brushtail possum. *Behavioral Ecology*
1130 *and Sociobiology*, **66**(1): 47-55.
- 1131 **Norton DA. 2009.** Species invasions and the limits to restoration: learning from the New
1132 Zealand experience. *Science*, **325**(5940): 569-571.
- 1133 **Öhman A. 2000.** In G. Fink (Ed.), *Encyclopedia of stress* (Vol. 2), Academic Press.
- 1134 **Otto SP. 2018.** Adaptation, speciation and extinction in the Anthropocene. *Proceedings of the*
1135 *Royal Society B*, **285**(1891): 20182047.
- 1136 **Palme R. 2019.** Non-invasive measurement of glucocorticoids: Advances and problems.
1137 *Physiology and Behaviour*, **199**: 229-243.
- 1138 **Palmer MA, Menninger HL, Bernhardt E. 2010.** River restoration, habitat heterogeneity and
1139 biodiversity: a failure of theory or practice?. *Freshwater Biology*, **55**: 205-222.
- 1140 **Parsons MH, Blumstein DT. 2010.** Familiarity breeds contempt: kangaroos persistently avoid
1141 areas with experimentally deployed dingo scents. *PLoS ONE*, **5**(5): 5e10403.
- 1142 **Parsons MH, Apfelbach R, Banks PB, Cameron EZ, Dickman CR, Frank AS, Jones ME,**
1143 **McGregor IS, McLean S, Müller-Schwarze D, Sparrow EE, Blumstein DT. 2017.**

- 1144 Biologically meaningful scents: a framework for understanding predator–prey research across
1145 disciplines. *Biological Reviews*. **93**(1): 98-114. doi: 10.1111/brv.12334.
- 1146 **Parris KM. 2016.** *Ecology of urban environments*. John Wiley and Sons.
- 1147 **Patten MA, Burger JC. 2018.** Reserves as double-edged sword: Avoidance behavior in an
1148 urban-adjacent wildland. *Biological Conservation*, **218**: 233-239.
- 1149 **Pianka ER. 2011.** *Evolutionary ecology*. Eric R. Pianka.
- 1150 **Pickett ST, Cadenasso ML, Grove JM, Nilon CH, Pouyat RV, Zipperer WC, Costanza R.**
1151 **2001.** Urban ecological systems: linking terrestrial ecological, physical, and socioeconomic
1152 components of metropolitan areas. *Annual Review of Ecology and Systematics*, **32**(1): 127-157.
- 1153 **Pitkanen A. 2000.** Connectivity of the rat amygdaloid complex. *The amygdala: A functional*
1154 *analysis*. Oxford University Press, Oxford: 31-115.
- 1155 **Polis GA, Barnes JD, Seely MK, Henschel JR, Enders MM. 1998.** Predation as a major cost
1156 of reproduction in Namib Desert tenebrionid beetles. *Ecology*, **79**(7): 2560-2566.
- 1157 **Price CJ, Banks PB. 2018.** Habitat augmentation for introduced urban wildlife: the use of piles
1158 of railway sleepers as refuge for introduced black rats *Rattus rattus*. *Australian Zoologist*, **39**(3):
1159 513-519.
- 1160 **Prugh LR, Stoner CJ, Epps CW, Bean WT, Ripple WJ, Laliberte AS, Brashares JS. 2009.**
1161 The rise of the mesopredator. *Bioscience*, **59**(9): 779-791.
- 1162 **Rayner MJ, Hauber ME, Imber MJ, Stamp RK, Clout MN. 2007.** Spatial heterogeneity of
1163 mesopredator release within an oceanic island system. *Proceedings of the National Academy of*
1164 *Sciences USA*, **104**(52): 20862-20865.
- 1165 **Rehnus M, Wehrle M, Palme R. 2014.** Mountain hares *Lepus timidus* and tourism: stress
1166 events and reactions. *Journal of Applied Ecology*, **51**(1): 6-12.
- 1167 **Rescorla RA, Solomon RL. 1967.** Two-process learning theory: Relationships between
1168 Pavlovian conditioning and instrumental learning. *Psychological Review*, **74**(3): 151.
- 1169 **Rieucan G, Vickery WL, Doucet GJ. 2009.** A patch use model to separate effects of foraging
1170 costs on giving-up densities: an experiment with white-tailed deer (*Odocoileus virginianus*).
1171 *Behavioral Ecology and Sociobiology*, **63**(6): 891-897.
- 1172 **Riley SP, Sauvajot RM, Fuller TK, York EC, Kamradt DA, Bromley C, Wayne RK. 2003.**
1173 Effects of urbanization and habitat fragmentation on bobcats and coyotes in southern California.
1174 *Conservation Biology*, **17**(2): 566-576.
- 1175 **Riley S, Brown J, Sikich J, Schoonmaker C, Boydston E. 2014.** Wildlife friendly roads: the
1176 impacts of roads on wildlife in urban areas and potential remedies. In: McCleery R, Moorman C,
1177 Peterson M. (eds) *Urban Wildlife Conservation*. Springer, Boston, MA, pp. 323- 360.

- 1178 **Ripple WJ, Estes JA, Beschta RL, Wilmers CC, Ritchie EG, Hebblewhite M, Berger J,**
1179 **Elmhagen B, Letnic M, Nelson MP, Schmitz OJ, Smith DW, Wallach AD, Wirsing AJ.**
1180 **2014.** Status and ecological effects of the world's largest carnivores. *Science*, **343**(6167):
1181 1241484.
- 1182 **Ritchie EG, Johnson CN. 2009.** Predator interactions, mesopredator release and biodiversity
1183 conservation. *Ecology Letters*, **12**(9): 982-998.
- 1184 **Rodríguez A, Andrén H, Jansson G. 2001.** Habitat-mediated predation risk and decision
1185 making of small birds at forest edges. *Oikos*, **95**(3): 383-396.
- 1186 **Rodrigues SM, LeDoux JE, Sapolsky RM. 2009.** The influence of stress hormones on fear
1187 circuitry. *Annual Review of Neuroscience*, **32**: 289-313.
- 1188 **Rodríguez A, Hausberger M, Clergeau P. 2010.** Flexibility in European starlings' use of social
1189 information: experiments with decoys in different populations. *Animal Behaviour*, **80**(6): 965-
1190 973.
- 1191 **Romero LM. 2004.** Physiological stress in ecology: lessons from biomedical research. *Trends in*
1192 *Ecology & Evolution*, **19**(5): 249-255.
- 1193 **Romero LM, Dickens MJ, Cyr NE. 2009.** The reactive scope model—a new model integrating
1194 homeostasis, allostasis, and stress. *Hormones and Behavior*, **55**(3), 375-389.
- 1195 **Roozendaal B. 2000.** Glucocorticoids and the regulation of memory
1196 consolidation. *Psychoneuroendocrinology*, **25**(3): 213-238.
- 1197 **Rosen JB, Schulkin J. 1998.** From normal fear to pathological anxiety. *Psychological*
1198 *Review*, **105**(2): 325-350.
- 1199 **Rosen JB. 2004.** The neurobiology of conditioned and unconditioned fear: a neurobehavioral
1200 system analysis of the amygdala. *Behavioral and Cognitive Neuroscience Reviews*, **3**(1): 23-41.
- 1201 **Rosen JB, Schulkin J. 2004.** Adaptive fear, allostasis, and the pathology of anxiety and
1202 depression. *Allostasis, Homeostasis and the Costs of Physiological Adaptation*: 164-227.
- 1203 **Santini L, González-Suárez M, Russo D, Gonzalez-Voyer A, von Hardenberg A, Ancillotto**
1204 **L. 2019.** One strategy does not fit all: determinants of urban adaptation in mammals. *Ecology*
1205 *Letters*, **22**(2): 365-376.
- 1206 **Santos X, Mateos E, Bros V, Brotons L, De Mas E, Herraiz JA, Herrando S, Miño À,**
1207 **Olmo-Vidal JM, Quesada J, Ribes J, Sabaté S, Sauras-Yera T, Serra A, Vallejo VR,**
1208 **Viñolas A. 2014.** Is response to fire influenced by dietary specialization and mobility? A
1209 comparative study with multiple animal assemblages. *PLoS ONE*, **9**(2): e88224.
- 1210 **Salo P, Korpimäki E, Banks PB, Nordström M, Dickman CR. 2007.** Alien predators are more
1211 dangerous than native predators to prey populations. *Proceedings of the Royal Society of London*
1212 *B: Biological Sciences*, **274**(1615): 1237-1243.

- 1213 **Sapolsky RM, Romero LM, Munck AU. 2000.** How do glucocorticoids influence stress
1214 responses? Integrating permissive, suppressive, stimulatory, and preparative actions. *Endocrine*
1215 *Reviews*, **21**(1): 55-89.
- 1216 **Say-Sallaz E, Chamaillé-Jammes S, Fritz H, Valeix M. 2019.** Non-consumptive effects of
1217 predation in large terrestrial mammals: Mapping our knowledge and revealing the tip of the
1218 iceberg. *Biological Conservation*, **235**:36–52. DOI: 10.1016/j.biocon.2019.03.044.
- 1219 **Schell CJ, Young JK, Lonsdorf EV, Santymire RM, Mateo JM. 2018.** Parental habituation to
1220 human disturbance over time reduces fear of humans in coyote offspring. *Ecology and*
1221 *Evolution*, **8**(24), 12965-12980.
- 1222 **Schmidt K, Kuijper DP. 2015.** A “death trap” in the landscape of fear. *Mammal Research*,
1223 **60**(4): 275-284.
- 1224 **Schmitz OJ, Beckerman AP, O’Brien KM. 1997.** Behaviorally mediated trophic cascades:
1225 effects of predation risk on food web interactions. *Ecology*, **78**:1388–1399.
- 1226 **Schmitz OJ, Krivan V, Ovadia O. 2004.** Trophic cascades: the primacy of trait-mediated
1227 indirect interactions. *Ecology Letters*, **7**(2): 153-163.
- 1228 **Schmitz OJ. 2008.** Effects of predator hunting mode on grassland ecosystem function. *Science*,
1229 **319**(5865): 952-954.
- 1230 **Schulkin J, Thompson BL, Rosen JB. 2003.** Demythologizing the emotions: adaptation,
1231 cognition, and visceral representations of emotion in the nervous system. *Brain and Cognition*,
1232 **52**(1): 15-23.
- 1233 **Sergio F, Caro T, Brown D, Clucas B, Hunter J, Ketchum J, McHugh K, Hiraldo**
1234 **F. 2008.** Top predators as conservation tools: ecological rationale, assumptions, and
1235 efficacy. *Annual Review of Ecology, Evolution and Systematics* **39**: 1– 19.
- 1236 **Seyle H. 1936.** A syndrome produced by diverse noxious agents. *Nature* (London), **138**: 32-39.
- 1237 **Shannon G, McKenna MF, Angeloni LM, Crooks KR, Fristrup KM, Brown E, Warner**
1238 **KA, Nelson MD, White C, Briggs J, McFarland S. 2016.** A synthesis of two decades of
1239 research documenting the effects of noise on wildlife. *Biological Reviews*, **91**(4): 982-1005.
- 1240 **Sheriff MJ, Dantzer B, Delehanty B, Palme R, Boonstra R. 2011.** Measuring stress in
1241 wildlife: techniques for quantifying glucocorticoids. *Oecologia*, **166**(4): 869-887.
- 1242 **Sherman LJ. 1984.** The effects of patchy food availability on nest-site selection and movement
1243 patterns of reproductively active female meadow voles *Microtus*
1244 *pennsylvanicus*. *Ecography*, **7**(3): 294-299.
- 1245 **Shochat E, Lerman S, Fernández-Juricic E. 2010.** Birds in urban ecosystems: population
1246 dynamics, community structure, biodiversity, and conservation. *Urban ecosystem ecology*,
1247 (urbanecosysteme):75-86.

- 1248 **Shrader AM, Brown JS, Kerley GI, Kotler BP. 2008.** Do free-ranging domestic goats show
1249 ‘landscapes of fear’? Patch use in response to habitat features and predator cues. *Journal of Arid*
1250 *Environments*, **72**: 1811–1819.
- 1251 **Sinclair ARE, Arcese P. 1995.** Population consequences of predation-sensitive foraging: The
1252 Serengeti wildebeest. *Ecology*, **76**(3): 882-891.
- 1253 **Smit B, Spaling H. 1995.** Methods for cumulative effects assessment. *Environmental Impact*
1254 *Assessment Review*, **15**(1): 81-106.
- 1255 **Smith JA, Suraci JP, Clinchy M, Crawford A, Roberts D, Zanette LY, Wilmers CC. 2017.**
1256 Fear of the human “super predator” reduces feeding time in large carnivores. *Proceedings of the*
1257 *Royal Society B: Biological Sciences*, **284**:20170433. DOI: 10.1098/rspb.2017.0433.
- 1258 **Smith JA, Donadio E, Pauli JN, Sheriff MJ, Middleton AD. 2019.** Integrating temporal
1259 refugia into landscapes of fear: prey exploit predator downtimes to forage in risky places.
1260 *Oecologia*, **189**:883–890. DOI: 10.1007/s00442-019-04381-5.
- 1261 **Snell-Rood EC, Wick N. 2013.** Anthropogenic environments exert variable selection on cranial
1262 capacity in mammals. *Proceedings of the Royal Society B: Biological Sciences*, **280**(1769):
1263 20131384.
- 1264 **Soso SB, Koziel JA, Johnson A, Lee YJ, Fairbanks WS. 2014.** Analytical methods for
1265 chemical and sensory characterization of scent-markings in large wild mammals: a review.
1266 *Sensors*, **14**: 4428–4465.
- 1267 **Spencer EE, Crowther MS, Dickman CR. 2014.** Risky business: do native rodents use habitat
1268 and odor cues to manage predation risk in Australian deserts?. *PLoS ONE*, **9**(2): e90566.
- 1269 **Stankowich T. 2008.** Ungulate flight responses to human disturbance: a review and meta-
1270 analysis. *Biological Conservation*, **141**(9): 2159-2173.
- 1271 **Sterling P, Eyer J. 1988.** Allostasis: a new paradigm to explain arousal pathology. In
1272 *‘Handbook of Life Stress, Cognition and Health’*(Eds K. Fisher and J. Reason.): 629–649.
- 1273 **Stillfried M, Fickel J, Börner K, Wittstatt U, Heddergott M, Ortmann S, Kramer-Schadt S,**
1274 **Frantz AC. 2017.** Do cities represent sources, sinks or isolated islands for urban wild boar
1275 population structure? *Journal of Applied Ecology*, **54**(1): 272-281.
- 1276 **Suraci JP, Clinchy M, Zanette LY, Wilmers CC. 2019.** Fear of humans as apex predators has
1277 landscape-scale impacts from mountain lions to mice. *Ecology Letters*. DOI: 10.1111/ele.13344.
- 1278 **Tablado Z, Jenni L. 2017.** Determinants of uncertainty in wildlife responses to human
1279 disturbance. *Biological Reviews*, **92**: 216-233.
- 1280 **Taylor-Brown A, Booth R, Gillett A, Mealy E, Ogbourne SM, Polkinghorne A, Conroy GC.**
1281 **2019.** The impact of human activities on Australian wildlife. *PLoS ONE*, **14**(1): e0206958.

- 1282 **Tews J, Brose U, Grimm V, Tielbörger K, Wichmann MC, Schwager M, Jeltsch F. 2004.**
1283 Animal species diversity driven by habitat heterogeneity/diversity: the importance of keystone
1284 structures. *Journal of Biogeography*, **31**(1): 79-92.
- 1285 **Therivel R, Ross B. 2007.** Cumulative effects assessment: does scale matter? *Environmental*
1286 *Impact Assessment Review*, **27**(5): 365-385.
- 1287 **Threlfall CG, Williams NS, Hahs AK, Livesley SJ. 2016.** Approaches to urban vegetation
1288 management and the impacts on urban bird and bat assemblages. *Landscape and Urban*
1289 *Planning*, **153**: 28-39.
- 1290 **Tigas LA, Van Vuren DH, Sauvajot RM. 2002.** Behavioral responses of bobcats and coyotes
1291 to habitat fragmentation and corridors in an urban environment. *Biological Conservation*, **108**(3):
1292 299-306.
- 1293 **Tooby J, Cosmides L. 1990.** The past explains the present: emotional adaptations and the
1294 structure of ancestral environments. *Ethology and Sociobiology*, **11**(4-5): 375-424.
- 1295 **Støen OG, Ordiz A, Evans AL, Laske TG, Kindberg J, Frøbert O, Swenson JE, Arnemo**
1296 **JM. 2015.** Physiological evidence for a human-induced landscape of fear in brown bears (*Ursus*
1297 *arctos*). *Physiology & Behavior*, **152**(A): 244-248.
- 1298 **Sweitzer RA. 1996.** Predation or starvation: consequences of foraging decisions by porcupines
1299 (*Erethizon dorsatum*). *Journal of Mammalogy*, **77**(4): 1068-1077.
- 1300 **Thomson RL, Tomás G, Forsman JT, Broggi J, Mönkkönen M. 2010.** Predator proximity as
1301 a stressor in breeding flycatchers: mass loss, stress protein induction, and elevated
1302 provisioning. *Ecology*, **91**(6): 1832-1840.
- 1303 **Threlfall CG, Law B, Banks PB. 2012.** Influence of landscape structure and human
1304 modifications on insect biomass and bat foraging activity in an urban landscape. *PLoS ONE*,
1305 **7**(6): e38800.
- 1306 **Towerton AL, Penman TD, Kavanagh RP, Dickman CR. 2011.** Detecting pest and prey
1307 responses to fox control across the landscape using remote cameras. *Wildlife Research*, **38**(3):
1308 208-220.
- 1309 **Travers M, Clinchy M, Zanette L, Boonstra R, Williams TD. 2010.** Indirect predator effects
1310 on clutch size and the cost of egg production. *Ecology Letters*, **13**(8): 980-988.
- 1311 **Trussell GC, Ewanchuk PJ, Matassa CM. 2006.** Habitat effects on the relative importance of
1312 trait-and density-mediated indirect interactions. *Ecology Letters*, **9**(11): 1245-1252.
- 1313 **Turcotte Y, Desrochers A. 2003.** Landscape-dependent response to predation risk by forest
1314 birds. *Oikos*, **100**(3): 614-618.

- 1315 **Van Buskirk J, Müller C, Portmann A, Surbeck M. 2002.** A test of the risk allocation
1316 hypothesis: tadpole responses to temporal change in predation risk. *Behavioral Ecology*, **13**(4):
1317 526-530.
- 1318 **Van Der Merwe M, Brown JS. 2008.** Mapping the landscape of fear of the cape ground squirrel
1319 (*Xerus inauris*). *Journal of Mammalogy*, **89**(5): 1162-1169.
- 1320 **Vijayakrishnan S, Kumar MA, Umapathy G, Kumar V, Sinha A. 2018.** Physiological stress
1321 responses in wild Asian elephants *Elephas maximus* in a human-dominated landscape in the
1322 Western Ghats, southern India. *General and Comparative Endocrinology*. DOI:
1323 10.1016/j.ygcen.2018.05.009.
- 1324 **Wallace KJ, Rosen JB. 2000.** Predator odor as an unconditioned fear stimulus in rats: elicitation
1325 of freezing by trimethylthiazoline, a component of fox feces. *Behavioral Neuroscience*, **114**(5):
1326 912-922.
- 1327 **Wallace KJ, Rosen JB. 2001.** Neurotoxic lesions of the lateral nucleus of the amygdala
1328 decrease conditioned fear but not unconditioned fear of a predator odor: comparison with
1329 electrolytic lesions. *Journal of Neuroscience*, **21**(10): 3619-3627.
- 1330 **Warburton B, Norton BG. 2009.** Towards a knowledge-based ethic for lethal control of
1331 nuisance wildlife. *Journal of Wildlife Management*, **73**(1): 158-164.
- 1332 **Weissburg M, Smee DL, Ferner MC. 2014.** The sensory ecology of nonconsumptive predator
1333 effects. *The American Naturalist*, **184**(2): 141-157.
- 1334 **Werner EE, Peacor SD. 2003.** A review of trait-mediated indirect interactions in ecological
1335 communities. *Ecology*, **84**(5): 1083-1100.
- 1336 **Western D. 2001.** Human-modified ecosystems and future evolution. *Proceedings of the*
1337 *National Academy of Sciences USA*, **98**(10): 5458-5465.
- 1338 **Willems EP, Hill RA. 2009.** Predator-specific landscapes of fear and resource distribution:
1339 effects on spatial range use. *Ecology*, **90**(2): 546-555.
- 1340 **Wingfield JC, Kitaysky AS. 2002.** Endocrine responses to unpredictable environmental events:
1341 stress or anti-stress hormones? *Integrative and Comparative Biology*, **42**(3): 600-609.
- 1342 **Woinarski JC, Burbidge AA, Harrison PL. 2015.** Ongoing unraveling of a continental fauna:
1343 decline and extinction of Australian mammals since European settlement. *Proceedings of the*
1344 *National Academy of Sciences USA*, **112**(15): 4531-4540.
- 1345 **Yates FE, Russell SM, Maran JW. 1971.** Brain-adenohypophysial communication in
1346 mammals. *Annual Review of Physiology*, **33**(1): 393-444.
- 1347 **Zanette LY, White AF, Allen MC, Clinchy M. 2011.** Perceived predation risk reduces the
1348 number of offspring songbirds produce per year. *Science*, **334**(6061): 1398-1401.

1349 **Zbyryt A, Bubnicki JW, Kuijper DPJ, Dehnhard M, Churski M, Schmidt K, Wong B.**
 1350 **2017.** Do wild ungulates experience higher stress with humans than with large carnivores?
 1351 *Behavioral Ecology*. DOI: 10.1093/beheco/axx142.

1352

Figure 1

A Venn diagram showing two main categories of stressor.

On the left are stressors that occur naturally in ecosystems, such as native predators, social and breeding interactions, availability of refuge and burrow/den microhabitats, disease / parasite prevalence, and availability of food and water. On the right are introduced stressors, primarily those arising from anthropogenic disturbances, such as: urban developments, human activity, introduced toxins and introduced predators. Where both categories of stressor occur together simultaneously, as in many urban environments, cumulative stress impacts can result in homeostatic overload or failure (as defined by Romero, Dickens, and Cyr, 2009). In these situations populations may be at particular risk of collapse and conservation action will be most urgent.




Figure 2

A conservation management approach that outlines the key steps for assessing if cumulative stress impacts are occurring between stressors that occur naturally in ecosystems and those that are introduced.

The circled areas indicate where conservation management initiatives may be used to mitigate these effects through management of vegetation complexity or supplementation materials such as water stations or carved 'tree-hollow' cavities.

Determine the status of the target population & the quality of the habitat		Determine how many additional stressors are present		Determine the intensity of the stressors present	
Is the target population obviously active?	No <input type="checkbox"/>	Are there any introduced pollutants?	Yes <input type="checkbox"/>	If across both columns more than two questions have been ticked, then the cumulative impacts are either: - Additive and an individual may be in a state of reactive homeostasis - Synergistic and an individual may be in a state of homeostatic overload - If exposure is chronic then an individual may experience homeostatic failure	
Is the target population's activity decreasing?	Yes <input type="checkbox"/>	Are there any fear inducing stressor of:	Yes <input type="checkbox"/>		
Are there adequate resources of food, water, and shelter?	No <input type="checkbox"/>	- Native predation pressures?	Yes <input type="checkbox"/>		
Does the habitat provide avenues to fulfil social interactions, including mating?	No <input type="checkbox"/>	- Introduced predation pressures?	<input type="checkbox"/>		
Has there been or is there likely to be any environmental disturbances, including stochastic weather events?	Yes <input type="checkbox"/>	- Human activity?	Yes <input type="checkbox"/>		
		Are any of these additional stressors likely to be persistent, without management (including the absence of an apex predator)?	Yes <input type="checkbox"/>		
If more than one box was ticked the population may be considered vulnerable to additional stressors		If more than one box was ticked the stressors may have cumulative effects (additive or synergistic)		Stressors that are circled are areas where conservation management may be used to mitigate the effects	

Figure 3

Graphical Abstract

Why wildlife conservation needs to measure: *Fear and stressing in predator-prey ecology: considering the twin stressors of predators and people on mammals*

