

The prediction of swim start performance based on squat jump force-time characteristics (#42630)

1

First submission

Guidance from your Editor

Please submit by **4 Dec 2019** for the benefit of the authors (and your \$200 publishing discount) .



Structure and Criteria

Please read the 'Structure and Criteria' page for general guidance.



Custom checks

Make sure you include the custom checks shown below, in your review.



Raw data check

Review the raw data. Download from the [materials page](#).



Image check

Check that figures and images have not been inappropriately manipulated.

Privacy reminder: If uploading an annotated PDF, remove identifiable information to remain anonymous.

Files

Download and review all files from the [materials page](#).

3 Figure file(s)

4 Table file(s)

1 Raw data file(s)

! Custom checks

Human participant/human tissue checks



Have you checked the authors [ethical approval statement](#)?



Does the study meet our [article requirements](#)?



Has identifiable info been removed from all files?



Were the experiments necessary and ethical?



Structure and Criteria

Structure your review

The review form is divided into 5 sections. Please consider these when composing your review:

1. BASIC REPORTING
2. EXPERIMENTAL DESIGN
3. VALIDITY OF THE FINDINGS
4. General comments
5. Confidential notes to the editor

 You can also annotate this PDF and upload it as part of your review

When ready [submit online](#).

Editorial Criteria

Use these criteria points to structure your review. The full detailed editorial criteria is on your [guidance page](#).

BASIC REPORTING

-  Clear, unambiguous, professional English language used throughout.
-  Intro & background to show context. Literature well referenced & relevant.
-  Structure conforms to [Peerj standards](#), discipline norm, or improved for clarity.
-  Figures are relevant, high quality, well labelled & described.
-  Raw data supplied (see [Peerj policy](#)).

EXPERIMENTAL DESIGN

-  Original primary research within [Scope of the journal](#).
-  Research question well defined, relevant & meaningful. It is stated how the research fills an identified knowledge gap.
-  Rigorous investigation performed to a high technical & ethical standard.
-  Methods described with sufficient detail & information to replicate.

VALIDITY OF THE FINDINGS

-  Impact and novelty not assessed. Negative/inconclusive results accepted. *Meaningful* replication encouraged where rationale & benefit to literature is clearly stated.
-  All underlying data have been provided; they are robust, statistically sound, & controlled.
-  Speculation is welcome, but should be identified as such.
-  Conclusions are well stated, linked to original research question & limited to supporting results.

Standout reviewing tips

3



The best reviewers use these techniques

Tip

Support criticisms with evidence from the text or from other sources

Example

Smith et al (J of Methodology, 2005, V3, pp 123) have shown that the analysis you use in Lines 241-250 is not the most appropriate for this situation. Please explain why you used this method.

Give specific suggestions on how to improve the manuscript

Your introduction needs more detail. I suggest that you improve the description at lines 57- 86 to provide more justification for your study (specifically, you should expand upon the knowledge gap being filled).

Comment on language and grammar issues

The English language should be improved to ensure that an international audience can clearly understand your text. Some examples where the language could be improved include lines 23, 77, 121, 128 – the current phrasing makes comprehension difficult.

Organize by importance of the issues, and number your points

1. Your most important issue
2. The next most important item
3. ...
4. The least important points

Please provide constructive criticism, and avoid personal opinions

I thank you for providing the raw data, however your supplemental files need more descriptive metadata identifiers to be useful to future readers. Although your results are compelling, the data analysis should be improved in the following ways: AA, BB, CC

Comment on strengths (as well as weaknesses) of the manuscript

I commend the authors for their extensive data set, compiled over many years of detailed fieldwork. In addition, the manuscript is clearly written in professional, unambiguous language. If there is a weakness, it is in the statistical analysis (as I have noted above) which should be improved upon before Acceptance.

The prediction of swim start performance based on squat jump force-time characteristics

Shiqi Thng^{Corresp., 1, 2}, Simon Pearson², Evelyne Rathbone¹, Justin Keogh^{1, 3, 4, 5}

¹ Faculty of Health Sciences and Medicine, Bond University, Gold Coast, QLD, Australia

² Queensland Academy of Sport, Nathan, QLD, Australia

³ Sports Performance Research Centre New Zealand, Auckland University of Technology, Auckland, New Zealand

⁴ Cluster for Health Improvement, Faculty of Science, , Health, Education and Engineering, University of the Sunshine Coast, Sippy Downs, QLD, Australia

⁵ Kasturba Medical College, Mangalore, Manipal Academy of Higher Education, Manipal, Karnataka, India

Corresponding Author: Shiqi Thng

Email address: sthng@bond.edu.au

Background: Close margins come between success in a sprint swimming event, with winning margins as close as tenths or hundredths of a second. Thus, any improvements made in the swimming start can have positive implications on overall swimming performance. However, it is still somewhat unclear what are the primary physiological characteristics underpinning swim start performance. The primary aim of this study was to develop a multiple regression model to determine key lower body force-time predictors using the squat jump for swim start performance as assessed by time to 5 m and 15 m in national and international level swimmers. A secondary aim was to determine if any differences exist between males and females in jump performance predictors for swim start performance.

Methods: A total of 38 males (age 21 ± 3.1 years, height 1.83 ± 0.08 m, body mass 76.7 ± 10.2 kg), 34 females (age 20.1 ± 3.2 years, height 1.73 ± 0.06 m, body mass 64.8 ± 8.4 kg) who had competed at either an elite ($n = 31$) or national level ($n = 41$) participated in this study. All tests were performed on the same day, with participants performing bodyweight squat jumps on a force platform followed by a swim start performance test using an instrumented starting block.

Results: Stepwise multiple linear regression with quadratic fitting identified concentric impulse and concentric impulse² as statistically significant predictors for time to 5 m ($R^2 = 0.659$) in males. With time to 15 m, concentric impulse, age and concentric impulse² were statistically significant predictors for males ($R^2 = 0.807$). For a quick time to 5 m and 15 m in males, a minimum concentric impulse of 200 – 230 N.s appears required, with any additional impulse production not being associated with a reduction in swim start times for most male swimmers. Concentric impulse, Reactive Strength Index modified and concentric mean power were identified as statistically significant predictors for female swimmers to time to 5 m ($R^2 = 0.689$). With time to 15 m, concentric impulse, body mass, concentric rate of power development and Reactive Strength Index modified ($R^2 = 0.841$) were identified as statistically significant predictors for females.

Discussion: The results of this study highlight the importance of lower body power and strength for swim start performance, although being able to produce greater than 200 or 230 N.s concentric impulse in squat jump did not necessarily increase swim start performance over 5 m and 15 m, respectively. Swimmers who can already generate greater these levels of concentric impulse may benefit more from improving their rate of force development and/or technical aspects of the swim start performance. The sex-related differences in key force-time predictors suggest that male and female swimmers may require

individualised strength and conditioning programs and regular monitoring of performance.

1 The prediction of swim start performance based on squat jump force-time characteristics

Shiqi Thng^{1,2}, Simon Pearson², Evelyne Rathbone¹, Justin W.L. Keogh^{1,3,4,5}

¹ Faculty of Health Sciences and Medicine, Bond University, Gold Coast, QLD, Australia

² Queensland Academy of Sport, Nathan, QLD, Australia

³ Sports Performance Research Centre New Zealand, AUT University, Auckland, New Zealand

⁴ Cluster for Health Improvement, Faculty of Science, Health, Education and Engineering,
University of the Sunshine Coast, Sippy Downs. QLD, Australia

⁵ Kasturba Medical College, Mangalore, Manipal Academy of Higher Education, Manipal,
Karnataka, India

Corresponding Author

Shiqi Thng

Bond University, Faculty of Health Science and Medicine, Gold Coast, Australia

Ph. +61 7 5595 4489

E: sthng@bond.edu.au

ORCID: 0000-0002-3215-1458

2

3

4 Abstract

5 Background: Close margins come between success in a sprint swimming event, with winning
6 margins as close as tenths or hundredths of a second. Thus, any improvements made in the
7 swimming start can have positive implications on overall swimming performance. However, it is
8 still somewhat unclear what are the primary physiological characteristics underpinning swim
9 start performance. The primary aim of this study was to develop a multiple regression model to
10 determine key lower body force-time predictors using the squat jump for swim start performance
11 as assessed by time to 5 m and 15 m in national and international level swimmers. A secondary
12 aim was to determine if any differences exist between males and females in jump performance
13 predictors for swim start performance.

14 Methods: A total of 38 males (age 21 ± 3.1 years, height 1.83 ± 0.08 m, body mass 76.7 ± 10.2
15 kg), 32 females (age 20.1 ± 3.2 years, height 1.73 ± 0.06 m, body mass 64.8 ± 8.4 kg) who had
16 competed at either an elite ($n = 31$) or national level ($n = 41$) participated in this study. All tests
17 were performed on the same day, with participants performing bodyweight squat jumps on a
18 force platform followed by a swim start performance test using an instrumented starting block.

19 Results: Stepwise multiple linear regression with quadratic fitting identified concentric impulse
20 and concentric impulse² as statistically significant predictors for time to 5 m ($R^2 = 0.659$) in
21 males. With time to 15 m, concentric impulse, age and concentric impulse² were statistically
22 significant predictors for males ($R^2 = 0.807$). For a quick time to 5 m and 15 m in males, a
23 minimum concentric impulse of 200 – 230 N.s appears required, with any additional impulse
24 production not being associated with a reduction in swim start times for most male swimmers.
25 Concentric impulse, Reactive Strength Index modified and concentric mean power were
26 identified as statistically significant predictors for female swimmers to time to 5 m ($R^2 = 0.689$).
27 With time to 15 m, concentric impulse, body mass, concentric rate of power development and
28 Reactive Strength Index modified ($R^2 = 0.841$) were identified as statistically significant
29 predictors for females.

30 Discussion: The results of this study highlight the importance of lower body power and strength
31 for swim start performance, although being able to produce greater than 200 or 230 N.s
32 concentric impulse in squat jump did not necessarily increase swim start performance over 5 m
33 and 15 m, respectively. Swimmers who can already generate greater these levels of concentric
34 impulse may benefit more from improving their rate of force development and/or technical
35 aspects of the swim start performance. The sex-related differences in key force-time predictors
36 suggest that male and female swimmers may require individualised strength and conditioning
37 programs and regular monitoring of performance.

38 Introduction

39 Swim start performance has been identified as a determining factor for success, especially in
40 sprint distance events, as it is the part of the race that the swimmer is travelling at the fastest
41 velocity [1, 2]. While the exact nature of starts may differ between the four swimming strokes,
42 there are three primary phases that contribute towards the overall start performance. The block
43 phase requires a quick reaction to the starting signal and a large take-off velocity that is primarily

horizontal in direction [3]. The subsequent flight phase is an example of projectile motion, whereby the swimmer becomes airborne and finishes when they contact the water [4, 2]. The flight phase is followed by the underwater phase, in which swimmers attempt to maintain a streamlined position with their arms outstretched in front of the head to minimise velocity loss while also performing propulsive undulatory leg kicks (except in breaststroke) until their head resurfaces before the 15 m mark [5]. The block, flight, and underwater phase account for approximately 11 %, 5 %, and 84 % respectively of the total start time [4]. The average velocity during the underwater phase is highly dependent on the take-off velocity acquired in the block phase, the horizontal distance obtained in the flight phase, as well as the degree of streamlining and effectiveness of the undulatory leg kicks during the underwater phase [6].

As close margins often exist between medallists in sprint swimming events, being able to identify areas to achieve marginal gains in performance by tenths or even hundredths of a second can make a difference in overall performance [7]. Previous research has highlighted a key component of swim start performance is the ability to produce high forces off the starting block [8, 9]. In a recent systematic review of eight cross-sectional studies, Thng et al. observed significant correlations between vertical jump and lower body strength scores to swim start performance in swimmers of a variety of standards, with these correlations typically higher for the jump than strength tests [10]. Specifically, near perfect correlations ($r > 0.90$) between jump height or take-off velocity and swim start performance were observed in the eight studies. These results highlight the importance of lower body power and strength as an important component of swim start performance. However, out of the 8 cross-sectional studies identified in the systematic review [10], only one study [11] utilised the OSB11 start block (OMEGA, Zurich, Switzerland) that is currently used in competitive swimming. The OSB11 start block which was introduced by the International Swimming Federation in 2010 has an angled kick plate at the rear of the block that allows the swimmer to adopt a kick start technique. Honda, Sinclair, Mason, and Pease [12] have identified that the angled kick plate on the OSB11 start block is capable of significantly improving both time to 5 m and 7.5 m, with a further 0.04 s improvement obtained in the kick start compared to the track start technique performed on the previous starting block. This is attributed to an increase in horizontal force application and subsequent take-off velocity results from the additional contribution of the rear leg on the kick plate. This view of Honda and colleagues [12] was consistent with the findings of Slawson et al. [4] who observed higher peak horizontal and vertical force generation with the OSB11 start blocks in elite swimmers, with these forces significantly correlated to a better start performance as assessed by block time, take-off velocity and flight distance.

In addition, all of the studies described in the systematic review by Thng et al. [10] only involved correlational analyses. While correlations describe the nature of a relationship between two variables, other statistical approaches such as multiple linear regression may provide more information regarding what power and strength (hereafter referred to as force-time characteristics) of jumping performance may best predict swim start performance in high performance swimmers. The lack of research using the OSB11 start block and kick start technique in these correlation studies needs to be addressed, as this relative lack of ecological validity with the start technique used in seven of the eight published studies may limit the generalisability to contemporary high-performance swimming. Another limitation of the

previous literature is the small sample sizes of recreational to sub-elite swimmers ($n = 7 - 27$) and the relative focus on male swimmers at the expense of their female counterparts. This is a concern as previous research [13, 14] have established differences in force and power capabilities between males and females in other athletic activities. For example, a number of studies have observed that males are able to produce higher velocities at the same percent of one repetition maximum and have a greater rate of force development and countermovement jump height than females [13-16]. Rice et al. [13] concluded that this greater jump height observed in males compared to females can be attributed to larger concentric impulse and thus greater velocity throughout most of the concentric phase at take-off in the countermovement jump. Further, the higher rate of force development and ability to produce greater velocities at the same percentage of one repetition maximum in males may be a result of greater muscle thickness and cross-sectional area, greater percentage of fast-twitch muscle fibres, greater concentration of anabolic hormones and higher neural activity during muscle contractions compared to females [17]. From a practical standpoint, these sex-related differences in force-time characteristics suggests there might need to be some potential differences in aspects of athletic monitoring and strength and conditioning programs between high-performance male and female swimmers.

Therefore, the primary objective of this study was to develop a multiple regression model to determine key lower body force-time predictors for swim start performance using the squat jump in high performance swimmers. Considering the potential sex differences in force-time characteristics during jumping, a secondary aim was to determine if differences exists between males and females in jump performance predictors for swim start performance.

Materials & Methods

Study Design

A cross-sectional study design was used to quantify the relationship between squat jump (SJ) force-time variables to swim start performance as assessed by times to 5 m and 15 m in national and international level swimmers. All tests were performed on the same day, with participants first performing SJ testing on the force platform, followed by a swim start performance test.

Participants

Participants were 38 males (age 21 ± 3.1 years, height 1.83 ± 0.08 m, body mass 76.7 ± 10.2 kg), 34 females (age 20.1 ± 3.2 years, height 1.73 ± 0.06 m, body mass 64.8 ± 8.4 kg) who had competed at either an elite ($n = 31$) or national level ($n = 41$) in front crawl, butterfly or breaststroke. Backstroke was excluded due to the start being initiated from within the water, instead of on the elevated OSB11 starting block. Elite level swimmers comprised of swimmers who had competed internationally in either the Olympics, Commonwealth Games or World Championships. National level swimmers comprised of swimmers that regularly competed at national championships. Swimmers were required to have at least 1 year of land-based resistance training experience under the supervision of a strength and conditioning coach. All participants gave informed consent to participate in the study, which was conducted in accordance with the Declaration of Helsinki and approved by Bond University Human Research Ethics Committee (0000016006), The University of Queensland Human Research Ethics Committee (HMS17/41) and Swimming Australia Ltd.

Methodology

Squat Jump test

Prior to the SJ test, participants completed a dynamic lower body warm-up under the supervision of a strength and conditioning coach. Following the warm-up, participants were given two practice jumps before the test was conducted. Jumps were performed on a force platform (ForceDecks FD4000, London, United Kingdom), with a sample rate of 1000 Hz. Participants started in an upright standing position with their hands on their hips. They were then instructed to adopt a squat position using a self-selected depth that was held for 3 seconds before they attempted to jump as high as possible [18]. A successful trial was one that did not display any small amplitude countermovement at the start of the jump phase on the force trace [19]. All participants were asked to perform three maximal intensity SJ with a 30-second passive rest in between each effort.

The SJ trial with the highest jump height was kept for data analysis. Jump height was determined by the conventional impulse-momentum method ($\text{Jump Height} = v^2/2g$, where v = velocity at take-off and g = gravitational acceleration) [20]. Ground reaction force data from the SJs were analysed using the commercially available ForceDecks software (ForceDecks, London, United Kingdom). Out of the variables provided by ForceDecks, 46 variables, excluding any left-to right asymmetry variables were initially extracted for use in further analysis.

Swim start

After completing a self-selected warm-up based on their usual pre-race warm-up routine, participants then performed three maximal effort swim starts (time to 15 m) with their main swim stroke (front crawl, butterfly, or breaststroke). Trials were started as per competition conditions and swimmers were instructed to swim to a distance past the 15 m mark, in order to ensure that representative values at the 15 m distance were obtained [21]. Two-minutes of passive recovery was given between each trial [22]. The start with the fastest time to 5 m for each individual were selected for further analysis. Starts performed using the front crawl stroke were also used for additional analysis based on the fastest time to 15 m. It was necessary to reduce this secondary analysis to a single stroke due to the influence of stroke type on 15 m times, where the greatest number of athletes performed the front crawl ($n = 50$).

Swim start performance were collected using a Kistler Performance Analysis System – Swimming (KPAS-S, Kistler Winterthur, Switzerland), which utilises a force instrumented starting block, constructed to match the dimensions of the Omega OSB11 block (KPAS-S Type 9691A1; Kistler Winterthur, Switzerland). Time to 5 m and 15 m were collected using five calibrated high speed digital cameras collecting at 100 frames per second, synchronised to the instrumented starting block using the KPAS-S. One camera was positioned 0.95 m above the water and 2.5 m perpendicular to the direction of travel to capture the start and entry of swimmer into the water, while the other three cameras were positioned 1.3 m underwater at 5 m, 10 m and 15 m perpendicular to the swimmer to capture the time to 15 m (Figure 1) [22]. The times to 5 m and 15 m were defined as the time elapsed from the starting signal until the apex of the swimmers' head passed the respective distances [22]. An Infinity Start System (Colorado Time Systems, Loveland, Colorado, USA) provided an audible starting signal to the athletes as well as an electronic start trigger to the KPAS-S system.

Please insert Figure one about here

Statistical Analysis

Descriptive statistics are reported as mean \pm SD for normally distributed continuous variables and frequency (%) for categorical variables. Normality was checked using histograms, normal Q-Q plots and the Shapiro-Wilk test. A principal component analysis (PCA) was carried out on 46 jump variables extracted from ForceDecks force platform (ForceDecks, London, United Kingdom) on 72 participants to reduce dimensionality. A second PCA was conducted to explore the new dataset of 32 jump performance variables and identify the principal components (PC) summarising the primary force-time variables. The decision on a suitable number of PCs to retain in each PCA required eigenvalues of 1.0 or greater (Kaiser criterion) and was supported by the use of a scree plot.

Multiple linear regression models using a stepwise regression method were initially performed to identify the potential predictors of the outcome variables of time (s) to 5 m and 15 m. Analyses were carried out on the entire dataset, and also on the data split by sex. Second order polynomial models were also investigated - the quadratic models were preferred to the linear models for males as they fitted the data better. Collinearity diagnostics were used to avoid the problem of multicollinearity. The assumptions of normality, linearity and homoscedasticity of residuals were verified. Results of the regression modelling are presented in terms of unstandardized coefficients, the 95% CI and p-values, along with the R^2 and standard error of estimate. Data were analysed with statistical software R version 3.5.3 and SPSS version 23.0.0 (SPSS Inc., Chicago, IL). P-values less than 0.05 were deemed to indicate statistical significance.

Results

Seventy-two swimmers, comprising 38 males and 34 females were included in this study. The physical characteristics of the participants are described in Table 1. While all 72 participants completed the time to 5 m, only 50 participants performed their swim start to 15 m using front crawl technique. Statistically significant differences among males and females were observed in a number of variables (Table 1), with males significantly heavier, taller and faster to 5 m and 15 m than females.

Please insert Table one about here

In the first PCA analysis on the 46 jump variables extracted from ForceDecks force platform (ForceDecks, London, United Kingdom), four PCs which explained 82 % of the variance were identified. Thirty-two most influential jump variables were identified from this initial PCA. A secondary PCA was run to explore the new dataset of 32 jump performance variables. The first three components, which explained 93 % of the variance, were retained. From this set, 15 variables were identified as potential predictors in subsequent regression models (Table 2). The results revealed that Component 1 accounting for 67.5 % of the variance, was of predominantly kinetic component. Component 2 accounting for 17.1 % of the variation, was predominantly a

time-dependent kinematic component. Lastly, Component 3 accounted for 8.5 % of the variation, with the highest load attributed to bodyweight.

Please insert Table two about here

Linear stepwise multiple regression analyses were performed to predict time to 5 m (see Figure 2 and Table 3) and time to 15 m (see Figure 3 and Table 4) in the overall sample of males and females as well as male and female subgroups.

Time to 5 m

Please insert Figure two about here

The scatterplot in Figure 2 shows a quadratic relationship between concentric impulse and time to 5 m in males ($R^2 = 0.693$). For a fast time to 5 m for males, a minimum concentric impulse production of around 180 - 200 N.s is required. While the generalised model then shows no additional reduction in time to 5 m with a higher concentric impulse for most swimmers, there are some outlier individuals who continue to derive additional performance benefit from an increased concentric impulse up to approximately 230 N.s. The relationship between concentric impulse and time to 5 m observed in females was linear ($R^2 = 0.487$), but this relationship was affected by other factors outlined in Table 3.

Please insert Table three about here

Concentric impulse was a statistically significant predictor in all three regression models (Table 3). The best prediction equations for time to 5 m in females and males were as follows:

Females: $T5\text{ m (s)} = 2.103 - 0.003 (\text{concentric impulse}) - 0.209 (\text{RSImod}) + 0.0002 (\text{concentric mean power})$

Males: $T5\text{ m (s)} = 2.645 - 0.010 (\text{concentric impulse}) + 0.00002 (\text{concentric impulse})^2$

Time to 15 m

Please insert Figure three about here

The scatterplot in Figure 3 shows a quadratic relationship between concentric impulse and time to 15 m in males ($R^2 = 0.746$). For a fast time to 15 m in males, a minimum concentric impulse production of around 230 N.s is required. However, similar to Figure 2, the relationship between

concentric impulse and time to 15 m observed in females was linear ($R^2 = 0.651$) but this relationship was also affected by other factors outlined in Table 4.

Please insert Table four about here

Concentric impulse was also the main significant predictor in all three regression models of the time to 15 m (Table 4). The best regression models were as follows:

Females: $T_{15\text{ m}}(\text{s}) = 9.303 - 0.030(\text{concentric impulse}) + 0.035(\text{body mass}) + 0.0002(\text{concentric RPD}) - 1.714(\text{RSImod})$

Males: $T_{15\text{ m}}(\text{s}) = 11.188 - 0.033(\text{concentric impulse}) - 0.048(\text{age}) + 0.00007(\text{concentric impulse})^2$

Discussion

The present study revealed that several lower body force-time characteristics, in particular concentric impulse, were significantly related to swim start performance in national and international level swimmers. However, when these analyses were performed for each sex individually, several differences in the prediction of swim start performance were observed. These sex-related differences in key force-time characteristics suggest that strength and conditioning programs and regular monitoring of performance may need to be tailored to male and female swimmers.

In the swim start, swimmers have to apply large forces rapidly on the start block to maximise horizontal take-off velocity, which in turn allows them to travel farther horizontally in the air before entering the water [23]. This task demand is consistent with the impulse-momentum relationship, whereby an impulse (the product of force and time of force application) needs to be generated to cause a change in momentum (i.e. velocity) of the system [24]. An analysis by Tor et al. [25] of the above water parameters in the swim start have found that take-off velocity and time on block were key predictors of swim start performance as assessed by time to 15 m using the OSB11 start block. Strong positive correlations between peak forces in the countermovement jump and peak forces on the OSB11 start block have also been reported by Cossor and colleagues [26]. Thus, to be able to achieve a high take-off velocity, a swimmer needs to be able to apply high forces/impulses off the starting block. Given that the swim start is mainly a concentric only movement, the findings of the present study further emphasise the important association between a swimmers' ability to produce impulse in the SJ and swim start performance.

It was expected that the current study would demonstrate a stronger prediction to 5 m than 15 m in the swim start. This hypothesis was based on how the movement pattern in the SJ is similar to the initial push-off in the block phase as well as the findings of Garcia-Ramos et al. [27] and Benjanuvatra et al. [8], who reported a significant correlation in take-off velocity [27] and jump height [8] in the SJ to 5 m ($r = -0.56$ and $r = -0.92$ respectively) but not 15 m. In contrast to this initial hypothesis, the current study demonstrated that the SJ force-time variables explained a greater amount of variance in time to 15 m than time to 5 m. Results of the current study were

also consistent with Garcia-Ramos et al. [11] who observed that the correlations between jump height and swim start performance were greater for the time to 15 m ($r = -0.67$) than time to 5 m ($r = -0.55$) using the kick start technique. Such equivalence in the literature was surprising, but it is possible that these contrasting findings from the current study to the limited literature could be attributed to a variety of between study differences, including the swim start technique and start block, as well as the sample size and homogeneity of participants included in the previously published studies. The current study utilised the kick start technique on the OSB11 start block, whereas Benjanuvatra et al. [8] and Garcia-Ramos et al. [27] utilised the grab start and track start, respectively. In addition, both of these studies included only female swimmers and had substantially smaller sample sizes ($n = 20$ and $n = 7$), whereas the current study utilised a mix of male and female swimmers, with a larger sample size ($n = 72$). As previously mentioned, the underwater phase is a key parameter in swim start performance, as a swimmer spends the highest percentage of the start in the underwater phase [25, 28]. Garcia-Ramos et al. [27] have suggested that swimmers require high levels of lower body strength and power to maximise their underwater kick performance. Therefore, it is possible that the stronger prediction in time to 15 m than 5 m in this study and the study by Garcia-Ramos et al. [11] may reflect the commonality in lower body force-time characteristics required for the block phase with the kick start technique and the undulatory kicks performed during the underwater phase.

Another focus of this study was examining potential sex-related differences in the force-time characteristics that may underpin swim start performance in high-performance swimmers. While concentric impulse was the strongest predictor for time to 5 m and 15 m in both males and females, the current study identified some differences between the sexes with respect to the predictors of time to 5 m and 15 m. For a quick time to 5 m and 15 m in males, a minimum concentric impulse of 200 – 230 N.s, respectively appears required, with any additional impulse production not being associated with a reduction in swim start times for most male swimmers. However, it is worth noting that within the dataset, there appear to be some athletes whose performance sits outside the generalised trend, showing increased performance gains from additional concentric impulse about the level at which most individuals are deriving no further benefit (Fig. 2 and 3). Nevertheless, these findings tend to suggest that for male athletes capable of producing greater than 230 N.s of impulse, it might be most beneficial for their strength and conditioning program to focus on improving their rate of force development, as it is possible that developing this high level of impulse in a shorter block time is required to further improve their swim start performance.

In contrast to the results for the male swimmers, which had concentric impulse as the sole contributing force-time variable from squat jumps, swim start performance to 5 m and 15 m for females were also influenced by other factors such as RSImod, mean power and concentric RPD. A few possible explanations for the differing strategies could be attributed to maximal strength capacity, load-velocity and neuromuscular capability between both sexes. Although lower body muscular strength was not measured in the current study, maximal strength has been shown to be a limiting factor in jumping ability and other lower body measure of explosive strength [29, 30]. Previous research has demonstrated that males possess greater maximal strength [31] and ability to produce greater velocities at the same percentage of one repetition maximum than their female counterparts [16]. When comparing the force-time curves in the countermovement jump between sexes, previous research [31, 32] have reported that the male and female differences in countermovement jump height were attributed to force characteristics and not temporal

characteristics of the force-time curve. This suggests that both sexes possess similar abilities to express forces, but the primary difference in jumping ability was due to the rate and magnitude of force production during both peak eccentric and concentric force production. Thus, these sex related differences might explain some of the differing strategies in the present study.

Conclusion

In summary, this study has identified squat jump concentric impulse as a key lower body force-time characteristic that was significantly related to swim start performance in high-performance swimmers. As impulse is the product of the ground reaction force and time of force application, it is integral for a swimmer to have the requisite ability to generate a high level of concentric impulse in a relatively short amount of time. Previous studies [33-35] have suggested that there is a trade-off between time spent on the starting block and take-off velocity. This is due to the likelihood of greater impulses being produced with greater block times. From a practical standpoint, a possible strategy to increase impulse generated on the starting block without excessively increasing the time of force application is to increase muscular strength and rate of force development qualities of the lower body through heavy resistance training and plyometric training [36, 37]. Heavy resistance training has been shown to increase power production, rate of power development, rate of force development and increases in muscle fiber cross-sectional area and neuromuscular activity [38]. The inclusion of plyometric training may also allow for the transfer maximal strength to power production and rate of force development [30], and has been shown to elicit significant improvements in key swim start parameters such as time to 5 m, take-off velocity and impulse [7, 23, 39].

Due to the different strength of the prediction equations, it appears that male and female swimmers utilise somewhat differing strategies during the swim start. While it is unknown if this is predominantly a result of the differences in muscular strength and force producing capacity between sexes, our results highlight the need for strength and conditioning coaches to consider individualising training programs to enhance swim start performance and ultimately swimming performance between sexes. From a monitoring perspective, if a swimmer possesses the concentric impulse production required but has slow start times to 5 m and 15 m, improving rate force development and/or assessing technical factors such as angle of entry, degree of streamline, hydrodynamic drag and underwater propulsion as mentioned would be imperative to maximise strength transfer to the swim start and ultimately swimming performance [35]. Thus, swimmers should be concurrently performing lower body strength and power training and practice the swim start to optimise the transfer of their strength and conditioning program in improving swim start performance [40].

Acknowledgements

This work was supported by the Queensland Academy of Sport's Sport Performance Innovation and Knowledge Excellence Unit in conjunction with Bond University Faculty of Health Sciences and Medicine. The authors would like to acknowledge Dr. Mark Osborne for his support in preparing this manuscript.

References

1. Cossor J, Mason B. Swim start performances at the sydney 2000 olympic games. *International Symposium on Biomechanics in Sports*. 2001:70-3.
2. Tor E, Pease D, Ball K. Characteristics of an elite swimming start. *BMS*. 2014;1:257-63.
3. Garcia-Hermoso A, Escalante Y, Arellano R, Navarro F, Dominguez AM, Saavedra JM. Relationship between final performance and block times with the traditional and the new starting platforms with a back plate in international swimming championship 50-m and 100-m freestyle events. *J Sports Sci Med*. 2013;12(4):698-706.
4. Slawson SE, Conway PP, Cossor JM, Chakravorti N, West AA. The categorisation of swimming start performance with reference to force generation on the main block and footrest components of the omega osb11 start blocks. *J Sports Sci*. 2013;31(5):468-78.
5. Formicola D, Rainoldi A. A kinematic analysis to evaluate the start techniques' efficacy in swimming. *Sport Sci Health*. 2015;11(1):57-66.
6. Mason B, Alcock A, Fowlie J. A kinetic analysis and recommendations for elite swimmers performing the sprint start. In: Menzel HJ, Chagas, M. H., editor. *Proceedings of XXV International Symposium on Biomechanics and Sport*; Ouro Preto: Brazil2007. p. 192-5.
7. Bishop DC, Smith RJ, Smith MF, Rigby HE. Effect of plyometric training on swimming block start performance in adolescents. *J Strength Cond Res*. 2009;23(7):2137-43.
8. Benjanuvatra N, Edmunds K, Blanksby B. Jumping abilities and swimming grab-start performance in elite and recreational swimmers. *IJARE*. 2007;1(3):231-41.
9. García-Ramos A, Feriche B, de la Fuente B, Argüelles-Cienfuegos J, Strojnik V, Strumbelj B et al. Relationship between different push-off variables and start performance in experienced swimmers. *Eur J Sport Sci*. 2015;15(8):687-95.
10. Thng S, Pearson S, Keogh JW. Relationships between dry-land resistance training and swim start performance and effects of such training on the swim start: A systematic review. *Sports Med*. 2019:1-17.
11. Garcia-Ramos A, Padial P, de la Fuente B, Arguelles-Cienfuegos J, Bonitch-Gongora J, Feriche B. Relationship between vertical jump height and swimming start performance before and after an altitude training camp. *J Strength Cond Res*. 2016;30(6):1638-45.
12. Honda KE, Sinclair PJ, Mason BR, Pease DL. A biomechanical comparison of elite swimmers start performance using the traditional track start and the new kick start. *BMS*. 2010;11:94-6.
13. Rice PE, Goodman CL, Capps CR, Triplett NT, Erickson TM, McBride JM. Force-and power-time curve comparison during jumping between strength-matched male and female basketball players. *Eur J Sport Sci*. 2017;17(3):286-93.
14. McMahon J, Rej S, Comfort P. Sex differences in countermovement jump phase characteristics. *Sports*. 2017;5(1):8.
15. Laffaye G, Wagner PP, Tombleson TI. Countermovement jump height: Gender and sport-specific differences in the force-time variables. *J Strength Cond Res*. 2014;28(4):1096-105.
16. Torrejon A, Balsalobre-Fernandez C, Haff GG, Garcia-Ramos A. The load-velocity profile differs more between men and women than between individuals with different strength levels. *Sports Biomech*. 2019;18(3):245-55.
17. Alegre L, Lara A, Elvira J, Aguado X. Muscle morphology and jump performance: Gender and intermuscular variability. *Journal of Sports Medicine and Physical Fitness*. 2009;49(3):320.
18. Mitchell LJ, Argus CK, Taylor KL, Sheppard JM, Chapman DW. The effect of initial knee angle on concentric-only squat jump performance. *Res Q Exerc Sport*. 2017;88(2):184-92.
19. Sheppard JM, Doyle TL. Increasing compliance to instructions in the squat jump. *J Strength Cond Res*. 2008;22(2):648-51.
20. Heishman A, Brown B, Daub B, Miller R, Freitas E, Bemben M. The influence of countermovement jump protocol on reactive strength index modified and flight time: Contraction time in collegiate basketball players. *Sports*. 2019;7(2):37.

21. Barlow H, Halaki M, Stuelcken M, Greene A, Sinclair PJ. The effect of different kick start positions on omega osb11 blocks on free swimming time to 15m in developmental level swimmers. *Human Movement Science*. 2014;34:178-86.
22. Tor E, Pease D, Ball K. The reliability of an instrumented start block analysis system. *J Appl Biomech*. 2015;31(1):62-7.
23. Rebutini VZ, Pereira G, Bohrer R, Ugrinowitsch C, Rodacki AL. Plyometric long jump training with progressive loading improves kinetic and kinematic swimming start parameters. *J Strength Cond Res*. 2014;30(9):2392-8.
24. Schilling BK, Falvo MJ, Chiu LZ. Force-velocity, impulse-momentum relationships: Implications for efficacy of purposefully slow resistance training. *Journal of sports science & medicine*. 2008;7(2):299.
25. Tor E, Pease D, Ball K. Key parameters of the swimming start and their relationship to start performance. *J Sports Sci*. 2015;33(13):1313-21.
26. Cossor JM, Slawson SE, Shillabeer B, Conway PP, West AA, editors. Are land tests a good predictor of swim start performance? ISBS-Conference Proceedings Archive; 2011.
27. Garcia-Ramos A, Tomazin K, Feriche B, Strojnik V, de la Fuente B, Arguelles-Cienfuegos J et al. The relationship between the lower-body muscular profile and swimming start performance. *J Hum Kinet*. 2016;50(1):157-65.
28. Vantorre J, Seifert L, Fernandes RJ, Boas JP, Chollet D. Kinematical profiling of the front crawl start. *Int J Sports Med*. 2010;31(1):16-21.
29. Andersen LL, Aagaard P. Influence of maximal muscle strength and intrinsic muscle contractile properties on contractile rate of force development. *Eur J Appl Physiol*. 2006;96(1):46-52.
30. Suchomel TJ, Nimphius S, Bellon CR, Stone MH. The importance of muscular strength: Training considerations. *Sports Med*. 2018;48(4):765-85.
31. Sole CJ, Mizuguchi S, Sato K, Moir GL, Stone MH. Phase characteristics of the countermovement jump force-time curve: A comparison of athletes by jumping ability. *The Journal of Strength & Conditioning Research*. 2018;32(4):1155-65.
32. Beckham GK, Suchomel TJ, Sole CJ, Bailey CA, Grazer JL, Kim SB et al. Influence of sex and maximum strength on reactive strength index-modified. *J Sports Sci Med*. 2019;18(1):65-72.
33. Breed RV, McElroy G. A biomechanical comparison of the grab, swing and track starts in swimming. *J Hum Mov Stud*. 2000;39:277-93.
34. Takeda T, Sakai S, Takagi H, Okuno K, Tsubakimoto S. Contribution of hand and foot force to take-off velocity for the kick-start in competitive swimming. *J Sports Sci*. 2017;35(6):565-71.
35. Vantorre J, Chollet D, Seifert L. Biomechanical analysis of the swim-start: A review. *J Sports Sci Med*. 2014;13(2):223-31.
36. West DJ, Owen NJ, Cunningham DJ, Cook CJ, Kilduff LP. Strength and power predictors of swimming starts in international sprint swimmers. *J Strength Cond Res*. 2011;25(4):950-5.
37. Aagaard P, Simonsen EB, Andersen JL, Magnusson P, Dyhre-Poulsen P. Increased rate of force development and neural drive of human skeletal muscle following resistance training. *Journal of applied physiology*. 2002;93(4):1318-26.
38. Jakobsen MD, Sundstrup E, Randers MB, Kjær M, Andersen LL, Krstrup P et al. The effect of strength training, recreational soccer and running exercise on stretch-shortening cycle muscle performance during countermovement jumping. *Human movement science*. 2012;31(4):970-86.
39. Rejman M, Bilewski M, Szczepan S, Klarowicz A, Rudnik D, Mackala K. Assessing the impact of a targeted plyometric training on changes in selected kinematic parameters of the swimming start. *Acta Bioeng Biomech*. 2017;19(2):149-60.
40. Breed RV, Young WB. The effect of a resistance training programme on the grab, track and swing starts in swimming. *J Sports Sci*. 2003;21(3):213-20.

Figure 1

Overview of the camera set-up and instrumented starting block.




Figure 1. Overview of the camera set-up and instrumented starting block (Kistler Group, 2019)

Figure 2

Relationship between concentric impulse (N.s) against time to 5 m (s) across females and males. Poly means polynomial regression to order 2, i.e. quadratic.

The grey dotted line and diamond markers represent the linear relationship between concentric impulse (N.s) and time to 5 m in females. The dashed line with circular markers represents the quadratic relationship between concentric impulse (N.s) and time to 5 m in males.




Figure 2. Relationship between concentric impulse (N.s) against time to 5 m (s) across females and males. Poly means polynomial regression to order 2, i.e. quadratic.

Figure 3

Relationship between concentric impulse (N.s) against time to 15 m (s) across females and males. Poly means polynomial regression to order 2, i.e. quadratic.

The grey dotted line and diamond markers represent the linear relationship between concentric impulse (N.s) and time to 15 m in females. The dashed line with circular markers represents the quadratic relationship between concentric impulse (N.s) and time to 15 m in males.




Figure 3. Relationship between concentric impulse (N.s) against time to 15 m (s) across females and males. Poly means polynomial regression to order 2, i.e. quadratic.

Table 1 (on next page)

Physical characteristics of participants.

Table 1: Physical characteristics of participants

Variables	Males		Females	
	5 m (<i>n</i> = 38)	15 m (<i>n</i> = 26)	5 m (<i>n</i> = 34)	15 m (<i>n</i> = 24)
Age (years)	21.0 ± 3.1	21.2 ± 3.2*	20.1 ± 3.2	19.2 ± 3.2
Body mass (kg)	76.7 ± 10.2**	76.5 ± 11.0**	64.8 ± 8.4	64.2 ± 8.4
Height (m)	1.83 ± 0.08**	1.85 ± 0.08**	1.73 ± 0.06	1.73 ± 0.06
Time to 5 m (s)	1.48 ± 0.09**		1.65 ± 0.08	
Time to 15 m (s)		6.4 ± 0.44**		7.3 ± 0.5

All data is presented as means and standard deviations. **p* < 0.05; ***p* < 0.001 between males and females.

Table 2(on next page)

List of 15 most influential potential predictors of swim start performance identified from the PCA and their correlations with the principal components.

- 1 Table 2: List of 15 most influential potential predictors of swim start performance identified from the
- 2 PCA and their correlations with the principal components.

Potential predictors	Principal Component		
	PC1	PC2	PC3
Variation explained for each component	67.5 %	17.1 %	8.5 %
Bodyweight (BW)	-0.71	0.11	0.68
Concentric impulse	-0.88	0.31	0.34
Concentric mean force	-0.91	-0.09	0.39
Concentric mean power	-0.94	0.13	0.14
Concentric peak force	-0.92	-0.15	0.32
Concentric rate of power development (RPD)	-0.93	-0.31	0.04
Force at peak power	-0.92	-0.05	0.33
Peak power	-0.95	0.24	0.14
Reactive strength index modified (RSI _{mod})	-0.90	-0.12	-0.20
Take-off peak force	-0.92	-0.15	0.32
Concentric peak velocity	-0.77	0.55	-0.29
Concentric rate of force development (RFD) BW	-0.59	-0.75	-0.15
Concentric RFD	-0.72	-0.66	0.05
Jump height (impulse-momentum)	-0.75	0.56	-0.31
Velocity at peak power	-0.68	0.66	-0.27

3

Table 3(on next page)

Multiple linear regression models to predict swim start time (s) to 5 m performance in females, males and both females and males combined.

1 Table 3: Multiple linear regression models to predict swim start time (s) to 5 m performance in females,
2 males and both females and males combined.

		% contribution	Beta coefficient (95% CI)	<i>p</i> -value
All	Concentric Impulse (N.s)	70.4	-0.002 (-0.002 to -0.001)	< 0.001
	Sex (Females)	5.4	0.065 (0.028 to 0.102)	0.001
	RSImod (m/s)	1.5	-0.084 (-0.164 to -0.004)	0.040
	Constant		1.882 (1.790 to 1.974)	< 0.001
	<i>R</i>² (SEE)		0.773 (0.059)	
Females	Concentric Impulse (N.s)	51.6	-0.003 (-0.004 to -0.002)	< 0.001
	RSImod (m/s)	9.5	-0.209 (-0.315 to -0.104)	< 0.001
	Concentric Mean Power (W)	7.8	0.0002 (0.00004 to 0.0003)	0.010
	Constant		2.103 (1.986 to 2.219)	< 0.001
	<i>R</i>² (SEE)		0.689 (0.047)	
Males	Concentric Impulse (N.s)	53.6	-0.010 (-0.015 to -0.005)	< 0.001
	Concentric Impulse² (N.s)²	12.3	0.00002 (0.00001 to 0.00003)	0.001
	Constant		2.645 (2.167 to 3.124)	< 0.001
	<i>R</i>² (SEE)		0.659 (0.055)	

3 SEE = standard error of estimate

4

Table 4(on next page)

Multiple linear regression models to predict swim start time (s) to 15 m performance in females, males and both females and males combined.

1 Table 4: Multiple linear regression models to predict swim start time (s) to 15 m performance in females,
2 males and both females and males combined.

		% contribution	Beta coefficient (95% CI)	p value
All	Concentric Impulse (N.s)	76.1	-0.008 (-0.011 to -0.004)	< 0.001
	Age (years)	3.5	-0.052 (-0.087 to -0.018)	0.004
	Sex (female)	3.0	0.362 (0.151 to 0.572)	0.001
	Constant		9.074 (8.503 to 9.646)	< 0.001
	R² (SEE)		0.826 (0.278)	
Females	Concentric Impulse (N.s)	65.1	-0.030 (-0.041 to -0.020)	< 0.001
	Body mass (kg)	9.3	0.035 (0.006 to 0.064)	0.020
	Concentric RPD (W/s)	4.9	0.0002 (0. 00006 to 0.0003)	0.004
	RSImod (m/s)	4.8	-1.714 (-3.215 to -0.213)	0.027
	Constant		9.303 (8.398 to 10.208)	< 0.001
	R² (SEE)		0.841 (0.225)	
Males	Concentric Impulse (N.s)	66.6	-0.033 (-0.058 to -0.008)	0.011
	Age (years)	9.4	-0.048 (-0.086 to -0.010)	0.016
	Concentric Impulse² (N.s)²	4.7	0.00007 (0.000007 to 0.0001)	0.031
	Constant		11.188 (8.975 to 13.401)	< 0.001
	R² (SEE)		0.807 (0.205)	

3 SEE = standard error of estimate

4