Plant-Insect interactions from Middle Triassic (late Ladinian) of Monte Agnello (Dolomites, N-Italy) - Initial pattern and response to abiotic environmental perturbations

Torsten Wappler, Evelyn Kustatscher, Elio Dellantonio

The Paleozoic–Mesozoic transition is characterized by the most massive extinction of the Phanerozoic. Nevertheless, an impressive adaptive radiation of herbivorous insects occurred on gymnosperm-dominated floras not earlier than during the Middle to Late Triassic, penecontemporaneous with similar events worldwide, all which exhibit parallel expansions of generalized and mostly specialized insect herbivory on plants, expressed as insect damage on a various plant organs and tissues. The flora from Monte Agnello is distinctive, due to its preservation in subaerially deposited pyroclastic layers with exceptionally preserved details. Thus, the para-autochthonous assemblage provides insights into environmental disturbances, caused by volcanic activity, and how they profoundly affected the structure and composition of herbivory patterns. These diverse Middle Triassic biota supply extensive evidence for insect herbivore colonization, resulting in specific and complex herbivory patterns involving the frequency and diversity of 20 distinctive damage types (DTs). These DT patterns show that external foliage feeders, piercer-and-suckers, leaf miners, gallers, and oviposition culprits were intricately using almost all tissue types from the dominant host plants of voltzialean conifers (e.g., *Voltzia*), horsetails, ferns (e.g., *Neuropteridium*, *Phlebopteris*, *Cladophlebis* and *Thaumatopteris*), seed ferns (e.g., *Scytophyllum*), and cycadophytes (e.g., *Bjuvia* and *Nilssonina*).

2 For PeerJ

3

4 **Plant–Insect interactions from Middle Triassic (late Ladinian) of Monte Agnello (Dolomites, N-**
5 **Italy) – Initial pattern and response to abiotic environmental perturbations**

6

7 Torsten Wappler¹, Evelyn Kustatscher^{2,3} & Elio Dellantonio⁴

8

9 ¹Steinmann Institute, University of Bonn, Nussallee 8, 53115 Bonn, Germany, e-mail: twappler@uni-
10 bonn.de;

11 ²Naturmuseum Südtirol, Bindergasse 1, 39100 Bozen/Bolzano, Italy, e-mail:
12 Evelyn.Kustatscher@naturmuseum.it;

13 ³Department für Geo- und Umweltwissenschaften, Paläontologie und Geobiologie, Ludwig-
14 Maximilians-Universität and Bayerische Staatssammlung für Paläontologie und Geobiologie, Richard-
15 Wagner-Straße 10, 80333 München, Germany;

16 ⁴Museo Geologico delle Dolomiti, Piazza SS. Filippo e Giacomo 1, 38037 Predazzo (TN), e-mail:
17 Elio.Dellantonio@muse.it.

18

19

20 INTRODUCTION

21 Continental arthropods and vascular plants have been major elements of terrestrial ecosystems
22 worldwide for nearly 400 million years and their varied ectophytic and endophytic associations can
23 provide a unique and direct record of the plant-insect interactions in the past (e.g., Labandeira &
24 Currano 2013). In 2006, Labandeira proposed four pulses of herbivore expansion, where the observed
25 Palaeozoic arthropod herbivory patterns – covering the first two phases – are mainly expressed by

26 damage patterns caused by mites and apterygote and/or basal pterygote herbivores on pteridophyte and
 27 basal gymnospermous plant hosts and are profoundly different from those that originated after the end-
 28 Permian mass extinction (Labandeira 2006a; Labandeira 2006b). Preliminary work on plant-insect
 29 interactions from early to late Permian floras of the Southern Alps indicates a moderately diverse
 30 pattern of damage occurring in a variety of habitats prior to the P-Tr crisis (pers. observation T.
 31 Wappler, 2013), penecontemporaneous with similar events in U.S. Southwest (e.g., Schachat et al.
 32 2014), Gondwana (Adami-Rodrigues et al. 2004a; Adami-Rodrigues et al. 2004b; Cariglino &
 33 Gutiérrez 2011; Gallego et al. 2014; Iannuzzi & Labandeira 2008; Prevec et al. 2009; Slater et al.
 34 2012), or Cathaysia (e.g., Glasspool et al. 2003). Nevertheless, herbivory expansion 2 was profoundly
 35 disrupted by environmental perturbations at the P-Tr boundary. The Early Triassic has been
 36 traditionally viewed as an unusual time marked by suppressed origination rates and low diversity (e.g.,
 37 Benton & Emerson 2007) generally attributed to the effects of extreme environmental conditions
 38 inflicted on Early Triassic ecosystems (e.g., Looy et al. 1999; Grauvogel-Stamm & Ash 2005;
 39 Roopnarine et al. 2007; Tong et al. 2007) but taphonomical biases cannot be excluded at least for
 40 European floras (Kustatscher et al. 2014b). In general, records of Early Triassic insects or of insect
 41 damage on plants are scant worldwide (comp. Table 1), so little is known about the mechanics and
 42 timing of diversification of this ecologically important group following the end-Permian mass-
 43 extinction event (Kustatscher et al. 2014b; Labandeira and Currano, 2013). Shcherbakov (2008a) even
 44 concluded that the entire class of insects was strongly reduced in diversity at the P-Tr boundary but
 45 following the end-Permian biotic crisis insect faunas already contained many elements common to
 46 modern insects (e.g., Aristov et al. 2013; Béthoux et al. 2005; Shcherbakov 2008b; Lukashevich et al.
 47 2010; Żyła et al. 2014; Haig et al. 2015) building the nucleus for the onset of the third pulse of
 48 herbivore expansion, coupled with an impressive adaptive radiation of herbivorous insects. Their
 49 associations with plants became significantly diverse being major elements for keystone communities

50 in terrestrial ecosystems worldwide (e.g., Ash 2014; Grauvogel-Stamm & Kelber 1996; Kustatscher et
 51 al. 2014b; Labandeira 2006a; Labandeira 2006b; Labandeira & Currano 2013; McLoughlin 2011;
 52 Moisan et al. 2012; Pott et al. 2008; Scott et al. 2004). Simultaneously, a major three-phased floral
 53 change has been proposed for Europe and probably worldwide (e.g., Grauvogel-Stamm & Ash 2005).
 54 The first stage lasted from the Induan to early Anisian, which in Europe is characterized by a
 55 “survival” interval dominated by the lycopsid *Pleuromeia* Corda ex Giebel 1853 and conifers coupled
 56 with relatively low levels of plant-insect interactional diversity (Kustatscher et al. 2014b); this is
 57 followed by a "recovery" interval characterized by the resurgence of lycopsids, sphenophytes, ferns,
 58 cycadophytes, conifers, ginkgophytes and seed ferns. The second stage occurred from the late Anisian
 59 to the Carnian. The third covers the Norian and Rhaetian stages, which is pivotal to understanding the
 60 evolution of trophically modern ecosystems (e.g., Benton 2010; Labandeira 2006b; Labandeira &
 61 Currano 2013).
 62 Thus, the late Middle Triassic (Ladinian) floras of the Dolomite Region in the Southern Alps of
 63 northeastern Italy provide an intriguing window into the early evidence for Herbivore Expansion 3.
 64 Ladinian floras from the Dolomites have been extensively studied in recent years (e.g., Kustatscher et
 65 al. 2014a; Kustatscher & Van Konijnenburg-van Cittert 2005 and references therein), evidencing a
 66 dominance of conifers (*Voltzia*, *Pelourdea*), while cycadophytes, seed ferns, ferns, horsetails, and
 67 lycopsids are much rarer. Nevertheless, the flora from Monte Agnello is markedly distinct from other
 68 Ladinian floras of the Dolomites by its higher diversity and abundance in cycadophytes, seed ferns and
 69 ferns. It is currently the best documented and most diverse late Middle Triassic biota in the Alps
 70 documenting a rich vascular plant record, including moderate levels of external foliage feeding,
 71 piercing-and-sucking, galling, and ovipositional damage.
 72 Of particular importance, from a taphonomic viewpoint, the Dolomites were subject to significant
 73 volcanic activity, beginning in the late Ladinian. Consequently, conditions for exceptional preservation

were high. Although most of the volcanic complexes were submarine, locally, such as in the area of Predazzo, subaerial eruptive centers existed (Hoernes 1912; Leonardi 1967), which alter the natural environment to variable extents and initiate very different effects on community composition, structure, function, and successional turnover on local and regional scales (e.g., Walker & Wardle 2014). This makes the Monte Agnello ideal for examining the response that such environmental perturbation had on community structures and offers the possibility to study the ecological expansion of interactional diversity recorded from the varied habitats.

Geological and paleontological setting

Monte Agnello (Fig. 1) represents an area that was marginally influenced by the Ladinian volcanic activity of the Predazzo volcano and is characterized by a well-preserved stratigraphic succession (e.g., Kustatscher et al. 2014a). The 250 meter thick volcanic succession is composed of “explosion breccia” at the base, followed by lava breccia, and alternations of lava flows and tuffs (Calanchi et al. 1977; Calanchi et al. 1978; Lucchini et al. 1982). The “explosion breccia” comprises lithic fragments (calcareous, volcanic and metamorphic fragments, clastic rocks, isolated crystals), related to the Permo-Triassic volcano-sedimentary succession and the metamorphic basement (Vardabasso 1930). The lithic fragments of the breccia are bound by carbonate and/or chlorite-serpentine cement (Calanchi et al. 1977). The thickness of this “explosion breccia” varies between 25 meters at Monte Agnello and 10 meters at Censi. The volcanic succession accumulated mostly in a subaerial environment, and is related to explosive phreatic activity (e.g., bomb sags, antidunes, accretionary lapilli; Calanchi et al. 1977; Lucchini et al. 1982).

The flora is preserved in the tuff lenses at the base of the “explosion breccia” of the volcanic succession at Predazzo (Kustatscher et al. 2014a), which hinders an appropriate stratigraphic correlation between the different sites. Considering that they are related to one or perhaps a few

phreatomagmatic events within the restricted time frame of late Ladinian volcanism the possible time difference between the single localities is however very reduced. The flora is composed of a large number of fronds, stems and reproductive organs of sphenophytes, ferns, seed ferns, cycadophytes and conifers. Due to the preservation in tuff layers, the organic material is missing and sometimes the remains are preserved only as impressions. Several stem fragments belong to the sphenophytes. The ferns are represented by Osmundaceae (*Neuropteridium elegans* (Brongniart) Schimper in Schimper and Schenk, 1879), Matoniaceae (*Phlebopteris fiemmensis* Kustatscher et al., 2014a) and Dipteridaceae (*Thaumatopteris* sp.). For the latter two families it is the oldest fossil occurrence to date for the Northern Hemisphere. Additional ferns of unknown botanical affinity are *Cladophlebis ladinica* Kustatscher et al., 2014a, *Cladophlebis* sp. (Osmundaceae and/or the Dicksoniaceae) and *Chiropteris monteagnellii* Kustatscher et al., 2014a (Dipteridaceae?). The seed ferns are represented by leaf fragments of *Scytophyllum bergeri* Bornemann, 1856. The cycadophyte leaf fragments probably belong to the genera *Bjuvia* Florin, 1933, *Taeniopteris* Brongniart, 1828 and/or *Macrotaeniopteris* Schimper, 1869 as well as *Nilssonia* Brongniart, 1828 and *Apoldia* Wesley, 1958. The conifers are represented by shoots of *Voltzia* Brongniart, 1828 and *Pelourdea* Seward, 1917 leaves. These plants grew probably during a humid spell, recently proposed for the late Ladinian of the Dolomites (Preto et al. 2010 and references therein).

MATERIAL AND METHODS

Data collection

Fossil plant assemblages were quantitatively censused from multiple sites at the base of the „explosive breccia“, that crops out on the northwestern slope of Monte Agnello – Censi, overlying a carbonate platform of late Anisian to Ladinian age (Sciliar Dolomite). About 684 specimens have been collected from eight distinctive sites denoted by the prefixes MA 1 – MA 8 (Fig. 1; Table 2). Sample size ranges

from 2 to 244 plant remains, depending primarily on the quality and accessibility of the fossils. For the quantitative study, each identifiable plant fossil was counted. Of the plant fossil specimens collected at Monte Agnello, all that were adequately preserved and exceeded a minimum size of 0.5 cm² were examined for insect damage. Parts and counterparts were matched whenever possible to avoid duplication. When possible, all specimens were assigned to a known species or plant morphotype. All analyzed specimens are housed at the Museo Geologico delle Dolomiti, Predazzo. Specimens occurring on the same rock slab are identified by different letters following the catalogue number whereas capital letters indicate parts and counterparts of the same specimen.

The most recent approach toward understanding the patterns of herbivory in the fossil record involves quantification of both the richness and intensity of insect damage (Wilf & Labandeira 1999; Labandeira et al. 2002, 2007; Kustatscher et al. 2014b). The richness of herbivory is determined first by establishing a classification system of distinctive, diagnosable damage types, or DTs, that can be used generally in studies of herbivore damage to plants. DTs then are grouped into functional feeding groups (FFG). Eight functional feeding groups are present in the Monte Agnello flora [(i) external foliage feeding, subdivided into hole, margin, surface feeding and skeleotization; (ii) piercing and sucking; (iii) oviposition, though not truly a feeding interaction but rather egg-laying that leaves a significant record of plant damage; (iv) mining and (v) galling]. To date, over 290 fossil DTs have been identified (pers. comm. C.C. Labandeira). Finally the DTs are ranked by their host specificity (HS), ranging from 1 for generalists to 3 for high host-plant specialization, which then allows non-generalized DTs (e.g., those with HS of 2 and 3) to be analyzed separately.

Each foliar element was photographed using a Canon EOS 30D camera with a Canon EF-S 60 mm f/2.8 macro lens or a Nikon Coolpix E4500. All photographs were optimized using Adobe Photoshop CS6 and Adobe Lightroom 5.

Institutional abbreviations

MGP, Museo Geologico delle Dolomiti, Predazzo. Specimens occurring on the same rock slab are identified by different capital letters following the catalogue number.

Quantitative analysis

Quantitative analyses of insect damage were done using R version 3.1.0 (www.r-project.org). For damage diversity analyses, sample size was standardized by selecting random subsets of foliar elements without replacement and calculating the damage diversity for the subsample. Subsets of the data were subjected to rarefaction using an analytic method detailed below, which extends the solution found by Wappler et al. (2012) to cases where individuals may belong to multiple classes and allows the explicit reconstruction of probability distributions for the rarefied sample (Heck et al. 1975). This process was repeated 5000 times, and the results were averaged to obtain the standardized damage diversity for the bulk flora and four single sub-localities (MA1, MA5, MA7, MA8). The remaining sub-localities were removed from the census because the target sample size of at least 40 specimens was not reached. The standard deviations (SD) for the resamples were calculated to provide sample error bars.

RESULTS

Damage on the bulk Monte Agnello flora

Of the 684 plant remains examined from the Monte Agnello flora, 83, or 12.13 %, exhibit some sort of damage represented by 20 different damage types. The taxa or morphotypes examined were represented by foliage, axes, stem fragments, fructifications, and dispersed seeds (Table 2). A total of 95 damage type occurrences were observed throughout the bulk flora: 45 on cycadophytes (representing 36.5% of all specimens), 37 on seed ferns (8.0%), ten on conifers (44.3%), and three on

ferns (7.6%) (Table 3), suggesting that selective feeding by insect herbivores preferentially targeted particular seed plants. This pattern of selectivity was also recognized within the early late Permian (Wuchiapingian) of the Gröden/Val Gardena Sandstone from the Bletterbach Gorge of the Dolomites (Northern Italy) (pers. observation T. Wappler, 2013). Herbivory recorded for the Monte Agnello sites represents nearly all of the fundamental modes of herbivory, excluding fungal infection, which was not observed (see Gunkel & Wappler 2015). Multiple DTs or functional feeding groups were only recorded in 1.6 % of the plant remains whereas the majority were only damaged in one way (~11 %). Seven distinctive functional feeding groups have been detected on the foliar elements from Monte Agnello, most of which occur on particular plant hosts. Types of the external foliage feeding constitute 78.9 % of all DT occurrences and preferentially occurred on the seed fern *S. bergeri* and consists of the exophytic consumption of live plant tissues, subdivided into skeletonization and margin-, hole- and surface feeding; this is the most common ensemble of Triassic damage types (Labandeira & Prevec 2014; pers. observation T. Wappler, 2013) (Fig. 2). Those of the galling FFG provided 11.5 % of all DT occurrences and are more or less evenly distributed among conifers, ferns and seed ferns (Figs. 3C-D, G). Gallings represents the most biologically complex of all major interactions, and represents arthropod-induced abnormal cell proliferation that can occur on all major plant organs (e.g., Kustatscher et al. 2014b; Scott et al. 2004); examples are widely known (e.g., Stone & Schönrogge 2003). Oviposition, though not a feeding interaction, comprised 5.2 % of all DT occurrences; examples are common (Ghosh et al. 2015; McLoughlin 2011) (Figs. 3E-F, I). Minor levels of insect damage were present for piercing-and-sucking (2.1 % of all DT occurrences; Fig. 3J) and mining (1.1 %; Fig. 3H) FFGs. Leafminers construct distinct leaf mines, most of which are quite conspicuous and represent a form of endophagous herbivory in which a herbivore targets and feeds on fluid tissues such as phloem, mesophyll or epidermal cell protoplasts (Sinclair & Hughes 2010); examples are uncommon and the possible mining structure on

the pteridosperm *Scytophyllum bergeri* (Fig. 3H) indicates that the origin and diversification of the leaf-mining habit occurred about 92 million years before the first appearance of fossil angiosperms (Ash 1997; Gnaedinger et al. 2014; Kustatscher et al. 2014b; McLoughlin 2011; Moisan et al. 2012; Pott et al. 2008).

Damage on individual species

Among the 28 taxa represented at Monte Agnello less than half indicate some kind of damage, whereas, three – *Scytophyllum bergeri*, *Bjuvia* cf. *dolomitica* and *Nilssonina* sp. – are the most herbivorized taxa (71,6 % of all DT occurrences) but only representing one-third of the flora (Table 2). The most abundant plant species are the conifers *Voltzia* sp. (Fig. 3C) and *Voltzia* sp. 2, which have the lowest damage frequency (2.44–2.94 %) of the common Monte Agnello taxa. Ferns are nearly equally as diverse as the seed ferns but damage frequency is at least ten times less abundant than among the seed ferns (Table 3; Fig. 3D). Sphenophytes displayed no signs of insect-mediated herbivory but the small number of sampled leaves open the possibility that more collecting and study may yet reveal damage to this group also.

Damage at distinct sub-localities

Plant material is generally preserved at the base of the “explosion breccia” at an angle to the bedding rather than compacted into a single horizon. Transport distance, therefore, must have been short, and burial was likely rapid. Thus the fossil leaf assemblages must be considered as para-autochthonous (e.g., Hanley et al. 2007). Minimal transport allows us to document considerable changes in species composition and insect folivory over short distances and recognize possible heterogeneity in the structure and composition of the source plant communities and their associated herbivores. Large-scale disturbances may profoundly alter the composition and structure of plant communities and are rarely

uniform in their influence on vegetation (Kustatscher et al. 2014a). Variations of floral composition and insect herbivore damage at the four sub-localities (MA1, MA5, MA7, MA8) censused are shown in Fig. 4 and Table 4. MA1 has the highest floral diversity (22 spp.), followed by MA5 (16 spp.) and MA7 (15 spp.). MA8 is an extremely low-diversity flora (7 spp.). Interestingly, all sites are strongly dominated by a single plant group representing in all cases over half of the characteristic plant material at that site. The most abundant plant lineage at MA5–MA8 is conifers, whereas at MA1 57% of the taphocoenosis is composed of cycadophytes. However, when analyzing herbivory on individual host groups at the four sub-localities, total damage frequency and external foliage feeding is overwhelmingly found on seed-fern hosts (Fig. 4), except MA7 where the preferred host-plants are cycadophytes.

228

DISCUSSION

Volcanogenic deposits can preserve spatio-temporal biotic patterns at levels of resolution not commonly represented in the fossil record. Consequently, the plant-insect assemblages recognized in this study appear compositionally and ecologically unique (Currano et al. 2011; Dale et al. 2005). The para-autochthonous early late Ladinian flora of the Monte Agnello (Dolomites, N-Italy) offers insights into the patterns of arthropod herbivory during the beginning of the third pulse of herbivore expansion (*sensu* Labandeira 2006a; Labandeira 2006b; Labandeira & Currano 2013: fig. 1). It also provides insights into the way herbivores responded to environmental perturbation and the reorganization of community structure. Even though our data are preliminary the palaeoecological and temporal setting of the early late Ladinian flora of the Monte Agnello in the Dolomites supports three major conclusions that parallel those drawn from data known from intensively studied Gondwanan sites.

(1) *Dominance of seed plant herbivory*. The dominance of seed plant herbivory by local arthropod herbivores, particularly that known since the Permian across western Euamerica (e.g., Schachat

et al. 2014), Europe (Geyer & Kelber 1987; Bernardi et al. in prep.), Cathaysia (Glasspool et al. 2003), and in the extensive glossopterid-dominated floras across Gondwana (e.g., Cariglino & Gutiérrez 2011; McLoughlin 2011; Prevec et al. 2009) is also a conspicuous component of the late Anisian to Ladinian environments. This documents the persistence of the preferential targeting of selected groups of seed plants, like the cycadophytes in Monte Agnello, particularly by external foliage feeders. The pattern could be interpreted to support Feeny's apparency hypothesis (Feeny 1976), as seed plants were the most abundant and conspicuous, and therefore would have been the most apparent to herbivore consumption. However, for the Monte Agnello data, a more likely explanation favors increased herbivory on particular plants due to the anatomy of their leaves, suggesting that particular physical traits, like the scleromorphic structures of conifer taxa, reduce the palatability and digestibility of such plant material or act as a deterrent when more palatable plants are available (Labandeira & Anderson 2005).

(2) *Increase of interactional diversity and rise of the leaf-mining habit.* There is an increase in plant-insect interactional diversity during the Early to Late Triassic in eastern Euamerica and Gondwana regions (e.g., Kustatscher et al. 2014b; Scott et al. 2004), coupled with an increase in the diversity of FFGs, DTs, and associated herbivore behaviors observed at Monte Agnello, compared to insect damage from earlier known floras (e.g., Kustatscher et al. 2014b). Of particular importance is the presence of the leaf-mining habit in which holometabolous insect larvae consume the inner parenchymal, epidermal, vascular, or other tissues of a plant, leaving the outer wall of the epidermis undamaged (Hering 1951). The earliest documented leaf-mining fossil records have been reported from Kyrgyzstan, Austria, Australia and South Africa in deposits of Middle to Late Triassic age (comp. Table 1).

(3) *Volcanic activity and site-specific habitat differences.* The data presented here show that volcanogenic deposits are valuable for the creation and preservation of in situ sequential stages

of biotic change not commonly represented in the fossil record. These episodic volcanic activities directly influenced the evolution of the environment, spatial structure and temporal dynamics of the plant community and the herbivores associated with the plants, resulting in vegetational heterogeneity had impact on both the likelihood and strength of interactions between plants and insect herbivores (e.g., Agrawal et al. 2006; Currano et al. 2011). Therefore the heterogeneity among the sub-localities indicates that volcanic disturbance caused compositional and structural changes in the ecosystem during the time it occupied the site, which explain variations in plant physiognomy, plant and insect herbivore composition, and the overall paleoecology (Table 4). This conclusion is supported by (1) the spatial variability in the percentage of herbivorized plant host specimens, (2) the elevated number of DTs on each host plant, and (3) the differences in evenness and the relative abundance distributions of damage among the single sub-localities.

These conclusions warrant further verification from investigations of additional new sites to clarify patterns of arthropod herbivory during this crucial period of time where terrestrial ecosystems were beginning to become modern.

ACKNOWLEDGEMENTS

We thank Federica Angeli (Trento, Italy), Fulvio Boninsegna (Predazzo, Italy), Andrea Braitto (Daiano, Italy), Daniele Ferrari (Museo Geologico delle Dolomiti, Predazzo, Italy), Christian Fontana (Vigo di Fassa, Italy) and Guido Roghi (CNR and University of Padova, Italy) for their help during fieldwork and the preparation of the fossils. Special thanks goes to Daniele Ferrari for the photographs of the plants. The authors acknowledge Ellen Currano, University of Wyoming, an anonymous reviewer, and the editor William DiMichele for their constructive and encouraging comments.

290 REFERENCES

- 291 **Adami-Rodrigues K, de Souza PA, Iannuzzi R, and Pinto ID. 2004a.** Herbivoria em floras
292 gonduânicas do neopaleózoico do Rio Grande do Sul: análise quantitativa. *Revista Brasileira de*
293 *Paleontologia* **7**:93-102.
- 294 **Adami-Rodrigues K, Iannuzzi R, and Pinto ID. 2004b.** Permian plant-insect interactions from a
295 Gondwana flora of southern Brazil. *Fossils and Strata* **51**:106-125.
- 296 **Agrawal AA, Lau JA, and Hambäck PA. 2006.** Community heterogeneity and the evolution of
297 interactions between plants and insect herbivores. *The Quarterly Review of Biology* **81**:349-376.
- 298 **Anderson JM, and Anderson HM. 1983.** *Palaeoflora of southern Africa: Molteno Formation*
299 *(Triassic). Volume 1: Part 1. Introduction: Part 2. Dicroidium.* Rotterdam: Balkema.
- 300 **Anderson JM, and Anderson HM. 1985.** *Palaeoflora of Southern Africa: Prodrum of South*
301 *African Megaflores: Devonian to Lower Cretaceous: Rotterdam.* Rotterdam: Balkema.
- 302 **Anderson JM, and Anderson HM. 2003.** Heyday of the gymnosperms: systematics and biodiversity
303 of the Late Triassic Molteno fructifications. *Strelitzia* **15**:1-398.
- 304 **Aristov DS, Bashkuev AS, Golubev VK, Gorochoy AV, Karasev EV, Kopylov DS, Ponomarenko**
305 **AG, Rasnitsyn AP, Rasnitsyn DA, Sinitshenkova ND, Sukatsheva ID, and Vassilenko DV.**
306 **2013.** Fossil insects of the middle and upper Permian of European Russia. *Paleontological*
307 *Journal* **47**:641-832.
- 308 **Ash S. 1997.** Evidence of arthropod-plant interactions in the Upper Triassic of the southeastern United
309 States. *Lethaia* **29**:237-248.
- 310 **Ash S. 1999.** An Upper Triassic *Sphenopteris* showing evidence of insect predation from Petrified
311 Forest National Park, Arizona. *International Journal of Plant Sciences* **160**:208-215.
- 312 **Ash S. 2000.** Evidence of oribatid mite herbivory in the stem of a Late Triassic tree fern from Arizona.
313 *Journal of Paleontology* **74**:1065-1071.
- 314 **Ash S. 2001.** New cycadophytes from the Upper Triassic Chinle Formation of the southwestern United
315 States. *Paleobios* **21**:15-28.
- 316 **Ash S. 2005.** A new Triassic flora and associated invertebrate fossils from the basal beds of the Chinle
317 Formation, near Cameron, Arizona. *Paleobios* **25**:17-34.
- 318 **Ash S, and Savidge RA. 2004.** The bark of the Late Triassic *Araucarioxylon arizonicum* tree from the
319 Petrified Forest National Park, Arizona. *IAWA Journal* **25**:349-368.

- 320 **Ash SR. 2014.** Contributions to the Upper Triassic Chinle flora in the American southwest.
321 *Palaeobiodiversity and Palaeoenvironments* **94**:279-294.
- 322 **Benton MJ. 2010.** The origins of modern biodiversity on land. *Philosophical Transactions of the*
323 *Royal Society of London B Biological Sciences* **365**:3667-3679.
- 324 **Benton MJ, and Emerson BC. 2007.** How did life become so diverse? The dynamics of
325 diversification according to the fossil record and molecular phylogenetics. *Palaeontology*
326 **50**:23-40.
- 327 **Béthoux O, Papier F, and Nel A. 2005.** The Triassic radiation of the entomofauna. *Comptes Rendus*
328 *Palevol* **4**:609-621.
- 329 **Bornemann JG. 1856.** *Über organische Reste der Lettenkohlengruppe Thüringens - Ein Beitrag zur*
330 *Fauna und Flora dieser Formation.* Leipzig: W. Engelmann Verlag.
- 331 **Brongniart AT. 1828–1837.** *Histoire des végétaux fossiles, ou recherches botaniques et géologiques*
332 *sur les végétaux renfermés dans les divers couches du globe.* Paris, Amsterdam: I.
333 Dufour/d'Ocagne.
- 334 **Calanchi N, Lucchini F, and Rossi PL. 1977.** M. Agnello: un apparato vulcanico a condotto centrale
335 nelle Dolomiti. *Minerologica Petrografica Acta* **21**:221-229.
- 336 **Calanchi N, Lucchini F, and Rossi PL. 1978.** The Volcanic Rocks from the Mount Agnello Area
337 (Fiemme Valley, Italy): A contribution to the knowledge of the Mid-Triassic magmatism of the
338 Southern Alps. *Tschermaks mineralogische und petrographische Mitteilungen* **25**:131-143.
- 339 **Cariglino B, and Gutiérrez PR. 2011.** Plant-insect interactions in a Glossopteris flora from the La
340 Golondrina Formation (Guadalupian–Lopingian), Santa Cruz Province, Patagonia, Argentina.
341 *Ameghiniana* **48**:103-112.
- 342 **Creber GT, and Ash SR. 2004.** The Late Triassic *Schilderia adamanica* and *Woodworthia arizonica*
343 trees of the Petrified Forest National Park, Arizona, USA. *Palaeontology* **47**:21-38.
- 344 **Currano ED, Jacobs BF, Pan AD, and Tabor NJ. 2011.** Inferring ecological disturbance in the fossil
345 record: A case study from the late Oligocene of Ethiopia. *Palaeogeography,*
346 *Palaeoclimatology, Palaeoecology* **309**:242-252.
- 347 **Dale VH, Swanson FJ, and Crisafulli CM. 2005.** *Disturbance, survival, and succession:*
348 *understanding ecological responses to the 1980 eruption of Mount St. Helens.* New York,
349 Springer.

- 350 **Feeny P. 1976.** Plant apparency and chemical defense. In: Wallace JW, and Mansell RL, eds.
351 *Biochemical interaction between plants and insects*. New York: Plenum Press, 1-40.
- 352 **Feng Z, Su T, Yang JY, Chen YX, Wei HB, Dai J, Guo Y, Liu JR, and Ding JH. 2014.** Evidence
353 for insect-mediated skeletonization on an extant fern family from the Upper Triassic of China.
354 *Geology* **42**:407-410.
- 355 **Florin R. 1933.** Studien über die Cycadales des Mesozoicums nebst Erörterungen über die
356 Spaltöffnungsapparate der Bennettitales. *Kungliga Svenska Vetenskapsakademiens Handlingar,*
357 *ser. 3* **12**:1-119.
- 358 **Gallego J, Cúneo R, and Escapa I. 2014.** Plant–arthropod interactions in gymnosperm leaves from
359 the Early Permian of Patagonia, Argentina. *Geobios* **47**:101-110.
- 360 **Gallego OF, Gnaedinger S, Kirsten O, and Giovanelli S. 2003.** Primera cita de trazas fósiles de
361 insectos en hojas del Pérmico de Uruguay y Triásico de Chile. *Universidad Nacional Del*
362 *Nordeste, Comunicaciones Científicas y Tecnológicas Biológicas*, **B032**:1-4.
- 363 **Gallego OF, Gnaedinger S, Labandeira CC, Martins-Neto RG, and Kirsten O. 2004.** Permian and
364 Triassic insect traces on fossil leaves from Uruguay and Chile. *International Congress on*
365 *Ichnology ICHNIA Abstract Book* **1**:35.
- 366 **Geyer G, and Kelber K-P. 1987.** Flügel und Lebensspuren von Insekten aus dem Unteren Keuper
367 Mainfrankens. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen* **174**:331-355.
- 368 **Ghosh AK, Kar R, and Chatterjee R. 2015.** Leaf galls on *Dicroidium hughesii* (Feistmantel) Lele
369 from the Triassic of India—a new record. *Alcheringa* **39**:92-98.
- 370 **Giebel CG. 1853.** Referat über Spieker. *Zeitschrift für die gesamten Naturwissenschaften* **2**: 34.
- 371 **Glasspool I, Hilton J, Collinson ME, and Wang S-J. 2003.** Foliar herbivory in Late Palaeozoic
372 Cathaysian gigantopterids. *Review of Palaeobotany and Palynology* **127**:125-132.
- 373 **Gnaedinger SC, Adami-Rodrigues K, and Gallego OF. 2007.** Evidencias de trazas de oviposición de
374 insectos (Odonata) en hojas del Triásico de Chile. *Ameghiniana* **44**:94R.
- 375 **Gnaedinger SC, Adami-Rodrigues K, and Gallego OF. 2008.** Insect egg ovipositions on leaves from
376 the Upper Triassic from northern Chile. In: Boardman DR, editor. *Simpósio Brasileiro de*
377 *Paleobotânica e Palinologia Boletim de Resumos*. Florianópolis, SC, Brazil: Asociación
378 Latinoamericana de Paleobotánica y Palinología. p 86.

- 379 **Gnaedinger SC, Adami-Rodrigues K, and Gallego OF. 2014.** Endophytic oviposition on leaves
380 from the Late Triassic of northern Chile: Ichnotaxonomic, palaeobiogeographic and
381 palaeoenvironment considerations. *Geobios* **47**:221-236.
- 382 **Grauvogel-Stamm L, and Ash SR. 2005.** Recovery of the Triassic land flora from the end-Permian
383 life crisis. *Comptes Rendus Palevol* **4**:593-608.
- 384 **Grauvogel-Stamm L, and Kelber K-P. 1996.** Plant-insect interactions and coevolution during the
385 Triassic in western Europe. *Paleontologica Lombarda, NS* **5**:5-23.
- 386 **Gunkel S, and Wappler T. 2015.** Plant-insect interactions in the upper Oligocene of Enspel
387 (Westerwald, Germany), including an extended mathematical framework for rarefaction.
388 *Palaeobiodiversity and Palaeoenvironments*, doi: 10.1007/s12549-014-0176-6.
- 389 **Haig DW, Martin SK, Mory AJ, McLoughlin S, Backhouse J, Berrell RW, Kear BP, Hall R,**
390 **Foster CB, Shi GR, and Bevan JC. 2015.** Early Triassic (early Olenekian) life in the interior
391 of East Gondwana: mixed marine–terrestrial biota from the Kockatea Shale, Western Australia.
392 *Palaeogeography, Palaeoclimatology, Palaeoecology* **417**:511-533.
- 393 **Hanley ME, Lamont BB, Fairbanks MM, and Rafferty CM. 2007.** Plant structural traits and their
394 role in anti-herbivore defence. *Perspectives in Plant Ecology, Evolution and Systematics* **8**:157-
395 178.
- 396 **Heck KL, van Belle G, and Simberloff D. 1975.** Explicit calculation of the rarefaction diversity
397 measurement and the determination of sufficient sample size. *Ecology* **56**:1459-1461.
- 398 **Heer O. 1877.** *Die vorweltliche Flora der Schweiz*: Wurster.
- 399 **Hering EM. 1951.** *Biology of the leaf miners*. Dordrecht: Springer Netherlands.
- 400 **Hoernes R. 1912.** Zur Geologie von Predazzo. *Sitzungsberichte der königlichen Akademie der*
401 *Wissenschaft* **121**:3-31.
- 402 **Hsü J, Chu C-n, Chen Y, Tuan S-y, Hu Y-f, and Chu W-c. 1974.** New Genera and Species of the
403 Late Triassic Plants from Yungjen, Yunnan I. *Acta Botanica Sinica* **16**:266-278.
- 404 **Iannuzzi R, and Labandeira CC. 2008.** The oldest record and early history of insect folivory. *Annals*
405 *of the Entomological Society of America* **101**:79-94.
- 406 **Kelber K-P, and Geyer G. 1989.** Lebensspuren von Insekten an Pflanzen des Unteren Keupers.
407 *Courier Forschungsinstitut Senckenberg* **109**:165-174.

- 408 **Kustatscher E, Dellantonio E, and von Konijnenburg-van Cittert JHA. 2014a.** The ferns of the
 409 late Ladinian, Middle Triassic flora from Monte Agnello, Dolomites, N-Italy. *Acta*
 410 *Palaeontologica Polonica* **59**:741-755.
- 411 **Kustatscher E, Franz M, Heunisch C, Reich M, and Wappler T. 2014b.** Floodplain habitats of
 412 braided river systems: depositional environment, flora and fauna of the Solling Formation
 413 (Buntsandstein, Lower Triassic) from Bremke and Fürstenberg (Germany). *Palaeobiodiversity*
 414 *and Palaeoenvironments* **94**:237-270.
- 415 **Kustatscher E, and Van Konijnenburg-van Cittert JHA. 2005.** The Ladinian flora (Middle
 416 Triassic) of the Dolomites: Palaeoenvironmental reconstructions and palaeoclimatic
 417 considerations. *GeoAlp* **2**:31-51.
- 418 **Labandeira CC. 2006a.** The four phases of plant-arthropod associations in deep time. *Geologica Acta*
 419 **4**:409-438.
- 420 **Labandeira CC. 2006b.** Silurian to Triassic plant and hexapod clades and their associations: new data,
 421 a review, and interpretations. *Arthropod Systematics & Phylogeny* **64**:53-94.
- 422 **Labandeira CC, and Anderson JM. 2005.** Insect leaf-mining in Late Triassic gymnospermous floras
 423 from the Molteno Formation of South Africa. *Geological Society of America Abstracts with*
 424 *Programs* **37**:15.
- 425 **Labandeira CC, and Currano ED. 2013.** The Fossil Record of Plant-Insect Dynamics. *Annual*
 426 *Review of Earth and Planetary Sciences* **41**:287-311.
- 427 **Labandeira CC, Johnson KR, and Lang PJ. 2002.** Preliminary assessment of insect herbivory across
 428 the Cretaceous-Tertiary boundary: Major extinction and minimum rebound. In: Hartman JH,
 429 Johnson KR, and Nichols DJ, eds. *The Hell Creek Formation of the northern Great Plains*.
 430 Boulder, Colorado: Geological Society of America Special Paper, 297-327.
- 431 **Labandeira CC, and Prevec R. 2014.** Plant paleopathology and the roles of pathogens and insects.
 432 *International Journal of Paleopathology* **4**:1-16.
- 433 **Labandeira CC, Wilf P, Johnson KR, and Marsh F. 2007.** *Guide to Insects (and Other) Damage*
 434 *Types on Compressed Plant Fossils. Version 3.0.* Washington, D.C.: Smithsonian Institution.
- 435 **Leonardi P. 1967.** *Le Dolomiti: geologia dei monti tra Isarco e Piave.* Rovereto: Consiglio Nazionale
 436 delle Ricerche e della Giunta Provinciale di Trento.
- 437 **Linck O. 1949.** Fossile Bohrgänge (*Anobichnium simile* ngn sp.) an einem Keuperholz. *Neues*
 438 *Jahrbuch für Mineralogie, Geologie und Paläontologie B* **90**:180-185.

- 439 **Looy CV, Brugman WA, Dilcher DL, and Visscher H. 1999.** The delayed resurgence of equatorial
440 forests after the Permian–Triassic ecologic crisis. *Proceedings of the National Academy of*
441 *Sciences of the United States of America* **96**:13857-13862.
- 442 **Lucchini F, Rossi PL, and Simboli G. 1982.** Il magmatismo triassico dell’area di Predazzo. In:
443 Castellarin A, and Vai GB, eds. *Guida alla Geologia del Sudalpino centro-orientale*. Roma:
444 Società Geologica Italiana, Guide Geologiche Regionali, 221-229.
- 445 **Lukashevich ED, Przhiboro AA, Marchal-Papier F, and Grauvogel-Stamm L. 2010.** The oldest
446 occurrence of immature Diptera (Insecta), Middle Triassic, France. *Annales de la Société*
447 *Entomologique de France (Nouvelle série)* **46**:4-22.
- 448 **McLoughlin S. 2011.** New records of leaf galls and arthropod oviposition scars in Permian–Triassic
449 Gondwanan gymnosperms. *Australian Journal of Botany* **59**:156-169.
- 450 **Meller B, Ponomarenko AG, Vasilenko DV, Fischer TC, and Aschauer B. 2011.** First beetle elytra,
451 abdomen (Coleoptera) and a mine trace from Lunz (Carnian, Late Triassic, Lunz-am-See,
452 Austria) and their taphonomical and evolutionary aspects. *Palaeontology* **54**:97-110.
- 453 **Minello LF. 1994.** As “florestas petrificadas” da região de São Pedro do Sul e Mata, RS. III – análise
454 morfológica megascópica, afinidades e considerações paleoambientais. *Acta Geologica*
455 *Leopoldensia* **39**:75-91.
- 456 **Moisan P, Labandeira CC, Matushkina NA, Wappler T, Voigt S, and Kerp H. 2012.** Lycopsid-
457 arthropod associations and odonatopteran oviposition on Triassic herbaceous *Isoetites*.
458 *Palaeogeography Palaeoclimatology Palaeoecology* **344-345**:6-15.
- 459 **Nathorst AG. 1876.** Bidrag till Sveriges fossila Flora. *Kongliga Svenska Vetenskaps-Akademiens*
460 *Handlingar* **14**:1-82.
- 461 **Nathorst AG. 1878.** *Beiträge zur fossilen Flora Schwedens. Über einige rhätische Pflanzen von Pålsjö*
462 *in Schonen*. Stuttgart: E. Schweizerbart'sche Verlagshandlung (E. Koch).
- 463 **Pott C, Labandeira CC, Krings M, and Kerp H. 2008.** Fossil insect eggs and ovipositional damage
464 on bennettitalean leaf cuticles from the Carnian (Upper Triassic) of Austria. *Journal of*
465 *Paleontology* **82**:778-789.
- 466 **Preto N, Kustatscher E, and Wignall PB. 2010.** Triassic climates – State of the art and perspectives.
467 *Palaeogeography, Palaeoclimatology, Palaeoecology* **290**:1-10.

- 468 **Prevec R, Labandeira CC, Neveling J, Gastaldo RA, Looy CV, and Bamford M. 2009.** Portrait of
469 a Gondwanan ecosystem: A new late Permian fossil locality from KwaZulu-Natal, South
470 Africa. *Review of Palaeobotany and Palynology* **156**:454-493.
- 471 **Roopnarine PD, Angielczyk KD, Wang SC, and Hertog R. 2007.** Trophic network models explain
472 instability of Early Triassic terrestrial communities. *Proceedings of the Royal Society B:*
473 *Biological Sciences* **274**:2077-2086.
- 474 **Roselt G. 1954.** Ein neuer Schachtelhalm aus dem Keuper und Beiträge zur Kenntnis von
475 *Neocalamites meriani* Brongn. *Geologie* **3**:617-643.
- 476 **Rozefelds AC, and Sobbe I. 1987.** Problematic insect leaf mines from the Upper Triassic Ipswich
477 Coal Measures of southeastern Queensland, Australia. *Alcheringa* **11**:51-57.
- 478 **Schachat S, Labandeira CC, Gordon J, Chaney D, Levi S, Halthore MS, and Alvarez J. 2014.**
479 Plant–insect interactions from the Early Permian (Kungurian) Colwell Creek Pond, North-
480 Central Texas: the early spread of herbivory in riparian environments. *International Journal of*
481 *Plant Sciences* **175**:855-890.
- 482 **Schimper WP. 1869.** *Traité de Paléontologie végétale ou la flore du monde primitif dans ses rapports*
483 *avec les formations géologiques et la flore du monde actuel. Tome premier.* Paris: Bailliére J.B.
484 et Fils ed.
- 485 **Schimper WP, and Schenk A. 1879.** In: Zittel KA, editor. *Handbuch der Palaeontologie, Teil II.*
486 *Palaeophytologie. Lief. I.* Leipzig: R. Oldenbourg, 1-152.
- 487 **Scott AC, Anderson JM, and Anderson HM. 2004.** Evidence of plant-insect interactions in the
488 Upper Triassic Molteno Formation of South Africa. *Journal of the Geological Society, London*
489 **161**:401-410.
- 490 **Seward AC 1917.** *Fossil plants, III Pteridospermae, Cycadofilices, Cordaitales, Cycadophyta.*
491 Cambridge: University Press.
- 492 **Shcherbakov DE. 2008a.** On Permian and Triassic insect faunas in relation to biogeography and the
493 Permian–Triassic crisis. *Paleontological Journal* **42**:15-31.
- 494 **Shcherbakov DE. 2008b.** Insect recovery after the Permian/Triassic crisis. *Alavesia* **2**:125-131.
- 495 **Sinclair RJ, and Hughes L. 2010.** Leaf miners: The hidden herbivores. *Austral Ecology* **35**:300-313.
- 496 **Slater BJ, McLoughlin S, and Hilton J. 2012.** Animal–plant interactions in a Middle Permian
497 permineralised peat of the Bainmedart Coal Measures, Prince Charles Mountains, Antarctica.
498 *Palaeogeography, Palaeoclimatology, Palaeoecology* **363–364**:109-126.

- 499 **Stone GN, and Schönrogge K. 2003.** The adaptive significance of insect gall morphology. *Trends in*
500 *Ecology & Evolution* **18**:512-522.
- 501 **Strullu-Derrien C, McLoughlin S, Philippe M, Mørk A, and Strullu DG. 2012.** Arthropod
502 interactions with bennettitalean roots in a Triassic permineralized peat from Hopen, Svalbard
503 Archipelago (Arctic). *Palaeogeography, Palaeoclimatology, Palaeoecology* **348-349**:45-58.
- 504 **Tillyard RJ. 1922.** Mesozoic insects of Queensland. No. 9. Orthoptera, and additions to the
505 Protorthoptera, Odonata, Hemiptera and Plannipennia. *Proceedings of the Linnean Society of*
506 *New South Wales* **47**:447-470.
- 507 **Tong J, Zhang S, Zuo J, and Xiong X. 2007.** Events during Early Triassic recovery from the end-
508 Permian extinction. *Global and Planetary Change* **55**:66-80.
- 509 **Vardabasso S. 1930.** Carta geologica del territorio eruttivo di Predazzo e Monzoni nelle Dolomiti di
510 Fiemme e Fassa. Padova: R. Scuola d'Ingegneria.
- 511 **Walker LR, and Wardle DA. 2014.** Plant succession as an integrator of contrasting ecological time
512 scales. *Trends in Ecology & Evolution* **29**:504-510.
- 513 **Walker MV. 1938.** Evidence of Triassic insects in the Petrified Forest National Monument, Arizona.
514 *Proceedings of the United States National Museum* **85**:137-140.
- 515 **Wappler T, Labandeira CC, Rust J, Frankenhäuser H, and Wilde V. 2012.** Testing for the effects
516 and consequences of mid Paleogene climate change on insect herbivory. *PLoS ONE* **7**:e40744.
- 517 **Webb JA. 1982.** Triassic species of *Dictyophyllum* from eastern Australia. *Alcheringa* **6**:79-91.
- 518 **Wesley, A. 1958.** Contributions to the Knowledge of the flora of the Grey Limestones of Veneto, Part
519 II. A revision of the Flora fossilis formationis oolithicae of De Zigno. *Centro Studi Petr. Geol.*
520 *del CNR, Univ. Padova* **21**:1-57.
- 521 **Wilf P, and Labandeira CC. 1999.** Response of plant-insect associations to Paleocene-Eocene
522 warming. *Science* **284**:2153-2156.
- 523 **Żyła D, Wegierek P, Owocki K, and Niedźwiedzki G. 2013.** Insects and crustaceans from the latest
524 Early–early Middle Triassic of Poland. *Palaeogeography, Palaeoclimatology, Palaeoecology*
525 **371**:136-144.

526
527

528 Funding

529 The work has been supported by the Comune di Predazzo (Provincia di Trento). This study is part of
 530 the project “The Permian-Triassic ecological crisis in the Dolomites: extinction and recovery dynamics
 531 in Terrestrial Ecosystems” financed by the Promotion of Educational Policies, University and Research
 532 Department of the Autonomous Province of Bolzano – South Tyrol. T.W. is supported by a Heisenberg
 533 grant from the German Science Foundation (1496/8-1). The funders had no role in study design, data
 534 collection and analysis, decision to publish, or preparation of the manuscript.

535

536 **Competing Interests**

537 The authors declare there are no competing interests.

538 **Author Contributions**

- 539 • Torsten Wappler analyzed the data, wrote the paper, prepared figures, and reviewed drafts of the
 540 paper.
- 541 • Evelyn Kustatscher wrote the paper, prepared figures, and reviewed drafts of the paper.
- 542 • Elio Dellantonio contributed on the geology, reviewed drafts of the paper, and contributed with
 543 iconography.

Table 1 (on next page)

Arthropod damage on Triassic plants

List of published records of arthropod damage on Triassic plants.

2 Table 1. List of published records of arthropod damage on Triassic plants.

Study	Age	Formation and locality	Damage type
Nathorst (1876); Nathorst (1878)	Rhaetian (Late Triassic)	Pålsjö, Scania, Sweden	<ul style="list-style-type: none"> possible oviposition scars on <i>Podozamites</i>
Ghosh et al. (2015)	Norian / Rhaetian (Late Triassic)	Parsora Formation (Dhaurai Hill beds); South Rewa Gondwana Basin, central India	<ul style="list-style-type: none"> disc-like galls on <i>Dicroidium hughesii</i>
Walker (1938); Ash (1997); Ash (1999); Ash (2000); Ash (2001); Ash (2005); Ash & Savidge (2004); Ash (2014); Creber & Ash (2004)	Norian (Late Triassic)	Chinle Formation, Petrified Forest National Park, Arizona, USA	<ul style="list-style-type: none"> marginal and non-marginal feeding traces on <i>Cynepteris</i>, <i>Marcouia</i>, <i>Zamites</i>, <i>Sphenopteris</i>, <i>Macrotæniopteris</i>, <i>Dechellyia</i>, <i>Nilssoniopteris</i> possible oviposition scars and insect eggs on <i>Dechellyia</i>, <i>?Equisetites</i> coprolite-bearing borings in <i>Itopsidema</i>, <i>Araucarioxylon</i>, <i>Schilderia</i>
Adami-Rodrigues et al. (2008)	Norian (Late Triassic)	El Tranquilo Group, Laguna Colorada Formation; Santa Cruz, Argentina	<ul style="list-style-type: none"> specific and complex herbivory patterns of several FFG's
Feng et al. (2014); Hsü et al. (1974)	Keuper (Late Triassic)	District Yungjen, Yunnan, China	<ul style="list-style-type: none"> crescent-shape bite marks on <i>Mixopteris</i> intense skeletonization <i>Dictyophyllum nathorstii</i>
Gallego et al. (2003); Gallego et al. (2004); Gnaedinger et al. (2007); Gnaedinger et al. (2008); Gnaedinger et al. (2014)	Carnian-Norian (Late Triassic)	La Ternera Fm. (Quebrada La Cachivarita locality; La Ternera hill area, Copiapó Province, and the Las Breas Fm. (Punta del Viento locality, Vicuña, Elqui Province), Chile	<ul style="list-style-type: none"> oviposition scars on <i>Heidiphyllum</i>, <i>Pseudoctenis</i>, <i>Taeniopteris</i>
Strullu-Derrien et al. (2012)	Carnian (Late Triassic)	De Geerdalen Formation; Hopen Island, Svalbard Archipelago	<ul style="list-style-type: none"> aggregations of pellets or coprolites within bennettitalean roots gall-like structures within the cortical or pith tissues of the larger (probable bennettitalean) axes
Rozefelds & Sobbe (1987); Tillyard (1922); Webb (1982)	Carnian (Late Triassic)	Blackstone Formation, Ipswich Coal Measures Group; Sydney Basin, New South Wales, Australia	<ul style="list-style-type: none"> possible oviposition scars and insect eggs on <i>Nilssoniopteris</i> possible galls or eggs on <i>Dictyophyllum</i> mining structures on

Meller et al. (2011); Pott et al. (2008); Aschauer & Wappler unpubl. Data	Carnian (Late Triassic)	Lunz Formation; Lunz am See, eastern Northern Calcerous Alps, Austria	<i>Heidiphyllum</i> , <i>Ginkgoites</i> <ul style="list-style-type: none"> possible oviposition scars and insect eggs on <i>Nilssoniopteris</i> possible mining structures on <i>Nilssonia</i> marginal and non-marginal feeding traces on <i>Nilssoniopteris</i>, and other bennettitalean leaves
Moisan et al. (2012)	Carnian (Late Triassic)	Madygen Formation; Turkestan Mountains, southwestern Kyrgyzstan, Central Asia	<ul style="list-style-type: none"> oviposition scars on <i>Isoetites</i>
Anderson & Anderson (1983); Anderson & Anderson (1985); Anderson & Anderson (2003); Labandeira & Anderson (2005); Scott et al. (2004)	Carnian (Late Triassic)	Molteno Formation; Karoo Basin, KwaZulu- Natal, Eastern Cape and Northern Cape, South Africa	<ul style="list-style-type: none"> specific and complex herbivory patterns involving the frequency and diversity of 79 distinctive damage types (DTs) on about 220 whole-plant species (liverworts, lycopods, horsetails, ferns, cycads, peltasperms, corystosperms, hamshawvialeans, ginkgoaleans, voltzialean conifers, bennettitaleans, gnetophytes)
Linck (1949); Roselt (1954)	Carnian/Ladinian (Upper/Middle Triassic)	Bedheim, Germany	<ul style="list-style-type: none"> borings in <i>Dadoxylon</i> possible oviposition scars on <i>Equisetites</i>
Geyer & Kelber (1987); Kelber & Geyer (1989)	Upper Ladinian (Middle Triassic)	Lettenkohle of Alsace, France; Lower Keuper of Franconia, Germany	<ul style="list-style-type: none"> crescent-shape bite marks on <i>Schizoneura</i>, <i>Taeniopteris</i> possible oviposition scars and insect eggs on <i>Equisetites</i>
Heer (1877)	Ladinian (late Middle Triassic)	Neuwelt, Lettenkohle, Switzerland	<ul style="list-style-type: none"> possible oviposition scars on <i>Equisetites</i>
Minello (1994)	Ladinian (Middle Triassic)	Xinigua, Rio Grande do Sul, Santa Maria Formation (Rosario do Sul Group), Brazil	<ul style="list-style-type: none"> coprolite-bearing borings in <i>Araucarioxylon</i>
Grauvogel-Stamm & Kelber (1996)	Early Anisian (Early Middle Triassic)	Grès à Voltzia Formation; Grès-à-Voltzia, northern Vosges Mountains, France	<ul style="list-style-type: none"> crescent-shape bite marks on <i>Neuropteridium</i> possible eggs entangled in plant debris
McLoughlin (2011)	Anisian – Ladinian (Middle	Wivenhoe Hill, Esk Trough,	<ul style="list-style-type: none"> oviposition scars on

	Triassic)	Esk Formation; Queensland, Australia	<i>Taeniopteris</i>
McLoughlin (2011)	Olenekian – Anisian (late Early to early Middle Triassic)	Turrismetta Head, Sydney Basin; New South Wales, Australia	<ul style="list-style-type: none"> • gall on <i>Dicrodium</i>
Kustatscher et al. (2014b)	Olenekian (Lower Triassic)	Solling Formation; Bremke and Fürstenberg, Germany	<ul style="list-style-type: none"> • specific herbivory patterns involving the frequency and diversity of 8 distinctive damage types (DTs) • external feeding damage on <i>Tongchuanophyllum</i>, <i>Neuropteridium</i>, <i>Pelourdea</i> • mid-vein gall on <i>Tongchuanophyllum</i> • linear series of lenticular or ovoidal oviposition scars on <i>Tongchuanophyllum</i>

Table 2 (on next page)

Floral and insect damage composition late Ladinian flora from Monte Agnello, Dolomites, Italy.

Table 2. Floral and insect damage composition late Ladinian flora from Monte Agnello, Dolomites, Italy. DMG, percentage of damage; Spec. Spezialized damage; PS, Piercing and sucking; Ovi, Oviposition; FFG, Functional Feeding groups; DTO, Damage type occurrence.

- 2 Table 2. Floral and insect damage composition late Ladinian flora from Monte Agnello, Dolomites, Italy. DMG, percentage of damage;
3 Spec. Spezialized damage; PS, Piercing and sucking; Ovi, Oviposition; FFG, Functional Feeding groups; DTO, Damage type occurrence.

Species	# leaves	% DMG	% Spec	% Gall	% Mine	% External	% PS	% Ovi	DT s	#FFG s	DTO All	DTO Spec	DTO External	DTnumbers
<i>Bjuvia</i> cf. <i>dolomitica</i>	113	15.93	0.89	0.89		14.16		0.89	9	5	21	1	18	1;2;3;12;14;17;29;80;100
<i>Chiropteris</i> <i>montagnellii</i>	12													
<i>Cladophlebis</i> <i>ladinica</i>	24	4.17						4.17	1		1			101
<i>Cladophlebis</i> sp.	4													
Cone indet.	3													
<i>Elatocladus</i> sp.	1													
Equisetoid stem fragment	1													
Indet.	5													
<i>Neuropteridium</i> <i>elegans</i>	3													
<i>Nilssonia</i> cf. <i>neuberi</i>	40	10.00				10.00			1	1	4		4	12
<i>Nilssonia</i> sp.	34	23.53	2.91			20.59	2.94		6	3	10	1	9	1;2;7;12;13;128
<i>Pelourdea</i> sp.	4	25.00				25.00			1	1	1		1	12
<i>Phlebopteris</i> <i>fiemmensis</i>	6	33.33	16.67	16.67		16.67			2	2	2	1	1	2;80
? <i>Podozamites</i> sp.	17	17.65						17.65	2		3			72;100
<i>Pterophyllum</i> sp.	1													
<i>Radicites</i> sp.	1													
<i>Schizoneura</i> <i>paradoxa</i>	6													

<i>Scytophyllum bergeri</i>	55	54.55	7.27	3.64	1.82	49.09		8	4	37	4	34	3;5;12;13;14;40;63;80	
Seed	2													
<i>Sphenozamites</i> sp.	37	13.51	2.70			13.51		3	2	5	1	5	2;8;12	
Stem indet.	6													
<i>Taeniopteris</i> sp.	8	25.00				25.00		2	1	2		2	12;14	
<i>Thaumatopteris</i> sp.	3													
<i>Voltzia</i> sp. 1	84	3.57	1.19	2.38			1.1 9	2	2	3	1		48;121	
<i>Voltzia</i> sp. 2	41	2.44		2.44				1	1	1			121	
<i>Voltzia</i> sp. indet.	170	2.94		2.35		0.59		2	2	5		1	12;121	
Wood	3													
Total	684	12.14	1.32	1.61	0.15	9.36	0.2 9	0.73	20	7	95	9	75	1;2;3;5;7;8;12;13;14;17; 29;40;48;63;72;80;100;1 01;121;128

Table 3(on next page)

Floral and insect damage composition of the late Ladinian flora from Monte Agnello, Dolomites, Italy on higher classification level.

Floral and insect damage composition of the late Ladinian flora from Monte Agnello, Dolomites, Italy on higher classification level. DMG, percentage of damage; Spec. Spezialized damage; PS, Piercing and sucking; Ovi, Oviposition; FFG, Functional Feeding groups; DTO, Damage type occurrence.

2 Table 3. Floral and insect damage composition of the late Ladinian flora from Monte Agnello, Dolomites, Italy on higher classification
3 level. DMG, percentage of damage; Spec. Spezialized damage; PS, Piercing and sucking; Ovi, Oviposition; FFG, Functional Feeding
4 groups; DTO, Damage type occurrence.

Plant groups	# leaves	% DMG	% Spec	% Gall	% Mine	% External	% PS	% Ovi	DT s	#FFG s	DTO All	DTO Spec	DTO External	DTnumbers
Conifer	303	1.00	0.33	2.31		0.66	0.33		3	3	10	1	2	12;48;121
Cycadophytes	250	16.00	1.20	0.40		13.60	0.40	1.60	14	6	45	3	38	1;2;3;7;8;12;13;14;17;29;72;80;100;128
Indet.	16													
Ferns	52	5.77	1.92	1.92		1.92		1.92	3	2	3	1	1	2;80;101
Seed ferns	55	54.55	7.27	3.64	1.82	49.09			8	4	37	4	34	3;5;12;13;14;40;63;80
Sphenophytes	8													1;2;3;5;7;8;12;13;14;17;29;40;48;63;72;80;100;101;121;128
Total	684	12.13	1.32	1.61	0.15	9.36	0.29	0.73	20	7	95	9	75	

5

Table 4(on next page)

Floral diversity and evenness

Floral diversity and evenness

2 Table 4. Floral diversity and evenness.

Flora	N	S	Rarefied species diversity at 40 leaves	Rarefied external damage diversity at 40 leaves	Rarefied specialized damage diversity at 40 leaves	Pielou's J	Simpson D
MA1	244	22	12.89±1.52	2.30±1.13	0.63±0.69	0.89	1.89
MA5	236	16	9.03±1.41	3.19±1.08	0.68±0.75	0.73	1.83
MA7	125	15	9.38±1.47	1.29±0.49	na	0.63	1.70
MA8	44	7	6.85±0.36	0.93±0.25	na	0.75	1.71

3

1

Simplified geological map of the Monte Agnello area (Dolomites, N-Italy)

Figure 1. Simplified geological map of the Monte Agnello area (Dolomites, N-Italy), modified from Vardabasso (1930). MA1-MA8, fossil sites.

2

Examples of external foliage feeding at Monte Agnello (Dolomites, N-Italy).

Examples of external foliage feeding at Monte Agnello (Dolomites, N-Italy). A, *Scytophyllum bergeri* Bornemann, 1856 with intensively consumed leaf margins (DT12, 14) (MGP63/97). B, Hole feeding indicated by leaf removal on both sides of the primary veins (DT63) on *S. bergeri* Bornemann, 1856 (MGP196/39A-B). C-D, Hole feeding on a Sphenophyte (DT8) (MGP194/106), enlarged in D. E, Marginal feeding on the cycadophyte *Nilssonia* cf. *neuberi* Stur ex Pott et al. 2007 (DT12) (MGP191/6A). F, Excision of leaf to primary vein (DT14) on *Bjuvia* cf. *dolomitica* Wachtler & van Konijnenburg-van Cittert, 2000 (MGP181/11A). G, Removal or abrasion of surface tissues with a weak reaction rim (DT29) indicated by the dotted lines on *B. cf. dolomitica* Wachtler & van Konijnenburg-van Cittert, 2000 (MGP196/43). H, Cuspate excision (DT81) on *S. bergeri* Bornemann, 1856 (MGP171/28), enlarged in I. J-L, External foliage feeding on *B. cf. dolomitica* Wachtler & van Konijnenburg-van Cittert, 2000 (MGP195/69A), deep excision of leaf margin enlarged in K (DT12) and interveinal tissue removed in L (DT17). Scale bars: striped, 10 mm; solid, 5 mm; dotted, 1 mm.

3

Examples of internal foliage consumption at Monte Agnello (Dolomites, N-Italy).

Examples of internal foliage consumption at Monte Agnello (Dolomites, N-Italy).

A–B, Elliptical piercing and sucking punctures on the conifer *Voltzia* sp. 1 (MGP196/35), enlarged in B (DT48). C, Ellipsoidal, sessile bud gall from branchlet (DT121) on the unaffiliated *Voltzia* sp. 1 (MGP171/81). D, Small, hemispherical, thoroughly carbonized structures (DT80) on *Phlebopteris fiemmensis* Kustatscher et al., 2014 (MGP181/57C), indicated by arrows. E, Fern *Speirocarpus* sp. (MGP197/69B) showing lenticular-ovoidal foliar oviposition scars (DT101), indicated by arrows. F and I, Lenticular-ovoidal foliar oviposition scars (DT100) on the unaffiliated cycadophytes (MGP196/6; MGP196/7A). G, Undifferentiated galling structures (DT80) on a seed-fern (MGP63/94), indicated by arrows. H, Semilinear, frass-laden, mining structure with a smooth and rimmed margin (DT40) on *Scytophyllum bergeri* Bornemann, 1856 (MGP63/98A), asterisk indicates initial place of oviposition. J, Ellipsoidal scale impressions with roughened surface (DT128) on the cycadophyte *Nilssonia* cf. *neuberi* Stur ex Pott et al. 2007 (DT128) (MGP194/72A). Scale bars: striped, 10 mm; solid, 5 mm; dotted, 1 mm.

4

Plant and damage composition within the single sub-localities

Figure 4. Plant and damage composition within the single sub-localities. Pie charts showing the frequency specimen data by A. Host plant abundance (pooled in higher taxonomic ranks). B-D. Damage composition. MA1, MA5, MA7, MA8, fossil sites.