

# Pots vs trammel nets: a catch comparison study in a Mediterranean small-scale fishery

**Andrea Petetta** Corresp., Equal first author, 1, 2, **Claudio Vasapollo** Equal first author, 2, **Massimo Virgili** <sup>2</sup>, **Giada Bargione** <sup>1, 2</sup>, **Alessandro Lucchetti** Corresp. 2

<sup>1</sup> Department of Biological, Geological and Environmental Sciences, University of Bologna, Bologna, Italy, Italy

<sup>2</sup> Institute for Biological Resources and Marine Biotechnologies (IRBIM), Italian National Research Council (CNR), Ancona, Italy, Italy

Corresponding Authors: Andrea Petetta, Alessandro Lucchetti

Email address: andrea.petetta@irbim.cnr.it, alessandro.lucchetti@cnr.it

Passive bottom-set nets are the most widely used fishing gears in Mediterranean small-scale fisheries (SSFs). Trammel nets, in particular, have key advantages such as their ease of use and handling and high capture efficiency for numerous commercial species. However, they entail high discard rates (5-44% of the total catch) connected to high mortality, thus exerting an adverse impact on benthic communities, besides catching individuals of commercial species under the minimum conservation reference size (MCRS) and specimens of protected species. Fish pots are seen as alternative and a more sustainable gear type that allow reducing discards in SSFs. In this study, a collapsible pot was tested at three coastal sites in the north-western Adriatic Sea (GFCM GSA 17) to compare its catch efficiency with that of the local traditional trammel nets. Data analysis demonstrated a similar catch efficiency for the commercial species, both among sites and as a whole. Moreover, the trammel net caught a larger amount of discards, both in terms of species number and of CPUEw. The catch comparison study involved the two most abundant landed species, common cuttlefish *Sepia officinalis* and annular sea bream *Diplodus annularis*. The pots were more effective for *S. officinalis*, whereas the trammel net was more effective for the shorter length classes for *D. annularis*, which were mostly under the MCRS (12 cm). The innovative pots could provide a valuable alternative to the trammel nets traditionally used in the Adriatic Sea, at least in certain areas and periods. Their main advantages include that they do not require a different rigging and they can be used without bait, while their foldable design allows large numbers to be easily loaded on board SSF vessels. The results of this pilot study indicate that pots can achieve the objectives of reducing discards and bycatch in SSFs without penalizing the catch of commercial species.

# Pots vs trammel nets: a catch comparison study in a Mediterranean small-scale fishery

Andrea Petetta<sup>1,2</sup>, Claudio Vasapollo<sup>1</sup>, Massimo Virgili<sup>1</sup>, Giada Bargione<sup>1,2</sup>, Alessandro Lucchetti<sup>1</sup>

<sup>1</sup> Institute for Biological Resources and Marine Biotechnologies (IRBIM), National Research Council (CNR), Ancona, Italy

<sup>2</sup> Department of Biological, Geological and Environmental Sciences, University of Bologna, Bologna, Italy

Corresponding Authors:

Alessandro Lucchetti<sup>1</sup>, Andrea Petetta<sup>1,2</sup>

Largo Fiera della Pesca, Ancona, 1, 60125, Italy

Email addresses: [alessandro.lucchetti@cnr.it](mailto:alessandro.lucchetti@cnr.it); [andrea.petetta@irbim.cnr.it](mailto:andrea.petetta@irbim.cnr.it)

## Abstract

Passive bottom-set nets are the most widely used fishing gears in Mediterranean small-scale fisheries (SSFs). Trammel nets, in particular, have key advantages such as their ease of use and handling and high capture efficiency for numerous commercial species. However, they entail high discard rates (5-44% of the total catch) connected to high mortality, thus exerting an adverse impact on benthic communities, besides catching individuals of commercial species under the minimum conservation reference size (MCRS) and specimens of protected species. Fish pots are seen as alternative and a more sustainable gear type that allow reducing discards in SSFs. In this study, a collapsible pot was tested at three coastal sites in the north-western Adriatic Sea (GFCM GSA 17) to compare its catch efficiency with that of the local traditional trammel nets. Data analysis demonstrated a similar catch efficiency for the commercial species, both among sites and as a whole. Moreover, the trammel net caught a larger amount of discards, both in terms of species number and of CPUE<sub>w</sub>. The catch comparison study involved the two most abundant landed species, common cuttlefish *Sepia officinalis* and annular sea bream *Diplodus annularis*. The pots were more effective for *S. officinalis*, whereas the trammel net was more effective for the shorter length classes for *D. annularis*, which were mostly under the MCRS (12 cm). The innovative pots could provide a valuable alternative to the trammel nets traditionally used in the Adriatic Sea, at least in certain areas and periods. Their main advantages include that they do not require a different rigging and they can be used without bait, while their foldable design allows large numbers to be easily loaded on board SSF vessels. The results of this pilot study indicate that pots can achieve the objectives of reducing discards and bycatch in SSFs without penalizing the catch of commercial species.

# Introduction

Gillnets and trammel nets (or set/passive nets) are the most widely used fishing gears in Mediterranean small-scale fisheries (SSFs) (Lucchetti et al., 2015), which account for more than 70,000 vessels (FAO, 2018) and approximately 150,000 jobs. Set nets consist of netting panels hanging in the water column, where they are held perpendicular to the bottom by floaters and sinkers. Bottom-set fixed nets passively exploit the movements of target species (Gabriel et al., 2005). The fish swimming into them are caught by being gilled, tangled or wedged in gillnets, which are constituted of a single netting panel. On the contrary, the typical catching method of trammel nets is trapping the fish in a pocket of netting thanks to three panels: an inner panel with small mesh size and two outer panels with larger mesh size (Fabi et al., 2002). The success of the passive nets is due to their ease of use and handling (especially on small boats) (Dinçer & Bahar, 2008), high selectivity (especially gillnets) (Holt, 1963; Fabi et al., 2002) and their high capture efficiency for numerous commercial species (Amengual-Ramis et al., 2016). Their technical parameters, e.g. mesh size, netting twine, hanging ratio and net drop, vary widely in relation to the characteristics of target species and fishing areas (e.g. depth, seafloor), as do their selection properties (Stergiou et al., 2006; Lucchetti et al., 2017a). Although passive nets are considered as selective gears, they nonetheless produce a large amount of discards (Goncalves et al., 2007; Tzanatos & Somarakis, 2007) that range from 5% to about 40% of the total biomass caught (Tsagarakis, Palialexis & Vassilopoulou, 2014). SSFs discards consist of species with low commercial value, individuals that are found in poor condition and specimens under the Minimum Conservation Reference Size (MCRS; Regulation (EU), 2019/1241). The proportion of undersized individuals in the catch is variable and for some commercial species it can be quite high (20% for *Sparus aurata*, 28.2-74.8% for *Diplodus* spp., 93.8% for *Pagellus acarne* in the eastern Mediterranean, Tzanatos et al., 2008; large numbers of *Diplodus bellottii*, *Argyrosomus regius* in Cadiz and *Diplodus* spp., *Pagrus pagrus* in the Cyclades, Goncalves et al., 2007). Notably, species that are caught in excessively small amounts for the fishers' target market may also become discards (Goncalves et al., 2007). Moreover, set nets can be also responsible for the incidental catch of protected species such as sea turtles (Lucchetti and Sala, 2010; Casale, 2011; Lucchetti et al., 2017b) and elasmobranchs with no economic value (Morey et al., 2006; Saidi et al., 2016; Bradai, Saidi & Enajjar, 2018). The reduction of discards and bycatch has become a priority for fisheries worldwide, by means of measures to improve selectivity and to preserve the environment (FAO, 2011). The Common Fishery Policy, through the article 15 of Regulation (EU), 2013/1380, calls for the development of more selective technical solutions, to avoid the catch of unwanted species and sizes. Several solutions are being tested in the Mediterranean. They include: i) gear modifications to improve size and species selection (Lucchetti et al., 2015, 2017a); ii) time/area fishing closures to minimize bycatch (Lucchetti, Vasapollo & Virgili, 2017a); iii) mitigation devices to avoid catching some protected species (e.g. UV lights for sea turtles; Virgili et al., 2017); and iv) alternative and more sustainable fishing gears (Amengual-Ramis et al., 2016).

As regards the latter point, experimental pots developed in the past few years in certain areas have ensured catch efficiencies comparable to those of traditional set nets (Furevik & Hågensen, 1997; Iskandar et al., 2006; Furevik et al., 2008; Königson et al., 2015; Amengual-Ramis et al., 2016). Pots are passive gears to which fish, crustaceans and molluscs are attracted by bait or pasture, whereas cephalopods are caught because use them as a refuge or a site to spawn. Pots have several appealing features – in particular a minimal habitat impact and low manufacturing cost – which have led them to be classified as LIFE (low-impact fuel-efficient) gears (Suuronen et al., 2012). Moreover, bycatch can be minimized by acting on bait, mesh size, materials, and position/design of the entrance and the escape gap(s) (Furevik & Løkkeborg, 1994; Furevik & Hågensen, 1997; Boutson et al., 2009).

In Mediterranean SSFs traditional pots are locally employed to target molluscs and crustaceans (Grati et al., 2010; Amengueal-Ramis et al., 2016), ensuring high catch efficiency and low discard rates (0.8–6.6%; Fabi et al., 2001). They are usually deployed in specific seasons (e.g. the spawning period of *Sepia officinalis*, Melli et al., 2014), in circumscribed areas (e.g. north western Adriatic Sea for *Squilla mantis*, Grati et al., 2018; Gulf of Cádiz (Spain), Thracian Sea (Greece) and Gulf of Gabès (Tunisia) for *Octopus vulgaris*, Ezzeddine-Najai, 1992; Tsangridis, Sánchez & Ioannidou, 2002; Sobrino et al., 2011), or in replacement of other gears (e.g. for *Nephrops norvegicus* during trawl fishing closures in Croatian northern Adriatic waters, Brčić et al., 2018). A major disadvantage of traditional pots is their large volume, which entails that vessels can carry only a limited number of units per trip.

In the Mediterranean Sea, studies of alternative fishing gears such as innovative pots are still limited (ICES SGPOT, 2008, 2009; Pol et al., 2010) and mainly regard those targeted to cephalopods like *O. vulgaris* (Sbrana et al., 2008) and crustaceans such as *N. norvegicus*, *Plesionika* spp. (Colloca, 2002; Sartor et al., 2006) and *Palinurus elephas* (Amengual-Ramis et al., 2016).

Based on these considerations, a pilot study was devised to test a fully collapsible pot design and to compare it to a traditional set net in commercial fishing conditions. The main goals of the study were to evaluate the respective catch compositions and to assess the effectiveness of the pots in terms of their use and handling, discards and bycatch reduction.

## Materials & Methods

### Study area

The pilot study was conducted in FAO Geographical Sub-Area 17 (north-western Adriatic Sea) and involved three coastal areas (Marina di Ravenna, Portonovo and Senigallia), where depth ranges from 5 to 19 m (Figure 1). Specifications of bottom type and average depth of the three sites are listed in Table 1. In these areas, SSFs mostly employ gillnets and trammel nets to catch cuttlefish (*S. officinalis*), various fish species (e.g. *Solea solea*, *Lithognathus mormyrus*, *Diplodus* spp., *S. aurata*, *Sciaena umbra*, *Umbrina cirrhosa*, *Dicentrarchus labrax*) and crustaceans (*S. mantis*, *Paeneus kerathurus*) (Fabi & Grati, 2005). Moreover, artisanal pots are deployed in spring to specifically target cuttlefish.

# ***Fishing gears and experimental setup***

The trials were conducted on board local professional fishing vessels (Table 1). The characteristics of the traditional trammel nets (GTRs) employed and the fishing grounds were selected by the fishers, and similarly the fishing operations (e.g. fishing time) and the sorting of the catch were carried out following the fishers' procedures, without interferences from the scientists on board. Experiments to compare the GTRs and the foldable pots were carried out from April to August, in 2016 and 2017, and involved a total number of 42 fishing trials: 20 at Marina di Ravenna, 12 at Portonovo and 10 at Senigallia. To minimize differences due to patchy species distribution, GTRs and pots were deployed close to each other (a few tens of metres). The technical features of the GTRs are reported in Figure 2. The three netting panels were made of transparent polyamide multifilament: 210/4 mm multifilament and 36 mm mesh bar for the inner panel and 210/3 mm multifilament and 200 mm mesh bar for the outer panels. The net had a nominal height of 2.5 m (35 meshes), although its effective vertical opening in the water was around 1.5 m. The float line and lead line were in propylene (diameter, 8 and 10 mm, respectively); the float line was reinforced with external oval-shaped floats (diameter, 8 cm); the lead line weighed 120 g/m. The total length of the set nets used in each site was: 500 m at Marina di Ravenna and Portonovo, 300 m at Senigallia (Table 1). The GTRs were set early in the morning and hauled in the afternoon.

The pots (manufactured by Trapula<sup>LTD</sup>, Croatia; Figure 3) have a stainless-steel bar frame with a pentagonal shape and a single oval entrance. Two steel structures on the top and bottom allow folding them. A propylene rope 5 mm in diameter was externally reinforced with a nylon net (32 mm square-mesh bar). Flexible steel bars 2 mm in diameter allow manual adjustment of the opening. To establish whether catch efficiency was related to the volume of the chamber (Furevik & Løkkeborg, 1994), two different pot sizes were tested: a smaller pot (SP) measuring 40 x 100 cm (height x width) and a larger pot (LP) measuring 60 x 140 cm. The pots were attached to a main propylene line (namely a gang) 8 mm in diameter anchored to the seabed by 2 m plastic branch lines 5 mm in diameter, placed at 15 m intervals. The pots were set 15 m apart according to the traditional rigging used by the local fishers (Fabi et al., 2001). The number of pots of each type ranged from 9 to 20 per fishing trial. Soaking time (i.e. the period of time the pots were left on the bottom) depended on weather conditions and fishers' tactics (Table 1). They were commonly retrieved after 2-3 days, or more, in case of adverse weather conditions, to attract a wider range of commercial species other than cuttlefish, for which traditional pots are usually left 24 hours (Fabi, 2001). The pots were not baited, but several black plastic ribbons were attached to the frame, to attract cuttlefish.

The GTR and pot hauls were paired: each haul consisted in setting and retrieving the GTR, the gang with SPs and the gang with LPs.

## ***Data collection***

For each haul, the crew sorted onboard the catch, that was kept separate by gear (GTR, LP, SP). The total catch for each gear was thus divided into a landed catch (species with commercial value, not necessarily target species) and discards, i.e. species discarded for different reasons

(invertebrates and fish species with no commercial value, commercial individuals under the MCRS or in poor conditions). All individuals were identified to the lowest taxonomic level possible, counted, weighed to the lowest 0.1 g and measured to the nearest 0.5 cm for total length (TL; fish) or mantle length (ML; cephalopods) and to the nearest 0.5 mm for carapace length (CL; crustaceans).

# **Data analysis**

The catch per unit effort (CPUE) was calculated for GTRs and pots. The GTR catch was standardized for the number of individuals ( $CPUE_I$ ) and total catch weight ( $CPUE_W$ ; in kg) captured by 1000 m of net in 12 h, considering that fishers commonly set 3000 to 6000 m of net and haul it up after about half a day. The pot catch was standardized for the  $CPUE_I$  and  $CPUE_W$  (in kg) captured by 66.6 pots (i.e., the number of pots corresponded to 1000 m of set net considering 15 m distance between two subsequent pots, according to Fabi et al., 2001) in 24 h. This duration corresponds to the commercial fishing time of the traditional pots used in the area to target cuttlefish (Fabi, 2001). The Kruskal-Wallis H test ( $\chi^2$ ) was applied to seek differences between the  $CPUE_W$  of GTRs and pots. A non parametric test was adopted, because the data distributions were not normal and extremely skewed, with wide tails. If differences did emerge, a pairwise Wilcoxon's signed rank test based on Bonferroni correction for multiple comparisons was applied to establish the levels showing significantly different median values. Differences in the size of the individuals caught by the GTRs, LPs and SPs were explored by analysing the length frequency distribution (LFD) of the landed species. The catch efficiency of each pot type vs GTR was compared using generalized linear mixed models (GLMMs; Holst and Revill, 2009). The probability for an individual to be retained in a pot follows from:

$$Pr \left\{ \frac{Pot}{(Pot + GTR)} \right\} = \frac{1}{1 + e^{-(\beta_0 + \beta_1 \times length + \beta_2 \times length^2 + \beta_3 \times length^3)}}$$

A binomial error distribution was used to calculate the probability of the number of fish caught in a pot ( $CPUE_I$ ) given that they were caught by both gears based on 1-cm size classes. A probability value of 0.5 corresponds to equal catches in both gears. According to Holst and Revill (2009), a 3<sup>rd</sup> order polynomial would be adequate for most cases, although in some instances a 1<sup>st</sup> or 2<sup>nd</sup> order would be enough. The best binomial model was chosen based on the lowest Akaike's Information Criterion (AIC) value. A random term was added to the models. Since the GTR and pot hauls were paired, the catches for each site and for each gear were pooled and the term 'site' was used as a random intercept.

The most abundantly caught species, *S. officinalis* and *Diplodus annularis*, were selected for the catch comparison analysis. Since only SPs were set at Portonovo, the catch comparison of LPs included only the Senigallia and Marina di Ravenna. The models are illustrated graphically with a 95% confidence interval (CI) calculated with a bootstrap method using 999 simulations. The free software R (R Core Team, 2018) and the R packages nlme (Pinheiro et al., 2018) and lme4 (Bates et al., 2014) were used for the analyses.

# **Results**

Overall, the three gears caught 53 species, 38 of which belonged to the landed fraction (GTRs=30, LPs=15, and SPs=22) and 28 to the discard species (GTRs=25, LPs=5, and SPs=5), thus confirming that the pots were more species-selective than GTRs (Supplementary Tables 1 and 2).

As regards the landed species, cuttlefish (*S. officinalis*) was the most abundant in terms of biomass at all 3 sites for all 3 gears. Two other abundant species caught by all 3 gears were annular seabream (*D. annularis*), caught at Marina di Ravenna and Senigallia but not at Portonovo, and striped seabream (*L. mormyrus*), caught at all 3 sites with greater abundance at Senigallia and Portonovo. Additional landed species caught by GTRs were *S. solea*, *S. mantis* and *S. umbra* at Marina di Ravenna; *Liza aurata*, *Sarda sarda* and *Scophthalmus rhombus* at Senigallia; and *Mugil cephalus* at Portonovo. Other landed species caught by the pots were *S. umbra* (LP, SP) and *Conger conger* (LP) at Marina di Ravenna and *Dentex dentex* (SP) at Portonovo (Supplementary Table 1).

The mean biomass values (calculated as CPUE<sub>w</sub>) and 95% CIs of the landed catch are reported in Table 2. The CI values of the three gears did overlap, indicating the lack of significant differences among them, both at each site and as a whole. Standardization of the landed catch weight failed to highlight significant differences in the medians among GTRs, LPs and SPs, either within the 3 sites or as a whole ( $\chi^2 = 2.59$ ,  $df = 2$ ,  $p = 0.274$ ; Figure 4).

The discards of LPs and SPs were lower than those of the GTRs both in terms of species number and of CPUE<sub>w</sub>; in fact, they were close to zero both at Senigallia and at Portonovo (Supplementary Table 2). The GTRs captured large amounts of *Alosa fallax* and *Pteroplatytrygon violacea* at Marina di Ravenna; *A. fallax* and *Liocarcinus vernalis* at Senigallia and *Eriphia verrucosa*, *Hexaplex trunculus* and *Maja crispata* at Portonovo (Supplementary Table 2).

The CIs of the mean biomass values (CPUE<sub>w</sub>) of GTR discards did not overlap with those of LPs and SPs, whereas those of LPs and SPs did (Table 2). The discards showed significant differences in terms of standardized biomass ( $\chi^2 = 11.34$ ,  $df = 2$ ,  $p = 0.004$ , Figure 4) and were mostly caught by GTRs. Wilcoxon's pairwise test showed that the median differences between gears were significant for GTRs vs LPs and for GTRs vs SPs ( $p = 0.011$  and  $p = 0.016$ , respectively), whereas the medians of LPs and SPs were not significantly different.

The LFD of *S. officinalis* and *D. annularis* at each site and as a whole is reported in Figure 5.

The lines of LPs and SPs were mostly above those of GTRs, indicating a greater catch efficiency.

The catch comparison curves (Figure 6 and Table 3) demonstrate that for *S. officinalis*, pots (both dimensions) were more efficient than GTRs. SPs were more efficient than GTRs for most *S. officinalis* sizes, except for the larger ones (above the 25 cm size class), for which the efficiency of both gears were similar, as the lower CI exceeds the limit of 0.5 indicating equal proportion of individual catches between both gears. In contrast, LPs showed the same efficiency as GTRs for the smaller individuals (below the 11 cm size class) and were more efficient for the larger individuals. LPs and SPs showed overlapping CIs from the 10 cm size class, i.e. a similar catch efficiency. As regards *D. annularis*, the GTR reflected a greater efficiency than both LPs

and SPs at the smaller sizes, usually under the MCRS of the species (12 cm). As a result, the percentage in number of undersized individuals of *D. annularis* caught by GTR was 15.1% (7.1% for Marina di Ravenna; 16.4% for Senigallia), while it was lower for both pots: 7.3% for SPs (3.3% for Marina di Ravenna; 8% for Senigallia) and 3.7% for LPs (3.3% for Marina di Ravenna; 4.5% for Senigallia). Therefore, LPs and SPs both showed a high selectivity for larger fish, although they also caught some undersized individuals. However, there were no differences among the three gears in the catch efficiency for the larger *D. annularis* individuals.

## Discussion

This study was aimed at testing the catching efficiency of an innovative pot design in the north-western Adriatic Sea; in particular, it was evaluated the pot's ability to provide an alternative to traditional trammel nets and to reduce discards in SSFs. The comparison implied a different standardization (CPUE<sub>w</sub> at 12 hours for trammel nets and 24 hours for pots) that represented a compromise to maintain the standardization unit as close as possible between the two gears, taking into account that they operate in different ways and with a different fishing time. The main finding, from the catch comparison analysis of the two most abundant species caught (cuttlefish and annular sea bream), was that the catch rates of the two types of pot tested, which differed only in dimensions, were comparable to those of the trammel nets. These data are in line with the high efficiency of the experimental pots reported in the Barents Sea (Furevik et al., 2008) and the Baltic Sea (Königson et al., 2015), where pots showed similar, if not higher, catch rates than those of passive nets, at least in a period of the year. Interestingly, the innovative pots caught a larger number of commercial species than the traditional ones used in the area by artisanal fishers, which are not collapsible, with a different shape and entrance, and mainly target cuttlefish (Fabi et al., 2001).

Regarding the annular sea bream, the poor selectivity of the trammel nets found for this species has been previously reported for *Diplodus* spp. by Tzanatos et al. (2008), who estimated a percentage (in weight) of the undersized individuals caught of 28.2% for *D. annularis* and 74.8% for *D. sargus*, being even higher than the average percentage of this study (15%). In contrast, pots seemed to be able to avoid *D. annularis* juveniles (average percentage of around 7%). Unlike studies in other areas (Munro, 1974; Furevik & Løkkeborg, 1994; Hedgärde et al., 2016), which concluded that larger pots are more effective than small ones, in this study, pot size seemed not to affect catch efficiency. However, the number of pots that actually produced a catch, ranged between  $38.5 \pm 3.3\%$  for SPs to  $42.5 \pm 4.3\%$  for LPs, stressing the need for increasing catch efficiency by using attractive baits.

The cost of a collapsible pot as the one tested in this study ranges from €50 (SP) to €100 (LP), whereas 100 m of the traditional trammel net used for the sea trials is around €200. Assumed that 100 m of nets would correspond to more than 6 pots (as stated in the Materials and Methods section), the alternative gear is more expensive. Nevertheless, whereas the set nets usually last a single season and are then too damaged to be repaired, these kind of pots last up to two years. Another advantage of pots is that they afford a more limited access to the catch, making them



less subject to depredation by large predators than set nets (e.g. seal- and dolphin-safe fishing gear; Königson, 2011; Königson et al., 2015). Moreover, they provide a greater catch quality, because they generally do not damage the specimens caught (Suuronen et al., 2012; Olsen, 2014). In addition, even if trammel nets require a shorter fishing time than pots, fishers could set different pots gangs in order to alternate the retrieve, and thus to haul them daily.

With reference to discards, the greater amount caught by the trammel nets clearly produces a greater impact on the benthic community, since discard mortality is high (Suuronen et al., 2012). Moreover, the cleaning of the trammel net implies an additional time and labour on deck for fishers, since discards must be released or untangled manually (Sartor et al., 2018; Szyńska et al., 2018). In contrast, the removal of discards from pots can be done without significantly reducing fishing time and leaving high probability for the unwanted organisms to survive (Suuronen et al., 2012). Discarding is a major issue for fisheries management worldwide (Tsagarakis, Palialexis & Vassilopoulou, 2014). In the Mediterranean, the Common Fisheries Policy (CFP Regulation (EU), 2013/1380) has introduced the obligation to land ('discard ban') all the individuals of the species with minimum legal size (MCRS, e.g. those species reported in the Annex III of the Council Regulation (EC) No 1967/2006), thus emphasizing the need to reduce discards (Damalas, 2015). The landing obligation is a matter of concern among fishers, which are facing difficulties related to storing and bringing to land the former discard, due to limited hold space, and to sorting time or personnel increasing (Maynou et al., 2018). The introduction of a new and alternative technology in a fishery, such as innovative pots, could help to achieve this goal of discard reduction only if it is acceptable to both fishers and fishery policies. In this context, fish collapsible pots are revealed to be: practical (i.e. involving no major changes to common fishers' practices), cost-effective (i.e. easy to use and not expensive to maintain, no waste of time for cleaning the gear), efficient (i.e. large spectrum of species caught) and enforceable (i.e. easy to control by inspection authorities).

With reference to bycatch of sensitive and protected species, during our study the trammel nets caught five specimens of the pelagic stingrays (*Pteroplatytrigon violacea*), a frequent event in several Adriatic fisheries (Bonanomi et al., 2018). This Elasmobranch species, usually discarded due to its scarce commercial value, is considered as a 'Least Concern' species in the IUCN red list (Baum et al., 2016). Another bycatch species caught by the trammel nets was the twait shad (*A. fallax*), listed in Annexes II and V of the Habitats Directive (European Commission, 1992) as requiring close protection. In addition, the passive nets deployed in the central and northern Adriatic are responsible for the bycatch – maybe as many as thousands individuals a year (Lucchetti, Vasapollo & Virgili, 2017b) – of loggerhead sea turtles (*Caretta caretta*), which are listed in Annex IV of the Habitats Directive. In contrast, the pots did not capture any of these species, substantially due of the small size of the pot entrance compared with the larger size of animals such as stingrays and sea turtles. As regards *A. fallax*, its absence in the pots catch may depend on their position close to the bottom, whereas trammel nets can also intercept pelagic fish species (*A. fallax*, *Sardina pilchardus*, *Engraulis encrasicolus*, *Scomber japonicus* etc.) which for different reasons can be discarded (Goncalves et al., 2007).

## Conclusions

The innovative pots tested in this study seem to provide a sound alternative to the traditional trammel nets used in the Adriatic Sea, at least in spring and summer, as concerns the small-scale fishery targeting common cuttlefish. These pots do not require a different vessel rigging nor changes to the on board practices; moreover, they can be used without baits and their foldable design involves that they can be easily stored on board the typical artisanal boats used in Mediterranean SSFs. The findings of this pilot study, although not conclusive, clearly indicate that these alternative gears go some way towards reducing bycatch and discards in SSFs while maintaining the commercial catch. Similar tests should be extended to other areas and seasons, also using baits, to provide a clearer assessment of their efficiency.

## Acknowledgements

This study was authorized by the Italian Coast Guard. It was conducted with the contribution of the LIFE financial instrument of the European Community, TartaLife Project - Reduction of sea turtle mortality in professional fishing (LIFE12NAT/IT/000937). It does not necessarily reflect the European Commission's views and in no way anticipates future policy. This work was finalised under the project "Innovative technologies and sustainable use of fisheries and biological resources in the Mediterranean sea" (FISHMED-PHD). We are indebted to the personnel of CNR-IRBIM (Ancona) and CESTHA (Marina di Ravenna), to the crew of FVs Jessica (Portonovo), Zio Lino (Senigallia) and Nemo (Marina di Ravenna) for their help in fieldwork. We are grateful to Word Designs for the language revision ([www.silviamodena.com](http://www.silviamodena.com)).

## References

- Amengual-Ramis JF., Vázquez-Archdale M., Cánovas-Pérez C., Morales-Nin B. 2016. The artisanal fishery of the spiny lobster *Palinurus elephas* in Cabrera National Park, Spain: Comparative study on traditional and modern traps with trammel nets. *Fisheries Research* 179:23–32. DOI: 10.1016/j.fishres.2016.01.02.
- Bates D., Mächler M., Bolker B., Walker S. 2014. Fitting linear mixed-effects models using lme4. *arXiv preprint arXiv:1406.5823*.
- Baum J., Bianchi I., Domingo A., Ebert DA., Grubbs RD., Mancusi C., Piercy A., Serena F., Snelson FF. 2016. *Pteroplatytrygon violacea*. *The IUCN Red List of Threatened Species*.
- Bonanomi S., Pulcinella J., Fortuna CM., Moro F., Sala A. 2018. Elasmobranch bycatch in the Italian Adriatic pelagic trawl fishery. *PLOS ONE*. DOI: <https://doi.org/10.1371/journal.pone.0191647>.
- Boutson A., Mahasawasde ÆC., Mahasawasde S., Tunkijjanukij ÆS. 2009. Use of escape vents to improve size and species selectivity of collapsible pot for blue swimming crab *Portunus pelagicus* in Thailand. *Fisheries Science*:25–33. DOI: 10.1007/s12562-008-0010-z.
- Bradai MN., Saidi B., Enajjar S. 2018. Overview on Mediterranean Shark's Fisheries: Impact on the Biodiversity. *Marine ecology: Biotic and abiotic interactions*:211–230.
- Brčić J., Herrmann B., Ma M., Baranovi M., Krstulovi S., Frane Š. 2018. Size selection of *Nephrops norvegicus* (L.) in commercial creel fishery in the Mediterranean Sea. *Fisheries*

- Research 200:25–32. DOI: 10.1016/j.fishres.2017.12.006.
- Colloca F. 2002. Life cycle of the deep-water pandalid shrimp *Plesionika edwardsii* (Decapoda , Caridea) in the central Mediterranean sea. *Journal of Crustacean Biology* 22:775–783. DOI: 10.1651/0278-0372(2002)022.
- Damalas D. 2015. Mission impossible : Discard management plans for the EU Mediterranean fisheries under the reformed Common Fisheries Policy. *Fisheries Research* 165:96–99. DOI: 10.1016/j.fishres.2015.01.006.
- Dinçer AC., Bahar M. 2008. Multifilament Gillnet Selectivity for the Red Mullet (*Mullus barbatus*) in the Eastern Black Sea Coast of Turkey , Trabzon. *Turkish Journal of Fisheries and Aquatic Sciences* 359:355–359.
- Directive H. 1992. Council Directive 92/43/EEC of 21 May 1992 on the conservation of natural habitats and of wild fauna and flora. *Official Journal of the European Union* 206:7–50.
- Ezzeddine-Najai S. 1992. Biologie et pêche du poulpe *Octopus vulgaris* Cuvier, 1797 (Cephalopoda, Octopoda) du golfe de Gabès. *Bulletin de l'Institut National des Sciences et Technologies Océanographiques et de Pêche à Salammbô* 19:5–19.
- Fabi G. 2001. *Sepia officinalis*: impact of three set of fishing techniques in the Adriatic and the Ligurian Sea. Study contract No. 98/069. :119.
- Fabi G., Grati F. 2005. Small-scale fisheries in the maritime department of Ancona (Central Northern Adriatic Sea). *AdriaMed Tech*:64–84.
- Fabi G., Grati F., Lucchetti A., Scarcella G. 2001. Osservazioni preliminari sulle catture di *Sepia officinalis* con tre attrezzi da posta in Medio Adriatico. *Biologia Marina Mediterranea* 8:660–664.
- Fabi G., Sbrana M., Biagi F., Grati F., Leonori I., Sartor P. 2002. Trammel net and gill net selectivity for *Lithognathus mormyrus* (L., 1758), *Diplodus annularis* (L., 1758) and *Mullus barbatus* (L., 1758) in the Adriatic and Ligurian seas. *Fisheries Research* 54:375–388.
- FAO. 2011. *International Guidelines on bycatch management and reduction of discards*. Rome.
- FAO. 2018. *The state of Mediterranean and Black Sea fisheries 2018*.
- Furevik DM., Hågensen SP. 1997. Experiments with cod pots as an alternative to gill-nets in the Varanger fjord in April-June and October-December 1996. *Proceedings of the 7th Russian/Norwegian Symposium: Gear Selection and Sampling Gears*:121–132.
- Furevik DM., Humborstad O., Jørgensen T., Løkkeborg S. 2008. Floated fish pot eliminates bycatch of red king crab and maintains target catch of cod. *Fisheries Research* 92:23–27. DOI: 10.1016/j.fishres.2007.12.017.
- Furevik DM., Løkkeborg S. 1994. Fishing trials in Norway for torsk (*Brosme brosme*) and cod (*Gadus morhua*) using baited commercial pots. *Fisheries research* 19:219–229.
- Gabriel O., Lange K., Dahm E., Wendt T. 2005. *Von Brandt's Fish Catching Methods of the World*. Blackwell Publishing.
- Goncalves J., Stergiou KI., Hernando JA., Puente E., Moutopoulos DK., Arregi L., Soriguer MC., Vilas C., Coelho R., Erzini K. 2007. Discards from experimental trammel nets in southern European small-scale fisheries. 88:5–14. DOI: 10.1016/j.fishres.2007.06.017.
- Grati F., Fabi G., Azzurro E. 2018. Seasonal dynamics of small-scale fisheries in the Adriatic Sea. DOI: 10.12681/mms.2153.
- Grati F., Polidori P., Scarcella G., Fabi G. 2010. Estimation of basket trap selectivity for changeable nassa ( *Nassarius mutabilis* ) in the Adriatic Sea. *Fisheries Research* 101:100–107. DOI: 10.1016/j.fishres.2009.09.012.
- Hedgärde M., Berg CW., Kindt-Larsen L., Lunneryd SG., Königson S. 2016. Explaining the

- catch efficiency of different cod pots using underwater video to observe cod entry and exit  
behaviour. *The Journal of Ocean Technology* 11:67–90.
- Holst R., Revill A. 2009. A simple statistical method for catch comparison studies. *Fisheries Research* 95:254–259. DOI: 10.1016/j.fishres.2008.09.027.
- Holt SJ. 1963. A method for determining gear selectivity and its application. *ICNAF Spec. Publ.* 5:106–115.
- ICES. 2009. ICES SGPO REPORT 2009 Report of the Study Group on the Development of Fish Pots for Commercial Fisheries and Survey Purposes T. In: Ancona, 19–20.
- ICES SGPO REPORT 2008 Report of the Study Group on the Development of Fish Pots for Commercial Fisheries and Survey Purposes 2008. Tórshavn , Faroe Islands.
- Iskandar D., Suzuki Y., Shiode D., Hu F., Tokai T. 2006. Catchability difference of gill net and collapsible baited pot for Japanese rock crab. *Indian Journal of Fisheries* 12:107–112.
- Königson S. 2011. Seals and fisheries: A study of the conflict and some possible solutions. Department of Marine Ecology; Institutionen för marin ekologi.
- Königson SJ., Fredriksson RE., Lunneryd S-G., Strömberg P., Bergström UM. 2015. Cod pots in a Baltic fishery: are they efficient and what affects their efficiency? *ICES Journal of Marine Science* 72:1545–1554.
- Lucchetti A., Buglioni G., Conides A., Klaoudatos D., Sartor P., Sbrana M., SPEDICATO M-T., Stamatopoulos C. 2015. Technical measures without enforcement tools: is there any sense? A methodological approach for the estimation of passive net length in small scale fisheries. *Mediterranean Marine Science* 16:82–89.
- Lucchetti A., Carbonara P., Colloca F., Lanteri L., Teresa M., Sartor P. 2017. Ocean & Coastal Management Small-scale driftnets in the Mediterranean : Technical features , legal constraints and management options for the reduction of protected species bycatch. *Ocean and Coastal Management* 135:43–55. DOI: 10.1016/j.ocecoaman.2016.11.002.
- Lucchetti A., Sala A. 2010. An overview of loggerhead sea turtle (*Caretta caretta*) bycatch and technical mitigation measures in the Mediterranean Sea. *Reviews in Fish Biology and Fisheries* 20:141–161.
- Lucchetti A., Vasapollo C., Virgili M. 2017a. Sea turtles bycatch in the Adriatic Sea set net fisheries and possible hot - spot identification. *Aquatic Conservation: Marine and Freshwater Ecosystems*:1–10. DOI: 10.1002/aqc.2787.
- Lucchetti A., Vasapollo C., Virgili M. 2017b. An interview-based approach to assess sea turtle bycatch in Italian waters. *PeerJ* 5:e3151.
- Maynou F., del Mar Gil M., Vitale S., Giusto GB., Fouts A., Rangel M., Rainha R., Erzini K., Gonçalves JMS., Bentes L. 2018. Fishers’ perceptions of the European Union discards ban: perspective from south European fisheries. *Marine Policy* 89:147–153.
- Melli V., Riginella E., Nalon M., Mazzoldi C. 2014. From Trap to Nursery . Mitigating the Impact of an Artisanal Fishery on Cuttlefish Offspring. *PLOS ONE* 9. DOI: 10.1371/journal.pone.0090542.
- Morey G., Moranta J., Riera F., Grau AM., Morales-Nin B. 2006. Elasmobranchs in trammel net fishery associated to marine reserves in the Balearic Islands (NW Mediterranean). *Cybiu* 30:125–132.
- Munro JL. 1974. The mode of operation of Antillean fish traps and the relationships between ingress, escapement, catch and soak. *ICES Journal of Marine Science* 35:337–350.
- Olsen L. 2014. Baited pots as an alternative fishing gear in the Norwegian fishery for Atlantic cod (*Gadus morhua*). The Arctic University of Norway.

- Pinheiro J., Bates D., DebRoy S., Sarkar D. 2018. R Core Team (2018). nlme: Linear and nonlinear mixed effects models. R package version 3.1-137.
- Pol MV., He P., Winger PD. 2010. Proceedings of the International Technical Workshop on Gadoid Capture by Pots (GACAPOT). *Massachusetts Division of Marine Fisheries Technical Report* 40:107.
- R Core Team. 2018. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. 2012. URL <http://www.R-project.org>.
- Regulation (EU). 2013. No. 1380/2013 of the European Parliament and of the Council of 11 December 2013 on the Common Fisheries Policy, amending Council Regulations (EC) No 1954/2003 and (EC) No 1224/2009 and repealing Council Regulations (EC) No 2. :22–61.
- Regulation (EU). 2019. Regulation (EU) 2019/1241 of the European Parliament and of the Council of 20 June 2019 on the conservation of fisheries resources and the protection of marine ecosystems through technical measures. :97.
- Saidi B., Enajjar S., Bradai MN. 2016. Elasmobranch captures in shrimps trammel net fishery off the Gulf of Gabès (Southern Tunisia, Mediterranean Sea). *Journal of Applied Ichthyology* 32:421–426. DOI: 10.1111/jai.13061.
- Sartor P., Rossetti I., Vannucci A., Viva C. 2006. Pesca del gobbetto striato, Plesionika edwardsii (brandt, 1851), con nasse sperimentali nel Tirreno settentrionale. *Biologia Marina Mediterranea* 13:288–289.
- Sartor P., Veli DL., Carlo F De., Ligas A., Massaro A., Musumeci C., Sartini M., Rossetti I., Sbrana M., Viva C. 2018. Reducing unwanted catches of trammel nets : experimental results of the “ guarding net ” in the caramote prawn , Penaeus kerathurus , small-scale fishery of the Ligurian Sea ( western Mediterranean ). :131–140.
- Sbrana M., Rossetti I., Sartor P., Vannucci A., De Ranieri S. 2008. Sperimentazione di differenti tipologie di trappole per la pesca del polpo Octopus vulgaris (Cuvier 1797) nel Mar Ligure orientale. *Biologia Marina Mediterranea* 15:356–357.
- Sobrinho I., Juarez A., Rey J., Romero Z., Baro J. 2011. Description of the clay pot fishery in the Gulf of Cadiz ( SW Spain ) for Octopus vulgaris : Selectivity and exploitation pattern. *Fisheries Research* 108:283–290. DOI: 10.1016/j.fishres.2010.12.022.
- Stergiou KI., Moutopoulos DK., Soriguer MC., Puente E., Lino PG., Zabala C., Monteiro P., Errazkin LA., Erzini K. 2006. Trammel net catch species composition , catch rates and métiers in southern European waters : A multivariate approach. *Fisheries Research* 79:170–182. DOI: 10.1016/j.fishres.2006.03.003.
- Suuronen P., Chopin F., Glass C., Løkkeborg S., Matsushita Y., Queirolo D., Rihan D. 2012. Low impact and fuel efficient fishing — Looking beyond the horizon. *Fisheries Research* 119–120:135–146. DOI: 10.1016/j.fishres.2011.12.009.
- Szynaka MJ., Bentes L., Monteiro P., Rangel M., Erzini K. 2018. Reduction of by-catch and discards in the Algarve small-scale coastal fishery using a monofilament trammel net rigged with a guarding net. *Scientia Marina*:121–129.
- Tsagarakis K., Palialexis A., Vassilopoulou V. 2014. Mediterranean fishery discards: review of the existing knowledge. *ICES Journal of Marine Science* 71:1219–1234. DOI: 10.4135/9781412953924.n678.
- Tsangridis A., Sánchez P., Ioannidou D. 2002. Exploitation patterns of Octopus vulgaris in two Mediterranean areas. *Scientia Marina* 66:59–68.
- Tzanatos E., Somarakis S. 2007. Discarding practices in a Mediterranean small-scale fishing fleet (Patraikos Gulf , Greece). *Fisheries Management and Ecology*:277–285. DOI:

498 10.1111/j.1365-2400.2007.00556.x.  
 499 Tzanatos E., Somarakis S., Tserpes G., Koutsikopoulos C. 2008. Catch length analysis , relation  
 500 to minimum landing sizes and management implications from a Mediterranean small-scale  
 501 fishery ( Patraikos Gulf , Greece ). *Fisheries Research* 93:125–134. DOI:  
 502 10.1016/j.fishres.2008.03.003.  
 503 Virgili M., Vasapollo C., Lucchetti A. 2017. Can ultraviolet illumination reduce sea turtle  
 504 bycatch in Mediterranean set net fisheries ? Can ultraviolet illumination reduce sea turtle  
 505 bycatch in Mediterranean set net fisheries ? *Fisheries Research* 199:1–7. DOI:  
 506 10.1016/j.fishres.2017.11.012.  
 507

# **Table 1**(on next page)

Summary of the fishing trials carried out at the three sites (Marina di Ravenna, Senigallia, Portonovo)

AVG: Average; SD: Standard Deviation; LOA: Length all out; GT: Gross tonnage; GTR: Trammel net; LP: Large pots; SP: Small pots.

1 **Table 1. Summary of the fishing trials carried out at the three sites (Marina di Ravenna, Senigallia, Portonovo).**  
2 **AVG: Average; SD: Standard Deviation; LOA: Length all out; GT: Gross tonnage; GTR: Trammel net; LP: Large pots; SP:**  
3 **Small pots.**

	<b>Marina di Ravenna</b>	<b>Senigallia</b>	<b>Portonovo</b>
<b>Bottom type</b>	sandy-mud with scattered rocky outcrops	sandy-mud	rocky
<b>AVG depth [m] ± SD</b>	9.9 ± 2.5	10.7 ± 0.43	6.0 ± 0.57
<b>Vessel characteristics</b>	LOA 12.4 m; 10 GT; 350 kW	LOA 12.4 m; 6 GT; 130 kW	LOA 6.6 m; 1 GT; 100 kW
<b>Study period</b>	April-July 2017	April-June 2016	May-August 2016
<b>No. of trials</b>	20	10	12
<b>GTR length [m]</b>	500	300	500
<b>No. of LPs</b>	20	9-10	0
<b>No. of SPs</b>	20	19-20	20
<b>AVG GTR soak time [h] ± SD</b>	19.9 ± 3.3	17.4 ± 2.0	17.2 ± 1.8
<b>AVG LP soak time [h] ± SD</b>	91.4 ± 28.7	73.9 ± 9.1	-
<b>AVG SP soak time [h] ± SD</b>	87.3 ± 25.6	76.6 ± 10.1	90.1 ± 21.3



## Table 2 (on next page)

Mean biomass values ( $CPUE_w$ ) with standard errors and confidence intervals, in brackets, of the landed catch and of discards for the three gears.

GTR (trammel net), LP (large pot) and SP (small pot), at each site (top) and as a whole (bottom). NA (not available).

1 **Table 2: Mean biomass values ( $CPUE_W$ ) with standard errors and confidence intervals, in brackets, of the landed catch and of discards for**  
 2 **the three gears,**

3 **GTR (trammel net), LP (large pot) and SP (small pot), at each site (top) and as a whole (bottom). NA (not available).**

	Site	GTR $CPUE_W$	LP $CPUE_W$	SP $CPUE_W$
Landed catch	M. di Ravenna	$5.41 \pm 0.76$ (3.92 - 6.89)	$7.27 \pm 1.77$ (3.81 - 10.73)	$3.72 \pm 0.64$ (2.48 - 4.97)
	Senigallia	$4.34 \pm 0.14$ (2.16 - 6.52)	$2.41 \pm 0.35$ (0.23 - 4.59)	$3.08 \pm 0.17$ (2.20 - 3.97)
	Portonovo	$2.90 \pm 0.70$ (1.53 - 4.27)	-	$2.48 \pm 0.59$ (1.33 - 3.62)
Discards	M. di Ravenna	$0.77 \pm 0.20$ (0.38 - 1.17)	$0.14 \pm 0.04$ (0.07 - 0.21)	$0.07 \pm 0.05$ (0.02 - 0.17)
	Senigallia	$0.95 \pm 0.57$ (0.17 - 2.07)	-	$0.04 \pm 0.0$ (NA)
	Portonovo	$1.52 \pm 0.79$ (0.02 - 3.06)	-	$0.06 \pm 0.0$ (NA)
Landed catch	GTR	$4.35 \pm 0.51$ (3.35 - 5.34)		
	LP	$6.01 \pm 1.39$ (3.28 - 8.74)		
	SP	$3.21 \pm 0.36$ (2.49 - 3.93)		
Discards	GTR	$0.95 \pm 0.23$ (0.51 - 1.40)		
	LP	$0.14 \pm 0.04$ (0.06 - 0.21)		
	SP	$0.06 \pm 0.03$ (0.01 - 0.12)		

**Table 3**(on next page)

Estimates of the parameters of the GLMM calculated for catch comparison

SE: standard error. GTR: Trammel net; LP: Large pots; SP: Small pots.


1 **Table 3. Estimates of the parameters of the GLMM calculated for catch comparison. SE:**  
 2 **standard error. GTR: Trammel net; LP: Large pots; SP: Small pots.**

Species	Model		Parameter	Estimate	SE	p
<i>Sepia officinalis</i>	Linear	LP vs GTR	$\beta_0$	-0.61	0.45	0.171
			$\beta_1$	0.08	0.03	0.009
	Quadratic	SP vs GTR	$\beta_0$	4.07	1.47	0.006
			$\beta_1$	-0.46	0.20	0.022
			$\beta_2$	0.01	0.01	0.038
<i>Diplodus annularis</i>	Quadratic	LP vs GTR	$\beta_0$	-27	5.62	<0.001
			$\beta_1$	3.33	0.71	<0.001
			$\beta_2$	-0.98	0.02	<0.001
	Quadratic	SP vs GTR	$\beta_0$	-11.39	4.24	0.007
			$\beta_1$	1.35	0.56	0.017
			$\beta_2$	-0.04	0.02	0.048

3

# Figure 1


Map of the study area where the trials were performed in 2016 and 2017 (April-August).



# Figure 2


Scheme of the trammel net used in the study.

Inner panel: middle; outer panel: top and bottom. PA: polyamide; PP: propylene;  $\varnothing$ : diameter; E: hanging ratio.



## Figure 3

Picture of the 'Trapula' pots tested in the study. See text for dimensions.





# Figure 4

Overall CPUE<sub>w</sub> of landings and discards of the three gears tested in the study.

GTR: trammel nets; LP: large pots; SP: small pots. Red dots: mean CPUE<sub>w</sub>; red bars: confidence intervals (CIs). (A) Commercial; (B) Discard.






## Figure 5

Length frequency distributions (LFDs) of *Sepia officinalis* in each site (A,B,C) and as a whole (D) and LFDs of *Diplodus annularis* in each site (E,F) and as a whole (G).

A,E: Marina di Ravenna, B: Portonovo, C,F: Senigallia. Dashed lines represent GTR (trammel nets); continuous lines represent SP (small pots); dotted lines represent LP (large pots); vertical dotted lines represent MCRS of 12 cm of *D. annularis*.



# Figure 6

Catch comparison curves for *Sepia officinalis* (A) and *Diplodus annularis* (B), representing the GLMM proportions of the total catches of the three gears.

GTR: trammel nets; LP: large pots; SP: small pots. Dashed and dotted lines represent the mean proportions of LP and SP, respectively; the vertical dotted line represents MCRS of 12 cm of *D. annularis*. Interpretation: a value of 0.5 indicates an even split between GTRs and Pots, whereas a value of 0.25 indicates that the Pots caught 25% of all the fish of that length class whereas 75% were caught by the GTRs. Shaded areas: 95% confidence intervals.

