

# A little frog leaps a long way: Multiple invasions to the Indian Subcontinent discovered in the Oriental tiny frog genus *Microhyla* (Amphibia: Microhylidae) (#46140)

1

First submission

## Guidance from your Editor

Please submit by **19 Mar 2020** for the benefit of the authors (and your \$200 publishing discount) .



### Structure and Criteria

Please read the 'Structure and Criteria' page for general guidance.



### Custom checks

Make sure you include the custom checks shown below, in your review.



### Raw data check

Review the raw data.



### Image check

Check that figures and images have not been inappropriately manipulated.

Privacy reminder: If uploading an annotated PDF, remove identifiable information to remain anonymous.

## Files

Download and review all files from the [materials page](#).

10 Figure file(s)  
11 Table file(s)  
2 Raw data file(s)  
2 Other file(s)

## ! Custom checks

### DNA data checks

- ! Have you checked the authors [data deposition statement](#)?
- ! Can you access the deposited data?
- ! Has the data been deposited correctly?
- ! Is the deposition information noted in the manuscript?



# Structure and Criteria

## Structure your review

The review form is divided into 5 sections. Please consider these when composing your review:

1. BASIC REPORTING
2. EXPERIMENTAL DESIGN
3. VALIDITY OF THE FINDINGS
4. General comments
5. Confidential notes to the editor

 You can also annotate this PDF and upload it as part of your review

When ready [submit online](#).

## Editorial Criteria

Use these criteria points to structure your review. The full detailed editorial criteria is on your [guidance page](#).

### BASIC REPORTING

-  Clear, unambiguous, professional English language used throughout.
-  Intro & background to show context. Literature well referenced & relevant.
-  Structure conforms to [Peerj standards](#), discipline norm, or improved for clarity.
-  Figures are relevant, high quality, well labelled & described.
-  Raw data supplied (see [Peerj policy](#)).

### EXPERIMENTAL DESIGN

-  Original primary research within [Scope of the journal](#).
-  Research question well defined, relevant & meaningful. It is stated how the research fills an identified knowledge gap.
-  Rigorous investigation performed to a high technical & ethical standard.
-  Methods described with sufficient detail & information to replicate.

### VALIDITY OF THE FINDINGS

-  Impact and novelty not assessed. Negative/inconclusive results accepted. *Meaningful* replication encouraged where rationale & benefit to literature is clearly stated.
-  All underlying data have been provided; they are robust, statistically sound, & controlled.
-  Speculation is welcome, but should be identified as such.
-  Conclusions are well stated, linked to original research question & limited to supporting results.

# Standout reviewing tips

3



The best reviewers use these techniques

## Tip

**Support criticisms with evidence from the text or from other sources**

## Example

*Smith et al (J of Methodology, 2005, V3, pp 123) have shown that the analysis you use in Lines 241-250 is not the most appropriate for this situation. Please explain why you used this method.*

**Give specific suggestions on how to improve the manuscript**

*Your introduction needs more detail. I suggest that you improve the description at lines 57- 86 to provide more justification for your study (specifically, you should expand upon the knowledge gap being filled).*

**Comment on language and grammar issues**

*The English language should be improved to ensure that an international audience can clearly understand your text. Some examples where the language could be improved include lines 23, 77, 121, 128 – the current phrasing makes comprehension difficult.*

**Organize by importance of the issues, and number your points**

1. Your most important issue
2. The next most important item
3. ...
4. The least important points

**Please provide constructive criticism, and avoid personal opinions**

*I thank you for providing the raw data, however your supplemental files need more descriptive metadata identifiers to be useful to future readers. Although your results are compelling, the data analysis should be improved in the following ways: AA, BB, CC*

**Comment on strengths (as well as weaknesses) of the manuscript**

*I commend the authors for their extensive data set, compiled over many years of detailed fieldwork. In addition, the manuscript is clearly written in professional, unambiguous language. If there is a weakness, it is in the statistical analysis (as I have noted above) which should be improved upon before Acceptance.*

# A little frog leaps a long way: Multiple invasions to the Indian Subcontinent discovered in the Oriental tiny frog genus *Microhyla* (Amphibia: Microhylidae)

Vladislav A Gorin<sup>1</sup>, Evgeniya N Solovyeva<sup>2</sup>, Mahmudul Hasan<sup>3</sup>, Hisanori Okamiya<sup>4</sup>, D.M.S. Suranjan Karunarathna<sup>5</sup>, Parinya Pawangkhanant<sup>6</sup>, Anslem de Silva<sup>7</sup>, Watinee Juthong<sup>8</sup>, Luan Thanh Nguyen<sup>9</sup>, Chatmongkon Suwannapoom<sup>6</sup>, Alexander Haas<sup>10</sup>, David P Bickford<sup>11</sup>, Indraneil Das<sup>12</sup>, Nikolay A Poyarkov<sup>Corresp. 1, 13</sup>

<sup>1</sup> Faculty of Biology, Department of Vertebrate Zoology, Lomonosov Moscow State University, Moscow, Russia

<sup>2</sup> Zoological Museum, Lomonosov Moscow State University, Moscow, Russia

<sup>3</sup> Department of Fisheries, Bangamata Sheikh Fazilatunnesa Mujib Science & Technology University, Jamalpur, Bangladesh

<sup>4</sup> Department of Biological Science, Faculty of Science, Tokyo Metropolitan University, Tokyo, Japan

<sup>5</sup> Nature Explorations and Education Team, Moratuwa, Sri Lanka

<sup>6</sup> School of Agriculture and Natural Resources, University of Phayao, Phayao, Thailand

<sup>7</sup> Amphibia and Reptile Research Organization of Sri Lanka, Gampola, Sri Lanka

<sup>8</sup> Prince of Songkla University, Songkhla, Thailand

<sup>9</sup> Asian Turtle Program—Indo-Myanmar Conservation, Hanoi, Vietnam

<sup>10</sup> Center for Natural History, Universität Hamburg, Hamburg, Germany

<sup>11</sup> Biology Department, University of La Verne, La Verne, California, United States

<sup>12</sup> Institute of Biodiversity and Environmental Conservation, Universiti Malaysia Sarawak, Kota Samarahan, Malaysia

<sup>13</sup> Joint Russian-Vietnamese Tropical Research and Technological Center, Hanoi, Vietnam

Corresponding Author: Nikolay A Poyarkov  
Email address: n.poyarkov@gmail.com

Frogs of the genus *Microhyla* include some of the world's smallest amphibians and represent the largest radiation of Asian microhylids, currently encompassing 50 species, distributed across the Oriental biogeographic region. The small-sized genus *Microhyla* remains one of the taxonomically most challenging groups of Asian frogs and was found to be paraphyletic with respect to large-sized fossorial *Glyphoglossus* (Matsui et al., 2011). We present an updated phylogeny of the genus with a nearly complete taxon sampling including 48 nominal *Microhyla* species and several undescribed candidate species. Phylogenetic analyses of 3207 bp of both mtDNA and nuDNA data recovered three well-supported groups: the *Glyphoglossus* clade, Southeast Asian *Microhyla* II clade (includes *M. annectens* species group), and a diverse *Microhyla* I clade including all other species. Within the largest major clade of *Microhyla* are seven well-supported subclades, that we identify as *M. achatina*, *M. fissipes*, *M. berdmorei*, *M. superciliaris*, *M. ornata*, *M. butleri*, and *M. palmipes* species groups. Phylogenetic position of 12 poorly known *Microhyla* species is clarified for the first time. These phylogenetic results, along with molecular clock and ancestral area analyses, show the *Microhyla* – *Glyphoglossus* assemblage to have

originated in Southeast Asia in the middle Eocene just after the first hypothesized land connections between the Indian Plate and the Asian mainland. While *Glyphoglossus* and *Microhyla* II remained within their ancestral ranges, *Microhyla* I expanded its distribution generally east to west colonizing and diversifying through the Cenozoic. The Indian Subcontinent was colonized by members of five *Microhyla* species groups independently, starting with the end Oligocene – early Miocene that coincides with an onset of seasonally dry climates in South Asia. Body size evolution modeling suggests that four groups of *Microhyla* have independently achieved extreme miniaturization with adult body size below 15 mm. Three of the five smallest *Microhyla* species are obligate phytotelm-breeders and we argue that their peculiar reproductive biology may be a factor involved in miniaturization. Body size increases in *Microhyla* – *Glyphoglossus* seem to be associated with a burrowing adaptation to seasonally dry habitats. Species delimitation analyses suggest a vast underestimation of species richness and diversity in *Microhyla* and reveal 15 to 33 undescribed species. We revalidate *M. nepenthicola*, synonymize *M. pulverata* with *M. marmorata*, and provide insights to taxonomic statuses of a number of poorly known species. Further integrative studies, combining evidence from phylogeny, morphology, advertisement calls, and behavior will result in a better systematic understanding of this morphologically cryptic radiation of Asian frogs.

# A little frog leaps a long way: Multiple invasions to the Indian Subcontinent discovered in the Oriental tiny frog genus *Microhyla* (Amphibia: Microhylidae)

Vladislav A. Gorin<sup>1</sup>, Evgeniya N. Solovyeva<sup>2</sup>, Mahmudul Hasan<sup>3</sup>, Hisanori Okamiya<sup>4</sup>, D.M.S. Suranjan Karunarathna<sup>5</sup>, Parinya Pawangkhanant<sup>6</sup>, Anslem de Silva<sup>7</sup>, Watinee Juthong<sup>8</sup>, Luan Thanh Nguyen<sup>9</sup>, Chatmongkon Suwannapoom<sup>6</sup>, Alexander Haas<sup>10</sup>, David P. Bickford<sup>11</sup>, Indraneil Das<sup>12</sup>, Nikolay A. Poyarkov<sup>1,13\*</sup>

*1 – Faculty of Biology, Department of Vertebrate Zoology, Lomonosov Moscow State University, Moscow, Russia*

*2 – Zoological Museum, Lomonosov Moscow State University, Moscow, Russia*

*3 – Department of Fisheries, Bangamata Sheikh Fazilatunnesa Mujib Science & Technology University, Jamalpur, Bangladesh*

*4 – Department of Biological Science, Faculty of Science, Tokyo Metropolitan University, Tokyo, Japan*

*5 – Nature Explorations and Education Team, Moratuwa, Sri Lanka*

*6 – School of Agriculture and Natural Resources, University of Phayao, Phayao, Thailand*

*7 – Amphibia and Reptile Research Organization of Sri Lanka, Gampola, Sri Lanka.*

*8 – Prince of Songkla University, Songkhla, Thailand*

*9 – Asian Turtle Program—Indo-Myanmar Conservation, Hanoi, Vietnam*

*10 – Universität Hamburg, Center for Natural History, Hamburg, Germany*

*11 – Biology Department, University of La Verne, La Verne, California, USA*

*12 – Institute of Biodiversity and Environmental Conservation, Universiti Malaysia Sarawak, Kota Samarahan, Malaysia*

*13 – Joint Russian-Vietnamese Tropical Research and Technological Center, Hanoi, Vietnam*

*\* – corresponding author; E-mail: [n.poyarkov@gmail.com](mailto:n.poyarkov@gmail.com)*

## Abstract

Frogs of the genus *Microhyla* include some of the world's smallest amphibians and represent the largest radiation of Asian microhylids, currently encompassing 50 species, distributed across the

Oriental biogeographic region. The small-sized genus *Microhyla* remains one of the taxonomically most challenging groups of Asian frogs and was found to be paraphyletic with respect to large-sized fossorial *Glyphoglossus* (Matsui *et al.*, 2011). We present an updated phylogeny of the genus with a nearly complete taxon sampling including 48 nominal *Microhyla* species and several undescribed candidate species. Phylogenetic analyses of 3207 bp of both mtDNA and nuDNA data recovered three well-supported groups: the *Glyphoglossus* clade, Southeast Asian *Microhyla* II clade (includes *M. annectens* species group), and a diverse *Microhyla* I clade including all other species. Within the largest major clade of *Microhyla* are seven well-supported subclades, that we identify as *M. achatina*, *M. fissipes*, *M. berdmorei*, *M. superciliaris*, *M. ornata*, *M. butleri*, and *M. palmipes* species groups. Phylogenetic position of 12 poorly known *Microhyla* species is clarified for the first time. These phylogenetic results, along with molecular clock and ancestral area analyses, show the *Microhyla* – *Glyphoglossus* assemblage to have originated in Southeast Asia in the middle Eocene just after the first hypothesized land connections between the Indian Plate and the Asian mainland. While *Glyphoglossus* and *Microhyla* II remained within their ancestral ranges, *Microhyla* I expanded its distribution generally east to west colonizing and diversifying through the Cenozoic. The Indian Subcontinent was colonized by members of five *Microhyla* species groups independently, starting with the end Oligocene – early Miocene that coincides with an onset of seasonally dry climates in South Asia. Body size evolution modeling suggests that four groups of *Microhyla* have independently achieved extreme miniaturization with adult body size below 15 mm. Three of the five smallest *Microhyla* species are obligate phytotelm-breeders and we argue that their peculiar reproductive biology may be a factor involved in miniaturization. Body size increases in *Microhyla* – *Glyphoglossus* seem to be associated with a burrowing adaptation to seasonally dry habitats. Species delimitation analyses suggest a vast underestimation of species richness and diversity in *Microhyla* and reveal 15 to 33 undescribed species. We revalidate *M. nepenthicola*, synonymize *M. pulverata* with *M. marmorata*, and provide insights to taxonomic statuses of a number of poorly known species. Further integrative studies, combining evidence from phylogeny, morphology, advertisement calls, and behavior will result in a better systematic understanding of this morphologically cryptic radiation of Asian frogs.

**Subjects:** Biodiversity, Biogeography, Evolutionary Studies, Taxonomy, Zoology

**Keywords:** molecular phylogeny, biogeography, miniaturization, Narrow-mouthed frogs, Southeast Asia, Microhylinae, species delimitation, Indian collision, cryptic species, *Glyphoglossus*

# Introduction

The tectonic collision between the Indian subcontinent (ISC) and the Eurasian landmass during the Early Cenozoic is widely recognized as a key event that caused significant geologic and climatic changes, such as the rise of the Himalaya, uplift of the Tibetan plateau, and a general drying of Central Asia (Harrison *et al.*, 1992; An *et al.*, 2001; Guo *et al.*, 2002; Molnar, 2005; Solovyeva *et al.*, 2018). However, this tectonic event also induced a major biotic exchange between the ISC and Eurasia and is widely regarded as a major driver of biotic diversification (Wilkinson *et al.*, 2002; Roelants *et al.*, 2004; Li *et al.*, 2013; Garg & Biju, 2019). Numerous studies demonstrated that Late Cretaceous floral and faunal elements reached and colonized tropical Asia from Gondwanaland via the northward drifting ISC (Dayanandan *et al.*, 1999; Klaus *et al.*, 2010, 2016; Kamei *et al.*, 2012; Morley, 2018), and many works regard the Indian Plate as an important source of Southeast Asian biodiversity, the so called “out-of-India” hypothesis (Bossuyt & Milinkovitch, 2001; Conti *et al.*, 2002; Gower *et al.*, 2002; Biju & Bossuyt, 2003; Sparks, 2003; Dutta *et al.*, 2004; Karanth, 2006; Datta-Roy & Karanth, 2009). However, a set of phylogenetic studies of different animal groups proposed an alternative “out-of-Eurasia” biogeographic hypothesis, suggesting a Southeast Asian origin of these lineages with further dispersal and colonization of ISC during its collision with Eurasian landmass (Raxworthy *et al.*, 2002; Renner, 2004; Köhler & Glaubrecht, 2007; Van der Meijden *et al.*, 2007; Macey *et al.*, 2008; Grismer *et al.*, 2016; Garg & Biju, 2019).

The tropical areas of South and Southeast Asia include four biogeographic regions recognized as global biodiversity hotspots (Myers *et al.*, 2000; Che *et al.*, 2010; De Bruyn *et al.*, 2014). Understanding the processes that sculpted this diversity is hampered by lack of comprehensive geological models and ongoing debate on timing and topography of the ISC–Eurasian collision (Acton, 1999; Van Hinsbergen *et al.*, 2011, 2012; Ali & Aitchison, 2008, 2012). Recent geologic models suggest land bridges connected ISC and modern Southeast Asia since the early Eocene (ca. 55–35 MYA), well before collision of the Indian plate with Eurasia (30–25 MYA) (Aitchison *et al.*, 2007; Aitchison & Ali, 2012; Hall, 2012; Ding *et al.*, 2017).



Several phylogenetic studies corroborate existence of pre-collision faunal exchanges between the ISC and Southeast Asia, demonstrating that they likely went in both directions: “out-of-India” and “out-of-Eurasia” (Klaus *et al.*, 2010; Li *et al.*, 2013; Grismer *et al.*, 2016; Garg & Biju, 2019). Combining data on tectonics, climatic history, and phylogenetics has proved to be a powerful instrument for examining patterns of diversification within clades and understanding processes involved in the assembly of high biodiversity in regions like South and Southeast Asia (e.g. De Bruyn *et al.*, 2014).

Frogs of the family Microhylidae are some of the most species rich groups of Anura, comprising 688 species in 13 subfamilies (Frost, 2020). Because of their transcontinental circumtropical distribution, microhylids are a promising test case for biogeography studies (Savage, 1973; Van Bocxlaer *et al.*, 2006; Van der Meijden *et al.*, 2007; Kurabayashi *et al.*, 2011). Among the 13 currently recognized subfamilies of microhylids, the subfamily Microhylinae is widely distributed in South, Southeast, and East Asia currently including eight genera with nearly a hundred species (Garg & Biju, 2019; Frost, 2020). Their phylogenetic relationships and historical biogeography have been discussed in several studies (Van Bocxlaer *et al.*, 2006; Frost *et al.*, 2006; Matsui *et al.*, 2011; de Sá *et al.*, 2012; Peloso *et al.*, 2016; Feng *et al.*, 2017). The most recent analysis of genus-level phylogeny within the Microhylinae by Garg & Biju (2019) suggested their origin on the ISC during early Paleocene with dispersal to the Asian mainland via several Eocene land bridges connecting the ISC with Southeast Asia. Following accretion of India and Eurasia in the Oligocene/Miocene, some Microhylinae lineages that diversified in Southeast Asia could have recolonized the ISC (Garg & Biju, 2019). However, phylogenetic relationships and historical biogeography within Microhylinae genera remain poorly resolved.

The genus *Microhyla* is the most species rich genus in the Microhylinae, currently comprising 50 recognized species (Poyarkov *et al.*, 2014, 2019; Biju *et al.*, 2019). Over half of this diversity has been described within the last 15 years (27 species, see Frost, 2020), yet *Microhyla* remains one of the most taxonomically challenging groups of Asian frogs. *Microhyla* is also the only Asian microhylid genus with a wide distribution over South, Southeast, and East Asia (see Fig. 1), making it an ideal model for studies on Asian biogeography. Monophyly of *Microhyla* was questioned by Matsui *et al.* (2011), based on analysis of mitochondrial DNA (mtDNA) markers, but was later corroborated by multilocus phylogenetic analyses (Peloso *et al.*,

2016; Tu et al., 2018). Most species of *Microhyla* are small-sized terrestrial frogs, while several diminutive species approach the lower body-size limit for vertebrates and represent some of the world's tiniest amphibians (Das & Haas, 2010; Poyarkov et al., 2014). In miniaturized groups of amphibians, a high proportion of cryptic diversity and rampant homoplasies are often recorded, muddying our estimates of diversity and evolutionary relationships (e.g. Rovito et al., 2013; Parra-Olea et al., 2016; Rakotoarison et al., 2017). Molecular phylogenetic analyses, optimally combined with behavioral and acoustic data, offer the best hope for clarifying diversity, species boundaries, and evolutionary relationships in many groups of Microhylidae, including the genus *Microhyla* (Hasan et al., 2014a; Garg et al., 2019; Poyarkov et al., 2018a, 2019).

Phylogenetic relationships among members of the genus *Microhyla* have been discussed in several recent studies (Matsui et al., 2011; Peloso et al., 2016; Tu et al., 2018; Nguyen et al., 2019; Biju et al., 2019). However, they were generally based on limited sampling (< 60% of recognized diversity), while few works have provided insights on biogeographic origin, patterns of distribution, and possible routes of colonization for the genus (Vineeth et al., 2018; Garg et al., 2019; Poyarkov et al., 2019). Though the greatest species diversity of *Microhyla* is observed in Southeast Asia (up to nine sympatric species in Indochina, see Fig. 1), some studies suggested the possibility of an Indian origin for the genus and several species groups (Garg et al., 2019; Garg & Biju, 2019). Extensive taxon sampling of all known members of the genus *Microhyla*, a more robustly resolved phylogeny, and sound divergence age estimates are important to understand historical distribution and diversification in this radiation of Asian frogs (Garg et al., 2019). Thus far, however, a comprehensive phylogenetic investigation with dating estimates within the genus *Microhyla* is lacking.

Herein, we identify unrecognized diversity and examine phylogenetic relationships among most species of *Microhyla* based on extensive geographic and taxonomic sampling, including 48 of 50 nominal species (96% of recognized taxa); phylogenetic information for 12 species and a number of undescribed species is reported for the first time. We use the resulting phylogeny (based on both mtDNA and nuDNA markers) to test biogeographic hypotheses in space and time and provide a scenario for *Microhyla* diversification. Our study provides the first nearly complete phylogeny for the genus *Microhyla* and links the Indian Subcontinent with Sundaland, Indo-Burma, and East Asia, thereby allowing a better understanding of biogeographic history and diversification of the group. We also evaluate miniaturization and

simulate body size evolution across different lineages of the genus, providing novel insights into morphological evolution in *Microhyla*.

## Materials and methods

### Taxon sampling

We used tissue samples stored in herpetological collections of Zoological Museum of Moscow University (ZMMU; Moscow, Russia); Zoological Institute, Russian Academy of Sciences (ZISP; St. Petersburg, Russia); Vertebrate Zoology Department, Biological Faculty, Moscow State University (ZPMSU; Moscow, Russia); Amphibian Research Center, Hiroshima University (IABHU; Higashihiroshima, Japan); Danum Valley Conservation Area, Specimen collection (RMBR; Sabah, Malaysia); Department of Fisheries, Bangamata Sheikh Fazilatunnesa Mujib Science & Technology University (DFBSFMSTU; Jamalpur, Bangladesh); School of Agriculture and Natural Resources, University of Phayao (AUP; Phayao, Thailand); and the Institute of Biodiversity and Environmental Conservation, Universiti Malaysia Sarawak, (UNIMAS; Sarawak, Malaysia).

We analyzed 122 tissue samples representing 40 nominal species of *Microhyla*, 14 species have not been included in previous phylogenetic analyses. In our analysis, we also included GenBank sequences from 78 specimens of approximately 37 nominal *Microhyla* species, 29 other Microhylidae representatives, and five non-microhylid outgroups used for rooting the phylogenetic tree and divergence times estimation (information summarized in Table S1). In total, we obtained molecular genetic data for 199 samples representing 48 nominal *Microhyla* species (Table S1). Geographic location of sampled populations is presented in Fig. S1. For alcohol-preserved voucher specimens stored in museum collections, muscle tissue was removed, preserved in 96% ethanol, and stored subsequently at -70°C.

### DNA extraction, amplification and sequencing

For molecular phylogenetic analyses, total genomic DNA was extracted from ethanol-preserved femoral muscle tissue using standard phenol-chloroform-proteinase K (final concentration 1 mg/ml) extraction procedures with consequent isopropanol precipitation (protocols followed *Sambrook & Russell, 2001*; *Swofford et al., 1996*).

For mtDNA, we amplified sequences covering fragments of 12S rRNA, tRNA<sup>Val</sup>, and 16S rRNA mtDNA genes to obtain an up to 2478 bp-length continuous fragment of mtDNA. The

16S rRNA gene has been widely applied in biodiversity surveys in amphibians (*Vences et al., 2005; Vieites et al., 2009*) and 12S rRNA + 16S rRNA data have been used in several important studies on Microhylinae phylogeny (*Matsui et al. 2011; Peloso et al. 2016*). These fragments have also proven to be particularly useful in taxonomic studies of the genus *Microhyla* and closely-related taxa (e.g. *Hasan et al., 2012, 2014a, 2014b; Howlader et al., 2015; Matsui, 2011; Matsui et al., 2011, 2013; Wijayathilaka et al., 2016*).

For nuDNA, we amplified a 729 bp-long fragment of brain-derived neurotrophic factor gene (*BDNF*). This marker was recently successfully applied in biodiversity and phylogenetic studies of Indian *Microhyla* species (see *Garg et al., 2019; Garg & Biju, 2019; Biju et al., 2019*). Primers used in PCR and sequencing were taken from the literature or designed by us and summarized in Table S2.

The PCR conditions for amplifying mtDNA fragments included an initial denaturation step of 5 min at 94°C, and 40 cycles of denaturation for 1 min at 94°C, primer annealing step for 1 min with TouchDown program from 65°C reducing 1°C every cycle to 55°C, and extension step for 1 min at 72°C, and the final extension step for 5 min at 72°C. The PCR conditions for amplifying *BDNF* gene followed *Van der Meijden et al. (2007)* and included an initial denaturation step of 5 min at 94°C followed with 32 cycles of denaturation for 1 min at 94°C, primer annealing step for 1 min at 50°C, and extension for 1 min at 72°C, with final extension step for 5 min at 72°C. Sequence data collection and visualization were performed on an ABI 3730xl automated sequencer (Applied Biosystems). PCR purification and cycle sequencing were done commercially through Evrogen® (Moscow, Russia). All unique sequences were deposited in GenBank (see Table S1).

### Phylogenetic analyses

In addition to newly obtained sequences, we added 107 sequences of Microhylidae available in GenBank to the final alignments; sequences of *Rhacophorus schlegelii*, *Alytes dickhilleni*, *A. muletensis*, *Blommersia transmarina* and *B. wittei* were selected as outgroup taxa for phylogenetic inference and were also used for time-tree calibration. Details on taxonomy, localities, GenBank accession numbers, and associated references for all examined specimens are summarized in Table S1.

Nucleotide sequences were initially aligned in MAFFT v.6 (Kato et al., 2002) with default parameters, subsequently checked by eye in BioEdit 7.0.5.2 (Hall, 1999), and adjusted as needed.

Phylogenetic tree reconstructions were performed with the following two data sets:

(1) A mtDNA dataset joining 12S rRNA and 16S rRNA for all examined samples, used for assessment of species groups and estimation of cryptic diversity within *Microhyla* (230 sequences, including 199 sequences of *Microhyla*);

(2) A concatenated mtDNA + nuDNA dataset, joining long 12S rRNA – 16S rRNA mtDNA fragment and *BDNF* gene sequences for 118 selected samples representing all major lineages within *Microhyla* (as revealed by initial analysis of mtDNA ), used for obtaining a more robust phylogenetic hypothesis, time-tree estimation, and ancestral range reconstruction for *Microhyla*.

The optimum partitioning schemes for alignments were identified with PartitionFinder 2.1.1 (Lanfear et al., 2012) using the *greedy* search algorithm under an AIC criterion, and are presented in Table S3. Phylogenetic trees were reconstructed with maximum likelihood (ML) and Bayesian algorithms (BI). ML trees were generated in IQ-TREE (Nguyen et al., 2015). Confidence in tree topology for ML analysis was assessed by 1000 bootstrap replications for ML analysis (ML BS). Bayesian inference (BI) was performed in MrBayes v3.1.2 (Ronquist & Huelsenbeck, 2003) with two simultaneous runs, each with four chains for 200 million generations. We checked that the effective sample sizes (ESS) were all above 200 by exploring likelihood plots using TRACER v1.6 (Rambaut & Drummond, 2007). The initial 10% of trees were discarded as burn-in. Confidence in tree topology was assessed by posterior probability for Bayesian analysis (BI PP) (Huelsenbeck & Ronquist, 2001). We *a priori* regarded tree nodes with ML BS values 75% or greater and BI PP values over 0.95 as sufficiently resolved (Huelsenbeck & Hillis, 1993; Felsenstein, 2004). For clarity, ML BS values between 75% and 50% (BI PP between 0.95 and 0.90) were regarded as tendencies and below 50% (BI PP below 0.90) were considered unresolved.

### Species delimitation

We examined putative species boundaries beyond those currently recognized by taxonomists based on two different species delimitation methods: The Automatic Barcode Gap Discovery (ABGD; Puillandre et al., 2012) and the Generalized Mixed Yule-Coalescent model

(GMYC; *Pons et al., 2006*). These methods enable the delimitation of independently-evolving species based on genetic data (*Fujita et al., 2012; Dellicour & Flot, 2015; Eyer et al., 2017*) and do not require *a priori* hypotheses of putative species groupings, thereby limiting potential bias in species delimitation.

The ABGD method is a single-gene approach to statistical detection of barcode gaps in a pairwise genetic distance distribution (*Puillandre et al., 2012*). Barcode gaps, presumably occurring between intra- and interspecific distances, were used to partition the 16S rRNA dataset into species hypotheses (initial partition). Resulting inferences were then recursively applied to yield finer partitions (recursive partitions) until no further partitioning was possible. ABGD analysis was run on the 16S rRNA dataset through a web-based interface (<http://www.abi.snv.jussieu.fr/public/abgd/>) using default parameters (10 steps of intraspecific divergence prior from  $P_{min} = 0.001$  to  $P_{max} = 0.10$ ,  $X = 2$ ).

The GMYC method is also a single-gene approach to identifying species ‘boundaries’ associated with shifts in branching rates between intra- and interspecies branching events on a time-calibrated ultrametric tree (*Pons et al., 2006; Fujisawa & Barraclough, 2013*). We used a Bayesian implementation of this method (bGMYC; *Reid & Carstens, 2012*), which was applied to the 16S rRNA data. We obtained the distribution of ultrametric phylogenetic trees of 16S rRNA haplotypes with BEAST v1.8.4 (*Drummond et al., 2012*), then used 100 random phylogenetic trees as an input for subsequent bGMYC analysis. We ran bGMYC for 50,000 generations with burn-in 40,000 and a thinning parameter of 100. We summarized results of bGMYC analyses in a matrix of pairwise co-assignment probabilities for each haplotype, shown as a heatmap (not presented).

In addition, both inter- and intraspecific uncorrected genetic *p*-distances were calculated using MEGA 6.1 (*Tamura et al., 2013*).

### **Divergence times estimation**

Molecular divergence dating was performed in BEAST v1.8.4, including the concatenated mtDNA + nuDNA dataset. We tested the molecular clock assumption separately for mtDNA and nuDNA markers with hierarchical likelihood ratio tests using PAML v4.7 (*Yang, 2007*). Following results of these tests, we used a strict clock model for BDNF and uncorrelated lognormal relaxed clock models for the mtDNA markers. The clock rate for BDNF was set to one. We used the same models and partitioning scheme as in the ML analysis. The Yule model

was set as the tree prior and the piecewise constant population size model was assumed; default priors were used for all other parameters. Two runs of 100 million generations were conducted in BEAST v1.8.4. Parameter convergence was assessed in Tracer with the first 10% of generations discarded as burn-in. We used TreeAnnotator v1.8.0 (part of the BEAST package) to generate the maximum clade credibility tree.

Because no reliable paleontological data have been reported for the Microhylinae, we used the three recently estimated calibration priors for this subfamily obtained from recent large-scale phylogenies of microhylids (Kurabayashi *et al.*, 2011), and a fossil record of Gastrophryninae (Sanchiz, 1998; Holman, 2003; de Sá *et al.*, 2012) as primary calibration points. We also applied two additional calibration points widely used in divergence time estimates of Anura: maximum age of the split between *Blommersia wittei* and *B. transmarina* from the Comoro islands at 15 MYA (Vences *et al.*, 2003), and the minimum age of *Alytes muletensis* – *A. dickhilleni* split at 5 MYA (Fromhage *et al.*, 2004). Calibration points and priors are summarized in Table S4.

# **Ancestral area reconstruction**

To infer a biogeographic history of *Microhyla*, a model-testing approach was applied using the ML tree of Lagrange (Ree *et al.*, 2005; Ree & Smith, 2008) in RASP 3.2 (Yu *et al.*, 2015). Species occurrences were categorized according to nine biogeographic areas, modified from Turner *et al.* (2001), Wood *et al.* (2012) and Chen *et al.* (2018), reflecting patterns of endemism in *Microhyla* (see Fig. 2A): (A) Mainland East Asia; (B) Eastern Indochina; (C) Western Indochina; (D) Indian Subcontinent; (E) Malayan Peninsula; (F) Sumatra + Java + Bali; (G) Borneo and adjacent Philippine islands; (H) Sri Lanka; and (I) East Asian islands (Taiwan + the Ryukyus) (for details on biogeographic regions definition and references see Supplemental Information 1). Maximum range-size was set to three areas, as no extant species occurs in more than three biogeographical regions. Matrices of modern distributions of taxa/ area are given in Table S5. Transitions between discrete states (ranges) along tree branches were modeled as a function of time, thus enabling ML estimation of the ancestral states at cladogenic events. A Lagrange analysis found the most likely ancestral area(s) at a node, the split of areas in the two descendant lineages, and calculated probabilities of these most likely areas for each node (Ree & Smith, 2008). Analyses used two models (Matzke, 2013): Lagrange Dispersal-Extinction-Cladogenesis (DEC; Ree & Smith, 2008), and the ML version of Statistical Dispersal-Vicariance

Analysis (S-DIVA; *Ronquist, 1997*). We assessed model fit using the Akaike Information Criterion (AIC) and Akaike weights.

Given the tremendous geological complexity of the region through time, we applied the following time and dispersal constraints to the analyses. We set four periods, corresponding to the main stages of gradual ISC movements northwards, formation of the first land bridges between the ISC and Southeast Asia, and final accretion between the ISC and the Asian mainland based on data from recent geologic models (based on *Hall, 2012*; *Ding et al., 2017*; *Morley, 2018*; see Fig. 3 for schematic paleogeographic maps). The following time periods were designated: (1) 100–57 MYA corresponds to complete isolation of the ISC from Eurasia, (2) 57–50 MYA marks the first assumed land connections between India and modern-day Sumatra; (3) during 50–35 MYA the ISC likely continued counter-clockwise movement northwards, forming land bridges with modern-day Indo-Burma; and (4) 35–0 MYA corresponds to the firm collision and formation of a stable land connection between the ISC and Eurasia. Transition matrices between biogeographic regions for each time period are presented in Table S6.

### Body size evolution

To assess evolution of body size and miniaturization in *Microhyla*, we used weighted squared-change parsimony (*Maddison, 1991*) as implemented in Mesquite v3.31 (*Maddison & Maddison, 2017*). We compiled data on maximum snout-vent length (SVL) of males and females for each *Microhyla* species as reported in literature and/or based on examination of available voucher specimens. SVL data for each *Microhyla* species are provided in Table S7.

## Results

### Taxon sampling, data collection and sequence characteristics

The complete aligned matrix of mtDNA data comprised 2478 bp, included 206 samples, representing 48 species of *Microhyla* (96% of the currently recognized species), five species (of nine currently recognized species) of the phylogenetically closely related genus *Glyphoglossus*, and 24 samples from outgroup taxa (see Table S1).

The concatenated mtDNA + nuDNA dataset comprised 3207 bp, including 118 samples from 100 ingroup and 18 outgroup taxa. Information on length of obtained fragments and their variability are given in Table S8.

### Phylogenetic relationships and species groups in *Microhyla*



The complete mtDNA-based genealogy of *Microhyla* and *Glyphoglossus* is presented in Figs. 4 and 5; a simplified collapsed tree is shown in Fig. S2. Bayesian Inference and Maximum Likelihood analyses of concatenated mtDNA + nuDNA data resulted in a topology that was generally congruent with the mtDNA-based genealogy, but with higher support for most nodes (Fig. 6). Overall, since the mtDNA + nuDNA phylogenetic tree was generally better resolved and showed higher nodal support values than the mtDNA tree, we relied on the mtDNA + nuDNA topology for inferring phylogenetic relationships and biogeographic history of the genus *Microhyla*.

The BI- and ML-analyses of the mtDNA data resulted in ingroup nodes the majority of which received high values of PP and BS support (Figs. 4 and 5). Observed topological patterns within the *Microhyla* – *Glyphoglossus* assemblage were generally congruent across analyses and agreed well with earlier phylogenies for the group (Matsui *et al.*, 2005, 2011, 2013; Matsui, 2011; Hasan *et al.*, 2012, 2014a; Howlader *et al.*, 2015; Peloso *et al.*, 2016; Wijayathilaka *et al.*, 2016; Seshadri *et al.*, 2016; Yuan *et al.*, 2016; Khatiwada *et al.*, 2017; Tu *et al.*, 2018; Garg *et al.*, 2019; Nguyen *et al.*, 2019; Poyarkov *et al.*, 2019; Biju *et al.*, 2019), although with generally higher node support values in our study.

All analyses unambiguously supported the monophyly of the *Microhyla* – *Glyphoglossus* assemblage; however, the basal node of this radiation was not sufficiently resolved in all analyses (Fig. 6 and Fig. S2). In agreement with Matsui *et al.* (2011), the genus *Microhyla sensu lato* was subdivided into two major deeply divergent groups: *Microhyla* I and *Microhyla* II, while monophyly of the genus with respect to *Glyphoglossus* was not supported according to mtDNA data (see Fig. S2). Though basal divergence in the *Microhyla* – *Glyphoglossus* clade based on mtDNA + nuDNA data was also not strongly supported (Fig. 6), the topology suggesting monophyly of *Microhyla* I + *Microhyla* II agreed well with results of recent multilocus phylogenies for this group (Peloso *et al.*, 2016; Tu *et al.*, 2018).

All species of *Microhyla* I, *Microhyla* II, and *Glyphoglossus* clades were consistently assigned to one of the nine generally strongly supported matrilineages (Figs. 4 and 5, A–I); same phylogenetic groupings were also revealed in the “total evidence” analysis (Fig. 6).

*Microhyla* I (subclade AI of Matsui *et al.*, 2011) included seven major clades and 43 putative species of tiny to mid-sized terrestrial frogs (Figs. 4 and 5, A–G):

- (A) Clade A corresponded to *M. achatina* species group and got only moderate levels of monophyly support in mtDNA-genealogy (0.90/61, hereafter node support values are given for BI PP / ML BS, respectively) (Fig. 4). Genealogical relationships within this group were poorly resolved. Phylogenetic positions of *M. fodiens* from central Myanmar, the *M. heymonsi* complex and *M. pineticola* from Indochina were unresolved. Other species form a strongly supported monophyly (1.0/98), which is further subdivided into two subclades: (A1) comprising species from Sundaland (*M. borneensis*, *M. nepenthicola*, *Microhyla* sp. 1 from Sabah, *M. malang*, *M. orientalis*, *M. mantheyi*) and southern Vietnam (*M. minuta*) (0.97/68), and (A2) with species from Sundaland (*M. achatina*, *M. gadjahmadai*), Myanmar (*M. irrawaddy*, *Microhyla* sp. 4 from northern Myanmar), and southern India (*M. kodial*) (1.0/87). ~~Monophyly of Clade A got no support in the “total evidence” analysis~~ (0.72/50; Fig. 6), while the clade including all members of *M. achatina* species group except *M. fodiens* got moderate support (0.91/62; Fig. 6).
- (B) Clade B corresponded to *M. fissipes* species group (1.0/100) and consisted of two well-supported subclades (Fig. 4): (B1) included species from Indochina and southern mainland China and Taiwan (*M. fissipes*, *M. mukhlesuri*), and species from eastern India, Bangladesh and the Andaman Islands (*M. mymensinghensis*, *M. chakrapanii*) (1.0/100); (B2) encompassing species from mainland China (*M. mixtura*, *M. beilunensis*, *M. fanjingshanensis*) and the Ryukyus (*M. okinavensis*, *Microhyla* sp. 3 from Yaeyama Archipelago) (1.0/98). The “total evidence” analysis suggested sister group relationships between *M. fanjingshanensis* and *M. okinavensis* (0.96/55; Fig. 6).
- (C) Clade C included the *M. berdmorei* complex and *M. pulchra* from Indochina and southern China, as well as *M. picta* from southernmost Vietnam (1.0/100). Clade C was recovered as sister clade to a clade A+B (1.0/90 for mtDNA, and 1.0/96 for “total evidence” datasets, respectively; see Figs. 5, 6).
- (D) Clade D encompassed species from Sri Lanka (*M. zeylanica*, *M. karunaratnei*) and southern India (*M. laterite*, *M. sholigari*, *M. darreli*) (D1, see Fig. 5), but also included species from northeastern India (*M. eos*), western Thailand (*Microhyla* sp. 2 from Tenasserim) and Thai-Malay Peninsula (*M. superciliaris*). The “total evidence” analysis strongly suggested sister group relationships between clades D and F (0.96/67; Fig. 6).

(E) Clade E included species with distribution on the Indian Subcontinent and Sri Lanka, including *M. ornata* complex (E1, *M. ornata*, *M. nilphamariensis*, *M. taraiensis*) and *M. rubra* complex members (E2, *M. rubra*, *M. mihintalei*). MtDNA data suggested a tendency for joining the Clades D and E in a monophylum (0.91/75; Fig. 5), but it was not supported by the “total evidence” analysis, placed Clade E as a sister group to the clade A+B+C, though with no support (0.70/37; Fig. 6). The monophyly of the clade joining matriline A–E was moderately supported by mtDNA (Fig. 5), but not by the mtDNA+nuDNA datasets (Fig. 6).

(F) Clade F corresponds to *M. butleri* species group and joined the *M. butleri* complex from southern China and Southeast Asia with *M. aurantiventris* from central Vietnam (1.0/100). Clade F was strongly supported as a sister lineage with respect to Clade E based on the “total evidence” analysis (0.96/67) (Fig. 6).

(G) Clade G was represented by the *M. palmipes* species complex from Java and Sumatra; its phylogenetic position of was poorly supported; the “total evidence” analysis suggested sister group relationships with respect to clade E+F but with moderate support (0.92/41; Fig. 6).

*Microhyla* II (subclade AIII of Matsui *et al.*, 2011) was represented by a single clade (H) and included nine nominal species of tiny to small-sized terrestrial or semi-arboreal frogs (Fig. 5, H):

(H) Clade H corresponds to *M. annectens* species group and was further subdivided in two subclades: (H1) included species from Borneo (*M. perparva*, *M. petrigena*), the Thai-Malay Peninsula (*M. annectens*), and Annamite Mountains of Indochina (*M. annamensis*, *M. marmorata* and *M. pulverata*) (1.0/88); and (H2) joined species from central (*M. nanapollexa*) and southern parts of Annamite Mountains (*M. arboricola*, *M. pulchella*). *M. marmorata* was recovered paraphyletic, with respect to *M. pulverata* (1.0/100; Fig. 5).

Finally, the large-sized fossorial frogs of the genus *Glyphoglossus* (subclade AII of Matsui *et al.*, 2011) was represented in our analysis by five of the nine recognized species; *G. capsus* from Borneo was recovered as a sister group to the clade that joined all other species from Indochina and Malay Peninsula (Fig. 5) (0.98/74).

# Species delimitation analyses

The BI matrilineal genealogy for the 48 nominal *Microhyla* species provided an initial assessment of species-level relationships (Figs. 4 and 5). Uncorrected genetic  $p$ -distances in 16S rRNA mtDNA gene within and among *Microhyla* species are given in Table S9. In addition to currently recognized taxa, our genealogy also depicted at least four lineages of *Microhyla* that likely represent unrecognized species (*Microhyla* spp. 1–4), and a number of potentially cryptic lineages within species complexes (intraspecific genetic differences  $p > 1.5\%$ , e.g., *M. malang*, *M. achatina*, *M. gadjahmadai*, *M. heymonsi*, *M. okinavensis*, *M. berdmorei*, *M. butleri*, *M. palmipes*, *M. petrigena*, *M. perparva* and *M. arboricola*; see Table S9). Generally, interspecific divergences were  $p > 3.0\%$ , but in some cases, intraspecific divergences exceeded interspecific divergences (Table S9).

To assess the number of putative species-level lineages within the genus *Microhyla*, we implemented two alternative approaches to species delimitation via tree-based bGMYC and distance-based ABGD analyses. These methods show varying performance depending on sample and population sizes, speciation rates, and other parameters, with bGMYC showing a tendency to oversplit, while ABGD often overlumps putative species; however, when these methods agree, the resulting delimitation gains plausibility (Dellicour & Flot, 2018). The results of species delimitation analyses are presented in Figs. 4 and 5, and summarized in Table S10. The combined results of both bGMYC and ABGD resolved all described species of *Microhyla*, except for *M. pulverata* that was indistinguishable from *M. marmorata*. This result is also corroborated by the mtDNA-genealogy (Fig. 5) and genetic divergence data (Table S9). Both analyses suggest that species diversity within *Microhyla* is greatly underestimated: for 48 recognized *Microhyla* species included in our genealogy, the bGMYC analysis recovered 81, and ABGD recovered 63 species-level lineages (Table S10). In most cases, analyses had congruent results, however in 14 species, bGMYC proposed more groups than were recovered by ABGD. Nonetheless, species delimitation analyses strongly indicate the presence of many unrecognized species-level lineages by further partitioning *M. heymonsi* (into 7 to 8 species), *M. bermodrei* (3 to 4 species), *M. malang* (3 species), *M. butleri* (2 to 4 species), and *M. gadjahmadai*, *M. okinavensis*, *M. palmipes*, *M. perparva*, *M. petrigena*, *M. achatina* and *M. arboricola* (each with 2 species) (Figs. 4 and 5; Table S10). For nine other taxa, the analyses gave incongruent results, with bGMYC splitting and ABGD lumping these disparate taxa (*M. beilunensis*, *M. chakrapanii*,

*M. mantheyi*, *M. mixtura*, *M. mukhlesuri*, *M. orientalis*, *M. superciliaris*, *Microhyla* sp. 3 and *Microhyla* sp. 4) (Table S10).

### Divergence times estimation

Estimated node-ages and the 95% highest posterior density (95% HPD) for the main nodes are summarized in detail in Table S11. The resulting BEAST chronogram for the ingroup is presented in Fig. 2; the general time-tree with node support and tree-calibration points is detailed in Fig. S3. Our analysis proposed that the most recent common ancestor (MRCA) of *Microhyla* and *Glyphoglossus* originated between late Paleocene and early Eocene, ca. 50.8 Ma (44.1–57.0). The group *Microhyla* + *Glyphoglossus* radiated within a comparatively narrow time period dated to the middle Eocene ca. 43.8 Ma (38.7–49.1). Diversification within the genus-level endemic radiations of *Microhyla* I, *Microhyla* II, and *Glyphoglossus* clades started in early to mid-Oligocene (from 35 to 25 Ma).

### Historical biogeography

RASP results are presented in Table S12. Ancestral areas and biogeographic processes (vicariance, dispersal, and colonization routes) reconstructed from our data are shown in Fig. 2. Paleogeography of South and Southeast Asia from early Paleocene to the Oligocene is shown in Fig. 3. According to both the DEC and the S-DIVA model, the MRCA of *Microhyla* + *Glyphoglossus* (node 12, Fig. 2) most likely inhabited Eastern Indochina. Eastern Indochina was also reconstructed as an ancestral range for *Microhyla* II and *Glyphoglossus* lineages (nodes 18 and 12, respectively, Fig. 2), and *Microhyla* I likely originated in western Indochina (node 30, Fig. 2). *Microhyla* II expanded its range to Borneo and the Malay Peninsula, but *Microhyla* I dispersed more widely to all biogeographic regions within the modern range of the genus, including at least five independent dispersal events from western Indochina to the Indian Subcontinent (Fig. 2). Results of our analyses suggest numerous cases of dispersal from the Asian mainland to islands including Sundaland, but only a single case of reverse dispersal (from Borneo to the Malay Peninsula, see Fig. 2).

### Body size evolution modeling

Results of ancestral state reconstructions of maximum SVL evolution for each taxon of *Microhyla* and *Glyphoglossus* are shown in Fig. 7. Maximum adult SVL differed substantially for males and females, so data were analyzed separately for each sex. *Glyphoglossus* (adult male SVL 30.0–95.0 mm) are generally larger compared to *Microhyla* (adult male SVL 10.6–35.0

mm); the MRCA of the *Microhyla* + *Glyphoglossus* assemblage is reconstructed as a mid-sized frog (25.0–30.0 mm SVL for males, 30.0–35.0 mm for females). Most species of *Microhyla* were found to be smaller (male SVL roughly 11.5–25.0 mm) than their common ancestor, with five cases of miniaturization (male SVL 11.5–15.0 mm). However, both *Glyphoglossus* and *M. berdmorei* species groups of *Microhyla* I have subsequently increased their body size (up to 105 mm in *Glyphoglossus molossus*, up to 35 mm in *M. berdmorei*) (Fig. 7).

## Discussion

### Updated phylogenetic relationships of *Microhyla*

In phylogenetic systematics, extensive taxon sampling increases accuracy and support of evolutionary relationships (Zwickl & Hillis, 2002). Herein, we present an updated phylogenetic study of the genus *Microhyla*, with the most complete taxon sampling including 48 of the 50 currently recognized species. The absent taxa in our study are *M. fusca* and *M. darevskii* – two enigmatic species from central and southern Vietnam. Further studies of museum specimens and increased field survey efforts are required to clarify their taxonomic status and phylogenetic affinities. We also did not sample *Microhyla maculifera* described from Danum Valley in Sabah, Borneo; it might correspond to *Microhyla* sp. 1 in our analysis, collected from its type locality.

The first phylogenetic study of Asian microhylids by Matsui *et al.* (2011) demonstrated paraphyly of *Microhyla* with respect to *Glyphoglossus* (at that time including *Calluella*). Subsequent multilocus phylogenetic studies (Peloso *et al.*, 2016; Tu *et al.*, 2018; Garg & Biju, 2019) strongly suggested sister group relationships between *Glyphoglossus* and *Microhyla sensu lato*, but still recognized presence of two deeply divergent lineages within *Microhyla*. Our time-tree suggests that divergence between the two major clades of *Microhyla* I and *Microhyla* II happened soon after the basal split within the *Microhyla* + *Glyphoglossus* assemblage during the middle Eocene and similar divergence time estimates were obtained by a recent analysis by Garg & Biju (2019). Robust phylogenies coupled with examination of external morphological and osteological characters are required to assess the evolutionary differences among the three subclades of the *Microhyla* + *Glyphoglossus* assemblage, likely warranting recognition as three distinct genera.

The first classifications of the genus *Microhyla* into species groups were based exclusively on morphological characters (Parker, 1934; Dubois, 1987; Fei *et al.*, 2005) but have



not been supported by more recent molecular data (Matsui *et al.*, 2011; Garg *et al.*, 2019). Matsui *et al.* (2011) assessed genealogical relationships among 20 *Microhyla* species and proposed recognition of five distinct species groups within the genus. Recently Garg *et al.* (2019) proposed a new scheme for grouping *Microhyla* species and recognized six species groups based on more extensive sampling of 33 species. Our phylogenetic hypothesis mostly agrees with these earlier proposed phylogenies of the genus, and a much more extensive taxon sampling allows us to revise species groups more accurately in *Microhyla* (Fig. 6).

*Microhyla achatina* group (Clade A, Figs. 4 and 6; part of group AId2 of Matsui *et al.*, 2011) includes species mostly from Southeast Asia (*M. achatina*, *M. gadjahmadai*, *M. heymonsi*, *M. pineticola*, *M. minuta*, *M. mantheyi*, *M. orientalis*, *M. malang*, *M. nepenthicola*, *M. borneensis*, *Microhyla* sp. 1), but also from Myanmar (*M. irrawaddy*, *M. fodiens*, *Microhyla* sp. 4) and southern India (*M. kodial*). This group includes tiny (male SVL 10.6 mm, Das & Haas, 2010) to mid-sized (male SVL 29.1 mm, Poyarkov *et al.*, 2019) frogs with lateral nostrils; dorsal skin shagreened to prominently granular skin; small disks on digits usually present and typically bearing terminal grooves; toe webbing rudimentary or absent; inner metatarsal tubercle rounded or oval-shaped, outer metatarsal tubercle rounded and small, or large, shovel-shaped (in *M. fodiens*, Poyarkov *et al.*, 2019); usually with a mid-dorsal line or skinfold and a light streak extending from posterior corner of eye to axilla (Garg *et al.*, 2019). The phylogenetic position of *M. minuta*, *M. pineticola*, and *M. borneensis sensu stricto* was assessed for the first time. Vietnamese *M. minuta* was recovered as a member of subgroup A1, joining species from Peninsular Malaysia (*M. mantheyi*), Java + Bali (*M. orientalis*), and Borneo (*M. malang*, *M. nepenthicola*, *M. borneensis*, *Microhyla* sp. 1). Our study also supports previously reported placement of south Indian *M. kodial* in one clade with Myanmar species *M. irrawaddy* and *M. sp. 4* (Poyarkov *et al.*, 2019); this clade has sister group relationships to the *M. achatina* + *M. gadjahmadai* clade from Java and Sumatra (subgroup A2). Unfortunately, phylogenetic positions of the *M. heymonsi* complex and *M. pineticola* remain unresolved. The morphologically different semi-fossorial *M. fodiens* from central Myanmar, previously identified as *M. rubra* (Wogan *et al.*, 2008; Peloso *et al.*, 2016), or *M. cf. berdmorei* (Garg *et al.*, 2019), clearly belongs to this group and forms a highly divergent lineage (Poyarkov *et al.*, 2019).

*Microhyla fissipes* group (Clade B, Figs. 4 and 6; part of group AId2 of Matsui *et al.*, 2011) joins species from East Asia (*M. fissipes*, *M. mixtura*, *M. beilunensis*, *M. okinavensis*, *M.*

*fanjingshanensis* and *Microhyla* sp. 3), Indochina, and eastern India (*M. mukhlesuri*, *M. mymensinghensis*, *M. chakrapanii*). We also agree with Garg *et al.* (2019) in recognizing this species group as distinct from the *M. achatina* group, though monophyly of the latter is only moderately supported (Fig. 4). Members of this group are in general morphologically similar to *M. achatina* species group, but can be distinguished from the latter by finger tips lacking disks, toe tips rounded or bearing tiny disks lacking terminal grooves; by inner metatarsal tubercle elongated, outer metatarsal tubercle small, rounded; and by a dark band running from canthus rostralis posteriorly towards groin and posterior parts of belly (Garg *et al.*, 2019). Our phylogeny recovered two well-supported subgroups within the *M. fissipes* group corresponding to Indochinese (B1) and East Asian (B2) taxa (Fig. 6) (Yuan *et al.*, 2016), supported sister relationships of *M. chakrapanii* and *M. mymensinghensis* (Garg *et al.*, 2019), and suggested sister group relationships of *M. fanjingshanensis* and *M. okinavensis* (Fig. 6).

*Microhyla berdmorei* group (Clade C, Figs. 5 and 6; group AId1 of Matsui *et al.*, 2011) encompasses the largest *Microhyla* species and, according to our data, includes the wide-ranging *M. berdmorei* complex (SVL up to 45.6 mm; Southeast Asia), *M. pulchra* (SVL up to 36.5 mm; Indochina and southern China), and stout-bodied semi-fossorial *M. picta* from southern Vietnam (SVL up to 33.4 mm). The phylogenetic position of *M. picta* was hypothesized here for the first time, and was recovered as sister taxon to *M. pulchra*. Members of the *M. berdmorei* species group exhibit considerable morphological differentiation: smooth to sparsely granular skin on dorsum; finger tips rounded, toe tips rounded or bearing tiny disks with or without terminal grooves; toe webbing from rudimentary to complete reaching toe disks; inner metatarsal tubercle oval, outer metatarsal tubercle from small to large, shovel-shaped (in *M. picta*); lacking mid-dorsal line or skinfold; with a light streak extending from posterior corner of eye to axilla; and a characteristic bright-yellow coloration of groin and posterior parts of belly. Based only on morphological characteristics, the unsampled *M. darevskii* (absent in our phylogeny), likely belongs to this species group.

*Microhyla superciliaris* group (Clade D; part of group AIdc of Matsui *et al.*, 2011) joins small-sized species from South and Southeast Asia, and includes a well-supported clade of southern Indian and Sri Lankan species (Figs. 5 and 6, D1, *M. zeylanica*, *M. laterite*, *M. sholigari*, *M. karunaratnei* and *M. darreli*; *M. zeylanica* group of Garg *et al.*, 2019), three species from northeastern India (*M. eos*), western Thailand (*Microhyla* sp. 2) and the Malay



Peninsula (*M. superciliaris*). Morphologically members of this group are small-sized (male SVL under 21.5 mm), have dorsal orientation of nostrils; smooth to granular skin on dorsum; finger disks rounded or with weak disks lacking terminal grooves; toe disks having terminal grooves; inner metatarsal tubercle oval-shaped, outer metatarsal tubercle small and rounded; toe webbing reduced or well-developed; mid-dorsal skinfold generally present; with a light streak from posterior eye corner to axilla; and often with contrasting black and white blotches on belly. Garg *et al.* (2019) proposed recognizing *M. zeylanica* species group for taxa inhabiting peninsular India + Sri Lanka, but Biju *et al.* (2019) reported *M. eos* from northeastern India as a sister lineage of this clade, refraining from assigning this species to any species group. Herein, we propose recognizing the *M. superciliaris* species group for taxa inhabiting Southeast Asia, northeastern and southern India, and Sri Lanka.

*Microhyla ornata* group (Clade E; group Alb of Matsui *et al.*, 2011) is comprised of species exclusively occurring in the Indian Subcontinent, subdivided into two groups, E1 (*M. ornata*, *M. nilphamariensis*, and *M. taraiensis*; corresponds to *M. ornata* group of Garg *et al.*, 2019), and E2 (*M. rubra*, *M. mihintalei*; corresponds to *M. rubra* group of Garg *et al.*, 2019) (Figs. 5 and 6). Morphologically *M. ornata* group includes small to mid-sized species with lateral nostrils; shagreened to granular dorsal skin; tips of digits lacking disks and terminal grooves; toe webbing rudimentary; inner and outer metatarsal tubercles present, latter may be enlarged; middorsal line or skinfold present; body flanks with dark band from nostrils to groin; and a light streak from posterior eye corner to axilla. Though Garg *et al.* (2019) proposed recognizing the more robust, semi-fossorial species *M. rubra* and *M. mihintalei* as a distinct *M. rubra* species group, we do not follow their scheme since phylogenetic relationships within Clade E are well resolved and subclades E1 and E2 are closely related (Matsui *et al.*, 2011). In addition, adaptations to burrowing lifestyle are not unique for *M. rubra* but are found in other lineages of *Microhyla* as well, such as *M. fodiens* of Clade A, and *M. picta* of Clade C.

*Microhyla butleri* group (Clade F; part of group A1c of Matsui *et al.*, 2011; see Figs. 5 and 6) includes the *M. butleri* complex from Southeast Asia and southern China and the closely related *M. aurantiventris* from central Vietnam. Morphologically members of this group show a dorso-lateral nostril position; prominently granular dorsal skin; presence of weak disks on digits bearing terminal grooves; moderately developed webbing on toes; inner and outer metatarsal tubercles small; middorsal line or skinfold present; characteristic ‘teddy-bear’-shaped dark

marking on dorsum edged with light color; body flanks lacking dark band from nostrils to groin; and a light eye-axilla streak present (Nguyen *et al.*, 2019).

*Microhyla palmipes* group (Clade G; group AIIa of Matsui *et al.*, 2011) presently includes a single species *M. palmipes*, from Java, Sumatra, and adjacent offshore islands in Indonesia, and according to our data, likely represents a species complex (see below). Morphological data on *M. palmipes* are scarce; they are small-sized frogs (male SVL 16.0 mm) with lateral nostrils; shagreened skin on dorsum; weak disks on digits lacking terminal grooves; moderately developed webbing on toes; inner and outer metatarsal tubercles small; middorsal line absent; dark markings on flanks and a light eye-axilla streak present (Bain & Nguyen, 2004; Poyarkov *et al.*, 2014).

Finally, the *M. annectens* group (Clade H of *Microhyla* II; group AIII of Matsui *et al.*, 2011; Figs. 5 and 6) encompasses tiny (male SVL 13.2 mm) to mid-sized (male SVL to 21.6 mm) frogs from Southeast Asia, and according to our phylogeny, is subdivided into two subgroups: H1 comprising species from Malayan Peninsula (*M. annectens*), Annamite Mountains in Vietnam (*M. annamensis*, *M. marmorata*, and *M. pulverata*), and Borneo (*M. petrigena* and *M. perparva*); and H2 including species from mountains of central and southern Vietnam (*M. arboricola*, *M. pulchella*, *M. nanapollexa*). *M. marmorata* was found to be paraphyletic with respect to *M. pulverata*. Morphologically, *M. annectens* group members are characterized by a relatively short body; lateral position of nostrils; sparsely granular to tubercular dorsal skin; complete toe webbing with well developed flattened and slightly expanded disks on digits bearing terminal grooves; inner metatarsal tubercle present, and outer metatarsal tubercle present or absent. Further morphological studies are required to examine morphological differentiation among *M. annectens* group members. Phylogenetic positions of *M. annamensis*, *M. marmorata*, *M. pulverata*, *M. arboricola*, and *M. pulchella* are for the first time reported in the present study.

Our updated phylogeny reveals several lineages of *Microhyla* that likely represent undescribed species: *Microhyla* sp. 1 from Sabah, Malaysia (corresponds to *Microhyla* sp. 1 of Matsui *et al.*, 2011), *Microhyla* sp. 2 from western Thailand (previously not reported), *Microhyla* sp. 3 from Yaeyama Archipelago (previously referred as *M. okinavensis*), and *Microhyla* sp. 4 from northern Myanmar (reported as *Microhyla* sp. A by Mulcahy *et al.*, 2018). Our study also recovered significant diversity within wide-ranging species complexes that might comprise

undescribed cryptic species (N = 31). This suggests that the taxonomy of the genus *Microhyla* still remains largely incomplete.

### **Indian Collision and historical biogeography of *Microhyla***

The origin of Asian microhylids, including the subfamily Microhylinae, is connected with a break-up of Gondwana and the Indian Collision (*Van Bocxlaer et al., 2006; Van der Meijden et al., 2007*). Most likely, ancestors of Asian Microhylidae subfamilies diverged and diversified on the Indian Plate during its long isolation and northward drifting in late Cretaceous and Paleocene (*Bossuyt & Milinkovitch, 2001; Kurabayashi et al., 2011; de Sá et al., 2012*) (see Fig. 3A). The basal divergence of the subfamily Microhylinae most likely took place on the Indian Plate prior to its first contact with Eurasia and the ISC is regarded as the original source of Microhylinae diversity (*Garg & Biju, 2019*). However, Southeast Asia (not the ISC) presently harbors the largest number of Microhylinae lineages and species (*Frost, 2020*).

Several recent biogeographic studies suggested that collision of the ISC with the Asian mainland was a more complicated process than previously conceived, implicating early opportunities for faunal exchange between the ISC and present-day Southeast Asia (*Klaus et al., 2010; Li et al., 2013; Grismer et al., 2016*). The “Out of India” scenario, with early dispersal from the ISC to Sundaland via brief land connection in early Eocene, has also been proposed for the Microhylinae (*Garg & Biju, 2019*). Recent progress in tectonic plate modeling further corroborates the possibility for biotic exchange between the ISC and Sundaland via the Incertus Arc land bridge, starting 55–50 Ma (Fig. 3B), although exact timing and configuration of the landmasses remains under debate (*Hall, 2012; Ding et al., 2017*). Interestingly, paleoclimate reconstructions suggest that modern-day megathermal angiosperm-dominated tropical forests also originated in the ISC. They later dispersed from there and became established across Sundaland starting about 50 Ma (*Morley, 2018*) coinciding with the onset of a perhumid climate in Southeast Asia (Fig. 3B).

The present time tree analysis indicates that the ancestral radiation of *Microhyla* + *Glyphoglossus* into three main lineages (*Microhyla* I, *Microhyla* II and *Glyphoglossus*) happened during a short time frame in the middle Eocene (ca. 43.8 Ma), slightly later than previous estimates (48.7 Ma; *Garg & Biju, 2019*). Our biogeographic analysis strongly suggests that the *Microhyla* + *Glyphoglossus* MRCA, as well as the *Microhyla* I + *Microhyla* II ancestor, inhabited Eastern Indochina (Fig. 2), which was connected to Sundaland (Fig. 3C). Thus, our

results support the Southeast Asian origin of the *Microhyla* – *Glyphoglossus* assemblage in contrast to the hypothesis by Garg & Biju (2019), that suggested the dispersal of *Microhyla* from the ISC into Asia from Oligocene to Miocene.

The *Microhyla* II clade remained largely within its ancestral area with most members of the group inhabiting Eastern Indochina and a few species dispersing to Borneo and the Malay Peninsula (Fig. 2). Compared to other *Microhyla* species groups, members of the *Microhyla* II clade are generally small and associated with perhumid montane evergreen forests or tropical rainforests; they do not occur in lowland seasonally dry areas. In fact, in Indochina, their distribution is restricted to mountainous areas (Parker, 1934; Poyarkov et al., 2014).

On the contrary, members of the *Microhyla* I clade dispersed widely and achieved a pan-Oriental distribution (see Figs. 1 and 2). Members of this clade are diverse ecologically and morphologically. They vary in body size from the smallest to the largest *Microhyla* taxa, occupy habitats that include open seasonally dry savannahs (Parker, 1934), and several species within the *Microhyla* I clade evolved adaptations towards digging and aestivation (*M. rubra*, *M. mihintalei*, *M. picta*, and *M. fodiens*; see Poyarkov et al., 2019). The MRCA of *Microhyla* I is hypothesized to inhabit western Indochina, the same region reconstructed as ancestral for all major internal nodes within *Microhyla* I, and for a number of included lineages (*M. heymonsi*, *M. fissipes*, *M. berdmorei*, and *M. superciliaris* species groups), respectively (Fig. 2).

Drifting of the ISC northwards led to a collision of the Indian plate with Eurasia from the Oligocene to the Miocene (Aitchison & Ali, 2012; Hall, 2012) (Fig. 3D). At the same time, the uplift of the Himalayas, coinciding with the middle Miocene thermal maximum, initiated the subsequent Miocene strengthening of the Indian monsoon and entailed the expansion of seasonally dry conditions across the northern parts of the Indian peninsula and Indochina. This resulted in the disappearance of closed tropical forests over much of the ISC (Morley, 2018) (Fig. 3D). Starting in the Oligocene, Indochina became the source of evergreen and seasonally dry floral elements that dispersed to the ISC with ongoing climate aridification (Morley, 2000). These conditions presumably facilitated colonization of the Indian Subcontinent by *Microhyla* I lineages.

Our biogeographic analysis reveals at least five independent cases of *Microhyla* I dispersal from Western Indochina to the ISC (see Fig. 2). Two of these took place in Late Oligocene–Early Miocene: *M. superciliaris* species group (29.2–22.5 Ma, Fig. 2, 1) and *M.*

*ornata* species group (31.4–18.8 Ma, Fig. 2, 2). Both lineages underwent significant diversification in the ISC and reached as far south as Sri Lanka. Three other cases of the ISC colonization by *Microhyla* I include more recent dispersal events in Late Miocene – Pliocene by *M. berdmorei* (6.9–4.1 Ma, Fig. 2, 3) and *M. fissipes* species groups (*M. mymensinghensis*, 7.2–5.1 Ma, Fig. 2, 4) to northeastern India and Bangladesh, with the only case of dispersal to southern peninsular India being the species of *M. achatina* group in the Middle Miocene (*M. kodial*, 16.2–8.9 Ma, Fig. 2, 5). The confusing phylogenetic position of *M. kodial* within the Southeast Asian *M. achatina* species group originally inspired the hypothesis that this might be a result of a human-mediated dispersal and introduction (Vineeth et al., 2018). However, subsequent discovery of its sister species *M. irrawaddy* and *Microhyla* sp. 4 in central Myanmar have made the hypothesis of natural dispersal of the *M. achatina* species group members from Southeast Asia to the ISC more plausible. Interestingly, *M. irrawaddy* inhabits seasonally dry savannah areas, with minimal rainfall (Poyarkov et al., 2019). Hence, progressing aridification of the northern and central parts of the ISC starting with late Miocene (Deepak & Karanth, 2018) could have created suitable habitats facilitating dispersal of *M. kodial* ancestors. Generally, western Indochina played an important role for the *Microhyla* I clade (Fig. 2) as this territory likely represents a “stepping stone” area connecting Southeast Asia and the ISC (Fig. 3D).

Diversification within *Microhyla* species group-level endemic radiations started in the Late Oligocene–Early Miocene and initiated multiple dispersals from Asian mainland to present-day islands and archipelagos (Fig. 2). These include multiple colonizations of Sundaland from both Indochina and the Malay Peninsula (by *M. annectens*, *M. palmipes*, *M. berdmorei* and *M. achatina* species groups; Fig. 2). This is not surprising, since these territories are believed to have been a single landmass throughout most of Cenozoic (Cannon et al., 2009; Woodruff, 2010; Hall, 2012). *Microhyla superciliaris* and *M. ornata* species groups experienced at least four independent dispersal events from southern India to Sri Lanka, corroborating results of recent studies, and suggesting a complex history of dispersals between these regions (Harikrishnan et al., 2012; Pyron et al., 2013; Agarwal et al., 2017; Karunarathna et al., 2019). Our study confirms placement of *M. chakrapanii* from Andaman Islands inside the Southeast and East Asian *M. fissipes* species group as sister species of *M. mymensinghensis* (see Garg et al., 2019). This confirms the faunal similarity of the Andamans with Southeast Asia rather than with peninsular India (see Das, 1994, 1999). Finally, members of *M. achatina*, *M. butleri*, and *M.*

*fissipes* species groups have dispersed several times from the Asian mainland to East Asian islands: Taiwan (*M. fissipes*, *M. heymonsi*, and *M. butleri*), and two times independently colonized the Ryukyus (*M. okinavensis* and *Microhyla* sp. 3). These results also corroborate data that suggested faunal exchanges between Eurasian continent and East Asian islands (e.g. *Ota*, 1998; *You et al.*, 2015; *Yuan et al.*, 2016; *Wang et al.*, 2017; *Nguyen et al.*, 2020) and require further study (*Lee et al.*, 2016; *Tominaga et al.*, 2019).

Interestingly, our phylogeny suggests numerous dispersal events from the Asian mainland to islands, with almost no dispersals back to the mainland (except for the *M. annectens* ancestor that is hypothesized to have dispersed from Borneo to Malay Peninsula, see Fig. 2). This also supports results of *de Bruyn et al.* (2014) who demonstrated that colonization events from younger Asian islands are comparatively rare, rather showing increased levels of immigration events as compared to Indochina or Borneo. Further studies are required to elucidate the role of islands in producing and preserving diversity in *Microhyla* frogs.

# **Implications for body size evolution in *Microhyla***

Miniaturization is a widespread and interesting morphological and ecological phenomenon in amphibians (*Hanken*, 1985; *Hanken & Wake*, 1993; *Rieppel*, 1996). It is common in several groups of frogs (*Clarke*, 1996; *Lehr & Coloma*, 2008) and reaches extremes in the Microhylidae (*Kraus*, 2011; *Rittmeyer et al.*, 2012; *Oliver et al.*, 2017; *Rakotoarison et al.*, 2017; *Scherz et al.*, 2019). The smallest Microhylinae and the smallest terrestrial vertebrate in Asia belong to *Microhyla* and include representatives of two different lineages within the genus: *Microhyla* I (*M. nepenthicola*, adult male size from 10.6 mm; see *Das & Haas*, 2010) and *Microhyla* II (*M. perparva*, males 10.5–11.9 mm, and *M. arboricola*, adult male size from 13.2 mm; see *Inger & Frogner*, 1979; *Poyarkov et al.*, 2014). Osteological consequences of miniaturization in *Microhyla* are not well studied yet diminutive species in *Microhyla* II clade show partial (*M. arboricola*) or almost complete (*M. perparva*, *M. nanapollexa*) reduction of the first finger. Similar patterns have also been reported in other miniature microhylids (e.g. *Kraus*, 2011; *Rakotoarison et al.*, 2017), however, patterns and drivers of body size evolution in *Microhyla* remain poorly understood.

In *Microhyla*, males generally tend to be smaller than females (see Table S7), yet our analyses revealed generally similar patterns of body size evolution in both sexes (Fig. 7). According to the most plausible scenario, the common ancestor of the *Microhyla* +

*Glyphoglossus* assemblage was a mid-sized frog (male SVL 25–30 mm, female SVL 30–35 mm). Body size increased in the *Glyphoglossus* clade (up to 105 mm SVL), was slightly reduced in the *Microhyla* I clade, and significantly reduced in the *Microhyla* II clade ancestors (see Fig. 7). *Microhyla* II members are all small-sized (< 25 mm SVL in males) and three species of this group reach extreme miniaturization (< 15 mm SVL in males; *M. arboricola*, *M. perparva*, *M. nanapollexa*). *Microhyla* I clade shows significant variation in body size: the *M. berdmorei* species group and, to a lesser extent, several species in other lineages (including *M. fodiens*, *M. rubra*, *M. mihintalei*, *M. aurantiventris*) demonstrate increased body sizes, while members of two lineages within *M. achatina* (*M. nepenthicola* + *M. borneensis* + *Microhyla* sp. 1) and *M. superciliaris* species groups (*M. superciliaris* + *Microhyla* sp. 2) are diminutive (Fig. 7).

Our analyses suggest that adult body size has independently changed several times in the evolution of the *Microhyla* + *Glyphoglossus* assemblage. At least two lineages show an increase in body size, while four other lineages demonstrate extreme miniaturization (Fig. 7). Significant increases in body size in these frogs seem to be connected with a fossorial life style. For example, all members of *Glyphoglossus* as well as some large *Microhyla* (*M. picta*, *M. fodiens*, *M. rubra* and *M. mihintalei*) have stout body habitus and are excellent burrowers (Poyarkov *et al.*, 2019). Most of these species inhabit open seasonally dry habitats at low elevations and burrowing is an important strategy for aestivation during dry periods. Increased body size might not only facilitate digging, but is also advantageous due to lower surface-area to volume ratios, hence leading to less evaporative water loss during aestivation (Tracy *et al.*, 2010).

Evolutionary causes for extreme miniaturization in frogs remain under debate (Scherz *et al.*, 2019), but are probably connected with life history strategies, such as exploiting new food resources (Lehr & Coloma, 2008) or adaptation to specific microhabitats such as leaf-litter or moss (Kraus, 2011). Scherz *et al.* (2019) noted that one miniaturized species is arboreal and breeds in water-filled leaf-axil phytotelmata. Interestingly, at least the three smallest known *Microhyla* species are also obligatory phytotelm-breeders and reproduce in water-filled pitcher-plants (*M. nepenthicola*, see Das & Haas, 2010; and *M. borneensis*, see Parker, 1928) or water-filled tree hollows (*M. arboricola*, see Poyarkov *et al.*, 2014; Vassilieva *et al.*, 2017). Among all other *Microhyla* species, *M. arboricola* is the only known arboreal species with a unique reproductive mode: developing tadpoles of this species are obligately oophagous (Vassilieva *et al.*, 2017). *M. arboricola* has a reduced clutch size ( $16 \pm 8$  eggs), compared to other *Microhyla*

species (usually over 400 eggs per clutch) (*Vassilieva et al., 2017*). Egg size appears to be one of the main constraints for miniaturization in animals (*Polilov, 2015*). On one hand, reduced clutch size might favor the choice of phytotelmata for reproduction due to the absence or low density of predators in such habitats; on the other hand, diminutive body size might be advantageous for phytotelmic frogs because it allows them to exploit smaller phytotelmata than are available to larger frogs (*Scherz et al., 2019*). Breeding biology of *M. petrigena* and *M. nanapollexa* is not yet reported. However, our field observations suggest that the latter species also reproduces in tree hollows. Further studies might shed light on evolutionary interdependencies between phytotelm-breeding and extreme miniaturization in *Microhyla*.

### **Taxonomic implications and cryptic diversity in *Microhyla***

Diminutive frogs are recognized as a source of astonishingly high undescribed cryptic diversity at different taxonomic levels due to incomplete phylogenetic information and widespread homoplasies in morphology (*Hanken & Wake, 1993; Rittmeyer et al., 2012; Scherz et al., 2019*). Until recently most miniature frog groups also attracted little attention by taxonomists (e.g. *Rakotoarison et al., 2017*). This is also true for the genus *Microhyla* as the only available systematic study addressing the genus by *Matsui et al. (2011)* provided important insights on phylogeny and taxonomy of these frogs, but did not provide other insights. In accordance with results of *Matsui et al. (2011)*, our phylogeny indicates that morphology-based classification schemes of *Parker (1934)*, *Dubois (1987)*, and *Fei et al. (2005)* do not reflect actual phylogenetic relationships among *Microhyla* species, most likely due to high frequency of homoplasies both in adult and larval morphology. Further thorough morphological and osteological studies along with a robust phylogeny are required to diagnose supraspecific-level taxa within the *Microhyla* + *Glyphoglossus* assemblage.

Since the sampling used in *Matsui et al. (2011)* was incomplete, and the number of recognized *Microhyla* species has since increased almost two-fold, many questions of *Microhyla* taxonomy remain unaddressed. In the present paper, we used an updated and almost complete phylogeny of the genus along with species delimitation methods to resolve several long-standing areas of confusion in the genus *Microhyla*. Our analyses suggest that despite recent progress in *Microhyla* taxonomy, the current number of recognized *Microhyla* species is still greatly underestimated. Based on species delimitation analyses, 15 (ABGD estimate) to 33 lineages (bGMYC estimate) probably reflect new species requiring formal description (Table S10).



However, due to possible overlap in levels of intra- and interspecific divergence, species delineation in *Microhyla* based on genetic differentiation alone is problematic (Garg et al., 2019). An integrative approach including morphology and acoustics must be applied for further progress in *Microhyla* taxonomy (Rakotoarison et al., 2017).

Below we give a brief summary of taxonomic implications of our results. The smallest member of the genus, *M. nepenthicola*, was described by Das & Haas (2010) from Kubah, Sarawak, but was synonymized with *M. borneensis* by Matsui (2011) based on morphological examination of the *M. borneensis* holotype from Kidi (*sic* for Bidi), Sarawak, Borneo (Parker, 1928). Matsui (2011) did not include in his molecular analysis materials from the type locality of *M. borneensis*; however, his taxonomy was widely accepted (Frost, 2020). We analyzed topotypic *M. borneensis* specimen from the Bidi region (Fig. 4, 38), specifically, from the Deded Krian National Park, near Bau, western Sarawak, and show it be a sister species of *M. nepenthicola*, and sufficiently divergent from the latter in 16S rRNA sequences ( $p = 3.5\%$ , Table S9). The bGMYC analysis also supports distinctiveness of *M. nepenthicola* from *M. borneensis* (Fig. 4). Specimens from the former are widespread along sandstone massifs of western Sarawak, and are diagnosable in showing a distinct, hour-glass pattern, as opposed to that of the latter, from the limestone hills of the interior, whose dorsum bears an indistinct, greyish-brown, shagreened pattern; therefore, we propose to revalidate the species *M. nepenthicola* Das & Haas, 2010. Our data further suggest that *M. borneensis*, *M. nepenthicola*, and *Microhyla* sp. 1 from Sabah form a clade of morphologically similar and closely related taxa. Further studies are required to fully clarify morphological differences between *M. nepenthicola* and *Microhyla* sp. 1.

Significant genetic differentiation is revealed within several species of the *M. achatina* species group suggesting presence of cryptic diversity. Examples include *M. malang* (3 putative species: populations from Sarawak, Sabah and Kalimantan); *M. orientalis* (2 putative species: populations from Bali and Java; our study confirms the occurrence of *M. orientalis* in Java); *M. mantheyi* (2 putative species within Malayan Peninsula); *M. achatina* (2 putative species within Java); and *M. gadjahmadai* (2 putative species within Sumatra) (see Fig. 4). Genetically, the most diverse cryptic species complex in *Microhyla* is the *M. heymonsi* complex. Earlier studies already recognized presence of several deeply divergent intraspecific lineages within *M. heymonsi* (Garg et al., 2019). Our new analyses revealed 7–8 highly divergent ( $p > 3.0\%$ ) lineages from China and northern Vietnam, Thailand and Laos, Thailand and south Vietnam, Taiwan,

Myanmar, peninsular Malaysia, and Sumatra (Fig. 4). Taxonomic status of these lineages has yet to be assessed. Our study also confirms the clear distinctiveness of an undescribed species *Microhyla* sp. 4 from northern Myanmar, and the full species status of *M. minuta*, *M. pineticola*, *M. Irrawaddy*, and *M. fodiens*, respectively.

Within the *M. fissipes* species group, the bGMYC analysis indicated the presence of three cryptic species-level lineages within *M. mukhlesuri*, although the ABGD analysis lumped *M. mukhlesuri* with *M. fissipes* (Fig. 4). Significant genetic differentiation in mtDNA-markers was reported for *M. mukhlesuri* by Yuan *et al.* (2016), but they were not corroborated by nuclear DNA. Further integrative studies are required to assess variation within *M. mukhlesuri*. Deep species-level divergence was revealed between populations of *M. okinavensis* from Okinawa and Amami islands ( $p=2.0\%$ ); populations from Yaeyama Archipelago formerly treated as *M. okinavensis* grouped with *M. mixtura* and most likely represent an undescribed species, *Microhyla* sp. 3 (Tominaga *et al.*, 2019; Hasan *et al.*, 2014a). Shallow divergence was also found among populations of *M. chakrapanii* from different islands of the Andaman Archipelago (Fig. 4).

Substantial genetic divergences also uncovered cryptic species lineages within the *M. berdmorei* species complex (Fig. 5) (Hasan *et al.*, 2012). The bGMYC analysis suggested presence of four distinct groups from Malayan Peninsula, Malaysia + Sumatra + Borneo, Indochina, and Bangladesh. Populations from northern Thailand previously described as *M. fowleri* are grouped within the Indochinese lineage of *M. berdmorei*, suggesting synonymy of the former (see Matsui *et al.*, 2011). Further studies, including examination of topotype material for *M. berdmorei* (Myanmar) and *M. darevskii* (central Vietnam), are needed to estimate taxonomic statuses of these newly revealed lineages and extent of their distribution. Within the *M. superciliaris* group, our study confirmed occurrence of *M. superciliaris* in southern Thailand (Songkhla), however a deep divergence of this population from the *M. superciliaris* population in Malaysia, suggests necessity of reevaluating taxonomy of Thai populations (Fig. 5). We also report an undescribed species *Microhyla* sp. 2, occurring in western Thailand where genetic variation among examined populations (Suratthani and Phetchaburi) also suggests presence of two cryptic lineages.

For *M. butleri* species group, our work confirms deep divergence and full species status of the recently described *M. aurantiventris* (Nguyen *et al.*, 2019), but also reveals additional

undescribed lineages within *M. butleri* that we treat here as a species complex. From two (ABGD) to four (bGMYC) cryptic lineages were recovered within this complex, with the most divergent lineages being distributed in Taiwan + mainland China versus the rest of the species range in Indochina and the Malay Peninsula (Fig. 5). We included only two *M. palmipes* samples in our analysis that were notably divergent in 16S rRNA sequences ( $p=2.9\%$ ) and originated from Bali and Sumatra; likely they both represent distinct species.

For *M. annectens* species group, our species delimitation analyses reveal a number of cryptic and undescribed lineages. Our study confirms genetic distinctiveness of recently described *M. arboricola* and *M. pulchella* (Poyarkov *et al.*, 2014), as well as of *M. annamensis*, for which genetic information was not previously available. We also added to our analysis a number of populations of *M. marmorata* throughout the species range and for the first time, included specimens of *M. pulverata* (collected from ca. 10 km north of the type locality in Gia Lai Province, central Vietnam). Both species were described by Bain & Nguyen (2004) based on morphological evidence and the main characters considered to be diagnostic for these species were belly coloration (marbled in *M. marmorata* versus dusty in *M. pulverata*) and skin texture. Our genetic data reveal almost no genetic differentiation between samples of *M. marmorata* and the topotypic *M. pulverata* ( $p = 0.3\%$ , see Table S9), the latter are nested within *M. marmorata* radiation and do not form a clade (Fig. 5). Moreover, our observations showed that belly coloration is highly variable within *M. marmorata*, especially in the southern part of the species range, and cannot be used as a reliable diagnostic character. Due to the lack of genetic and morphological differentiation, we hereby formally treat *Microhyla pulverata* Bain & Nguyen, 2004 as a subjective junior synonym of *Microhyla marmorata* Bain & Nguyen, 2004. Some other species of *M. annectens* species group show deep intraspecific divergences in 16S rRNA sequences, such as *M. arboricola* ( $p = 2.2\%$  between populations from Dak Lak and Khanh Hoa provinces of Vietnam), *M. petrigena* ( $p = 3.1\%$  between populations from Sarawak and Sabah), and *M. perparva* ( $p = 4.3\%$  between populations from Sarawak and Indonesian Kalimantan) (Fig. 5). It is likely that these lineages represent distinct species, including several new taxa awaiting formal description.

## Conclusions

Herein, we provide an updated phylogenetic hypothesis for the genus *Microhyla*. An exhaustive taxonomic sampling for this group is challenging due to the high number of narrow-ranged or point-endemic species across South and Southeast Asia. In the present study, however, we examined mtDNA and nuDNA markers for 48 of 50 recognized *Microhyla* species (96%), including 12 nominal species and several undescribed candidate species that have not been examined phylogenetically before our work, thus providing the most comprehensive taxonomic sampling for *Microhyla* to date. Our data further highlight the importance of broad phylogenetic sampling and ground-level field research to gather an accurate picture of global biodiversity, phylogenetic relationships, and evolutionary patterns in cryptic groups such as microhylid frogs.

We recognize nine species groups within the *Microhyla* – *Glyphoglossus* assemblage (*M. achatina*, *M. fissipes*, *M. berdmorei*, *M. superciliaris*, *M. ornata*, *M. butleri*, *M. palmipes*, *M. annectens* species groups and *Glyphoglossus*), divided into three clades of probable generic-level differentiation: *Microhyla* I, *Microhyla* II and *Glyphoglossus*. Further integrative research combining phylogenetic and morphological lines of evidence is required to fully diagnose these recognized groups and test our taxonomic arrangement. The basal radiation of the *Microhyla* – *Glyphoglossus* assemblage is dated to the middle Eocene and likely took place in Southeast Asia. Following drifting of the Indian Plate northwards and formation of firm land bridges between the subcontinent and Asian mainland in Oligocene, ancestors of *Microhyla* colonized India several times from Southeast Asia and later diversified there. Our analysis also suggests that such dispersal occurred independently in five different species groups of the *Microhyla* II clade. Our results further corroborate the growing set of evidence for early-Eocene land connections between the Indian Subcontinent and Southeast Asia. Progressing aridification since the late Oligocene – Miocene likely facilitated dispersal of Southeast Asian biotic elements to India including the ancestral lineage-genus *Microhyla*. Our study further highlights the importance of Indochina not only as a cradle of autochthonous amphibian diversity and a key evolutionary hotspot for the herpetofauna (Bain & Hurley, 2011; Geissler et al., 2015; de Bruyn et al., 2014), but also as a stepping stone facilitating dispersal between Sundaland, the Indian Subcontinent, and East Asia (e.g. Wood et al., 2010; Chen et al., 2017; Suwannapoom et al., 2018; Poyarkov et al., 2018b). Further phylogenetic studies across different faunal groups with Indo-Southeast Asian affinities are required to clarify impact of complex paleogeography and paleoclimate history on formation of extant biodiversity in Asia.

Comprising the smallest tetrapods in Asia, frogs in the genus *Microhyla* represent a promising model group for studies of evolutionary drivers and constraints of miniaturization in vertebrates. Our study suggests that four groups of *Microhyla* independently achieved extreme miniaturization with adult body sizes <15 mm. Evolution of body size in *Microhyla* – *Glyphoglossus* assemblage seems to be driven by natural history: the largest body sizes are observed in burrowing species adapted to aestivation during the dry season, while three of the five smallest known *Microhyla* species occur only in perhumid montane forests and are phytotelm-breeders. Further research is required on how reproductive ecology in phytotelmata, often leading to reduction of clutch size, facilitates extreme miniaturization in *Microhyla*.

The present work clearly indicates a vast underestimation of diversity and species richness of *Microhyla*. We revalidate *M. nepenthicola* as a valid species, synonymize *M. pulverata* with *M. marmorata*, confirm species-level differentiation for a number of taxa not included in earlier phylogenies, and reveal a large number of cryptic lineages representing putative undescribed species. Alternative approaches to species delimitation suggest that at least 15 to 33 lineages of *Microhyla* likely correspond to species-level differentiation. Further integrative studies combining genetic, morphological, and acoustic parameters are essential for a better understanding of evolutionary relationships and taxonomy within this morphologically cryptic and diverse radiation of Asian frogs.

## Acknowledgements

The authors are grateful to the following colleagues who took part in the fieldwork, collected material and discussed the results, assembled and facilitated obtaining samples, as well as providing access to collections and other materials: S. Shonleben, V. F. Orlova, R. A. Nazarov, N. B. Ananjeva, N. L. Orlov, A. B. Vassilieva, E. A. Galoyan, K. D. Milto, D. Gabadage, M. Botejue, M. Madawala, Than Zaw, T. V. Nguyen, S. N. Nguyen, K.-H. Lee, Y.-P. Lin, H.-Y. Tseng, S.-F. Yang, O. S. Bezman-Moseiko, T. Neang, H. N. Nguyen, T. Igawa, S. Komaki, S.-M. Lin, Y. Lee, J. Lee, Y. M. Pui, K. V. Minin, A. A. Vedenin, J. Che, S. Howard, and P. Geissler. We thank A. M. Fetisova and E. S. Popov for providing important information and references on tectonics, paleogeography, and climate evolution in Asia. We are grateful to S. Shonleben, V. D. Kretova, T. V. Duong, S. Komaki, A. S. Dubrovskaya, P. V. Yushchenko, S. S. Idiatullina, and

984 A. N. Kuznetsov for tremendous help during our work on this project. We are also grateful to  
985 anonymous reviewers for useful comments on an earlier version of the manuscript.

986

## 987 **References**

- 988 **Acton GD. 1999.** Apparent polar wander of India since the Cretaceous with implications for  
989 regional tectonics and true polar wander. *In: Radhakrishna T, Piper JDA (Eds.) The Indian*  
990 *Subcontinent and Gondwana: a Palaeomagnetic and Rock Magnetic Perspective. Memoires of*  
991 *Geological Society of India*, **44**:129–175.
- 992 **Agarwal I, Biswas S, Bauer AM, Greenbaum E, Jackman TR, de Silva A, Batuwita S. 2017.**  
993 Cryptic species, taxonomic inflation, or a bit of both? New species phenomenon in Sri Lanka as  
994 suggested by a phylogeny of dwarf geckos (Reptilia, Squamata, Gekkonidae, *Cnemaspis*).  
995 *Systematics and Biodiversity* **15**:1–13.
- 996 **Aitchison JC, Ali JR, Davis AM. 2007.** When and where did India and Asia collide? *Journal of*  
997 *Geophysical Research: Solid Earth* **112**:B05423.
- 998 **Aitchison JC, Ali JR. 2012.** India-Asia collision timing. *Proceedings of the National Academy of*  
999 *Sciences* **109**(40):E2645–E2646.
- 1000 **Ali JR, Aitchison JC. 2008.** Gondwana to Asia: Plate tectonics, paleogeography and the biological  
1001 connectivity of the Indian subcontinent from the Middle Jurassic through latest Eocene (166-35  
1002 Ma). *Earth-Science Reviews* **88**:145–166.
- 1003 **Ali JR, Aitchison JC. 2012.** Comment on “Restoration of Cenozoic deformation in Asia and the  
1004 size of Greater India” by D. J. J. Van Hinsbergen et al. *Tectonics* **31**:TC4006.
- 1005 **An Z, Kutzbach JE, Prell WL, Porter SC. 2001.** Evolution of Asian monsoons and phased uplift  
1006 of the Himalaya–Tibetan Plateau since late Miocene times. *Nature* **411**:62–66.  
1007 <https://doi.org/10.1038/35075035>
- 1008 **Bain RH, Hurley MM. 2011.** A biogeographic synthesis of the amphibians and reptiles of  
1009 Indochina. *Bulletin of the American Museum of Natural History* 2011(360):1–38.
- 1010 **Bain RH, Nguyen QT. 2004.** Three new species of narrow-mouth frogs (genus: *Microhyla*) from  
1011 Indochina, with comments on *Microhyla annamensis* and *Microhyla palmipes*. *Copeia*  
1012 **2004**(3):507–524.
- 1013 **Biju SD, Bossuyt F. 2003.** New frog family from India reveals an ancient biogeographical link with  
1014 the Seychelles. *Nature* **425**(6959):711–714.

- 101 **Biju SD, Garg S, Kamei RG, Maheswaran G. 2019.** A new *Microhyla* species (Anura:  
1016 Microhylidae) from riparian evergreen forest in the eastern Himalayan state of Arunachal  
1017 Pradesh, India. *Zootaxa* **4674**:100–116.
- 101 **Bossuyt F, Milinkovitch MC. 2001.** Amphibians as indicators of early Tertiary “out-of-India”  
1019 dispersal of vertebrates. *Science* **292(5514)**:93–95.
- 102 **Cannon CH, Morley RJ, Bush ABG. 2009.** The current refugial rainforests of Sundaland are  
1021 unrepresentative of their biogeographic past and highly vulnerable to disturbance. *Proceedings of*  
1022 *the National Academy of Sciences* **106**:11188–11193.
- 102 **Che J, Zhou WW, Hu JS, Yan F, Papenfuss TJ, Wake DB, Zhang YP. 2010.** Spiny frogs (Paini)  
1024 illuminate the history of the Himalayan region and Southeast Asia. *Proceedings of the National*  
1025 *Academy of Sciences* **107(31)**:13765–70.
- 102 **Chen J-M, Poyarkov NA, Suwannapoom C, Lathrop A, Wu Y-H, Zhou W-W, Yuan Z-Y, Jin**  
1027 **J-Q, Chen H-M, Liu H-Q, Nguyen TQ, Nguyen SN, Duong TV, Eto K, Nishikawa K,**  
1028 **Matsui M, Orlov NL, Stuart BL, Brown RM, Rowley JJJ, Murphy RW, Wang Y-Y, Che J.**  
1029 **2018.** Large-scale phylogenetic analyses provide insights into unrecognized diversity and  
1030 historical biogeography of Asian leaf-litter frogs, genus *Leptolalax* (Anura: Megophryidae).  
1031 *Molecular Phylogenetics and Evolution* **124**:162–171.
- 103 **Chen JM, Zhou WW, Poyarkov NA, Stuart BL, Brown RM, Lathrop A, Wang YY, Yuan ZY,**  
1033 **Jiang K, Hou M, Chen HM. 2017.** A novel multilocus phylogenetic estimation reveals  
1034 unrecognized diversity in Asian horned toads, genus *Megophrys* sensu lato (Anura:  
1035 Megophryidae). *Molecular Phylogenetics and Evolution* **106**:28–43.
- 103 **Clarke BT. 1996.** Small size in amphibians — its ecological and evolutionary implications.  
1037 *Symposia of the Zoological Society of London*, **69**:201–224.
- 103 **Conti E, Eriksson T, Schönerberger J, Sytsma KJ, Baum DA. 2002.** Early Tertiary out-of-India  
1039 dispersal of Crypteroniaceae: Evidence from phylogeny and molecular dating. *Evolution*  
1040 **56(10)**:1931–1942.
- 104 **Das I, Haas A. 2010.** New species of *Microhyla* from Sarawak: Old World’s smallest frogs crawl  
1042 out of miniature pitcher plants on Borneo (Amphibia: Anura: Microhylidae). *Zootaxa* **2571**:37–  
1043 52.
- 104 **Das I. 1994.** A check-list of the amphibians and reptiles of Andaman and Nicobar Islands. *Journal*  
1045 *of the Andaman Science Association* **10(1)**:44-49.

- 104**Das I. 1999.** Biogeography of the amphibians and reptiles of the Andaman and Nicobar Islands. In:  
1047 **H. Ota** (Ed.). *Tropical island herpetofauna. Origin, current diversity and conservation.* pp:43–  
1048 77. Elsevier Science B.V., Amsterdam.
- 104**Datta-Roy A, Karanth KP. 2009.** The out-of-India hypothesis: What do molecules suggest?  
1050 *Journal of biosciences* **34**(5):687–697.
- 105**Dayanandan S, Aston PS, Williams SM, Primack RB. 1999.** Phylogeny of the tropical tree family  
1052 Dipterocarpaceae based on nucleotide sequences of the chloroplast *rbcL* gene. *American journal*  
1053 *of Botany* **86**:1182–1190.
- 105**De Bruyn M, Stelbrink B, Morley RJ, Hall R, Carvalho GR, Cannon CH, Van den Bergh G,**  
1055 **Meijaard E, Metcalfe I, Boitani L, Maiorano L, Shoup R, von Rintelen T. 2014.** Borneo and  
1056 Indochina are major evolutionary hotspots for Southeast Asian biodiversity. *Systematic Biology*  
1057 **63**:879-901.
- 105**de Sá RO, Streicher JW, Sekonyela R, Forlani MC, Loader SP, Greenbaum E, Richards S,**  
1059 **Haddad CF. 2012.** Molecular phylogeny of microhylid frogs (Anura: Microhylidae) with  
1060 emphasis on relationships among New World genera. *BMC Evolutionary Biology* **12**(1):241.
- 106**Deepak V, Karanth P. 2018.** Aridification driven diversification of fan-throated lizards from the  
1062 Indian subcontinent. *Molecular Phylogenetics and Evolution* **120**:53–62 DOI  
1063 10.1016/j.ympev.2017.11.016.
- 106**Dellicour S, Flot J-F. 2015.** Delimiting species-poor datasets using single molecular markers: a  
1065 study of barcode gaps, haplowebs and GMYC. *Systematic Biology* **64**:900–908.
- 106**Dellicour S, Flot J-F. 2018.** The hitchhiker’s guide to single-locus species delimitation. *Molecular*  
1067 *Ecology Resources* **18**:1234–1246 DOI: 10.1111/1755-0998.12908
- 106**Ding L, Maksatbek S, Cai F, Wang H, Song P, Ji W, Xu Q, Zhang L, Muhammad Q, Upendra**  
1069 **B. 2017.** Processes of initial collision and suturing between India and Asia. *Science China Earth*  
1070 *Sciences* **60**(4):635–651.
- 107**Drummond AJ, Suchard MA, Xie D, Rambaut A. 2012.** Bayesian phylogenetics with BEAUti  
1072 and the BEAST 1.7. *Molecular Biology and Evolution* **29**:1969–1973.
- 107**Dubois A. 1987.** Miscellanea taxinomica batrachologica (II). *Alytes* **6**:1–9.
- 107**Dutta SK, Vasudevan K, Chaitra MS, Shanker K, Aggarwal R. 2004.** Jurassic frogs and the  
1075 evolution of amphibian endemism in the Western Ghats. *Current Science* **86**:211–216.



- 1076 **Eyer PA, Seltzer R, Reiner-Brodetzki T, Hefetz A. 2017.** An integrative approach to untangling  
1077 species delimitation in the *Cataglyphis bicolor* desert ant complex in Israel. *Molecular*  
1078 *phylogenetics and evolution* **115**:128–139.
- 1079 **Fei L, Ye C-Y, Jiang J, Xie F, Huang Y (Eds.) 2005.** *An Illustrated Key to Chinese Amphibians.*  
1080 Sichuan Publishing House of Science and Technology, Chengdu.
- 1081 **Felsenstein J. 2004.** *Inferring Phylogenies*. Vol. 2. Sunderland: Sinauer Associates, Inc., 465.
- 1082 **Feng Y, Blackburn DC, Liang D, Hillis DM, Wake DB, Cannatella DC, Zhang P. 2017.**  
1083 Phylogenomics reveals rapid, simultaneous diversification of three major clades of Gondwanan  
1084 frogs at the Cretaceous–Paleogene boundary. *Proceedings of the National Academy of Sciences*  
1085 **114**:5864–5870.
- 1086 **Fromhage L, Vences M, Veith M. 2004.** Testing alternative vicariance scenarios in Western  
1087 Mediterranean discoglossid frogs. *Molecular Phylogenetics and Evolution* **31**(1):308–322.
- 1088 **Frost DR, Grant T, Faivovich J, Bain RH, Haas A, Haddad CFB, de Sá RO, Channing A,**  
1089 **Wilkinson M, Donnellan SC, Raxworthy CJ, Campbell JA, Blotto BL, Moler PE, Drewes**  
1090 **RC, Nussbaum RA, Lynch JD, Green DM, Wheeler WC. 2006.** The amphibian tree of life.  
1091 *Bulletin of the American Museum of Natural History* **297**:1–370.
- 1092 **Frost DR. 2020.** Amphibian species of the world 6.0, an online reference. (2020-01). New York:  
1093 Darrel Frost and the American Museum of Natural History. Available at  
1094 <http://research.amnh.org/herpetology/amphibia/index.html> (accessed 15 January 2020).
- 1095 **Fujisawa T, Barraclough TG. 2013.** Delimiting species using single-locus data and the  
1096 Generalized Mixed Yule Coalescent approach: a revised method and evaluation on simulated  
1097 data sets. *Systematic Biology* **62**:707–724.
- 1098 **Fujita MK, Leaché AD, Burbrink FT, McGuire JA, Moritz C. 2012.** Coalescent based species  
1099 delimitation in an integrative taxonomy. *Trends in Ecology and Evolution* **27**:480–488.
- 1100 **Garg S, Biju SD. 2019.** New microhylid frog genus from Peninsular India with Southeast Asian  
1101 affinity suggests multiple Cenozoic biotic exchanges between India and Eurasia. *Scientific*  
1102 *Reports* **9**(1):1906.
- 1103 **Garg S, Suyesh R, Das A, Jiang J, Wijayathilaka N, Amarasinghe AT, Alhadi F, Vineeth KK,**  
1104 **Aravind NA, Senevirathne G, Meegaskumbura M, Biju SD. 2019.** Systematic revision of  
1105 *Microhyla* (Microhylidae) frogs of South Asia: a molecular, morphological, and acoustic  
1106 assessment. *Vertebrate Zoology* **69**(1):1–71.

- 110Geissler P, Hartmann T, Ihlow F, Rödder D, Poyarkov NA, Nguyen TQ, Ziegler T, Böhme W.  
1108 2015. The Lower Mekong: an insurmountable barrier to amphibians in southern Indochina?.  
1109 *Biological Journal of the Linnean Society* **114**(4):905–914.
- 111Gower DJ, Kupfer A, Oommen OV, Himstedt W, Nussbaum RA, Loader SP, Presswell B,  
1111 Müller H, Krishna SB, Boistel R, Wilkinson M. 2002. A molecular phylogeny of ichthyophiid  
1112 caecilians (Amphibia: Gymnophiona: Ichthyophiidae): out of India or out of South East Asia?.  
1113 *Proceedings of the Royal Society of London, Series B: Biological Sciences*. **269**(1500):1563-  
1114 1569.
- 111Grismer JL, Schulte JA, Alexander A, Wagner P, Travers SL, Buehler MD, Welton LJ, Brown  
1116 RM. 2016. The Eurasian invasion: phylogenomic data reveal multiple Southeast Asian origins  
1117 for Indian Dragon Lizards. *BMC Evolutionary Biology* **16**(1):43.
- 111Guo ZT, Ruddiman WF, Hao QZ, Wu HB, Qiao YS, Zhu RX, Peng SZ, Wei JJ, Yuan BY, Liu  
1119 TS. 2002. Onset of Asian desertification by 22Myr ago inferred from loess deposits in China.  
1120 *Nature* **416**(6877):159–163 DOI 10.1038/416159a.
- 112Hall R. 2012. Late Jurassic–Cenozoic reconstructions of the Indonesian region and the Indian Ocean.  
1122 *Tectonophysics* **570**:1–41.
- 112Hall TA. 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program  
1124 for Windows 95/98/NT. *Nucleotide* **41**:95–98.
- 112Hanken J, Wake DB. 1993. Miniaturization of body size: organismal consequences and  
1126 evolutionary significance. *Annual Review of Ecology and Systematics* **24**:501–519.
- 112Hanken J. 1985. Morphological novelty in the limb skeleton accompanies miniaturization in  
1128 salamanders. *Science* **229**:871–874.
- 112Harikrishnan S, Vasudevan K, De Silva A, Deepak V, Kar NB, Naniwadekar R, Lalremruata  
1130 A, Prasanna KR, Aggarwal RK. 2012. Phylogeography of *Dasia* Gray, 1830 (Reptilia:  
1131 Scincidae), with the description of a new species from southern India. *Zootaxa* **3233**(1):37–51.
- 113Harrison TM, Copeland P, Kidd WSF, Yin A. 1992. Raising Tibet. *Science* **255**(5052):1663–1670  
1133 DOI 10.1126/science.255.5052.1663.
- 113Hasan M, Islam MM, Khan MMR, Alam MS, Kurabayashi A, Igawa T, Kuramoto M, Sumida  
1135 M. 2012. Cryptic anuran biodiversity in Bangladesh revealed by mitochondrial 16S rRNA gene  
1136 sequences. *Zoological science* **29**(3):162–172.

- 1133 **Hasan M, Islam MM, Kuramoto M, Kurabayashi A, Sumida M. 2014a.** Description of two new  
1138 species of *Microhyla* (Anura: Microhylidae) from Bangladesh. *Zootaxa* **3755**:401–408.
- 1139 **Hasan M, Islam MM, Khan MMR, Igawa T, Alam MS, Djong HT, Kurniawan N, Joshy H,**  
1140 **Sen YH, Belabut DM, Kurabayashi A. 2014b.** Genetic divergences of South and Southeast  
1141 Asian frogs: a case study of several taxa based on 16S ribosomal RNA gene data with notes on  
1142 the generic name *Fejervarya*. *Turkish Journal of Zoology* **38**(4):389–411.
- 1143 **Holman JA. 2003.** *Fossil frogs and toads of North America*. In: Farlow JO (Ed.) Indiana university  
1144 press. Bloomington: Indiana 246 pp.
- 1145 **Howlader MSA, Nair A, Gopalan SV, Merilä J. 2015.** A new species of *Microhyla* (Anura:  
1146 Microhylidae) from Nilphamari, Bangladesh. *PloS One* **10**(3):e0119825:1–18.
- 1147 **Huelsenbeck JP, Hillis DM. 1993.** Success of phylogenetic methods in the four-taxon case.  
1148 *Systematic Biology* **42**(3):247–264 DOI 10.1093/sysbio/42.3.247.
- 1149 **Huelsenbeck JP, Ronquist F. 2001.** MrBayes: Bayesian inference of phylogenetic trees.  
1150 *Bioinformatics* **17**:754–755.
- 1151 **Inger, R. F., and K. J. Frogner. 1979.** New species of narrow-mouth frogs (genus *Microhyla*) from  
1152 Borneo. *Sarawak Museum Journal* **27**:311–322.
- 1153 **Kamei RG, Mauro DS, Gower DJ, Van Bocxlaer I, Sherratt E, Thomas A, Babu S, Bossuyt F,**  
1154 **Wilkinson M, Biju SD. 2012.** Discovery of a new family of amphibians from northeast India  
1155 with ancient links to Africa. *Proceedings of the Royal Society B: Biological Sciences*.  
1156 **279**(1737):2396-2401.
- 1157 **Karanth KP. 2006.** Out-of-India: Gondwanan origin of some tropical Asian biota. *Current Science*  
1158 **90**:789–792.
- 1159 **Karunarathna S, Poyarkov NA, de Silva A, Madawala M, Botejue M, Gorin VA, Surasinghe T,**  
1160 **Gabadage D, Ukuwela KD, Bauer AM. 2019.** Integrative taxonomy reveals six new species of  
1161 day geckos of the genus *Cnemaspis* Strauch, 1887 (Reptilia: Squamata: Gekkonidae) from  
1162 geographically isolated hill forests in Sri Lanka. *Vertebrate Zoology* **64**:247–298.
- 1163 **Katoh K, Misawa K, Kuma K, Miyata T. 2002.** MAFFT: a novel method for rapid multiple  
1164 sequence alignment based on fast Fourier transform. *Nucleic Acids Research* **30**(14):3059–3066.
- 1165 **Khatiwada JR, Shu GC, Wang SH, Thapa A, Wang B, Jiang J. 2017.** A new species of the genus  
1166 *Microhyla* (Anura: Microhylidae) from Eastern Nepal. *Zootaxa* **4254**(2):221–239.

- 1166 **Klaus S, Morley RJ, Plath M, Zhang YP, Li JT. 2016.** Biotic interchange between the Indian  
1168 subcontinent and mainland Asia through time. *Nature Communications* **7**:12132.
- 1169 **Klaus S, Schunart CD, Streit B, Pfenninger M. 2010.** When Indian crabs were not yet Asian -  
1170 biogeographic evidence for Eocene proximity of India and Southeast Asia. *BMC Evolutionary*  
1171 *Biology* **10**:1–9.
- 1172 **Köhler F, Glaubrecht M. 2007.** Out of Asia and into India: on the molecular phylogeny and  
1173 biogeography of the endemic freshwater gastropod *Paracrostoma* Cossmann, 1900  
1174 (Caenogastropoda: Pachychilidae). *Biological Journal of the Linnean Society* **91(4)**:627–651.
- 1175 **Kraus F. 2011.** At the lower size limit for tetrapods, two new species of the miniaturized frog genus  
1176 *Paedophryne* (Anura, Microhylidae). *ZooKeys* **154**:71–88 DOI 10.3897/zookeys.154.1963
- 1177 **Kurabayashi A, Matsui M, Belabut DM, Yong HS, Ahmad N, Sudin A, Kuramoto M, Hamidy**  
1178 **A, Sumida M. 2011.** From Antarctica or Asia? New colonization scenario for Australian-New  
1179 Guinean narrow mouth toads suggested from the findings on a mysterious genus  
1180 *Gastrophrynoides*. *BMC Evolutionary Biology* **11(1)**:175.
- 1181 **Lanfear R, Calcott B, Ho SYW, Guindon S. 2012.** PartitionFinder: combined selection of  
1182 partitioning schemes and substitution models for phylogenetic analyses. *Molecular Biology and*  
1183 *Evolution* **29(6)**:1695–1701 DOI 10.1093/molbev/mss020.
- 1184 **Lee KH, Shaner PJ, Lin YP, Lin SM. 2016.** Geographic variation in advertisement calls of a  
1185 Microhylid frog – testing the role of drift and ecology. *Ecology and Evolution* **6(10)**:3289–3298.
- 1186 **Lehr E, Coloma LA. 2008.** A minute new Ecuadorian Andean frog (Anura: Strabomantidae,  
1187 *Pristimantis*). *Herpetologica* **64(3)**:354–367.
- 1188 **Li JT, Li Y, Klaus S, Rao DQ, Hillis DM, Zhang YP. 2013.** Diversification of rhacophorid frogs  
1189 provides evidence for accelerated faunal exchange between India and Eurasia during the  
1190 Oligocene. *Proceedings of the National Academy of Sciences* **110(9)**:3441–3446.
- 1191 **Macey JR, Kuehl JV, Larson A, Robinson MD, Ugurtas IH, Ananjeva NB, Rahman H, Javed**  
1192 **HI, Osman RM, Doumma A, Papenfuss TJ. 2008.** Socotra Island the forgotten fragment of  
1193 Gondwana: Unmasking chameleon lizard history with complete mitochondrial genomic data.  
1194 *Molecular Phylogenetics and Evolution* **49(3)**:1015–1018.
- 1195 **Maddison WP, Maddison DR. 2017.** Mesquite: a modular system for evolutionary analysis.  
1196 Version 3.31. Available at <http://mesquiteproject.org>.

- 1197 **Maddison WP. 1991.** Squared-change parsimony reconstructions of ancestral states for continuous-  
1198 valued characters on a phylogenetic tree. *Systematic Zoology* **40(3)**:304–314 DOI  
1199 10.2307/2992324.
- 1200 **Matsui M, Hamidy A, Belabut DM, Ahmad N, Panha S, Sudin A, Khonsue W, Oh HS, Yong**  
1201 **HS, Jiang JP, Nishikawa K. 2011.** Systematic relationships of Oriental tiny frogs of the family  
1202 Microhylidae (Amphibia, Anura) as revealed by mtDNA genealogy. *Molecular Phylogenetics*  
1203 *and Evolution* **61(1)**:167–176.
- 1204 **Matsui M, Hamidy A, Eto K. 2013.** Description of a new species of *Microhyla* from Bali,  
1205 Indonesia (Amphibia, Anura). *Zootaxa* **3670(4)**:579–590.
- 1206 **Matsui M, Ito H, Shimada T, Ota H, Saidapur SK, Khonsue W, Tanaka-Ueno T, Wu GF. 2005.**  
1207 Taxonomic relationships within the pan-oriental narrowmouth toad *Microhyla ornata* as revealed  
1208 by mtDNA analysis (Amphibia, Anura, Microhylidae). *Zoological Science* **22(4)**:489–495.
- 1209 **Matsui M. 2011.** Taxonomic revision of one of the Old World's smallest frogs, with description of a  
1210 new Bornean *Microhyla* (Amphibia, Microhylidae). *Zootaxa* **2814(1)**:33–49.
- 1211 **Matzke NJ. 2013.** Probabilistic historical biogeography: new models for founder-event speciation,  
1212 imperfect detection, and fossils allow improved accuracy and model-testing. *Frontiers of*  
1213 *Biogeography* **5(4)**.
- 1214 **Molnar P. 2005.** Mio-Pliocene growth of the Tibetan Plateau and evolution of East Asian climate.  
1215 *Palaeontologia Electronica* **8(1)**:1–23.
- 1216 **Morley RJ. 2000.** *Origin and evolution of tropical rain forests*, Wiley, Chichester 362 pp.
- 1217 **Morley RJ. 2018.** Assembly and division of the South and South-east Asian flora in relation to  
1218 tectonics and climate change. *Journal of Tropical Ecology* **34(4)**:209–234.
- 1219 **Mulcahy DG, Lee JL, Miller AH, Chand M, Thura MK, Zug GR. 2018.** Filling the BINs of life:  
1220 Report of an amphibian and reptile survey of the Tanintharyi (Tenasserim) Region of Myanmar,  
1221 with DNA barcode data. *ZooKeys* **757**:85–152.
- 1222 **Myers N, Mittermeier RA, Mittermeier CG, Da Fonseca GA, Kent J. 2000.** Biodiversity  
1223 hotspots for conservation priorities. *Nature* **403(6772)**:853.
- 1224 **Nguyen LT, Poyarkov NA, Nguyen TT, Nguyen TA, Tran VH, Gorin VA, Murphy RW,**  
1225 **Nguyen SN. 2019.** A new species of the genus *Microhyla* Tschudi, 1838 (Amphibia: Anura:  
1226 Microhylidae) from Tay Nguyen Plateau, Central Vietnam. *Zootaxa* **4543**:549–580.

- 1222 **Nguyen LT, Schmidt HA, Von Haeseler A, Minh BQ. 2015.** IQ-TREE: a fast and effective  
1228 stochastic algorithm for estimating maximum-likelihood phylogenies. *Molecular Biology and*  
1229 *Evolution* **32(1)**:268–274.
- 1230 **Nguyen TV, Duong TV, Luu KT, Poyarkov NA. 2020.** A new species of *Kurixalus* (Anura:  
1231 Rhacophoridae) from northern Vietnam with comments on the biogeography of the genus.  
1232 *Journal of Natural History in press* DOI 10.1080/00222933.2020.1728411
- 1233 **Oliver PM, Iannella A, Richards SJ, Lee MSY. 2017.** Mountain colonisation, miniaturisation and  
1234 ecological evolution in a radiation of direct-developing New Guinea frogs (*Choerophryne*,  
1235 Microhylidae). *PeerJ* **5**:e3077. DOI 10.7717/peerj.3077
- 1236 **Ota H. 1998.** Geographic patterns of endemism and speciation in amphibians and reptiles of the  
1237 Ryukyu Archipelago, Japan, with special reference to their paleogeographic implications.  
1238 *Researches on Population Ecology* **40**:189–204.
- 1239 **Parker HW. 1928.** The brevipitid frogs of the genus *Microhyla*. *Annals and Magazine of Natural*  
1240 *History*, Series 10 **2**:473–499.
- 1241 **Parker HW. 1934.** *A Monograph of the Frogs of the Family Microhylidae*. British Museum  
1242 (Natural History), London, (1) + viii + 208 pp.
- 1243 **Parra-Olea G, Rovito SM, García-París M, Maisano JA, Wake DB, Hanken J. 2016.** Biology of  
1244 tiny animals: three new species of minute salamanders (Plethodontidae: *Thorius*) from Oaxaca,  
1245 Mexico. *PeerJ* **4**:e2694.
- 1246 **Peloso PL, Frost DR, Richards SJ, Rodrigues MT, Donnellan S, Matsui M, Raxworthy CJ,**  
1247 **Biju SD, Lemmon EM, Lemmon AR, Wheeler WC. 2016.** The impact of anchored  
1248 phylogenomics and taxon sampling on phylogenetic inference in narrow-mouthed frogs (Anura,  
1249 Microhylidae). *Cladistics* **32(2)**:113–140.
- 1250 **Polilov AA. 2015.** Small is beautiful: features of the smallest insects and limits to miniaturization.  
1251 *Annual Review of Entomology* **60**:103–121.
- 1252 **Pons J, Barraclough TG, Gomez-Zurita J, Cardoso A, Duran DP, Hazell S, Kamoun S, Sumlin,**  
1253 **WD, Vogler AP. 2006.** Sequence-based species delimitation for the DNA taxonomy of  
1254 undescribed insects. *Systematic Biology* **55**:595–609.
- 1255 **Poyarkov NA, Gorin VA, Zaw T, Kretova VD, Gogoleva SI, Pawangkhanant P, Che J. 2019.**  
1256 On the road to Mandalay: contribution to the *Microhyla* Tschudi, 1838 (Amphibia: Anura:

- 1257 Microhylidae) fauna of Myanmar with description of two new species. *Zoological Research*  
1258 **40(4)**:244–276 DOI 10.24272/j.issn.2095-8137.2019.044.
- 1259 **Poyarkov NA, Nguyen TV, Duong TV, Gorin VA, Yang JH. 2018a.** A new limestone-dwelling  
1260 species of *Micryletta* (Amphibia: Anura: Microhylidae) from northern Vietnam. *PeerJ* **6**:e5771  
1261 DOI 10.7717/peerj.5771
- 1262 **Poyarkov NA, Suwannapoom C, Pawangkhanant P, Aksornneam A, Duong TV, Korost DV,**  
1263 **Che J. 2018b.** A new genus and three new species of miniaturized microhylid frogs from  
1264 Indochina (Amphibia: Anura: Microhylidae: Asterophryinae). *Zoological Research* **39(3)**:130–  
1265 155.
- 1266 **Poyarkov NA, Vassilieva AB, Orlov NL, Galoyan EA, Tran DTA, Le DTT, Kretova VD,**  
1267 **Geissler P. 2014.** Taxonomy and distribution of narrow-mouth frogs of the genus *Microhyla*  
1268 Tschudi, 1838 (Anura: Microhylidae) from Vietnam with descriptions of five new species.  
1269 *Russian Journal of Herpetology* **21**:89–148.
- 1270 **Puillandre N, Lambert A, Brouillet S, Achaz G. 2012.** ABGD, Automatic Barcode Gap Discovery  
1271 for primary species delimitation. *Molecular Ecology* **21**:1864–1877.
- 1272 **Pyron RA, Kandambi HKD, Hendry CR, Pushpamal V, Burbrink FT, Somaweera R. 2013.**  
1273 Genus-level molecular phylogeny of snakes reveals the origins of species richness in Sri Lanka.  
1274 *Molecular Phylogenetics and Evolution* **66**:969–975 DOI 10.1016/j.ympev.2012.12.004
- 1275 **Rakotoarison A, Scherz MD, Glaw F, Köhler J, Andreone F, Franzen M, Glos J, Hawlitschek**  
1276 **O, Jono T, Mori A, Ndriantsoa SH. 2017.** Describing the smaller majority: integrative  
1277 taxonomy reveals twenty-six new species of tiny microhylid frogs (genus *Stumpffia*) from  
1278 Madagascar. *Vertebrate Zoology* **67(3)**:271–398.
- 1279 **Rambaut A, Drummond AJ. 2007.** Tracer v1. 5. Available at <http://beast.bio.ed.ac.uk/Tracer>.
- 1280 **Raxworthy CJ, Forstner MRJ, Nussbaum RA. 2002.** Chameleon radiation by oceanic dispersal.  
1281 *Nature* **415**:784–787.
- 1282 **Ree RH, Moore BR, Webb CO, Donoghue MJ. 2005.** A likelihood framework for inferring the  
1283 evolution of geographic range on phylogenetic trees. *Evolution* **59(11)**:2299–2311 DOI  
1284 10.1111/j.0014-3820.2005.tb00940.x.
- 1285 **Ree RH, Smith SA. 2008.** Maximum likelihood inference of geographic range evolution by  
1286 dispersal, local extinction, and cladogenesis. *Systematic Biology* **57(1)**:4–14 DOI  
1287 10.1080/10635150701883881.

- 1288**Reid NM, Carstens BC. 2012.** Phylogenetic estimation error can decrease the accuracy of species  
1289 delimitation: a Bayesian implementation of the general mixed Yule-coalescent model. *BMC*  
1290 *Evolutionary Biology* **12**:196.
- 129**Renner SS. 2004.** Tropical trans-Atlantic disjunctions, sea surface currents, and wind patterns.  
1292 *International Journal of Plant Sciences* **165**:23–33.
- 129**Rieppel O. 1996.** Miniaturization in tetrapods: consequences for skull morphology. *Symposia of the*  
1294 *Zoological Society of London* **69**:47–61.
- 129**Rittmeyer EN, Allison A, Grundler MC, Thompson DK, Austin CC. 2012.** Ecological guild  
1296 evolution and the discovery of the world’s smallest vertebrate. *PloS One* **7(1)**:e29797. DOI  
1297 10.1371/journal.pone.0029797
- 129**Roelants K, Jiang J, Bossuyt F. 2004.** Endemic ranid (Amphibia: Anura) genera in southern  
1299 mountain ranges of the Indian subcontinent represent ancient frog lineages: Evidence from  
1300 molecular data. *Molecular Phylogenetics and Evolution* **31(2)**:730–740.
- 130**Ronquist F, Huelsenbeck JP. 2003.** MrBayes 3: Bayesian phylogenetic inference under mixed  
1302 models. *Bioinformatics* **19(12)**:1572–1574 DOI 10.1093/bioinformatics/btg180.
- 130**Ronquist F. 1997.** Dispersal-vicariance analysis: a new approach to the quantification of historical  
1304 biogeography. *Systematic Biology* **46**:195–203.
- 130**Rovito SM, Parra-Olea G, Hanken J, Bonett RM, Wake DB. 2013.** Adaptive radiation in  
1306 miniature: the minute salamanders of the Mexican highlands (Amphibia: Plethodontidae:  
1307 *Thorius*). *Biological Journal of the Linnean Society* **109**:622–643.
- 130**Russell DW, Sambrook J. 2001.** *Molecular cloning: a laboratory manual*. Cold Spring Harbor,  
1309 NY: Cold Spring Harbor Laboratory.
- 131**Sanchiz B. 1998.** *Encyclopedia of paleoherpetology. Part 4. Salientia*. München: Pfeil 275 pp.
- 131**Savage JM. 1973.** The geographic distribution of frogs: patterns and predictions: *In: Evolutionary*  
1312 *biology of the anurans*. University of Missouri Press, Missouri: 351–445.
- 131**Scherz MD, Hutter CR, Rakotoarison A, Riemann JC, Rödel MO, Ndriantsoa SH, Glos J,**  
1314 **Roberts SH, Crottini A, Vences M, Glaw F. 2019.** Morphological and ecological convergence  
1315 at the lower size limit for vertebrates highlighted by five new miniaturised microhylid frog  
1316 species from three different Madagascan genera. *PloS One* **14(3)**.



- 131 **Seshadri KS, Singal R, Priti H, Ravikanth G, Vidisha MK, Saurabh S, Pratik M, Gururaja KV.**  
1318 **2016.** *Microhyla laterite* sp. nov., a new species of *Microhyla* Tschudi, 1838 (Amphibia: Anura:  
1319 Microhylidae) from a laterite rock formation in South West India. *PloS One* **11(3)**:e0149727.
- 132 **Solovyeva EN, Lebedev VS, Dunayev EA, Nazarov RA, Bannikova AA, Che J, Murphy RW,**  
1321 **Poyarkov NA. 2018.** Cenozoic aridization in Central Eurasia shaped diversification of toad-  
1322 headed agamas (*Phrynocephalus*; Agamidae, Reptilia). *PeerJ* **6**:e4543 DOI 10.7717/peerj.4543
- 1323 **Sparks JS. 2003.** Molecular phylogeny and biogeography of the Malagasy and South Asian cichlids  
1324 (Teleostei: Perciformis: Cichlidae). *Molecular Phylogenetics and Evolution* **30**:599–614.
- 1325 **Suwannapoom C, Sumontha M, Tunprasert J, Ruangsuwan T, Pawangkhanant P, Korost DV,**  
1326 **Poyarkov N.A. 2018.** A striking new genus and species of cave-dwelling frog (Amphibia:  
1327 Anura: Microhylidae: Asterophryinae) from Thailand. *PeerJ* **6**:e4422 DOI 10.7717/peerj.4422
- 1328 **Tamura K, Stecher G, Peterson D, Filipski A, Kumar S. 2013.** MEGA6: molecular evolutionary  
1329 genetics analysis version 6.0. *Molecular Biology and Evolution* **30(12)**:2725–2729.
- 1330 **Tominaga A, Matsui M, Shimoji N, Khonsue W, Wu CS, Toda M, Eto K, Nishikawa K, Ota H.**  
1331 **2019.** Relict distribution of *Microhyla* (Amphibia: Microhylidae) in the Ryukyu Archipelago:  
1332 High diversity in East Asia maintained by insularization. *Zoologica Scripta* **48(4)**:440–453.
- 1333 **Tracy CR, Christian KA, Tracy CR. 2010.** Not just small, wet, and cold: effects of body size and  
1334 skin resistance on thermoregulation and arboreality of frogs. *Ecology* **91(5)**:1477–1484. DOI  
1335 10.1890/09-0839.1
- 1336 **Tu N, Yang M-H, Liang D, Zhang P. 2018.** A large-scale phylogeny of Microhylidae inferred from  
1337 a combined dataset of 121 genes and 427 taxa. *Molecular Phylogenetics and Evolution* **126**:85–  
1338 91.
- 1339 **Turner H, Hovenkamp P, Van Welzen PC. 2001.** Biogeography of Southeast Asia and the West  
1340 Pacific. *Journal of Biogeography* **28**:217–230.
- 1341 **Van Bocxlaer I, Roelants K, Biju SD, Nagaraju J, Bossuyt F. 2006.** Late Cretaceous vicariance in  
1342 Gondwanan amphibians. *PloS One* **1(1)**:e74.
- 1343 **Van der Meijden A, Vences M, Hoegg S, Boistel R, Channing A, Meyer A. 2007.** Nuclear gene  
1344 phylogeny of narrow-mouthed toads (Family: Microhylidae) and a discussion of competing  
1345 hypotheses concerning their biogeographical origins. *Molecular Phylogenetics and Evolution*  
1346 **44**:1017–1030.


- 1344 **Van Hinsbergen DJ, Kapp P, Dupont - Nivet G, Lippert PC, DeCelles PG, Torsvik TH. 2011.**  
1348 Restoration of Cenozoic deformation in Asia and the size of Greater India. *Tectonics* **30(5)**:  
1349 TC5003.
- 1350 **Van Hinsbergen DJ, Lippert PC, Dupont-Nivet G, McQuarrie N, Doubrovine PV, Spakman W,**  
1351 **Torsvik TH. 2012.** Greater India Basin hypothesis and a two-stage Cenozoic collision between  
1352 India and Asia. *Proceedings of the National Academy of Sciences* **109(20)**:7659–7664.
- 1353 **Vassilieva AB, Trounov VL, Poyarkov NA, Galoyan EA. 2017.** The phytotelm tadpoles of  
1354 *Microhyla arboricola* (Anura: Microhylidae) from Vietnam, with comments on reproductive  
1355 biology and development. *Zootaxa* **4247(4)**:413–428.
- 1356 **Vences M, Thomas M, Bonett RM, Vieites DR. 2005.** Deciphering amphibian diversity through  
1357 DNA barcoding: chances and challenges. *Philosophical Transactions of the Royal Society of*  
1358 *London Series B: Biological Sciences* **360(1462)**:1859–1868.
- 1359 **Vences M, Vieites DR, Glaw F, Brinkmann H, Kosuch J, Veith M, Meyer A. 2003.** Multiple  
1360 overseas dispersal in amphibians. *Proceedings of the Royal Society of London Series B:*  
1361 *Biological Sciences*. **270(1532)**:2435–2442.
- 1362 **Vieites DR, Wollenberg KC, Andreone F, Köhler J, Glaw F, Vences M. 2009.** Vast  
1363 underestimation of Madagascar's biodiversity evidenced by an integrative amphibian inventory.  
1364 *Proceedings of the National Academy of Sciences* **106(20)**:8267–8272.
- 1365 **Vineeth KK, Radhakrishna UK, Godwin RD, Anwasha S, Rajashekhar KP, Aravind NA. 2018.**  
1366 A new species of *Microhyla* Tschudi, 1838 (Anura: Microhylidae) from West Coast of India: an  
1367 integrative taxonomic approach. *Zootaxa* **4420**:151–179.
- 1368 **Wang YH, Hsiao YW, Lee KH, Tseng HY, Lin YP, Komaki S, Lin SM. 2017.** Acoustic  
1369 differentiation and behavioral response reveals cryptic species within *Buergeria* treefrogs (Anura,  
1370 Rhacophoridae) from Taiwan. *PloS One* **12(9)**.
- 1371 **Wijayathilaka N, Garg S, Senevirathne G, Karunarathna N, Biju S, Meegaskumbura M. 2016.**  
1372 A new species of *Microhyla* (Anura: Microhylidae) from Sri Lanka: an integrative taxonomic  
1373 approach. *Zootaxa* **4066(3)**: 331–342.
- 1374 **Wilkinson M, Sheps J, Oommen OV, Cohen BL. 2002.** Phylogenetic relationships of Indian  
1375 caecilians (Amphibia: Gymnophiona) inferred from mitochondrial rRNA gene sequences.  
1376 *Molecular Phylogenetics and Evolution* **23(3)**:401–407.

- 1377 **Wogan GOU, Vindum JV, Wilkinson JA, Koo MS, Slowinski JB, Win H, Thin T, Kyi SW, Oo**  
1378 **SL, Lwin KS, Shein AK. 2008.** New country records and range extensions for Myanmar  
1379 amphibians and reptiles. *Hamadryad* **33**:83–96.
- 1380 **Wood PL, Heinicke MP, Jackman TR, Bauer AM. 2012.** Phylogeny of bent-toed geckos  
1381 (*Cyrtodactylus*) reveals a west to east pattern of diversification. *Molecular Phylogenetics and*  
1382 *Evolution* **65**(3):992–1003.
- 1383 **Woodruff DS. 2010.** Biogeography and conservation in Southeast Asia: how 2.7 million years of  
1384 repeated environmental fluctuations affect today’s patterns and the future of the remaining  
1385 refugial-phase biodiversity. *Biodiversity and Conservation* **19**(4):919–941.
- 1386 **Yang Z. 2007.** PAML 4: phylogenetic analysis by maximum likelihood. *Molecular Biology and*  
1387 *Evolution* **24**(8):1586–1591 DOI 10.1093/molbev/msm088.
- 1388 **You CW, Poyarkov NA, Lin SM. 2015.** Diversity of the snail-eating snakes *Pareas* (Serpentes,  
1389 *Pareatidae*) from Taiwan. *Zoologica Scripta* **44**(4):349–361.
- 1390 **Yu Y, Harris AJ, Blair C, He X. 2015.** RASP (Reconstruct Ancestral State in Phylogenies): a tool  
1391 for historical biogeography. *Molecular Phylogenetics and Evolution* **87**:46–49.
- 1392 **Xuan ZY, Suwannapoom C, Yan F, Poyarkov NA, Nguyen SN, Chen HM, Chomdej S,**  
1393 **Murphy RW, Che J. 2016.** Red River barrier and Pleistocene climatic fluctuations shaped the  
1394 genetic structure of *Microhyla fissipes* complex (Anura: Microhylidae) in southern China and  
1395 Indochina. *Current Zoology* **62**(6):531–543.
- 1396 **Zwickl DJ, Hillis DM. 2002.** Increased taxon sampling greatly reduces phylogenetic error.  
1397 *Systematic Biology* **51**:588–598. DOI 10.1080/10635150290102339

# Figure 1

Distribution and species richness of Oriental tiny frogs of the genus *Microhyla*.


Heatmap indicates approximate number of sympatrically co-occurring species (from 1 to 9); the highest species richness is observed in southern Vietnam and Malayan Peninsula. Photo by Nikolay A. Poyarkov.



# Figure 2

Biogeographic history of *Microhyla*.

A – Biogeographic regions used in the present study; B – BEAST chronogram on the base of 3207 bp-long mtDNA+nuDNA dataset with the results of ancestral area reconstruction in RASP. For biogeographic areas definitions, species occurrence data and transition matrices see Supplemental Information 1 and Tables S5–S6. Node colors correspond to the respective biogeographic areas; values inside node icons correspond to node numbers (see Fig. S3 and Table S11 for divergence time estimates); values near nodes indicate marginal probabilities for ancestral ranges (S-DIVA analysis); icons illustrate vicariant and dispersal events (see Legend). Red arrows from 1 to 5 correspond to the dispersals to the Indian Subcontinent by *Microhyla* II lineages.




# Figure 3

Paleogeography and climate of South and Southeast Asia, 60–25 Ma.

Tectonic reconstructions modified from *Hall (2012)*; paleoclimate reconstructions based on *Morley (2018)*. Solid arrows indicate directions of plant dispersals (dark-green – perhumid floral elements, light-green – seasonal wet / seasonal dry elements) (from *Morley, 2018*); red arrows show probable areas of Microhylinae diversification and ways of their dispersal. A – K/T boundary to early Paleocene: the isolated Indian subcontinent (ISC) is drifting northwards cradling perhumid tropical biota, Southeast Asia (SEA) has primarily seasonal wet or seasonal dry climate, no land connection between SEA and ISC, basal radiation of Microhylinae in the ISC; B – Paleocene to early Eocene: the ISC and SEA are at the same latitude within same perhumid climate zone, first land connections between India and Sundaland via Incertus Arc, dispersal of Microhylinae from the ISC to SEA; C – Middle Eocene: land connection between the ISC and mainland Southeast Asia (modern-day Myanmar), basal radiation of *Microhyla* – *Glyphoglossus* assemblage in SEA; D – Oligocene: India drifts into northern high-pressure zone and seasonally dry climates predominate across the ISC and SEA, *Microhyla* II lineages colonize the ISC from SEA.








# Figure 4

Updated mtDNA-genealogy of the *Microhyla* - *Glyphoglossus* assemblage (full tree, part 1).


BI genealogy of *Microhyla* and *Glyphoglossus* samples examined in this study reconstructed from 2478 bp of mtDNA fragment. Values at nodes correspond to BI PP/ML BS, respectively; numbers at tips correspond to sample numbers summarized in Supplementary Table S1. Colors and letters (A-I) correspond to species groups of the *Microhyla* - *Glyphoglossus* assemblage. Yellow and red bars present the results of species delimitation analyses from bGMYC and ABGD algorithms, respectively. Frog photos are given in one scale, numbers near thumbnails correspond to species: 1 - *Microhyla nepenthicola*; 2 - *M. borneensis*; 3 - *M. malang*; 4 - *M. orientalis*; 5 - *M. mantheyi*; 6 - *M. minuta*; 7 - *M. achatina*; 8 - *M. irrawaddy*; 9 - *M. heymonsi*; 10 - *M. pineticola*; 11 - *M. fodiens*; 12 - *M. fissipes*; 13 - *M. mukhlesuri*; 14 - *M. chakrapanii*; 15 - *M. okinavensis*; 16 - *M. berdmorei* (Vietnam); 17 - *M. berdmorei* (Malaysia); 18 - *M. picta*; 19 - *M. pulchra*; 20 - *M. zeylanica*; 21 - *M. sholigari*; 22 - *M. karunaratnei*; 23 - *Microhyla* sp. 2; 24 - *M. ornata*; 25 - *M. mihintalei*; 26 - *M. butleri*; 27 - *M. aurantiventris*; 28 - *M. palmipes*; 29 - *M. annamensis*; 30 - *M. marmorata*; 31 - *M. pulverata*; 32 - *M. annectens*; 33 - *M. petrigena*; 34 - *M. perparva*; 35 - *M. pulchella*; 36 - *M. arboricola*; 37 - *Glyphoglossus molossus*; 38 - *G. guttulatus*. Photos by Nikolay A. Poyarkov, Indraneil Das, Vladislav A. Gorin, Parinya Pawangkhanant, Luan Thanh Nguyen, and Evgeniya N. Solovyeva.



# Figure 5

Updated mtDNA-genealogy of the *Microhyla* - *Glyphoglossus* assemblage (full tree, part 2).

BI genealogy of *Microhyla* and *Glyphoglossus* samples examined in this study reconstructed from 2478 bp of mtDNA fragment. Values at nodes correspond to BI PP/ML BS, respectively; numbers at tips correspond to sample numbers summarized in Supplementary Table S1. For legend, see Fig. 4. Photos by Nikolay A. Poyarkov, Indraneil Das, Vladislav A. Gorin, Parinya Pawangkhanant, Luan Thanh Nguyen, and Evgeniya N. Solovyeva.






# Figure 6

Maximum Likelihood tree for the “total evidence” analysis of the 3207 bp-long concatenated mtDNA + nuclear DNA dataset.

Values at nodes correspond to BEAST PP / ML BS / BI PP, respectively; black and white circles correspond to well-supported ( $BI\ PP \geq 0.95$ ;  $ML\ BS \geq 90$ ) and moderately supported ( $0.95 > BI\ PP \geq 0.90$ ;  $90 > ML\ BS \geq 75$ ) nodes, respectively; no circles indicate unsupported nodes. Color marking of species groups in *Microhyla* species complex corresponds to Figs. 4–5, but not to Fig. 2. For legend, see Fig. 4. Photos by Nikolay A. Poyarkov, Indraneil Das, Vladislav A. Gorin, Parinya Pawangkhanant, Luan Thanh Nguyen, and Evgeniya N. Solovyeva.



# Figure 7

Body size evolution among members of the *Microhyla* - *Glyphoglossus* assemblage.

See Table S7 for SVL data. Color of branches corresponds to average SVL in males (A) and females (B) (see legend).

