

# Acrodont tooth implantation via severe ankylosis as an adaptation of strong bite force (#45825)

1

First submission

## Guidance from your Editor

Please submit by **29 Feb 2020** for the benefit of the authors (and your \$200 publishing discount) .



### Structure and Criteria

Please read the 'Structure and Criteria' page for general guidance.



### Raw data check

Review the raw data.



### Image check

Check that figures and images have not been inappropriately manipulated.

Privacy reminder: If uploading an annotated PDF, remove identifiable information to remain anonymous.

## Files

Download and review all files from the [materials page](#).

3 Figure file(s)  
2 Table file(s)  
1 Raw data file(s)  
2 Other file(s)



# Structure and Criteria

## Structure your review

The review form is divided into 5 sections. Please consider these when composing your review:

1. BASIC REPORTING
2. EXPERIMENTAL DESIGN
3. VALIDITY OF THE FINDINGS
4. General comments
5. Confidential notes to the editor

 You can also annotate this PDF and upload it as part of your review

When ready [submit online](#).

## Editorial Criteria

Use these criteria points to structure your review. The full detailed editorial criteria is on your [guidance page](#).

### BASIC REPORTING

-  Clear, unambiguous, professional English language used throughout.
-  Intro & background to show context. Literature well referenced & relevant.
-  Structure conforms to [Peerj standards](#), discipline norm, or improved for clarity.
-  Figures are relevant, high quality, well labelled & described.
-  Raw data supplied (see [Peerj policy](#)).

### EXPERIMENTAL DESIGN

-  Original primary research within [Scope of the journal](#).
-  Research question well defined, relevant & meaningful. It is stated how the research fills an identified knowledge gap.
-  Rigorous investigation performed to a high technical & ethical standard.
-  Methods described with sufficient detail & information to replicate.

### VALIDITY OF THE FINDINGS

-  Impact and novelty not assessed. Negative/inconclusive results accepted. *Meaningful* replication encouraged where rationale & benefit to literature is clearly stated.
-  All underlying data have been provided; they are robust, statistically sound, & controlled.
-  Speculation is welcome, but should be identified as such.
-  Conclusions are well stated, linked to original research question & limited to supporting results.

# Standout reviewing tips

3



The best reviewers use these techniques

## Tip

**Support criticisms with evidence from the text or from other sources**

## Example

*Smith et al (J of Methodology, 2005, V3, pp 123) have shown that the analysis you use in Lines 241-250 is not the most appropriate for this situation. Please explain why you used this method.*

**Give specific suggestions on how to improve the manuscript**

*Your introduction needs more detail. I suggest that you improve the description at lines 57- 86 to provide more justification for your study (specifically, you should expand upon the knowledge gap being filled).*

**Comment on language and grammar issues**

*The English language should be improved to ensure that an international audience can clearly understand your text. Some examples where the language could be improved include lines 23, 77, 121, 128 – the current phrasing makes comprehension difficult.*

**Organize by importance of the issues, and number your points**

1. Your most important issue
2. The next most important item
3. ...
4. The least important points

**Please provide constructive criticism, and avoid personal opinions**

*I thank you for providing the raw data, however your supplemental files need more descriptive metadata identifiers to be useful to future readers. Although your results are compelling, the data analysis should be improved in the following ways: AA, BB, CC*

**Comment on strengths (as well as weaknesses) of the manuscript**

*I commend the authors for their extensive data set, compiled over many years of detailed fieldwork. In addition, the manuscript is clearly written in professional, unambiguous language. If there is a weakness, it is in the statistical analysis (as I have noted above) which should be improved upon before Acceptance.*

# Acrodont tooth implantation via severe ankylosis as an adaptation of strong bite force

Kelsey M Jenkins<sup>Corresp., 1</sup>, Jack O Shaw<sup>1</sup>

<sup>1</sup> Geology and Geophysics, Yale University, New Haven, United States of America

Corresponding Author: Kelsey M Jenkins  
Email address: kelsey.jenkins@yale.edu

Extant and extinct reptiles exhibit numerous combinations of tooth implantation and attachment. Tooth implantation ranges from those possessing roots and lying within a socket (thecodonty), to teeth lying against the lingual wall of the jawbone (pleurodonty), to teeth without roots or sockets that are attached to the apex of the marginal jawbones (acrodonty). Attachment may be ligamentous or via ankylosis (i.e., fusion). Adaptive reasonings are proposed as an underlying driver for evolutionary changes in some forms of tooth implantation and attachment. However, a substantiated adaptive hypothesis is lacking for the state of acrodonty via severe ankylosis that is seen in several lineages of Lepidosauria, a clade that is plesiomorphically pleurodont. We hypothesize that acrodonty via severe ankylosis, as seen in *Sphenodon punctatus* and acrodontan lizards, is an adaptation either resulting from or allowing for a stronger bite force. We use bite force data gathered from the literature to show that those taxa possessing acrodont dentition attached via severe ankylosis possess a stronger bite force than those taxa with pleurodont dentition. Furthermore, we discuss the evolution of acrodonty and potential behaviors related to strong bite force that influenced the evolution of acrodonty within Rhynchocephalia, Acrodonta, and Trogonophidae. Implications for the evolution of different forms of tooth implantation and attachment include ancient behavioral shifts.

# **Acrodont Tooth Implantation Via Severe Ankylosis as an Adaptation of Strong Bite Force**

Kelsey M. Jenkins<sup>1\*</sup>, Jack O. Shaw<sup>1</sup>

<sup>1</sup> Department of Geology and Geophysics, Yale University, New Haven, Connecticut, USA

Corresponding Author:

Kelsey M. Jenkins\*

Department of Geology and Geophysics, Yale University, New Haven, Connecticut, USA

Email address: kelsey.jenkins@yale.edu

## **Abstract**

Extant and extinct reptiles exhibit numerous combinations of tooth implantation and attachment. Tooth implantation ranges from those possessing roots and lying within a socket (thecodonty), to teeth lying against the lingual wall of the jawbone (pleurodonty), to teeth without roots or sockets that are attached to the apex of the marginal jawbones (acrodonty). Attachment may be ligamentous or via ankylosis (i.e., fusion). Adaptative reasonings are proposed as an underlying driver for evolutionary changes in some forms of tooth implantation and attachment. However, a substantiated adaptive hypothesis is lacking for the state of acrodonty via severe ankylosis that is seen in several lineages of Lepidosauria, a clade that is plesiomorphically pleurodont. We hypothesize that acrodonty via severe ankylosis, as seen in

*Sphenodon punctatus* and acrodontan lizards, is an adaptation either resulting from or allowing for a stronger bite force. We use bite force data gathered from the literature to show that those taxa possessing acrodont dentition attached via severe ankylosis possess a stronger bite force than those taxa with pleurodont dentition. Furthermore, we discuss the evolution of acrodonity and potential behaviors related to strong bite force that influenced the evolution of acrodonity within Rhynchocephalia, Acrodonta, and Trogonophidae. Implications for the evolution of different forms of tooth implantation and attachment include ancient behavioral shifts.

## Introduction

Acrodont tooth implantation, where the tooth rests at the summit of the tooth-bearing bone, evolved multiple times within Lepidosauria. It appears at least twice within squamate reptiles, as seen in acrodontan lizards (Romer, 1956) and trogonophid amphisbaenians (Gans, 1960), and once within Rhynchocephalia (Jenkins et al., 2017). In Acrodonta and *Sphenodon punctatus*, the only living representative of Rhynchocephalia, the dentition is severely ankylosed (i.e., fused) to the adjacent bone. In those taxa, teeth and surrounding tissues have been investigated thoroughly via histological studies (Cooper & Poole, 1973; Smirina & Ananjeva, 2007; Kieser et al., 2009, 2011; Haridy, 2018), CT data (Dosedělová et al., 2016), and developmental research (Buchtová et al., 2013). However, the evolution of acrodont tooth implantation is seldom discussed in an adaptive context.

Smith (1958) suggested that acrodonity and thecodonty (i.e., where the tooth is implanted in a socket) are traits associated with anchoring permanent dentition. However, the fossil record shows that some reptiles possessing acrodont dentition replaced their teeth regularly (Gow, 1977; Haridy, LeBlanc & Reisz, 2018). Furthermore, it is unknown if acrodonity and severe ankylosis are truly associated with anchoring permanent dentition, or if those combined traits somehow

inhibit tooth replacement. While it is possible that thecodonty evolved to anchor permanent dentition in mammals, it is well known that toothed archosaurs, which also possess thecodont implantation, replace their teeth with some regularity (e.g., Edmund, 1962; D’Emic et al., 2019).

There are a suite of traits commonly associated with acrodont tooth implantation. Reduced tooth counts, severe tooth wear, and a loss of the alveolar foramen are also commonly associated with acrodont tooth implantation (Augé, 1997; Zaher & Rieppel, 1999; Haridy, 2018). A slowing or lack of tooth replacement, called monophyodonty, is also commonly associated with acrodont tooth implantation (Smith, 1958; Cooper et al., 1970). Even with monophyodont dentition, additional teeth are typically still added to the posterior end of the tooth row throughout ontogeny, as is the plesiomorphic condition within Reptilia (Robinson, 1976; Gow, 1977; Rieppel, 1992; Reynoso, 2003). Severe ankylosis accompanies acrodonty in acrodontan squamates and in *Sphenodon punctatus*. Typically, ankylosis is severe in those taxa and the tooth and bone lose identity from one another (Fig. 1). Herein, we call those combined traits acrodonty via severe ankylosis (AVSA). AVSA caused some to erroneously propose that *S. punctatus* lacks teeth entirely, instead possessing a serrated jawbone (Mlot, 1997). On the contrary, severe wear may obscure the anterior dentition in older, acrodont, monophyodont lepidosaurs, causing them to appear to be absent.

The ancestral state of tooth implantation and attachment in the reptile line involved a tooth set in a shallow socket (i.e., subthecodonty) attached via ankylosis. However, reptiles have since explored all forms of tooth implantation (acrodonty, pleurodonty, and thecodonty) and attachment (ankylosis and gomphosis) in varying combinations (Bertin et al., 2018). Adaptive interpretations are frequently used to explain why reptiles may stray from the ancestral state. For example, dentine infoldings, called plicidentine, evolved independently multiple times within

Reptilia, and it is interpreted to be a mechanism to strengthen tooth attachment in kinetic-feeding predators (Maxwell, Caldwell & Lamoureux, 2011). Even the loss of teeth seems to be a response to the evolution of other adaptive structures, like the keratinous beak in birds and turtles (Davit-Béal, Tucker & Sire, 2009). The evolution of AVSA as seen in acrodontans and *Sphenodon punctatus* lacks any adaptive hypothesis. Here we suggest that AVSA is an adaptation associated with strong bite force.

Anecdotal evidence suggests that acrodont taxa possess a strong bite force: *S. punctatus* is said to possess a powerful, painful, and ‘vice-like’ bite (Robb, 1977; Daugherty and Cree, 1990), and one of the authors (KMJ) notes from personal experience that the veiled chameleon, *Chamaeleo calyptratus*, also possesses a painful bite and is reluctant to release its victim. Bite-force analyses also indicate agamids have a stronger bite than *S. punctatus*, relative to body size (Schaerlaeken et al., 2008; Jones and Lappin, 2009). The literature concerning bite force in lepidosaurs is numerous and implies a multitude of benefits for increased bite force. For instance, increase bite force is thought to improve prey capture and handling, particularly for the consumption of hard-bodied prey (Herrel et al., 1999; Herrel et al., 2001; Verwajen, Van Damme & Herrel, 2002; Meyers et al., 2018). High bite force may also aid in territory defense and dominance (Herrel, Meyers & Vanhooydonck, 2001; Lailvaux et al., 2004; Huyghe et al., 2005; Husak et al., 2006; Jones & Lappin, 2009), and mating success (Lappin & Husak, 2005; Husak et al., 2009; Herrel et al., 2010). Higher bite force in lizards is often accompanied by skeletal correlates in the cranium and increased mass of the adductor musculature compared to those with lower bite force (Herrel, McBrayer & Larson, 2007; Fabre et al., 2014).

We hypothesize that taxa possessing acrodont dentition also possess a higher bite force, compared to those with pleurodont dentition, relative to body size. Furthermore, we suggest that

AVSA as seen in Acrodonta and *Sphenodon punctatus* protects teeth from the risk of breakage during strong biting and allows for effective feeding in older individuals with severely worn dentition. In order to test our hypothesis, we compared bite force among lepidosaurian taxa. We found that size-normalized bite force was significantly greater in acrodont lepidosaurs than pleurodont lepidosaurs. Furthermore, we discuss the evolution of AVSA within an adaptive context in response to high bite force.

## Materials & Methods

To assess the relationship between bite force and tooth implantation, we collected data from thirty-eight peer-reviewed papers that record lepidosaurian bite force data (Supplementary Files). Publications that do not record metrics used in these statistical analyses and were excluded. The metrics recorded are the means of snout-vent length (SVL), head depth (HD), and bite force (BF). SVL, HD, and BF were all log-transformed and inspected for normality, following previous studies (Erickson et al., 2004; Wroe et al., 2005; Sellers et al., 2017).

We focused on the relationship between SVL and BF, as SVL is the most commonly reported measure of size in reptiles. However, many squamate reptiles possess elongate body forms that are not necessarily correlated to cranial allometry, and thus may not strongly correlate with bite force. Because of this, we also standardized by head depth in separate analyses.

Multiple studies evaluating lepidosaur bite force suggest that head depth is a good predictor of bite force because it accommodates the adductor muscles (Herrel, de Grauw & Lemos-Espinal, 2001; Lappin, Hamilton & Sullivan, 2006; McBrayer & Anderson, 2007; Herrel et al., 2010).

Tooth implantation was assessed by the authors.

To examine differences in bite force between acrodont and pleurodont taxa, analyses of covariance (ANCOVA) were performed using both size variables (SVL and HD) as covariates.

To further compare bite force across taxa of significantly different body masses, we calculated normalized bite force (NBF) as the residuals of a linear regression fit to (1) bite force and SVL or (2) bite force and HD. We refer to these values as SVL-NBF and HD-NBF, respectively. Differences in NBF between tooth implantation groups were then assessed using one-sided Kolmogorov–Smirnov (KS) and Mann-Whitney U (MWU) tests. Given unequal sample sizes, we also tested jackknife variants of the two tests (1000 replicates) which generated the same results.

## Results

Bite force is higher in acrodont taxa than in pleurodont taxa when size standardized (Fig. 2). Raw bite force values ranged from 0.3 to 409.3 Newtons. SVL ranged 13.0–389.0 mm, and HD ranged 1.2–55.5 mm. SVL NBF ranged -2.14 to 1.76, whereas HD NBF ranged -2.85 to 3.93 (Table 1). ANCOVAs indicate significant differences between tooth implantation type groups after accounting for SVL and HD (Table 2). According to both KS and MWU tests acrodont SVL-NBF and HD-NBF values were significantly greater than pleurodonts (Table 2).

Linear regressions of bite force, SVL, and HD, indicate that the three variables are significantly and positively correlated ( $p\text{-value} < 0.05$ ). Given strong, linear log-log relationships amongst the three values, data were log-transformed for all analyses. This relationship was stronger between bite force and SVL ( $p\text{-value} \ll 0.05$ , Adj R-square = 0.729), compared to bite force and HD ( $p\text{-value} \ll 0.05$ , Adj R-square = 0.337). Given this, we focus on the relationship between SVL and bite force for the remainder of this paper.

To evaluate the proportion of the lepidosaurian tree that has been examined in terms of bite force, we tallied all known publications to the best of our knowledge that record lepidosaurian bite force (Fig. 3; Supplementary Files). This includes those publications that were

not included in the initial analyses that compare bite force between acrodont and pleurodont taxa. Species of seventeen lepidosaurian families were found to be represented, including four acrodont families and 13 pleurodont families. Dactyloidae was represented by the most species ( $n = 49$ ), while the families of Sphenodontidae, Varanidae, and Trogonophidae were only represented by one species each ( $n = 1$ ).

## Discussion

Thus far, anatomical research related to biteforce in lepidosaurian reptiles has focused primarily on cranial musculature and skeletal dimensions, namely head depth, length, and width. However, teeth are more intimately associated with biting and oral processing than the aforementioned elements. Dental morphology is often adapted for diet, with generalists possessing a more generalized dentition and specialists possessing more unique tooth morphologies. For example, large bulbous molariform teeth are seen in some durophagous lizards (Estes and Williams, 1984). This is a stray from most iguanian and gekkotan lizards that possess simple columnar teeth with one to three small cusps. It should come as no surprise that tooth implantation and attachment is also shaped by oral processing capabilities. However, there is still the question of whether (1) AVSA developed due to strong bite force, or if (2) strong bite force evolved in response to AVSA.

In the first scenario, AVSA is a response to increased bite force. This concept must take into account the morphology of a typical pleurodont tooth versus a tooth that is influenced by AVSA. A typical pleurodont tooth, like that of geckos and non-acrodontan iguanians, is roughly cylindrical, and possesses a tooth base that is much less wide than it is tall. Unworn teeth in acrodontans and *Sphenodon punctatus* are comparatively much shorter and possess a wide tooth base that is strongly adhered to the jaw. Under higher bite forces and larger amounts of oral

processing, a pleurodont tooth would be more susceptible to breakage. While pleurodont teeth are regularly replaced, high rates of breakage would still be detrimental to the animal's feeding capability. An acrodon tooth with a wide base and firm attachment is better suited to withstand higher bite forces without breakage. However that is not to say that breakage could not occur in taxa possessing AVSA. The extremely strong adherence of teeth can result in the occasional breakage of both tooth and bone in those taxa (Dosedělová et al., 2016).

In the second scenario, strong bite force is a response to AVSA. Lepidosaurs possessing AVSA are monophyodont and exhibit severe wear, particularly in the anterior dentition as seen in older individuals. Although those individuals have extremely worn teeth, they still manage to capture and consume prey. This is particularly the case for *Sphenodon punctatus*, which is known to capture and saw small birds with their teeth using propalinal jaw movement (Cartland-Shaw et al., 1998). If dentition is severely worn due to a lack of replacement, increased bite force would be crucial in allowing the jaws to clamp tightly onto prey. Thus, older individuals with few functional teeth can still forage and consume as needed. If strong bite force in *S. punctatus* and Acrodonta evolved as a mechanism to aid in territory defense or increased mating success, an older animal may still be successful even though it possesses severely worn teeth. At this time, we cannot favor one hypothesis over another. It is also possible that different lineages acquired AVSA under either scenario.

Of the taxa that were examined in previous publications, fewer species of lepidosaurs with acrodon tooth implantation have been studied in regard to bite force in comparison to those with pleurodont implantation (Fig. 3). Of the pleurodont taxa, 49 species of *Anolis* lizards (Dactyloidae) were the subjects of 20 publications that record bite force alone. Those taxa make up the largest proportion of pleurodont taxa analyzed here. The large number of *Anolis*-based

studies is partly because they are speciose and represent a particularly important model taxon for ecological studies in the Americas. Conversely, only 16 unique species of acrodont lepidosaurs belonging to four separate families are the subject of 17 publications that record bite force. A larger sampling of acrodont taxa need to be examined in order to lend further support to the hypothesis that acrodont taxa possess a stronger bite force than pleurodont taxa. While this is impossible for *Sphenodon punctatus*, which is the only living representative of Sphenodontidae, a larger sampling of different species of acrodontan lizards and trogonophid amphisbaenians may be subjected to bite force analyses. Representatives of 16 families of squamate lizards have been subjected to bite force analyses, which leaves a large portion of the squamate line understudied. This is particularly the case for particularly snakes, for which we could find no bite-force research.

### Can AVSA be reversed?

The transition from pleurodont to acrodont tooth implantation occurred independently several times within Lepidosauria, but only in Acrodonta and Rhynchocephalia is the ankylosing so extreme that the tooth-bone boundary is difficult to detect upon initial inspection. Stem Acrodonta do not possess the extensive ankylosis and apical tooth implantation that is seen in the crown group, although the roots of the teeth are much shorter than most other iguanians (Simões et al., 2015). All crown acrodontans possess some degree of acrodont tooth implantation accompanied by severe ankylosis. Within that clade, AVSA may be a fixed trait that lacks the plasticity to explore other forms of tooth implantation and attachment. It is also possible that there has been no selective pressure acted upon tooth implantation and attachment within Acrodonta that would drive members of the clade away from AVSA since it initially evolved.

While this may imply a potentially adaptive circumstance to the initial evolution of this trait, it cannot be excluded that this trait may not longer act as an adaptation in extant Acrodonta.

AVSA is persistent within Rhynchocephalia, but several transitions in tooth implantation and attachment occurred from an initially acrodont state (Jenkins et al., 2017). *Ankylosphenodon pachyostosis* possesses ‘ankylotheodont’ dentition, in which the tooth has deeply implanted roots, but is nonetheless ankylosed to the surrounding bone (Reynoso, 2000). One genus, *Sapheosaurus*, potentially lacks marginal dentition, although it is unknown if this is due to extensive wear or if this taxon was truly edentulous (Cocude-Michel, 1963). The tooth plates seen in *Oenosaurus muelheimensis* also represent an interesting derivation from the typical tooth seen within Rhynchocephalia (Rauhut et al, 2012). Nonetheless, the tooth implantation seen in *O. muelheimensis* is acrodont, and tooth attachment is extensive. Although acrodonty is widespread within the clade, tooth implantation seems to be a more plastic trait within Rhynchocephalia than it is within Acrodonta.

### Acrodonty in Amphisbaenia

Trogonophidae do not possess the severe ankylosing associated with AVSA, as seen in Gans (1960) and Gans and Montero (2008). The relatively lower bite force seen in *T. wiegmanni* compared to other acrodont taxa seen in our results was also likely impacted by the fact that *T. wiegmanni* is an elongate, serpentine-like form. Because of that, using SVL to standardize our results may not be meaningful in the case of this taxon. However, the other taxa examined in this study are not impacted by extremely elongate body plans. When bite-force is standardized by head depth, the same trend of greater acrodont bite force is more apparent for *T. wiegmanni*.

Trogonophidae is the only clade within Amphisbaenia to evolve acrodont tooth implantation. However, other amphisbaenians possess teeth with roots of varying lengths.

Overall, amphisbaenians possess shorter roots than seen in most iguanians and geckos. Tooth implantation in amphisbaenians is often described as ‘subacrodont’ or ‘subpleurodont’ to denote the stray from the ‘typical’ pleurodont tooth implantation seen in most other squamates (Estes, 1975; Yatkola, 1976; Sullivan, 1985; Charig and Gans, 1990; Kearney, Maisano & Rowe, 2004; Gans and Montero, 2008; Longrich et al., 2015; Čerňanský, Klembara & Müller., 2016). The evolution of tooth implantation and attachment in Amphisbaenia has not been explored further, but the trend towards dentition with shorter roots is intriguing. Bite-force experiments conducted on amphisbaenians could address the evolution of acrodonty within the clade. Further research on this, and on the dental tissues of amphisbaenians, is certainly warranted to address the evolution of tooth implantation and attachment in Amphisbaenia.

## Conclusions

Acrodonty via severe ankylosis seen in *Sphenodon punctatus* and acrodontan squamates is likely an adaptation shaped by strong bite forces. The appearance of AVSA within those lineages, as well as extinct lineages, has implications for the early evolution of this trait. This may be informative of the early evolution of rhynchocephalian and acrodontan behavioral shifts, such as increased territoriality or dietary shifts. Within Amphisbaenia, a range of tooth implantation states exist, grading from pleurodonty to acrodonty. Further research on the dental changes taking place within this clade may suggest a new hypothesis for the evolution of acrodonty or lend support to the hypothesis presented here. While large portions of the squamate line have been subjected to bite force analyses, a much larger portion of the lineage has not, particularly snakes. Greater variation in bite force may exist among the clade, with subsequent implications for dental evolution in terms of tooth implantation, attachment, and morphology. A

251 greater understanding of the evolution of dentition, particularly tooth implantation and  
252 attachment, will inform us of changes in behavior in ancient reptiles lineages.

## 253 **Acknowledgements**

254 We thank D. Meyer and R. Armfield for providing helpful comments on an early version  
255 of this manuscript. Access to specimens was aided by G. Watkins-Colwell.

256

# References

- Augé, M. 1997. A consideration of the phylogenetic significance of acrodonty. *Herpetological Journal* 7:111-113.
- Bertin, T.J.C., B. Thivichon-Prince, A.R.H. LeBlanc, M.W. Caldwell, and L. Viriot. 2018. Current Perspectives on tooth implantation, attachment, and replacement in Amniota. *Frontiers of Physiology* 6:1-20 DOI: 10.3389/fphys.2018.01630
- Buchtová, M., O. Zahradníček, S. Balková, and A.S. Tucker. 2013. Odontogenesis in the veiled chameleon (*Chamaeleo calyptratus*). *Archives of Oral Biology* 58:118-133.
- Cartland-Shaw, L.K., A. Cree, C.M. Skeaff, N.M. Grimmond. 1998. Differences in dietary and plasma fatty acids between wild and captive populations of a rare reptile, the tuatara (*Sphenodon punctatus*). *Journal of Comparative Physiology B* 168:569-580.
- Čerňanský, A., J. Klembara, and J. Müller. 2016. The new rare record of the late Oligocene lizards and amphisbaenians from Germany and its impact on our knowledge of the European terminal Palaeogene. *Palaeobiodiversity and Palaeoenvironments* 96:559-587.
- Charig, A.J., and C. Gans. 1990. Two new amphisbaenians from the Lower Miocene of Kenya. *Bulletin of the British Museum of Natural History* 46:19-36.
- Cocude-Michel, M. 1963. Les rhynchocéphales et les sauriens des calcaires lithographiques (Jurassique Supérieur) d'Europe occidentale. *Nouvells Archives du Museum d'Histoire Naturelle de Lyon* 7:1-187.
- Cooper, J.S., and D.F.G Poole. 1973. The dentition and dental tissues of the agamid lizard, *Uromastix*. *Journal of Zoology* 169:85-100 DOI: 10.1111/j.1469-7998.1973.tb04654.x

- 278 Cooper, J.S., D.F.G. Poole, and R. Lawson. 1970. The dentition of agamid lizards with special  
279 reference to tooth replacement. *Journal of Zoology* 162:85-98 DOI: 10.1111/j.1469-  
280 7998.1970.tb01259.x
- 281 Daugherty, C.H., and A. Cree. 1990. Tuatara: a survivor from the dinosaur age. *New Zealand*  
282 *Geographic* 6:66-86.
- 283 Davit-Béal, T., A.S. Tucker, and J-Y. Sire. 2009. Loss of teeth and enamel in tetrapods: fossil  
284 record, genetic data and morphological adaptations. *Journal of Anatomy* 214:477-501  
285 DOI: 10.1111/j.1469-7580.2009.01060
- 286 D’Emic, M.D., P.M. O’Connor, T.R. Pascucci, J.N. Gavras, E. Mardakhayava, and E.K. Lund.  
287 2019. Evolution of high tooth replacement rates in theropod dinosaurs. *PLoS ONE*  
288 14:e0224734.
- 289 Dosedělová, H., K. Štěpánková, T. Zikmund, H. Lesot, J. Kaiser, K. Novotny, J. Štembírek, Z.  
290 Knotek, O. Zahradníček, and M. Buchtova. 2016. Age-related changes in the tooth-bone  
291 interface area of acrodont dentition in the chameleon. *Journal of Anatomy* 229:356-368  
292 DOI: 10.1111/joa.12490
- 293 Edmund, A.G., 1962. Sequence and rate of tooth replacement in the crocodilia. *Contributions to*  
294 *the Life Science Division of the Royal Ontario Museum* 52:1-42.
- 295 Erickson, G.M., A.K. Lappin, T. Parker, and K.A. Vliet. 2004. Comparison of bite-force  
296 performance between long-term captive and wild American alligators (*Alligator*  
297 *mississippiensis*). *Journal of Zoology* 262:21-28.
- 298 Estes, R. 1975. Lower vertebrates from the Fort Union Formation, Late Paleocene, Big Horn  
299 Basin, Wyoming. *Herpetologica* 31:365-385.

- 300 Estes, R., and E.E. Williams. 1984. Ontogenetic variation in the molariform teeth of lizards.  
301 *Journal of Vertebrate Paleontology* 4:96-107.
- 302 Fabre, A-C., D.V. Andrade, K. Huyghe, R. Cornette, and A. Herrel. 2014. Interrelationships  
303 between bones, muscles, and performance: biting in the lizard *Tupinambis merianae*.  
304 *Evolutionary Biology* 41:518-527.
- 305 Gans, C. 1960. Studies on amphisbaenids (Amphisbaenia, Reptilia): 1. A taxonomic revision of  
306 the Trogonophinae, and a functional interpretation of the amphisbaenid adaptive pattern.  
307 *Bulletin of the American Museum of Natural History* 119:129-204.
- 308 Gans, C., and R. Montero. 2008. An atlas of amphisbaenian skull anatomy. *Biology of the*  
309 *Reptilia* 21:621-738.
- 310 Gow, C.E. 1977. Tooth function and succession in the Triassic reptile *Procolophon trigoniceps*.  
311 *Palaeontology* 20:695-704.
- 312 Haridy, Y. 2018. Histological analysis of post-eruption tooth wear adaptations, and ontogenetic  
313 changes in tooth implantation in the acrodontan squamate *Pogona vitticeps*. *PeerJ*  
314 6:e5923 DOI: 10.7717/peerj.5923
- 315 Haridy, Y., A.R. LeBlanc, and R.R. Reisz. 2018. The Permian reptiles *Opisthodontosaurus*  
316 *carrolli*: a model for acrodont tooth replacement and dental ontogeny. *Journal of*  
317 *Anatomy* 232:372-382 DOI: 10.1111/joa.12754
- 318 Herrel, A., L. Spithoven, R. Van Damme, and F. De Vree. 1999. Sexual dimorphism of head size  
319 in *Gallotia galloti*: testing the niche divergence hypothesis by functional analyses.  
320 *Functional Ecology* 13:289-297.
- 321 Herrel, A., E. de Grauw, and J.A. Lemos-Espinal. 2001. Head shape and bite performance in  
322 xenosaurid lizards. *Journal of Experimental Zoology* 290:101-107.

323 Herrel, A., J.J. Meyers, and B. Vanhooydonck. 2001. Correlations between habitat use and body  
 324 shape in a phrynosomatid lizard (*Urosaurus ornatus*): a population-level analysis.  
 325 *Biological Journal of the Linnean Society* 74:305-314.

326 Herrel, A., L.D. McBrayer, and P.M. Larson. 2007. Functional basis for sexual differences in  
 327 bite force in the lizard *Anolis carolinensis*. *Biological Journal of the Linnean Society*  
 328 91:111-119.

329 Herrel, A., J.A. Moore, E.M. Bredeweg, and N.J. Nelson. 2010. Sexual dimorphism, body size,  
 330 bite force and male mating success in tuatara. *Biological Journal of the Linnean Society*  
 331 100:287-292.

332 Husak, J.F., A.L. Lappin, and R.A. Van Den Bussche. 2009. The fitness advantage of a high-  
 333 performance weapon. 2009. The fitness advantage of a high-performance weapon.  
 334 *Biological Journal of the Linnean Society* 96:840-845.

335 Herrel, A., R. Van Damme, B. Vanhooydonck, and F. De Vree. 2001. The implications of bite  
 336 performance for diet in two species of lacertid lizards. *Canadian Journal of Zoology*  
 337 79:662-670.

338 Herrel, A., J.A. Moore, E.M. Bredeweg, and N.J. Nelson. 2010. Sexual dimorphism, body size,  
 339 and bite force and male mating success in tuatara. *Biological Journal of the Linnean*  
 340 *Society* 100:287-292.

341 Husak, J.F., A.K. Lappin, S.F. Fox, and J.A. Lemos-Espinal. 2006. Bite force performance  
 342 predicts dominance in male venerable collared lizards (*Crotaphytus antiquus*). *Copeia*  
 343 2006:301-306.

344 Huyghe, K., B. VanHooydonck, H. Sheers, M. Molina-Borja, and R. Van Damme. 2005.  
 345 Morphology, performance and fighting capacity in male lizards, *Gallotia galloti*.  
 346 *Functional Ecology* 19:800-807.

347 Jenkins, K.M., M.E.H. Jones, T. Zikmund, A. Boyde, and J.D. Daza. 2017. A review of tooth  
 348 implantation among rhynchocephalians (Lepidosauria). *Journal of Herpetology* 51:300-  
 349 306 DOI: 10.1670/16-146

350 Jones, M.E.H., and K.A. Lappin. 2009. Bite-force performance of the last rhynchocephalian  
 351 (Lepidosauria: *Sphenodon*). *Journal of the Royal Society of New Zealand* 39:71-83.

352 Kearney, M., J.A. Maisano, and T. Rowe. 2004. Cranial anatomy of the extinct amphisbaenian  
 353 *Rhinerua hatcherii* (Squamata, Amphisbaenia) based on high-resolution x-ray computed  
 354 tomography. *Journal of Morphology* 264:1-33.

355 Kieser, J.A., T. Tkatchenko, M.C. Dean, M.E.H. Jones, W. Duncan, and N.J. Nelson. 2009.  
 356 Microstructure of dental hard tissues and bone in the tuatara dentary, *Sphenodon*  
 357 *punctatus* (Diapsida: Lepidosauria: Rhynchocephalia). *Comparative Dental Morphology*  
 358 13:80-85.

359 Kieser, J.A., L-H. He, M.C. Dean, M.E.H. Jones, W.J. Duncan, M.V. Swain, and N.J. Nelson.  
 360 2011. Structure and compositional characteristics of caniniform dental enamel in the  
 361 tuatara *Sphenodon punctatus* (Lepidosauria: Rhynchocephalia). *New Zealand Dental*  
 362 *Journal* 107:44-50.

363 Lailvaux, S.P., A. Herrel, B. VanHooydonck, J.J. Meyers, and D.J. Irschick. 2004. Performance  
 364 capacity, fighting tactics and the evolution of life-stage male morphs in the green anole  
 365 lizard (*Anolis carolinensis*). *Proceedings of the Royal Society of London Series B*  
 366 271:2501-2508 DOI: 10.1098/rspb.2004.2891

- 367 Lappin, A.K., and J.F. Husak. 2005. Weapon performance, not size, determines mating success  
368 and potential reproductive output in the Collared Lizard (*Crotaphytus collaris*). *The*  
369 *American Naturalist* 166:426-436.
- 370 Lappin, A.K., P.S. Hamilton, and B.K. Sullivan. 2006. Bite-force performance and head shape in  
371 a sexually dimorphic crevice-dwelling lizard, the common chuckwalla [*Sauromalus ater*  
372 (=obesus)]. *Biological Journal of the Linnean Society* 88:215-222.
- 373 Longrich, N.R., J. Vinther, R.A. Pyron, D. Pisani, and J.A. Gauthier. 2015. Biogeography of  
374 worm lizards (Amphisbaenia) driven by end-Cretaceous mass extinction. *Proceedings of*  
375 *the Royal Society B* 282:20143034.
- 376 Maxwell, E.E., M.W. Caldwell, and D.O. Lamoureux. 2011. The structure and phylogenetic  
377 distribution of amniote plicidentine. *Journal of Vertebrate Paleontology* 31:553-561.
- 378 McBrayer, L.D., and R.A. Anderson. 2007. Sexual size dimorphisms and bite force in the  
379 Norther Alligator Lizard, *Elgaria coerulea* 41:554-559.
- 380 Meyers, J.J., K.C. Nishikawa, and A. Herrel. 2018. The evolution of bite force in horned lizards:  
381 the influence of dietary specialization. *Journal of Anatomy* 232:214-226.
- 382 Mlot, C. 1997. Return of the tuatara. *Science News* 152:300-301.
- 383 Rauhut, O.W.M., A.M. Heyng, A. López-Arbarello, and A. Hecker. 2012. A new  
384 rhynchocephalian from the Late Jurassic of Germany with a dentition that is unique  
385 amongst tetrapods. *PLoS ONE* 7:e46839.
- 386 Reynoso, V.H. 2000. An unusual sphenodontian (Reptilia: Diapsida) from the Tlayua Formation  
387 (Albian), Central Mexico. *Journal of Paleontology* 74:133-148.

Reynoso, V.H. 2003. Growth patterns and ontogenetic variation of the teeth and jaws of the Middle Jurassic sphenodontian *Cynosphenodon huizachalensis* (Reptilia: Rhynchocephalia). *Canadian Journal of Earth Science* 40:609-619.

Rieppel, O. 1992. The skull in a hatchling of *Sphenodon punctatus*. *Journal of Herpetology* 26:80-84.

Robb, J. 1977. *The Tuatara*. New Zealand: Meadowfield.

Robinson, P.L. 1976. How *Sphenodon* and *Uromastyx* grow their teeth and use them. *Morphology and Biology of Reptiles* 3:43-65.

Romer, A.S. 1956. *Osteology of the Reptiles*. Chicago: University of Chicago Press.

Schaerlaeken, V., A. Herrel, P. Aerts, and C.F. Ross. 2008. The functional significance of the lower temporal bar in *Sphenodon punctatus*. *The Journal of Experimental Biology* 211:3908-3914.

Sellers, K.C., K.M. Middleton, J.L. Davis, and C.M. Holliday. 2017. Ontogeny of bite force in a validated biomechanical model of the American alligator. *Journal of Experimental Biology* 220:2036-2046.

Simões, T.R., E. Wilner, M.W. Caldwell, L.C. Weinschütz, and A.W.A. Kellner. 2015. A stem acrodontan lizard in the Cretaceous of Brazil revises early lizard evolution in Gondwana. *Nature Communications* 6:8149.

Smirina, E.M., and N.B. Ananjeva. 2007. Growth layers in bones and acrodont teeth of the agamid lizard *Laudakia stoliczkana*. *Amphibia-Reptilia* 28:193-204.

Smith, H.M. 1958. Evolutionary lines in tooth attachment and replacement in reptiles: their possible significance in mammalian dentition. *Transactions of the Kansas Academy of Science* 61:216-225.

411 Sullivan, R.M. 1985. A new middle Paleocene (Torrejonian) rhineurid amphisbaenian  
 412 amphisbaenian, *Plesiorhineura tsentasi* new genus new species, from the San Juan Basin,  
 413 New Mexico. *Journal of Paleontology* 59: 1481-1485.

414 Verwaijen, D., R. Van Damme, and A. Herrel. 2002. Relationships between head size, bite force,  
 415 and prey handling efficiency and diet in two sympatric lacertid lizards. *Functional*  
 416 *Ecology* 16:842-850.

417 Wroe, S., C. McHenry, and J. Thomason. 2005. Bite club: comparative bite force in big biting  
 418 mammals and the prediction of predatory behaviour in fossil taxa. *Proceedings of the*  
 419 *Royal Society B* 272:619-625.

420 Yatkola, D.A. 1976. Mid-Miocene lizards from western Nebraska. *Copeia* 1976:645-654.

421 Zaher, H., and O. Rieppel. 1999. Tooth implantation and replacement in squamates, with special  
 422 reference to mosasaur lizards and snakes. *American Museum Novitates* 3271:1-19.

423

424 **Figure Legend**

425 **Figure 1: Acrodont tooth implantation via severe ankylosis as seen in two acrodontan**

426 **lizards (A & B) and a rhynchocephalian (C).** A) the chameleon *Fercifer oustaleti* YPM R

427 1214, B) the agamid *Agama agama* YPM R 17936, and C) the rhynchocephalian *Sphenodon*

428 *punctatus* YPM R 10647. D) Pleurodont tooth implantation as seen in *Ctenosaura* sp. YPM R

429 11060.

430 **Figure 2: Acrodont vs. Pleurodont bite force.** [Left panel] Log-transformed SVL and HD

431 (mm) versus log bite force (N). Grey line represents a linear regression. [Right panel] SVL- and

432 HD-normalized bite force distributions.


433 **Figure 3: Number of species analyzed for bite force by family, colored by tooth**

434 **implantation.**

# Figure 1

Acrodont tooth implantation via severe ankylosis as seen in two acrodontan lizards (A & B) and a rhynchocephalian (C).


A) the chameleon *Fercifer oustaleti* YPM R 1214, B) the agamid *Agama agama* YPM R 17936, and C) the rhynchocephalian *Sphenodon punctatus* YPM R 10647. D) Pleurodont tooth implantation as seen in *Ctenosaura* sp. YPM R 11060.



# Figure 2


Acrodont vs. Pleurodont bite force.

[Left panel] Log-transformed SVL and HD (mm) versus log bite force (N). Grey line represents a linear regression. [Right panel] SVL- and HD-normalized bite force distributions.



# Figure 3

Number of species analyzed for bite force by family, colored by tooth implantation.



**Table 1** (on next page)

Summary statistics. SVL = snout-vent length. HD = head depth. NBF = normalized bite force

Table

1:

Summ

ary

statisti

|            | SVL (mm) |        |        |       | HD (mm) |       |       |      | Raw bite force (N) |        |       |       | SVL NBF |      |       |      | HD NBF |      |       |      |
|------------|----------|--------|--------|-------|---------|-------|-------|------|--------------------|--------|-------|-------|---------|------|-------|------|--------|------|-------|------|
|            | Min.     | Max.   | Mean   | SD    | Min.    | Max.  | Mean  | SD   | Min.               | Max.   | Mean  | SD    | Min.    | Max. | Mean  | SD   | Min.   | Max. | Mean  | SD   |
| All        | 13.00    | 389.00 | 100.91 | 71.15 | 1.22    | 55.50 | 9.93  | 7.12 | 0.30               | 409.35 | 34.46 | 65.21 | -2.17   | 1.76 | 0.00  | 0.72 | -2.85  | 3.93 | 0.00  | 1.03 |
| Acrodont   | 45.34    | 250.00 | 125.22 | 71.33 | 4.00    | 33.40 | 12.19 | 8.96 | 5.08               | 238.30 | 52.20 | 65.40 | -2.17   | 1.30 | 0.26  | 1.00 | -0.38  | 1.28 | 0.51  | 0.51 |
| Pleurodont | 13.00    | 389.00 | 97.27  | 70.65 | 1.22    | 55.50 | 9.67  | 6.87 | 0.30               | 409.35 | 31.81 | 64.99 | -1.52   | 1.76 | -0.04 | 0.67 | -2.85  | 3.93 | -0.06 | 1.06 |

cs. SVL = snout-vent length. HD = head depth. NBF = normalized bite force



## **Table 2**(on next page)

Summary of statistical test results describing differences between bite force of taxa with different tooth implantation.

ANCOVAs tested covariation and raw bite force values. KS and MWU tests compared distributions of NBF values.

1

2 Table 2: Summary of statistical test results describing differences between bite force of taxa with  
3 different tooth implantation. ANCOVAs tested covariation and raw bite force values. KS and  
4 MWU tests compared distributions of NBF values.

| Variable               | Test   | Test statistic | P-value  | Jackknife<br>mean p-value<br>(10000<br>replicates) |
|------------------------|--------|----------------|----------|----------------------------------------------------|
| SVL (logged)           | ANCOVA | 4.32           | 0.041    |                                                    |
|                        | KS     | 0.38258        | 0.003748 | 0.0483                                             |
|                        | MWU    | 2160           | 0.003064 | 0.0313                                             |
| Head depth<br>(logged) | ANCOVA | 4.21           | 0.042    |                                                    |
|                        | KS     | 0.4803         | 0.002    | 0.0344                                             |
|                        | MWU    | 1461           | 0.001303 | 0.0246                                             |

5