

Nectar robbing in the trainbearers (*Lesbia*, Trochilidae)

Boris Igić^{1,2}, Ivory Nguyen¹ and Phillip B. Fenberg^{3,4}

¹ Department of Biological Sciences, University of Illinois at Chicago, Chicago, IL, USA

² Botany Department, The Field Museum of Natural History, Chicago, IL, USA

³ School of Ocean and Earth Sciences, National Oceanography Centre Southampton, University of Southampton, Southampton, UK

⁴ Department of Life Sciences, Natural History Museum, London, UK

ABSTRACT

Many flower visitors engage in floral larceny, a suite of so-called 'illegitimate' visits in which foragers take nectar without providing pollination services. The data on prevalence of illegitimate visits among hummingbirds, as well as the total proportion of foraging and diet that such visits comprise is broadly lacking. Here, we report the occurrence of nectar larceny in the two currently recognized species of trainbearers and analyze the proportion of plant visits categorized by mode of interaction as: robbing, theft, and/or pollination. We augment our original field observations using a trove of data from citizen science databases. Although it is difficult to distinguish primary vs. secondary robbing and theft vs. pollination, we conservatively estimate that ca. 40% of the recorded nectar foraging visits involve nectar robbing. Males appear to engage in robbing marginally more than females, but further studies are necessary to confidently examine the multi-way interactions among sex, species, mode of visitation, and other factors. Our results also indicate that the suggested relationship between serrations on bill tomia and traits such as nectar robbing or territorial defense may be complicated. We discuss the significance of these findings in the context of recent developments in study of nectar foraging, larceny, and pollination from both avian and plant perspectives.

Submitted 3 March 2020
Accepted 25 June 2020
Published 12 August 2020

Corresponding author
Boris Igić, boris@uic.edu

Academic editor
Ann Hedrick

Additional Information and
Declarations can be found on
page 11

DOI 10.7717/peerj.9561

© Copyright
2020 Igić et al.

Distributed under
Creative Commons CC-BY 4.0

Subjects Animal Behavior, Biodiversity, Ecology, Plant Science, Zoology

Keywords Nectar robbing, Hummingbirds, Feeding behavior, Floral larceny, Pollination, Pollination biology, Trainbearers, Plant defense

INTRODUCTION

A growing list of bird species across several clades are known to forage for nectar on so-called "illegitimate" flower visits, in which flower rewards are taken without the requisite provision of pollination services. Also termed 'floral larceny,' this mode of foraging is correspondingly gaining a broader appreciation as an important factor shaping the ecology and evolution of plant-animal interactions (*Lara & Ornelas, 2001; Irwin et al., 2010; Rojas-Nossa, Sánchez & Navarro, 2016; Boehm, 2018*). Although species such as flower piercers (*Diglossa*, Passeriformes) are widely known to depend on nectar larceny, there are many reports of illegitimate visits by hummingbirds (*Lara & Ornelas, 2001; Gonzalez & Loiselle, 2016*). Some morphological characteristics of hummingbirds,

OPEN ACCESS

including bill length and tomial serrations, are thought to be particularly closely associated with nectar larceny (Ornelas, 1994, but see Rico-Guevara et al., 2019).

Nearly all plants that provide nectar and pollen rewards experience larceny, most often from insects and vertebrates, and many defend against it (Irwin et al., 2010). The remarkable frequency of illegitimate visits necessitated the development and adoption of a more precise lexicon of larceny (Inouye, 1980), which attempts to separate it into canonical modes, partly in service of conceptual clarification useful in pursuit of identifying its ecological and evolutionary causes and consequences. Thus, “primary nectar robbers” mechanically create a hole in flower tissue through which they remove nectar, bypassing the floral opening. By contrast, “secondary nectar robbers” remove nectar utilizing openings previously fashioned by primary robbers. Deceptively resembling pollinators, “nectar thieves” also access nectar through the floral opening. Due to a mismatch between flower and thief morphology or behavior, however, pollination does not take place. Finally, some plant species are vulnerable to larceny by “base workers”, visitors who probe along the base of the flower (between the free petals) and obtain nectar, while still failing to effect pollination.

These modes of illegitimate nectaring visits may be difficult to distinguish from each other and/or legitimate visits—pollination—without careful observations and manipulations. Moreover, they can clearly quantitatively overlap so that, for example, thieving may merely reduce pollination efficiency without eliminating it, or individual birds can engage in a mix of primary and secondary robbing. While many studies report one or more modes of nectar foraging among hummingbirds (McDade & Kinsman, 1980; Roubik, Holbrook & Parra, 1985; Feinsinger et al., 1987; Ornelas, 1994), we lack assessments of their relative importance (Snow & Snow, 1980), which are critical for a more complete understanding of the incentives and behaviors that drive the observed patterns or plant–animal interactions.

Here, we present original observations and collect data from public photograph databases with a focus on documenting the frequency of modes of nectar foraging by Green- and Black-tailed Trainbearers (*Lesbia nuna* and *L. victoriae*). We discuss the significance of common occurrences of nectar robbing in the context of plant-pollinator co-evolution and bill morphology. The original observations took place in and around Ollantaytambo, Peru, on *Brugmansia sanguinea*, *Fuchsia boliviensis* and *Passiflora tripartita*. We coarsely document the proportion of pollination, theft and primary- and secondary-robbing visits to flowers, by leveraging images from citizen-science databases eBird, iNaturalist, and other photographed/reported occurrences in the literature. Our highly preliminary analyses find that floral larceny is surprisingly common and dominated by robbing (at least ca. 40%). Specifically, if secondary robbing is common among hummingbirds, as we suggest, assessment of the proposed associations between tomial serrations and mode of nectar foraging (Ornelas, 1994; Rico-Guevara et al., 2019) could be difficult.

MATERIALS AND METHODS

Species and site description

The genus *Lesbia* is currently comprised of two short-billed hummingbird species, Green-tailed Trainbearer, *Lesbia nuna* (Lesson, 1832), and Black-tailed Trainbearer,

Lesbia victoriae (Bourcier & Mulsant, 1846). Their geographic distributions largely overlap, ranging from Colombia to Peru in montane scrub and other semi-open habitats. The species are differentiated by color, specifically the extent of emerald green on dorsal side and tail (more in *L. nuna*). Black-tailed Trainbearers are generally larger, as well. Both species are strongly sexually dimorphic, with males in possession of a spectacularly long tail. The species are difficult to differentiate and confidently assign because of a great amount of geographic variation within each species (many subspecies names exist) and possible existence of undescribed species (*Weller & Schuchmann, 2004; Stiles, 2004*). These difficulties are especially apparent in attempts to identify individuals from single photographs, and we consequently conducted our analyses (see below) at the generic level.

Description of original observation

On 14 January 2019, we recorded and photographed a male Green-tailed Trainbearer, *Lesbia nuna*, which nectar-robbed several flowers of *Fuchsia boliviensis* (Onagraceae) in Ollantaytambo, Peru (13°15'44"S, 72°16'14"W), and then followed the same individual on a 50 m foraging bout, while it nectar-robbed flowers of *Brugmansia sanguinea* (Solanaceae)—a species with spectacular ca. 30 cm-long tubular crimson-to-yellow flowers—and then legitimately visiting *Salvia leucantha* (Lamiaceae). Both rusty and black-throated flowerpiercer (*Diglossa sittoides* and *D. brunneiventris*) occur at relatively high densities in the area (B. Igić, P.B. Fenberg, 2019, personal observations), are known to pierce both *Fuchsia* and *Brugmansia* species, and may have been responsible for the previously made calyx and petal hole at the base of *B. sanguinea* flower.

Literature searches

We examined the literature and internet resources to accumulate a database of additional instances of nectar collecting visits by *Lesbia* species. First, we used internet and Google Scholar exact keyword searches of the existing scientific literature, including: “*Datura*” or “*Brugmansia*” and “*Lesbia*” or “trainbearer”; “*Lesbia*” and “diet”. We employed “advanced search” filtering, screening out a variety of ribald nouns, to reduce the considerable volume of images with unrelated or offensive content. Documentation of any details on foraging ecology of the trainbearers is sparse. We found three sources either documenting or hinting at illegitimate flower visits by trainbearers. *Gould (1861, p. 15)* cited in *Ornelas (1994)*, contains a passing reference to piercing of a *Brugmansia sp.* flower by a *Lesbia sp.*, but the visit was originally inferred as insect gleanings, not nectar robbing, as later re-framed by *Ornelas (1994)*, and neither the species nor locations were identified. *Snow & Snow (1980)* documented feeding behavior for a species identified as *Lesbia nuna* in the vicinity of Bogotá, Colombia. The observed individuals—nearly all females—visited six plant species, and were thought to either utilize holes in corolla or probe around the base of the corolla split in *Castilleja fissifolia*. In a paper documenting secondary robbing behavior in Cinereous Conebill, *Vogt (2006)* mentions unpublished observations of secondary nectar robbing by a trainbearer. Additionally, although we could find no papers detailing their insect foraging in any detail, 21.6% of *Lesbia nuna* feeding records comprised of insect hawking (*Snow & Snow, 1980*), and gut contents of a single individual

included arthropods (Remsen, Stiles & Scott, 1986). Therefore, as generally holds for hummingbirds, the trainbearers' diet at a minimum contains arthropods and nectar. The proportion of time spent on each mode of nectaring is not known.

Foraging mode collection and scoring procedures

Consequently, we examined all identified species records for the Genus *Lesbia* in two public databases—iNaturalist (iNaturalist, 2019) and eBird (Sullivan et al., 2009), and in online searches, primarily of the photo service flickr (<https://www.flickr.com>). In each case, we made last visits or downloaded database records on 6 January 2020. For all resulting records, we applied the same methodology. Accessions, metadata, image locations and scored data used in our analyses are available in [Supplemental Materials](#). Those without any flowers were discarded, as were those in which birds were simply perched or flying near a flower. We then closely examined the remaining images, in which birds were hovering in close proximity (about a body length) to flowers, and facing them. For that subset of 180 records, we extracted the location, date, and comment metadata. We recorded individual sex (“m”, “f”, or “NA”), species identification (scoring specific epithets as “nuna”, “victoriae”, or “sp.”), estimated mode of visitation (primary nectar-robbing, secondary nectar-robbing, thieving, pollination, or “NA”), as well as our confidence in the estimate of the mode of interaction (“low” vs. “high”; to help facilitate critical re-examinations of our results). To ensure consistent scoring of nectar foraging mode, both B.I. and P.B.F. examined all 180 records and jointly agreed on marked scoring. Finally, with the help of several colleagues, we identified and recorded many (but not all) of at least 50 visited plant species.

Often, modes of visitation are unclear and we combined terms for the most accurate characterization of visitation. For example, primary vs. secondary nectar robbing and pollination vs. thieving are generally hard to distinguish, so we simply combined the likely modes of visitation (there were only two instances of clear thieving). This enables us to conservatively summarize modes of interactions as approximate ranges, without making unnecessary and possibly incorrect assumptions. We should note here that nearly all of the combined pollination/thieving visits appear to have been legitimate visits, but establishing the extent of pollen transfer is prohibitively demanding for most studies of modes of nectar feeding. Similarly, we recorded the species identifications, but found them to be uncertain for a number of individuals, and we consequently highlight the jointly presented feeding habits for all of *Lesbia*. Published (uploaded) observations were not a priori assigned or randomized in any way. Due to these limitations we simply report a number of pooled summary statistics. Moreover, we find it unlikely that they suffer of a strong bias with respect to visit mode, because the mode of visit is often unclear to a photographer until after the photo is taken. Instead, we initially found it more likely that there would be a photography bias in favor of photographing males, with their outstanding long tails (68 out of 114 records with foraging mode; exact binomial test, $p < 0.03$). We examine associations and interactions with sex in part due to this possibility. As pooling over the ignored variable(s) can be misleading, so we also perform limited three-way visual analyses, but caution against their over-interpretation. We coarsely

examine associations between species ID (*nuna/victoriae*), sex (f/m), and binarized mode of interaction with flowers (robbing/other), examining the null hypothesis that the three traits are mutually independent. For model selection, we used log-linear models implemented in R package MASS (*R Core Team, 2019*), and AIC scores from LRstats. The nature of our data collection and resulting exploratory analyses suggests that associations ought to be re-tested by obtaining a new set of data in advance of further testing.

Bill morphology

Short-billed hummingbirds and those with serrated tomlia are thought to be more likely to nectar-rob flowers with long corolla tubes, co-adapted with long-billed hummingbirds (*Lara & Ornelas, 2001*). We did not find a reference to presence/absence of serrated tomlia in trainbearers, so we scored this trait in 17 specimens of *Lesbia nuna* and four specimens of *Lesbia victoriae* at the Field Museum of Natural History. We used a dissecting microscope to visualize and score this trait on the available collections, as well as measure their bill lengths (tip to operculum; all performed by I.N.).

RESULTS

Field observation of robbing

In our field observations, we recorded a single adult male foraging bout, during which this individual visited flowers of three plant species. Approximately ten flowers of *Fuchsia boliviensis* were robbed (unclear if primary or secondary robbing took place) as well as two flowers of *Brugmansia sanguinea*, while five flowers of *Salvia boliviensis* were apparently legitimately visited (ruling out or confirming pollination success requires controlled experiments). Photographs collected in the course of observations were deposited in iNaturalist (<https://www.inaturalist.org/observations/37206000>; *iNaturalist, 2019*).

Database collection

We collected and examined more than 1,550 photographs of *Lesbia* and found floral visits for 180 foraging bouts (*Fig. 1; Supplemental Data Table*) on at least 42 flowering plant genera, identified thus far. It is unlikely that many of these instances are replicates, recording the same individual, although this is somewhat uncertain for the vicinity of Cerro de Monserrate in Bogotá, Colombia, with about a dozen documented foraging bouts.

We assigned the mode of visitation for 135 visits with some confidence, while the mode remains unknown for 45 visits, most often because pictured birds were hovering in the vicinity of a flower, immediately before or after nectaring (*Table 1*). The most common recorded mode was “pollination/thieving” (46 visits), a category that we scored conservatively (vaguely) because it is clearly continuous, virtually impossible to confidently score separately without functional study. Ideally, one would examine pollen transfer or effected seed set, after bagging and marking flowers. Our impression, however, is that the majority of these visits at least plausibly resulted in pollen transfer. The second most common mode was combined “nectar robbing” (primary and secondary robbing; 41 visits) indicating that an individual fed through a pierced hole in the side of the corolla.




Figure 1 (A) A nectar robbing visit by a black-tailed trainbearer *Lesbia victoriae* on *Brugmansia sanguinea*, whose flower was previously pierced by a species of *Diglossa* (photo by Diego Emerson Torres; used with permission); (B) bar plot illustrating the relative visit mode frequencies of *Lesbia* species. Of the total 135 visits whose mode is known (could be assigned), 60.7% were legitimate (combining pollination and pollination/thieving) and 39.3% illegitimate (combining all robbing and thieving). [Full-size !\[\]\(1679558f37f6db0dd8360a2a7e913e90_img.jpg\) DOI: 10.7717/peerj.9561/fig-1](https://doi.org/10.7717/peerj.9561/fig-1)

It was followed by “pollination” (36 visits), which indicated a fairly clear legitimate interaction, “secondary nectar robbing” (10 visits) and thieving (two visits). Although we could not confidently assign any visits to the “primary robbing” mode, this is entirely due to the one key limitation of our observational approach, reliance on scoring from photographs. If we conservatively lump all “pollination/thieving” and “pollination” observations as legitimate, such visits comprise approximately 60.7% of the total and illegitimate visits 39.3% (combining all robbing and thieving).

With a number of limitations in mind, we present a few preliminary analyses in an attempt to estimate the association of species ID, sex, and mode of visitation. When these factors are jointly considered in a three-way contingency table, we find that three variables are likely not mutually independent (AIC = 56.43, LR $\chi^2 = 13.55$, d.f. = 4, $p < 0.01$). Non-robbing visits by *Lesbia nuna* females and robbing visits by *Lesbia victoriae* males have the highest residuals. The lowest AIC model posits conditional independence (mode is independent of species, given sex; AIC = 48.78, LR $\chi^2 = 1.91$, d.f. = 2, $p = 0.39$), indicating

Table 1 Cross tabulation data of nectar foraging mode, sex, and species identity for *Lesbia* spp. “Other” denotes lumped pollination and thieving, and “Robbing” combines both modes of robbery. Sex identification relied on tail length and subtle color differences as a proxy, and species identification was in nearly all cases copied from the original source. Of 180 total foraging bouts captured, we assigned visit mode to 135, and each of visit (nectar foraging) mode, sex and species to 114 shown in the table.

		Other	Robbing
<i>L. nuna</i>	Female	18	5
	Male	12	7
<i>L. victoriae</i>	Female	16	7
	Male	23	26

that sex differences are greater than species differences. One broad interpretation is that the females generally rob less than the (territorial) males. We emphasize that this result is preliminary, and the associations may stem from a number of data limitations, especially confounding unmeasured variables, or those not considered a priori. For example, more *Lesbia victoriae* males were observed in urban parks where plant communities and robbing frequency may differ, compared to rural or natural areas. Moreover, a model with all two-way interactions, but without the three-way interaction of the saturated model, provides only slightly worse fit (δ AIC = 0.15). It is therefore presently unclear whether species or sex differences are more important in predicting nectar foraging modes.

Bill morphology

We examined 17 specimens of *Lesbia nuna* and found no serrated tomia. The imbalance of sex ratio (15 males) is concerning, but perhaps unimportant in this case, because serrations are more likely to be present in males (*Rico-Guevara et al., 2019*). The mean bill length for males was 15.9 ± 0.7 mm, and for females 14.2 ± 3.2 mm. Similarly, we found no tomial serrations in four specimens (three male, one female) of *Lesbia victoriae*. Their mean bill length for males was 14.7 ± 0.4 mm, and one female measured 11.0 mm.

DISCUSSION

Hummingbirds are generally thought to serve as legitimate flower visitors, and effect pollination in exchange for host plant nectar. There is, however, an increasing appreciation of the fact that nectar obtained during illegitimate visits—including both primary and secondary robbing, as well as thieving—may comprise a substantial portion of diets of many species (*Lara & Ornelas, 2001; Irwin et al., 2010; Boehm, 2018*). Presently, the frequency distribution of nectar larceny within and among bird individuals and species, and a variety of covariates (community composition, territoriality, seasonality, geography, etc.) is unclear. Most reports of hummingbird nectar foraging document the presence (or imply absence) of nectar robbing, and do not provide an estimate of the proportion of visits dedicated to illegitimate visits. This is in large part due to the difficulty of accruing this kind of data and, when larceny occurs, separating its modes.

Our study originated with a simple natural history observation of robbing by a trainbearer, which we augmented using a trove of citizen-science data from crowd-sourced

public databases. We believe that such estimates are a first step in improving our understanding of the dynamics between plants and pollinators, especially why and how (e.g., primary vs. secondary robbing) some individual birds or species steal more than others, how larceny affects plants, and how they respond to larceny. Below, we discuss the results and their immediate implications of this small-scale study and place them in the context of the literature on ecology and evolution of plant–hummingbird interactions.

We find that both currently described species of *Lesbia* steal nectar, mostly by robbing. Unambiguous illegitimate visitation constitutes a substantial proportion, around 40% of all nectar foraging visits. Although [Snow & Snow \(1980\)](#) surveyed a relatively small area, as a part of a study with a far greater scope, they found that robbing comprises 64% of all visits by *Lesbia nuna* in the northern Andes, and all illegitimate visits occurred on one of five visited plant species (*Castilleja fissifolia*, Orobanchaceae). Observations of nine species at this high-elevation site found that five species visited flowers illegitimately, and robbing constituted 17% of all flower visits (range 0–64%). Plant-side studies, which estimate the proportion of flowers robbed on a single plant species, uncover a wide range of robbery rates due to visiting hummingbirds. For example, [Maloof & Inouye \(2000\)](#) reviewed reports of visitation to eight plant species (by a variety of animal species), and found robbing ranged from 11% to 89%. Similarly, [Del Coro Arizmendi \(2001\)](#) found that total robbing in four highland Mexican species ranges from 9% to 58%, and [Navarro \(2001\)](#) found that 75% of *Macleania bullata* (Ericaceae) flowers contained pierced holes consistent with primary robbing. An unpublished report (D. Boose, cited in [Maloof & Inouye, 2000](#)) found that the Stripe-tailed Hummingbird (*Eupherusa exima*) robs 98% of *Razisea spicata* (Acanthaceae) flowers, but the mode is not stated. In each of these plant-side studies, however, the proportion of hummingbirds' visits comprised of each visit mode is unknown, and in some cases the primary robber is not a hummingbird.

Despite its critical importance for understanding the animal-side foraging incentives and plant-side pollination and defense strategies, primary and secondary robbing are difficult to disentangle ([Vogt, 2006](#); [Boehm, 2018](#)). [Snow & Snow \(1980\)](#) is the only hummingbird study we found, which lists data that enables a calculation of the relative frequencies of visit modes, including primary vs. secondary robbing. While we may have missed existing records, it appears that such estimates are rare at best. Confidently scoring the specific mode of robbing (primary vs. secondary) demands evidence bearing on the question whether “breaking-and-entering” took place, which in turn requires careful experimental data collection. For example, capturing the act of piercing of flowers or documenting entry through a pre-existing hole could enable unambiguous scoring of robbing mode. We find that secondary nectar robbing is relatively common, comprising between 7.4% and 37.8% of the visits ([Fig. 1](#)). This large range estimate reflects uncertainty (not variability) due to the limitations of our approach. As secondary robbing is critically dependent on primary robbing—an able robber must first make a hole—we examined additional photographs of many robbed plant species, and we found extensive flower damage as well as records of several species of flowerpiercers (*Diglossa*) commonly observed in the areas in close proximity to trainbearer robbing locations (B. Igić, P.B. Fenberg, 2019, personal observation). It is therefore likely that secondary robbing may

represent the majority of illegitimate visits by trainbearers, especially because their range overlaps with multiple species of flowerpiercers. Finally, it is possible that due to the difficulty of observation, this mode of nectar foraging is underestimated in other hummingbird species.

From the hummingbird perspective, if primary robbing is more likely to allow access to previously unvisited, relatively nectar-rich flowers, then perhaps secondary robbing ought to be less advantageous (Irwin *et al.*, 2010). Framed in this way, any secondary robbing may appear somewhat paradoxical. It is possible that, instead, one or more of the common underlying assumptions are flawed. Specifically, we can imagine that, for example, if they are faced with resource limitation (of flowers that are ordinarily legitimately exploited), trainbearers respond by robbing or thieving from inferior resources, a situation that may be exacerbated by territoriality. Although our study cannot strongly substantiate it, a trend of higher robbing frequency in males than females (complicated by association with species), may be related to such phenomena. On the other hand, the evolutionary game between hummingbirds, flowers, and all other interactors (whole pollinator guilds, herbivores, etc.) is likely substantially more complex (Maloof & Inouye, 2000), broadening the set of conditions for stable non-zero frequencies of secondary robbing. If we are to gain a more complete understanding of hummingbird foraging strategies, it will be critical to record these patterns and more closely scrutinize the ecological context, including target plants and other visitors (Richman, Irwin & Bronstein, 2017).

Floral larceny is often targeted at flowers of plants that have close co-evolutionary relationships with other legitimately pollinated animal species, such as the sword-billed hummingbird and its guild of long-flowered species (Soteras *et al.*, 2018). Some characteristics of hummingbirds, including short length and tomial serrations are hypothesized to be particularly closely associated with primary nectar robbing (Ornelas, 1994). The idea that the serration of bill tomia affects nectar robbing ability and efficiency remains broadly untested. Recent studies suggest that the presence of serrations is sexually dimorphic, with greater or exclusive expression in males, and that this feature could instead be shaped by a role in plumage preening or fighting, especially male territorial defense (Rico-Guevara *et al.*, 2019). Although additional field studies of nectar robbing and examination of bill morphologies for the presence of serrate tomia are needed to clarify their role, our data provide some insight. First, our data imply that, if secondary nectar robbing is more common than previously thought, it may be difficult to detect an association between serration and (combined) robbing observations. Serrations may well provide flower-robbing utility, but they do not appear necessary for secondary robbing. Second, if males perform primary robbing more frequently, then the existence of sexual dimorphism for tomial serrations is less problematic for the Ornelas hypothesis. Moreover, neither sexual dimorphism nor the proposed role in defense precludes simultaneous or exaptive function of serrated tomia in piercing. Consequently, it will be difficult to accumulate a great weight of evidence that exclusively support or detracts from this hypothesis. It is maybe instructive that many other specialized piercers, such as the Wedge-billed Hummingbird (*Schistes geoffroyi* and Purple-crowned fairy *Heliothryx barroti*), fail to present sexually dimorphic serrations (Rico-Guevara *et al.*, 2019).

A systematic recording of the nectar feeding mode, along with a number of predictors,

coupled with careful studies of feeding behavior and morphology are likely to be most persuasive in examining the validity and generality of the Ornelas hypothesis.

Predominance of different modes of larceny is likely to be associated with differing evolutionary responses from flowering plants, including variation in amount and/or time-dependence of nectar release, mating system (increased reliance on self-fertilization), and defenses, both physical (spines, thicker calyces or bracts) and chemical (secondary metabolites). A number of plant species visited by trainbearers are members of the nightshade family (Solanaceae), known for common production of alkaloids, which are broadly toxic to vertebrates. Perhaps not coincidentally, a great number and quantity of alkaloids are produced by species of *Brugmansia* and *Nicotiana*, and tissue at the base of flowers (calyx) contains some of the highest concentrations of such compounds (Saitoh, Noma & Kawashima, 1985; Alves, Sartoratto & Trigo, 2007). *Brugmansia* species have not been studied in equal detail to tobacco, but at least some plants in this genus can incapacitate juvenile humans, orders of magnitude larger than hummingbirds, following casual skin contact (Andreola et al., 2008). Moreover, given that there is a strong association between hummingbird pollination and self-fertility (Wolowski et al., 2013), stereotypically hummingbird-pollinated flowers may be more commonly robbed by both avian and hymenopteran robbers. This may be especially true for those flowers with relatively longer tubular corollas and large amounts of nectar (Lara & Ornelas, 2001; Rojas-Nossa, Sánchez & Navarro, 2016), precisely the syndrome commonly associated with hummingbird pollination.

CONCLUSION

Our observations and analyses point to additional needed work on systematics and ecology of *Lesbia*, especially their unsettled taxonomy, feeding ecology and energy budgets. More broadly, we sorely lack observational data to establish the relative frequencies of nectar feeding modes and manipulative field study to obtain direct evidence for nectar feeding dynamics across all hummingbirds and other nectar-feeding birds (Irwin et al., 2010). Our study documents the presence of larceny in trainbearers and, more specifically, shows that floral larceny is surprisingly common. These findings indicate that larceny, especially secondary robbing, may be more common among hummingbirds than is generally appreciated. Furthermore, we demonstrate that citizen science databases can be fruitfully employed to establish the presence of nectar robbing and coarsely examine its frequency. Such approaches can also generate hypotheses and guide an improved understanding of plant-side effects and responses, including pollination efficiency, the distribution of visitor interactions (Malooof & Inouye, 2000; Del Coro Arizmendi, 2001), plant incentives and defenses (as well as overall fitness) (Pelayo, Rengifo & Soriano, 2011), and ultimately better explain the dynamics in the co-evolutionary game between flowers and their visitors.

ACKNOWLEDGEMENTS

We are greatly indebted to iNaturalist & eBird projects, and all contributing natural history enthusiasts whose images and data enabled us to conduct this study. We thank Diego Emerson Torres for providing data and permission to use the image shown in Fig. 1.

Ben Marks & John Bates (FMNH) provided advice and access to bird collections, and Cathy Rushworth & Carl Rothfels provided generous help with plant species identification. Rainee Kaczorowski, an anonymous reviewer, and Mason-Gamer-Igic lab groups provided comments and edits, which improved the manuscript. Jackie Treehorn, a park ranger, fixed the cable.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This work was supported by a grant from the National Science Foundation NSF DEB-1655692 to Boris Igić and Royal Society Grant RG160700 to Phillip B. Fenberg. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:
National Science Foundation (NSF): DEB-1655692.
Royal Society: RG160700.

Competing Interests

The authors declare that they have no competing interests.

Author Contributions

- Boris Igić conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.
- Ivory Nguyen performed the experiments, authored or reviewed drafts of the paper, and approved the final draft.
- Phillip B. Fenberg conceived and designed the experiments, performed the experiments, authored or reviewed drafts of the paper, and approved the final draft.

Data Availability

The following information was supplied regarding data availability:
The raw data is available in the [Supplemental Files](#).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.9561#supplemental-information>.

REFERENCES

- Alves MN, Sartoratto A, Trigo JR. 2007. Scopolamine in *Brugmansia suaveolens* (Solanaceae): defense, allocation, costs, and induced response. *Journal of Chemical Ecology* 33(2):297–309 DOI 10.1007/s10886-006-9214-9.
- Andreola B, Piovan A, Da Dalt L, Filippini R, Cappelletti E. 2008. Unilateral mydriasis due to angel's trumpet. *Clinical Toxicology* 46(4):329–331 DOI 10.1080/15563650701378720.

- Boehm M. 2018.** Biting the hand that feeds you: wedge-billed hummingbird is a nectar robber of a sicklebill-adapted Andean bellflower. *Acta Amazonica* **48(2)**:146–150
DOI [10.1590/1809-4392201703932](https://doi.org/10.1590/1809-4392201703932).
- Del Coro Arizmendi M. 2001.** Multiple ecological interactions: nectar robbers and hummingbirds in a highland forest in Mexico. *Canadian Journal of Zoology* **79(6)**:997–1006
DOI [10.1139/z01-066](https://doi.org/10.1139/z01-066).
- Feinsinger P, Beach JH, Linhart YB, Busby WH, Murray KG. 1987.** Disturbance, pollinator predictability, and pollination success among Costa Rican cloud forest plants. *Ecology* **68(5)**:1294–1305 DOI [10.2307/1939214](https://doi.org/10.2307/1939214).
- Gonzalez O, Loiselle BA. 2016.** Species interactions in an andean bird–flowering plant network: phenology is more important than abundance or morphology. *PeerJ* **4(1)**:e2789
DOI [10.7717/peerj.2789](https://doi.org/10.7717/peerj.2789).
- Gould J. 1861.** *An introduction to the Trochilidae, or family of humming-birds*. London: Taylor & Francis.
- iNaturalist. 2019.** Available at <https://www.inaturalist.org> (accessed 25 December 2019).
- Inouye DW. 1980.** The terminology of floral larceny. *Ecology* **61(5)**:1251–1253
DOI [10.2307/1936841](https://doi.org/10.2307/1936841).
- Irwin RE, Bronstein JL, Manson JS, Richardson L. 2010.** Nectar robbing: ecological and evolutionary perspectives. *Annual Review of Ecology, Evolution, and Systematics* **41(1)**:271–292
DOI [10.1146/annurev.ecolsys.110308.120330](https://doi.org/10.1146/annurev.ecolsys.110308.120330).
- Lara C, Ornelas J. 2001.** Preferential nectar robbing of flowers with long corollas: experimental studies of two hummingbird species visiting three plant species. *Oecologia* **128(2)**:263–273
DOI [10.1007/s004420100640](https://doi.org/10.1007/s004420100640).
- Maloof JE, Inouye DW. 2000.** Are nectar robbers cheaters or mutualists? *Ecology* **81(10)**:2651–2661 DOI [10.1890/0012-9658\(2000\)081\[2651:ANRCOM\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2000)081[2651:ANRCOM]2.0.CO;2).
- McDade LA, Kinsman S. 1980.** The impact of floral parasitism in two neotropical hummingbird-pollinated plant species. *Evolution* **34(5)**:944–958
DOI [10.1111/j.1558-5646.1980.tb04033.x](https://doi.org/10.1111/j.1558-5646.1980.tb04033.x).
- Navarro L. 2001.** Reproductive biology and effect of nectar robbing on fruit production in *Macleania bullata* (Ericaceae). *Plant Ecology* **152(1)**:59–65 DOI [10.1023/A:1011463520398](https://doi.org/10.1023/A:1011463520398).
- Ornelas JF. 1994.** Serrate tomia: An adaptation for nectar robbing in hummingbirds? *Auk* **111(3)**:703–710.
- Pelayo RC, Rengifo C, Soriano PJ. 2011.** Avian nectar robbers of *passiflora mixta* (Passifloraceae): do they have a positive effect on the plant? *Interciencia* **36(8)**:587–592.
- R Core Team. 2019.** *R: a language and environment for statistical computing*. Vienna: The R Foundation for Statistical Computing. Available at <http://www.R-project.org/>.
- Remsen J, Stiles FG, Scott PE. 1986.** Frequency of arthropods in stomachs of tropical hummingbirds. *Auk* **103(2)**:436–441.
- Richman SK, Irwin RE, Bronstein JL. 2017.** Foraging strategy predicts foraging economy in a facultative secondary nectar robber. *Oikos* **126(9)**:1250–1257 DOI [10.1111/oik.04229](https://doi.org/10.1111/oik.04229).
- Rico-Guevara A, Rubega M, Hurme K, Dudley R. 2019.** Shifting paradigms in the mechanics of nectar extraction and hummingbird bill morphology. *Integrative Organismal Biology* **1(1)**:oby006 DOI [10.1093/iob/oby006](https://doi.org/10.1093/iob/oby006).
- Rojas-Nossa SV, Sánchez JM, Navarro L. 2016.** Effects of nectar robbing on male and female reproductive success of a pollinator-dependent plant. *Annals of botany* **117(2)**:291–297.

- Roubik DW, Holbrook NM, Parra GV. 1985.** Roles of nectar robbers in reproduction of the tropical treelet *quassia amara* (simaroubaceae). *Oecologia* **66**(2):161–167
DOI 10.1007/BF00379850.
- Saitoh F, Noma M, Kawashima N. 1985.** The alkaloid contents of sixty *Nicotiana* species. *Phytochemistry* **24**(3):477–480 DOI 10.1016/S0031-9422(00)80751-7.
- Snow DW, Snow BK. 1980.** Relationships between hummingbirds and flowers in the andes of colombia. *Bulletin of the British Museum of Natural History (Zoology)* **38**:105–139.
- Soteras F, More M, Ibanez AC, Del Rosario Iglesias M, Cocucci AA. 2018.** Range overlap between the sword-billed hummingbird and its guild of long-flowered species: an approach to the study of a coevolutionary mosaic. *PLOS ONE* **13**(12):e0209742
DOI 10.1371/journal.pone.0209742.
- Stiles FG. 2004.** Proposal (143) to South American classification committee split *Lesbia gouldii* from *Lesbia nuna*. Available at <http://www.museum.lsu.edu/~Remsen/SACCprop143.htm> (accessed 21 January 2020).
- Sullivan BL, Wood CL, Iliff MJ, Bonney RE, Fink D, Kelling S. 2009.** eBird: a citizen-based bird observation network in the biological sciences. *Biological Conservation* **142**(10):2282–2292
DOI 10.1016/j.biocon.2009.05.006.
- Vogt CA. 2006.** Secondary nectar robbing, a previously unsubstantiated foraging behavior of the Cinereous Conebill (*conirostrum cinereum*). *Ornitologia Neotropical* **17**(4):613–617.
- Weller AA, Schuchmann KL. 2004.** Biogeographic and taxonomic revision of the trainbearers *Lesbia* (Trochilidae), with the description of two new subspecies. *Ornithologischer Anzeiger* **43**(2):115–136.
- Wolowski M, Saad CF, Ashman T-L, Freitas L. 2013.** Predominance of self-compatibility in hummingbird-pollinated plants in the Neotropics. *Naturwissenschaften* **100**(1):69–79
DOI 10.1007/s00114-012-0995-0.