

Microbiome dynamics in the tissue and mucus of acroporid corals differ in relation to host and environmental parameters

Giulia M Marchioro ^{Equal first author, 1, 2, 3}, **Bettina Glasl** ^{Corresp., Equal first author, 3, 4, 5}, **Aschwin H Engelen** ², **Ester A Serrão** ², **David G Bourne** ^{3, 4, 5}, **Nicole S Webster** ^{3, 4, 6}, **Pedro R Frade** ²

¹ University of Algarve, Faro, Portugal

² CCMAR - Centre of Marine Sciences, University of Algarve, Faro, Portugal

³ AIMS@JCU, Townsville, Queensland, Australia

⁴ Australian Institute of Marine Science, Townsville, Queensland, Australia

⁵ College of Science and Engineering, James Cook University, Townsville, Australia

⁶ Australian Centre for Ecogenomics, University of Queensland, Brisbane, Queensland, Australia

Corresponding Author: Bettina Glasl

Email address: b.glasl@aims.gov.au

Corals are associated with diverse microbial assemblages; however, the spatial-temporal dynamics of intra-species microbial interactions are poorly understood. The coral-associated microbial community varies substantially between tissue and mucus microhabitats, however, the factors controlling the occurrence, abundance, and distribution of microbial taxa over time have rarely been explored for different compartments simultaneously. Here, we test 1) differentiation in microbiome diversity and composition between coral compartments (surface mucus and tissue) of two *Acropora* hosts (*A. tenuis* and *A. millepora*) common along inshore reefs of the Great Barrier Reef, as well as 2) the potential linkage between shifts in individual coral microbiome families and underlying host and environmental parameters. Amplicon based 16S ribosomal RNA gene sequencing of 136 samples collected over 14 months, revealed significant differences in bacterial richness, diversity and community structure among mucus, tissue and the surrounding seawater. Seawater samples were dominated by members of the Synechococcaceae and Pelagibacteraceae bacterial families. The mucus microbiome of *Acropora* spp. was dominated by members of Flavobacteriaceae, Synechococcaceae and Rhodobacteraceae and the tissue was dominated by Endozoicimonaceae. Mucus microbiome in both *Acropora* species was primarily correlated with seawater parameters including levels of chlorophyll *a*, ammonium, particulate organic carbon and the sum of nitrate and nitrite. In contrast, the correlation of the tissue microbiome to the measured environmental (i.e. seawater parameters) and host health physiological factors differed between host species, suggesting host-specific modulation of the tissue-associated microbiome to intrinsic and extrinsic factors. Furthermore, the correlation between

individual coral microbiome members and environmental factors provides novel insights into coral microbiome-by-environment dynamics and hence has potential implications for current reef restoration and management efforts (e.g. microbial monitoring / observatory programs).

Microbiome dynamics in the tissue and mucus of acroporid corals differ in relation to host and environmental parameters

Giulia M. Marchioro^{*,1,2,3}, Bettina Glasl^{*,3,4,5}, Aschwin H. Engelen², Ester A. Serrão², David G.
Bourne^{3,4,5}, Nicole S. Webster^{3,4,6}, Pedro R. Frade²

*these authors contributed equally to the study

¹ University of Algarve, Faro, Portugal

² CCMAR- Centre of Marine Sciences, University of Algarve, Faro, Portugal

³ AIMS@JCU, Townsville, Queensland, Australia

⁴ Australian Institute of Marine Science, Townsville, Queensland, Australia

⁵ College of Science and Engineering, James Cook University, Townsville, Queensland,
Australia

⁶ Australian Centre for Ecogenomics, University of Queensland, Brisbane, Queensland, Australia

Corresponding Author:

Bettina Glasl

Australian Institute of Marine Science, Townsville, Queensland, Australia

Email address: b.glasl@aims.gov.au

Abstract

Corals are associated with diverse microbial assemblages; however, the spatial-temporal dynamics of intra-species microbial interactions are poorly understood. The coral-associated microbial community varies substantially between tissue and mucus microhabitats, however, the factors controlling the occurrence, abundance, and distribution of microbial taxa over time have rarely been explored for different compartments simultaneously. Here, we test 1) differentiation in microbiome diversity and composition between coral compartments (surface mucus and tissue) of two *Acropora* hosts (*A. tenuis* and *A. millepora*) common along inshore reefs of the Great Barrier Reef, as well as 2) the potential linkage between shifts in individual coral microbiome families and underlying host and environmental parameters. Amplicon based 16S ribosomal RNA gene sequencing of 136 samples collected over 14 months, revealed significant differences in bacterial richness, diversity and community structure among mucus, tissue and the surrounding seawater. Seawater samples were dominated by members of the Synechococcaceae and Pelagibacteraceae bacterial families. The mucus microbiome of *Acropora* spp. was dominated by members of Flavobacteriaceae, Synechococcaceae and Rhodobacteraceae and the tissue was dominated by Endozoicimonaceae. Mucus microbiome in both *Acropora* species was primarily correlated with seawater parameters including levels of chlorophyll *a*, ammonium, particulate organic carbon and the sum of nitrate and nitrite. In contrast, the correlation of the tissue microbiome to the measured environmental (i.e. seawater parameters) and host health physiological factors differed between host species, suggesting host-specific modulation of the tissue-associated microbiome to intrinsic and extrinsic factors. Furthermore, the correlation between individual coral microbiome members and environmental factors provides novel insights into coral microbiome-by-environment dynamics and hence has potential implications for current reef restoration and management efforts (e.g. microbial monitoring / observatory programs).

Introduction

Coral microbiomes include the well-characterized endosymbiotic dinoflagellates of the family Symbiodiniaceae, and a vast diversity of bacteria and archaea (Bourne, Morrow & Webster, 2016; Frade et al., 2016a; Rowher et al., 2002). The microbiome has a fundamental role in the health and stability of the coral holobiont; it recycles nutrients, removes waste products and

defends against pathogens (Lema, Willis & Bourne, 2012; Morris et al., 2011; Rådecker et al., 2015; Rosado et al., 2019). The coral microbiome is influenced by a variety of intrinsic and extrinsic factors. Coral microbiomes are host species-specific and were thought to remain relatively stable over space and time (Frias-Lopez et al., 2002; Rowher et al., 2002). However, recent studies have proposed that spatial-temporal factors such as environmental parameters (Chen et al., 2011), depth (Glasl et al., 2017), geography (Hong et al., 2009; Littman et al., 2009a), seasonality (Ceh, van Keulen & Bourne, 2011; Chen et al., 2011; Hong et al., 2009; Koren & Rosenberg, 2006), coastal pollution (Klaus et al., 2007), and the physiological status of the host (Grottoli et al., 2018; Littman, Willis & Bourne, 2009b) can also influence the occurrence and relative abundance of microbial taxa. For instance, Li et al. (2015) reported a dynamic relationship between the community structure of coral-associated bacteria and the seasonal variation in environmental parameters such as dissolved oxygen and rainfall. Glasl et al. (2019a) showed that although host-associated microbiomes were five-times less responsive to the environment compared to the seawater microbiome, they were still affected by environmental factors (e.g. temperature, turbidity, and nutrient concentration).

The coral provides different microhabitats for its microbial associates, including the surface mucus layer, coral tissue, skeleton and gastrovascular cavity, each differing in microbial richness, diversity and community structure, often assessed through alpha- and beta-diversity metrics (Agostini et al., 2012; Engelen et al., 2018; Pollock et al., 2018; Sweet, Croquer & Bythell, 2011). Each microhabitat has a unique set of biochemical features and harbors a specific microbial community (Engelen et al., 2018; Pollock et al., 2018; Sweet, Croquer & Bythell, 2011). Hence, revealing microhabitat-specific host-microbiome associations and their specific sensitivities to environmental fluctuations is crucial to our understanding of coral holobionts. For example, the coral surface mucus layer is a polysaccharide-protein-lipid complex that provides an interface between the coral epithelium and the surrounding seawater (Brown & Bythell, 2005). Here microbes take advantage of a nutrient-rich medium and particular microbiome members found in the coral mucus overlap with both the tissue and the seawater microbial communities (Bourne & Munn, 2005; Brown & Bythell, 2005; Glasl, Herndl & Frade, 2016; Sweet, Croquer & Bythell, 2011). In contrast to the extracellular polymeric nature of the surface mucus layer, the coral tissue consists of two distinct layers (epidermis and gastrodermis) and a connective-tissue layer, the mesoglea (Muller-Parker, D'Elia & Cook, 2015). The coral tissue

harbors photosymbiotic dinoflagellates (family Symbiodiniaceae), that can provide up to 100% of energy required by their coral host (Muller-Parker, D'Elia & Cook, 2015). The Symbiodiniaceae community has been shown to vary in tandem with the bacterial community in early life stages of corals (Quigley et al., 2019) and this may be caused by the release of complex organic molecules such as the organosulfur compound dimethylsulfoniopropionate (DMSP; Bourne et al., 2013; Frade et al., 2016b). The coral tissue microbiome is mostly represented by bacteria belonging to the phyla Proteobacteria and Actinobacteria. For example, the gammaproteobacterial *Endozoicomonas* are abundant in the coral's endodermal tissue and are often considered 'true' coral symbionts (Bayer et al., 2013; Glasl et al., 2019b; Neave et al., 2016, 2017). When compared to the surface mucus layer, the microbial community in the tissue is significantly less dense and diverse (Bourne & Munn, 2005; Koren & Rosenberg, 2006), likely attributed to the more spatially stable and host controlled environment (Bourne & Munn, 2005), although divergent evidence suggests the mucus is less diverse than the tissue (Pollock et al., 2018). Furthermore, tissue-associated bacterial communities form aggregations within the coral cell layers, also referred to as coral-associated microbial aggregates (CAMAs), and are often co-localized near algal symbiont cells, highlighting potential metabolic interactions between symbionts (Wada et al., 2019).

In this study, we test the hypotheses that different coral compartments (surface mucus layer and tissue) of *Acropora* spp. harbor distinct microbial communities and that different intrinsic and extrinsic factors explain microbiome dynamics within these compartments. Furthermore, we aim to identify significant correlations of individual bacterial families associated with coral tissue and mucus with host-physiological and seawater parameters.

Materials & Methods

Sample collection

Samples of *Acropora millepora* (Ehrenberg, 1834), *Acropora tenuis* (Dana, 1846) and seawater were collected monthly, at Geoffrey Bay (Magnetic Island) in the Great Barrier Reef (Supplementary Figure S1), between February 2016 and March 2017, for amplicon based 16S ribosomal RNA (rRNA) gene sequencing along with environmental metadata. All samples were collected under the permit G16/38348.1 issued by the Great Barrier Reef Marine Park Authority.

Samples (n = 3 per sample type and per sampling event) for molecular analysis were collected as part of the Australian Microbiome Initiative and the sample procedure has previously been outlined by Glasl et al. (2019a). In brief, coral nubbins (approximately 5 cm tall) of both *Acropora* species were collected, rinsed with 0.2 µm filter-sterilized seawater and placed into cryogenic vials. Coral mucus from the same specimens was collected with sterile cotton swabs as previously described by Glasl, Herndl & Frade (2016). Seawater samples for molecular analysis were collected in sterile collapsible bags, pre-filtered through a 50 µm filter mesh to remove large particles, and subsequently filtered onto a 0.2 µm Sterivex filter (Millipore). Coral nubbins, mucus swabs and Sterivex filters were immediately snap frozen in liquid nitrogen after collection and stored at -80°C until further processing. To acquire environmental information, water and sediment samples were collected in duplicate for each sampling event as described in Glasl et al. (2019a) and further analyzed according to the standard procedures of Australian Institute of Marine Science (Devlin & Lourey, 2000). The environmental information processed includes common reef water quality measures such as salinity, particulate organic carbon, total suspended solids, concentrations of chlorophyll *a*, ammonium, the sum of nitrite and nitrate, particulate nitrogen, nitrite, total nitrogen, non-purgeable organic carbon, non-purgeable inorganic carbon, phosphate and silica as well as total organic carbon in the sediment, total organic nitrogen in the sediment and grainsize percentage of sediments < 0.63 µm, between 0.63 µm and 2 mm, and > 2 mm. Seawater temperatures and daylight hours were obtained from AIMS long-term monitoring temperature records (<http://eatlas.org.au>).

Sample preparation and genetic assays

Frozen coral tissue was airbrushed into a *ziploc* bag with phosphate-buffered saline (PBS) solution added until all tissue was removed from the skeletal fragment (PBS volume noted). The resulting tissue slurry was homogenized for 1 min at 12,500 rpm using a hand-held tissue homogenizer (Heidolph Silent Crusher M), pelleted (10 min at 16,000 rcf) and snap frozen in liquid nitrogen. DNA from the tissue and mucus samples was extracted using the DNeasy PowerBiofilm kit (QIAGEN). DNA extracts were sent on dry ice to the Ramaciotti Centre for Genomics (Sydney, Australia) for sequencing. The bacterial 16S rRNA gene was sequenced using the 27F (Lane, 1991) and 519R (Turner et al., 1999) primers on the Illumina MiSeq platform using a dual indexed 2 x 300 bp paired-end approach. Primers pairs were selected to

warrant comparability across datasets of the Australian Microbiome Initiative
(<https://www.australianmicrobiome.com>).

Sequence analysis

Sequencing data were analyzed as single nucleotide variants following the standardized platform of the Australian Microbiome Initiative (Brown et al., 2018). In brief, paired-end reads were merged using FLASH software (Magoc & Salzberg, 2011) and FASTA formatted sequences were extracted from FASTQ files. Sequences < 400 bp in length, and / or containing one or more N's, or homopolymer runs of > 8 bp were removed with MOTHUR (v1.34.1; Schloss et al., 2009). Sequences were de-replicated and ordered by abundance using USEARCH (64 bit v10.0.240; Edgar, 2010). Sequences with less than 4 representatives and Chimeras were removed, and the quality-filtered sequences were mapped to chimera-free zero-radius operational taxonomic units (zOTUs). A table containing the samples and their read abundances was created and the zOTUs were taxonomically classified with SILVA v132 database (Yilmaz et al., 2014) using MOTHUR's implementation of the Wang classifier (Wang et al., 2007) and a 60% Bayesian probability cut-off. This sequencing dataset has already been used in a previous contribution by the research group (Glasl et al., 2019a), but in the current study it is analyzed from a different perspective aiming at comparing temporal microbiome dynamics between two distinct coral compartments.

Chloroplasts and mitochondria derived reads were removed from the dataset and remaining data was rarefied to a sequencing depth of 3,500 reads per sample in R (R Core Team, 2015) using `subset_taxa()` function in the `phyloseq` package (McMurdie & Holmes, 2013). Read counts per sample were transformed into relative abundances.

Coral holobiont photopigment quantification

Photopigment (chlorophyll *a*) concentrations in the tissue of corals were quantified using a spectrophotometric approach (Glasl et al., 2019b). Tissue pellets were thawed on ice to avoid sample degradation and resuspended in 1 ml of 90% ethanol. Samples were sonicated for 1 min and centrifuged for 5 min at 10,000 rcf. Subsequently, 700 µl of the supernatant was removed and transferred to a new tube. The resuspension, sonication and centrifugation were repeated on the remainder of the pellet. The supernatant was recovered again, combined with the previous

extraction and mixed by inversion. Sample extract and 90% ethanol (blank read) were loaded in triplicate (200 μ l each) to a 96-well plate and the absorbance was recorded at 470, 632, 649, 665, 696 and 750 nm in a Cytation 3 multi-mode microplate reader (BioTek, Winooski, USA) and analyzed using the software Gen5 (BioTek, Winooski, USA). Blank corrected absorbance measures were used to calculate chlorophyll *a* concentrations (Supplementary Equation S1).

Coral protein quantification

Soluble protein concentrations of coral tissue samples were quantified using a colorimetric protein assay kit (Pierce BCA Protein Assay Kit; Glasl et al., 2019b). Tissue pellets were thawed on ice and resuspended in 1 ml PBS. The resuspension (25 μ l) was added to 200 μ l of working reagent from the kit in a 96-well plate. The plate was mixed thoroughly on a plate shaker for 30 seconds and then incubated at 37°C for 30 min. The plate was cooled down at room temperature. The absorbance was measured at 563 nm in a Cytation 3 multi-mode microplate reader (BioTek, Winooski, USA) and analyzed using the software Gen5 (BioTek, Winooski, USA). Measurements of the standards and samples were blank corrected to remove background absorbance. For each plate, a protein standard curve was obtained using bovine serum albumin (BSA) solution at concentrations between 25 and 2,000 μ g ml⁻¹.

Symbiodiniaceae cell counting

To determine cell numbers of Symbiodiniaceae in the coral tissue, the tissue pellet was thawed on ice, resuspended in 1 ml of 0.2 μ l filtered seawater and fixed in 2% formaldehyde (final concentration) to preserve the symbiont cells. The solution was passed through a syringe needle to reduce cell agglomeration and diminish the bias from cell clumps. Samples were then mixed for 1 min and 10 μ l of the homogenate was loaded onto a Neubauer haemocytometer (0.100 mm depth). Symbiodiniaceae cells were counted under 40 x magnification with an Olympus CX31 light microscope. In total, six independent haemocytometer loadings (24 squares each with 0.1 μ l volume) were used per sample to ensure robustness of density determinations.

Statistical analyses

Statistical analyses were performed using RStudio (v1.1.463). Analyses of microbial communities were performed on rarefied relative abundance data at zOTU level. zOTU richness

and Shannon-Weaver diversity were compared across host compartments, host species and reference seawater samples using non-parametric Analysis of Variance (Kruskal-Wallis test using function `kruskal.test`), followed by Dunn's test for multiple comparisons (function `dunn.test`). All p-values were adjusted using the Benjamini-Hochberg multiple comparison correction method to decrease the false discovery rate (Benjamini & Hochberg, 1995). A Venn diagram was constructed to describe the shared and unique zOTUs among mucus, tissue and seawater microbiomes using VennDiagram package (Chen & Boutros, 2011) and visualized using `eulerr` package (Larsson, 2020). Non-Metric Multidimensional Scaling (NMDS) was used to illustrate the microbial community structure among host species and host compartments based on Bray-Curtis dissimilarities (`phyloseq` package, McMurdie & Holmes, 2013). Permutational Multivariate Analysis of Variance (PERMANOVA, 999 permutations) was used to test for differences in microbial structure between host species and host compartments using the `adonis2` function of the `vegan` package (Oksanen et al., 2013).

Physiological variables were normalized (i.e., chlorophyll *a* normalized to protein content, chlorophyll *a* normalized to Symbiodiniaceae numbers, Symbiodiniaceae density normalized to protein content) following common procedures in coral physiology studies (Frade et al., 2008, Iglesias-Prieto & Trench, 1997). Due to fragmentation of the collected coral branches, coral surface area could not be measured. Environmental and physiological variables were standardized and checked for collinearity using the Pearson correlation coefficient. Redundant variables based on Pearson's correlation (> 0.7 or < -0.7 ; Dormann et al., 2013) were removed from the analysis. Non-correlated variables were then used in a Bray-Curtis distance-based Redundancy Analysis (db-RDA), which quantifies the impact of the explanatory variables on the microbiome (dis)similarities (Legendre & Anderson, 1999). zOTU relative abundance, environmental and physiological metadata were used for db-RDA using the `phyloseq` package (McMurdie & Holmes, 2013). The analysis tests the statistical relationship between microbial community composition and the environmental/physiological variables for each coral compartment and host species combination. A model selection tool (`ordiR2step` function in the `vegan` package, *sensu* Blanchet, Legendre & Borcard, 2008) was performed to select the best db-RDA model (i.e., the best explanatory variables) for variation in microbiome composition of each coral compartment (mucus and tissue) in each host species (Johnson & Omland, 2004). The significance of each explanatory variable was confirmed with an ANOVA-like permutational test

(function `permutest`) for dbRDA. The explanatory value (in %) of significant explanatory variables (e.g. environmental and physiological parameters) on each microbiome was assessed with Variation Partitioning Analysis of the `vegan` package (Oksanen et al., 2013). A correlation matrix (based on the default Pearson correlation) between the relative abundance of the 20 most abundant microbial families and significant environmental variables was generated using the R package `MicrobiomSeq` (Ssekagiri, Sloan & Ijaz, 2017), for which p-values were adjusted using the Benjamini-Hochberg multiple comparison correction (Benjamini & Hochberg, 1995).

Results

Composition of coral tissue and mucus microbiomes

The bacterial 16S rRNA genes of 136 samples, including coral tissue (n = 24 for *A. millepora*; n = 30 for *A. tenuis*), coral mucus layer (n = 24 for *A. millepora*; n = 28 for *A. tenuis*) and seawater (n = 30; used as reference samples) were sequenced and 12,051 zOTUs identified at single nucleotide variants.

zOTU richness differed significantly among mucus, tissue and seawater microbiomes (Kruskal-Wallis $Chi^2_{(2,133)} = 57.74$, $p = 2.89 \times 10^{-13}$), but not between seasons (see Supplementary Table S1). Coral zOTU richness differed between species (*A. millepora* vs *A. tenuis*; Kruskal-Wallis $Chi^2_{(1,134)} = 12.23$, $p = 0.00047$). Seawater harbored the richest microbial community (558 zOTU \pm 54.6), followed by the mucus (*A. millepora*, 220 zOTU \pm 188; *A. tenuis* 511 zOTU \pm 234) and tissue (*A. millepora*, 125 zOTU \pm 31.6; *A. tenuis*, 173 zOTU \pm 146; Supplementary Table S1). Alpha diversity based on Shannon Index also differed significantly among microbiomes from mucus, tissue and seawater (Kruskal-Wallis $Chi^2_{(2,133)} = 53.37$, $p = 2.57 \times 10^{-12}$), but not between seasons (see Supplementary Table S1). Coral zOTU Shannon differed between species (*A. millepora* vs *A. tenuis*; Kruskal-Wallis $Chi^2_{(1,134)} = 6.002$, $p = 0.01429$). Alpha diversity measures of mucus samples were not significantly different (Shannon Index: *A. millepora*, 4.18 ± 0.83 ; *A. tenuis*, 5.15 ± 0.69) from seawater samples (Shannon Index: 4.40 ± 0.209 ; Supplementary Table S1). In contrast, the tissue microbiome was dramatically different from the mucus and seawater microbiomes and harbored the lowest microbial diversity (Shannon Index: *A. millepora*, 3.35 ± 0.63 ; *A. tenuis*, 3.54 ± 0.84).

Sequences affiliated to the phyla Proteobacteria dominated the microbial community of all samples (average relative abundance \pm SD; mucus: $44.1 \pm 11.5\%$; tissue: $62.8 \pm 2\%$;

seawater: 39.6 ± 3.1), followed in dominance by Bacteroidetes (mucus: $27.5 \pm 13.0\%$; tissue: $9.6 \pm 10.9\%$; seawater: $12.0 \pm 11.4\%$) and Cyanobacteria (mucus: $14.4 \pm 9.0\%$; tissue: $9.8 \pm 11.0\%$; seawater: $38.5 \pm 4.0\%$). Mucus microbiomes for both *Acropora* species (Figure 1) were characterized mostly by members of the family Flavobacteriaceae (average relative abundance \pm SD; for *A. tenuis*: $17.3 \pm 9.1\%$; *A. millepora*: $17.3 \pm 12.7\%$), Synechococcaceae (*A. tenuis*: $12.3 \pm 7.8\%$; *A. millepora*: $13.1 \pm 10.2\%$) and Rhodobacteraceae (*A. tenuis*: $5.7 \pm 3.0\%$; *A. millepora*: $6.4 \pm 6.4\%$; Figure 1). In contrast, the Endozoicimonaceae family dominated the tissue microbiome (*A. tenuis*: $43.2 \pm 31.7\%$; *A. millepora*: $20.5 \pm 19.7\%$), with additional representation of Flavobacteriaceae (*A. tenuis*: $7.9 \pm 9.6\%$; *A. millepora*: $7.2 \pm 9.6\%$), Synechococcaceae (*A. tenuis*: $5.5 \pm 6.8\%$; *A. millepora*: $12.3 \pm 14.5\%$) and Rhodobacteraceae (*A. tenuis*: $6.5 \pm 10.4\%$; *A. millepora*: $5.3 \pm 8.5\%$; Figure 1) families. Seawater samples were mostly characterized by members of Synechococcaceae ($36.6 \pm 3.9\%$) and Pelagibacteraceae ($18.6 \pm 4.9\%$), but also by Rhodobacteraceae ($8.6 \pm 4.8\%$) and Flavobacteriaceae ($8.0 \pm 2.6\%$; Figure 1). Tissue and mucus microbiomes exclusively shared 1,193 zOTU (9.9%), mucus and seawater microbiomes exclusively shared 1,458 zOTU (12.1%), whereas the tissue and seawater microbiome shared only 66 zOTU (0.6%; Figure 2).

Microbial community composition (beta-diversity) significantly differed among mucus, tissue and seawater (Figure 3; PERMANOVA, $pseudo-F_{(2,126)} = 14.53$, $p = 0.001$), between *Acropora* species (PERMANOVA, $pseudo-F_{(1,126)} = 4.42$, $p = 0.001$), and between seasons (PERMANOVA, $pseudo-F_{(1,126)} = 1.90$, $p = 0.011$). Interaction between species and compartment was also significant (PERMANOVA, $pseudo-F_{(1,126)} = 3.07$, $p = 0.002$; other interactions were not significant; Supplementary Table S2).

Explanatory variables of coral tissue and mucus microbiomes

Physiological parameters of the tissue (i.e., chlorophyll *a* normalized to protein content, chlorophyll *a* normalized to Symbiodiniaceae numbers, Symbiodiniaceae density normalized to protein content) remained stable between host species.

Out of a total of 20 environmental variables measured for seawater and sediment, 6 variables were non-mutually collinear and were thus included in the db-RDA analysis. Selected variables were salinity, concentration of particulate organic carbon (POC), total suspended solids

(TSS), chlorophyll *a* (Chl*a*), ammonium (NH₄⁺) and the sum of nitrite and nitrate concentrations (i.e., NO₂⁻/NO₃⁻; Supplementary Table S3).

Environmental/physiological parameters investigated in this study explained a limited amount of variation in the microbial community of mucus and tissue of the two *Acropora* species studied (Figure 4). For example, seawater parameters explained 14% (Chl*a*, NH₄⁺ and NO₂⁻/NO₃⁻) and 10% (POC and NO₂⁻/NO₃⁻) of the compositional variability for the mucus microbiome in *A. tenuis* and *A. millepora*, respectively (ANOVA-like permutational test for dbRDA; Supplementary Table S4); NO₂⁻/NO₃⁻ was the only explanatory environmental variable common to the mucus microbiome of both *Acropora* species (5% of compositional variability explained in each species). In comparison, for the seawater microbiome, environmental parameters (NO₂⁻/NO₃⁻, TSS, POC, Salinity and Chl*a*) explained 32% of the compositional variability of the microbiome (Supplementary Figure S2), suggesting greater environmental sensitivity by the microbial community in the seawater compared to the coral-associated communities.

In contrast, tissue microbiomes of *A. millepora* and *A. tenuis* differed substantially in their response to environmental and/or to physiological parameters. While host physiology (i.e., Symbiodiniaceae density normalized to protein contents) and environment (TSS and Chl*a*) explained 6% and 10%, respectively, of the variation of the tissue microbiome in *A. tenuis*, in *A. millepora*, the compositional variation was solely explained (10%) by environmental parameters (NO₂⁻/NO₃⁻ and TSS; Variation Partitioning Analysis and ANOVA-like permutational test for dbRDA; Supplementary Table S4). TSS was the only explanatory environmental variable common to the tissue microbiomes of both *Acropora* species (total of 5% and 4% in *A. tenuis* and in *A. millepora*, respectively).

Correlation between bacterial families and environmental/physiological parameters

The relative abundance of Synechococcaceae in tissues of both *Acropora* species and the mucus of *A. tenuis* was negatively correlated with TSS ($p = 0.025$ - 0.039 ; Figure 5 and Supplementary Tables S5 and S6). In contrast, Synechococcaceae was positively correlated to total NO₂⁻/NO₃⁻ in both species (mucus of *A. tenuis*, $p = 0.002$, Supplementary Table S6; and tissue of *A. millepora*, $p = 0.024$, Supplementary Table S5). For *A. tenuis*, Synechococcaceae abundance in the tissue correlated negatively with the only significant physiological parameter (Symbiodiniaceae density normalized to protein contents; $p = 0.025$). In the mucus of *A. millepora*, the abundance of

Pirellulaceae was positively correlated with $\text{NO}_2^-/\text{NO}_3^-$ ($p = 0.035$) and negatively correlated with TSS ($p = 0.019$), while OCS155 was positively correlated to $\text{NO}_2^-/\text{NO}_3^-$ ($p = 0.015$). Proteobacteria from the mucus of *A. tenuis*, Pelagibacteraceae and Halomonadaceae, were both strongly negative correlated with chlorophyll *a* in the seawater (Pelagibacteraceae, $p = 0.013$; Halomonadaceae, $p = 0.008$). Additionally, Halomonadaceae correlated negatively with NH_4^+ ($p = 0.005$; Figure 5 and Supplementary Tables S5 and S6).

Tissue-associated Endozoicimonaceae showed a strong significant positive correlation with Symbiodiniaceae density normalized to protein content in *A. tenuis* ($p = 0.0003$). In contrast, in the tissue of *A. millepora*, Endozoicimonaceae were negatively correlated with $\text{NO}_2^-/\text{NO}_3^-$ ($p = 0.020$), whereas the abundance of Cryomorphaceae family was negatively correlated with TSS ($p = 0.020$; Figure 5, Supplementary Table S5).

Discussion

Microbial communities associated with corals are continually exposed to fluctuations in the surrounding environment and the physiology of their host. Previous studies have demonstrated changes in the coral microbiome in response to thermal stress (Ainsworth & Hoegh-Guldberg, 2009; Grottoli et al., 2018; Lee et al., 2015; Thurber et al., 2009), ocean acidification (Grottoli et al., 2018; Thurber et al., 2009), organic matter enrichment (Garren & Azam, 2012), bleaching events (Bourne et al., 2008) and other environmental and physiological factors (Glasl et al., 2019a; Guppy & Bythell, 2006; Kelly et al., 2014; Li et al., 2015; Pollock et al., 2018). However, the coral microbiome is not homogenous across the animal and an improved understanding of the sensitivity of the microorganisms inhabiting each coral compartment is needed. This study highlights compositional differences in the bacterial communities associated with coral mucus and coral tissue, as well as with the surrounding seawater, findings that are largely consistent with previous studies (Apprill, Weber & Santoro, 2016; Bourne & Munn, 2005; Engelen et al., 2018; Pollock et al., 2018; Sweet, Croquer & Bythell, 2011). Furthermore, the high similarity between mucus and seawater microbiomes (see Supplementary Tables S1 and S2, Figures 2 and 3) and the high dissimilarity between tissue and seawater microbiomes suggests that the mucus microbial community is more strongly influenced by the external environment than the tissue community. Similar results have been reported for other coral species in the Caribbean (*Orbicella faveolata*, *Diploria strigosa*, *Montastraea cavernosa*, *Porites porites* and *Porites*

astreoides), where mucus and seawater shared significantly more microbial taxa than those shared by tissue and seawater microbiomes (Apprill, Weber & Santoro, 2016). Our results also support that mucus microbiomes are richer and more diverse than tissue microbiomes, which is a pattern corroborated by many studies (Bourne & Munn, 2005; Koren & Rosenberg, 2006).

Despite the host species-specificity of the coral microbiomes, some bacterial taxa were ubiquitously associated with a particular coral compartment. For example, Flavobacteriaceae and Synechococcaceae dominated the mucus of both species, while Endozoicimonaceae dominated the tissue microbiome of both *Acropora* species. However, overall microbiome composition also showed some overlap between host compartments, consistent with previous reports of overlap between the mucus and tissue microbiomes of other coral species (Engelen et al., 2018; Sweet, Croquer & Bythell, 2011). This intersection is a natural feature of the coral holobiont as both compartments are within the same host and because the constituents of the surface mucus layer are originally produced inside the tissue (Bythell & Wild, 2011). The sharing of some microbial taxa between compartments may also arise due to methodological challenges associated with retrieving samples that are exclusively mucus or coral tissue (Sweet, Croquer & Bythell, 2011). Similar methodological limitations could also obscure differences between the mucus and seawater microbiomes (Brown & Bythell, 2005).

Explanatory factors of mucus microbiome variation

We hypothesized that the coral mucus microbiome, which is in direct contact with seawater, would be primarily correlated with seawater parameters, whereas the tissue microbiome would be most affected by the physiological state of the coral host. Mucus is highly hydrated: mucocyte cells release their secretions in a condensed form which then undergo a massive swelling upon hydration, forming a visco-elastic gel (Brown & Bythell, 2005). Surface mucus can therefore be influenced by the presence of nutrients dissolved in the surrounding seawater (Tanaka, Ogawa & Miyajima, 2010). As expected, environmental factors (i.e. seawater parameters) were influential in shaping the mucus microbiome of both species (*A. millepora* and *A. tenuis*), consistent with recent studies relating changes in the mucus microbiome with environmental perturbations (Li et al., 2015; Pollock et al., 2018). However, the extent of influence from environmental parameters (10% of variation) on the mucus microbiome was much lower than the influence of environment on the seawater microbiome (32% of variation), suggesting that other factors also play a role in

modulating the mucus microbiome. For instance, the surrounding environment may interact with host physiology and together they alter the bacterial community structure of the mucus. Mucus is a nutrient-rich medium fueled by the photosynthetic activity of the Symbiodiniaceae (Brown & Bythell, 2005) and therefore it is expected that some degree of variation in its chemical composition is explained by host-Symbiodiniaceae factors. For example, *A. millepora* and *A. tenuis* at the sampling site (Geoffrey Bay at Magnetic Island) associate with distinct Symbiodiniaceae (LaJeunesse et al., 2018; Ulstrup & van Oppen, 2003; van Oppen et al., 2001). *A. millepora* colonies were associated with *Durussdinium* (van Oppen et al., 2001) whereas *A. tenuis* harbored *Cladocopium* spp. (Ulstrup & van Oppen, 2003). Links between mucus chemical composition and microbiome community structure have been proposed (Tremblay et al., 2011). Physiological factors regulating the dynamics of production and release of the surface mucus layer could also contribute to regulating mucus microbial composition (Glasl, Herndl & Frade, 2016).

Fluctuations of NH_4^+ , $\text{NO}_2^-/\text{NO}_3^-$, *Chla* and POC in the surrounding seawater significantly correlated with the mucus microbiome variation in *Acropora* species. Li et al. (2015) and Chen et al. (2011) suggested that rainfall had a crucial effect on bacterial community variation in the coral microbiome, being mostly associated with an increase in the relative abundance of the *Bacilli* group (Chen et al., 2011; Li et al., 2015). In the present study, $\text{NO}_2^-/\text{NO}_3^-$ (and its collinear variables daylight, particulate nitrogen and grainsize of sediments; Supplementary Table S3) had the greatest influence on microbiome structure, being a significant factor for both studied species. The link between rainfall and increasing nutrients (such as $\text{NO}_2^-/\text{NO}_3^-$) is well established for inshore reefs (Fabricius, 2005). In the current study, higher amounts of particulate and dissolved nutrients (but a decrease in TSS), corresponded to an increase in mucus-associated Synechococcaceae, Pirellulaceae, OCS155 and Rhodobacteraceae and a decrease in Halomonadaceae. For instance, Synechococcaceae in the mucus was highly positively correlated with $\text{NO}_2^-/\text{NO}_3^-$ and negatively correlated with TSS. These findings corroborate previous work in which the abundance of free-living *Synechococcus* in shallow coastal waters decreased significantly under lower nutrient (especially nitrate) and higher TSS concentrations (Uysal & Köksalan, 2006).

Dissolved nutrients, such as nitrogen and phosphorus, can affect coral physiology and drive changes in the associated microbial community (Shaver et al., 2017; Thompson et al.,

2015). For example, organic-rich nutrients from terrestrial run-off negatively affect the health of corals and promote rapid growth of opportunistic heterotrophic bacteria (e.g. Vibrionales, Flavobacteriales and Rhodobacterales), thus affecting the overall composition of the coral microbiome (McDevitt-Irwin et al., 2017; Weber et al., 2012). In our study, the abundance of Flavobacteriaceae and Rhodobacteraceae in the mucus of *A. tenuis* correlated with TSS and NH_4^+ , respectively. The coral holobiont, including cyanobacteria related to *Synechococcus* spp. (Lesser et al., 2004), can also efficiently take up inorganic nitrogen, for example, as nitrogen is required by the photosynthesis production of its Symbiodiniaceae symbionts (Yellowlees, Rees & Leggat, 2008). In fact, NH_4^+ can be assimilated by both coral and its Symbiodiniaceae (Pernice et al., 2012), and recent work has implicated bacteria such as *Vibrio* and *Alteromonas* in the incorporation and translocation of NH_4^+ into coral tissues and associated Symbiodiniaceae (Ceh et al., 2013). Nitrifying members of the mucus microbiome, such as ammonium oxidizing bacteria (e.g., Pirellulaceae) and archaea, are fueled by NH_4^+ (Beman et al., 2007; Siboni et al., 2008; Yang et al., 2013), and $\text{NO}_2^-/\text{NO}_3^-$ can be respired by nitrate reducers putatively active in coral microbiomes (Siboni et al., 2008; Yang et al., 2013). Interestingly, Pirellulaceae abundances in the mucus of *A. millepora* positively correlated with concentrations of environmental $\text{NO}_2^-/\text{NO}_3^-$, the products of ammonium oxidation. These nitrogen-cycling processes mediated by microbes are highly dependent on oxygen availability, but because oxygen concentration in the mucus shows strong diel fluctuations (Shashar, Cohen & Loya, 1993), it is possible that both aerobic (e.g., nitrification) and anaerobic (e.g., denitrification) processes happen within the mucus layer at different times of the day. Temporal dynamics in the coral mucus microbiome are thus likely influenced by the individual and collective metabolic capabilities of the diverse assemblage of microbes and by nutrient availability in the surrounding waters.

Explanatory factors of tissue microbiome variation

The statistical relation between the coral tissue microbiome and the environmental and physiological parameters differed between coral species. Whereas the tissue microbiome of *A. tenuis* corresponded to both environment and host physiology, *A. millepora* correlated only with environmental parameters. This difference may be associated to specific features of each species, through which *A. millepora* could modulate the internal environment and create more stable

intra-tissue conditions than *A. tenuis* (e.g., via skeletal light modulation, host morphology and tissue thickness, *sensu* Enriquez, Mendez & Iglesias-Prieto, 2005). A non-mutually exclusive alternative explanation is the influence of the algal symbiont (Symbiodiniaceae) genotype associated to the host. Little et al. (2004) investigated Symbiodiniaceae communities associated with *A. millepora* and *A. tenuis* on Magnetic Island demonstrating that the coral-algal endosymbiotic relationship in *Acropora* spp. is distinct between species, dynamic and flexible (corals associate with different Symbiodiniaceae types at different life stages, for example), and contributes significantly to physiological attributes of the coral holobiont. For example, different algal genotypes can affect the nutrient availability (e.g. carbon and nitrogen) in the coral holobiont (Pernice et al., 2015; Bayliss et al., 2019). Environmental factors such as seawater temperature can also lead to temporal changes in the symbiont community (Cooper et al., 2011; Howells et al., 2012; Rocker, Willis & Bay, 2012). As the microbiome is strongly associated to the coral holobiont, any disturbance in the host-Symbiodiniaceae relationship may have indirect effects on the microbial composition and its response to environmental and physiological factors. Other studies demonstrate the influence of Symbiodiniaceae on the host microbial community and also support the idea that these two components of the coral holobiont are finely tuned (Glas et al., 2017; Grottoli et al., 2018; Littman, Bourne & Willis, 2010; Littman, Willis & Bourne, 2009b; Quigley et al., 2019). In the present study, Endozoicimonaceae were strongly positively correlated with the Symbiodiniaceae density in the tissue of *A. tenuis* and negatively correlated with $\text{NO}_2^-/\text{NO}_3^-$ in *A. millepora* (see Figure 5). These results are to some extent at odds with experimental results showing a stable dominance of Endozoicimonaceae in tissues of *Pocillopora verrucosa* irrespective of excess dissolved organic nitrogen and despite a bleaching phenomenon concomitant with structural changes in its Symbiodiniaceae community (Pogoreutz et al., 2018).

Besides the diversity of Symbiodiniaceae associated to each coral species, other factors can affect the coral and its response to environmental parameters, such as photochemical efficiency (Fv/Fm) and symbiont density (Cunning & Baker, 2014; Da-Anoy, Cabaitan & Conaco, 2019). For instance, Da-Anoy, Cabaitan & Conaco (2019) demonstrated a greater reduction of Fv/Fm in *A. tenuis* in response to elevated temperatures compared to *A. millepora* and the temperature responses of the corals did not directly correlate with their associated seemingly associated Symbiodiniaceae. This suggests that other species-specific physiological

factors could modulate the responses of the coral to the environment and, indirectly, influence the tissue-associated microbiome. One such factor is the way coral-associated microbial aggregates (CAMAs) are distributed throughout the tissue, which varies within populations and can vary among coral species (Work & Aeby, 2014; Wada et al., 2019).

Total suspended solids (TSS) was the only environmental parameter measured in the present study that significantly related to the tissue microbiome of both coral species. TSS can impact corals by limiting light availability for photosynthesis and decreasing Symbiodiniaceae densities, which can indirectly affect microbial communities (Fabricius, 2005; Pollock et al., 2014). High levels of suspended solids characterize the environment of inshore reefs such as those found around Magnetic Island. The decrease in TSS is strongly associated with an increase in the abundance of tissue-associated Synechococcaceae and Cryomorphaceae. Cryomorphaceae are typical copiotrophs in the phylum Bacteroidetes and their increase in the tissue of *A. millepora* could relate to declines in coral holobiont health.

Conclusions

This study highlights that microbiomes inhabiting different physical microniches within the coral holobiont differ in their linkage between host and environmental factors. Microbiomes of *Acropora* spp. differed significantly among host compartments (surface mucus layer and tissue) and species (*A. tenuis* and *A. millepora*). Seawater parameters had the greatest influence on the mucus microbiome in both species whereas the tissue microbiomes showed differential patterns to environmental/host-physiological parameters, suggesting host-specific modulation of the tissue microbiome. While further research is needed to unequivocally define the drivers of coral microbiome variation, by investigating temporal variation in water quality and coral health measures and correlating these with microbial community dynamics across distinct host compartments in closely related species, this study has identified several intrinsic and extrinsic factors that contribute to microbiome composition in corals.

Acknowledgements

We thank Michele Skuza, Neale Johnston and the AIMS water quality team for their help with analyzing the water quality samples. We thank Heidi Luter, Katarina Damjanovic and Joe

Gioffre for their assistance in the field. We also thank Sara Bell, Florita Flores and Carlos Alvarez for their expertise in the laboratory.

References

- Agostini, S., Suzuki, Y., Higuchi, T., Casareto, B. E., Yoshinaga, K., Nakano, Y., & Fujimura, H. (2012). Biological and chemical characteristics of the coral gastric cavity. *Coral Reefs*, 31(1), 147–156. <https://doi.org/10.1007/s00338-011-0831-6>
- Ainsworth, T. D., & Hoegh-Guldberg, O. (2009). Bacterial communities closely associated with coral tissues vary under experimental and natural reef conditions and thermal stress. *Aquatic Biology*, 4, 289–296. <https://doi.org/10.3354/ab00102>
- Ainsworth, T. D., Krause, L., Bridge, T., Torda, G., Raina, J.-B., Zakrzewski, M., Gates, R. D., Padilla-Gamiño, J. L., Spalding, H. L., Smith, C., Woolsey, E. S., Bourne, D. G., Bongaerts, P., Hoegh-Guldberg, O., & Leggat, W. (2015). The coral core microbiome identifies rare bacterial taxa as ubiquitous endosymbionts. *The ISME Journal*, 9(10), 2261–2274. <https://doi.org/10.1038/ismej.2015.39>
- Apprill, A., Weber, L. G., & Santoro, A. E. (2016). Distinguishing between Microbial Habitats Unravels Ecological Complexity in Coral Microbiomes. *MSystems*, 1(5), e00143-16, /msys/1/5/e00143-16.atom. <https://doi.org/10.1128/mSystems.00143-16>
- Bayer, T., Neave, M. J., Alsheikh-Hussain, A., Aranda, M., Yum, L. K., Mincer, T., Huguen, K., Apprill, A., & Voolstra, C. R. (2013). The Microbiome of the Red Sea Coral *Stylophora pistillata* Is Dominated by Tissue-Associated Endozoicomonas Bacteria. *Applied and Environmental Microbiology*, 79(15), 4759–4762. <https://doi.org/10.1128/AEM.00695-13>

- Bayliss, S.L.J., Scott, Z.R., Coffroth, M.A., & terHorst, C.P. (2019) Genetic variation in *Breviolum antillologorgium*, a coral reef symbiont, in response to temperature and nutrients. *Ecology and Evolution*, 9, 2803-2813. <https://doi.org/10.1002/ece3.4959>
- Beman, J. M., Roberts, K. J., Wegley, L., Rohwer, F., & Francis, C. A. (2007). Distribution and Diversity of Archaeal Ammonia Monooxygenase Genes Associated with Corals. *Applied and Environmental Microbiology*, 73(17), 5642–5647. <https://doi.org/10.1128/AEM.00461-07>
- Benjamini, Y., & Hochberg, Y. (1995). Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal statistical society: series B (Methodological)*, 57(1), 289-300.
- Blanchet, F. G., Legendre, P., & Borcard, D. (2008). Forward selection of explanatory variables. *Ecology*, 89(9), 2623–2632. <https://doi.org/10.1890/07-0986.1>
- Bourne, D. G., Dennis, P. G., Uthicke, S., Soo, R. M., Tyson, G. W., & Webster, N. (2013). Coral reef invertebrate microbiomes correlate with the presence of photosymbionts. *The ISME Journal*, 7(7), 1452–1458. <https://doi.org/10.1038/ismej.2012.172>
- Bourne, D. G., Morrow, K. M., & Webster, N. S. (2016). Insights into the Coral Microbiome: Underpinning the Health and Resilience of Reef Ecosystems. *Annual Review of Microbiology*, 70(1), 317–340. <https://doi.org/10.1146/annurev-micro-102215-095440>
- Bourne, D. G., & Munn, C. B. (2005). Diversity of bacteria associated with the coral *Pocillopora damicornis* from the Great Barrier Reef. *Environmental Microbiology*, 7(8), 1162–1174. <https://doi.org/10.1111/j.1462-2920.2005.00793.x>
- Bourne, D., Iida, Y., Uthicke, S., & Smith-Keune, C. (2008). Changes in coral-associated microbial communities during a bleaching event. *The ISME Journal*, 2(4), 350–363. <https://doi.org/10.1038/ismej.2007.112>

- 576
- 577 Brown, B., & Bythell, J. (2005). Perspectives on mucus secretion in reef corals. *Marine Ecology*
- 578 *Progress Series*, 296, 291–309. <https://doi.org/10.3354/meps296291>
- 579
- 580 Brown, M. V., van de Kamp, J., Ostrowski, M., Seymour, J. R., Ingleton, T., Messer, L. F.,
- 581 Jeffries, T., Siboni, N., Laverock, B., Bibiloni-Isaksson, J., Nelson, T. M., Coman, F.,
- 582 Davies, C. H., Frampton, D., Rayner, M., Goossen, K., Robert, S., Holmes, B., Abell, G.
- 583 C. J., ... Bodrossy, L. (2018). Systematic, continental scale temporal monitoring of
- 584 marine pelagic microbiota by the Australian Marine Microbial Biodiversity Initiative.
- 585 *Scientific Data*, 5(1), 180130. <https://doi.org/10.1038/sdata.2018.130>
- 586
- 587 Bythell, J. C., & Wild, C. (2011). Biology and ecology of coral mucus release. *Journal of*
- 588 *Experimental Marine Biology and Ecology*, 408(1–2), 88–93.
- 589 <https://doi.org/10.1016/j.jembe.2011.07.028>
- 590
- 591 Ceh, J., Kilburn, M. R., Cliff, J. B., Raina, J.-B., van Keulen, M., & Bourne, D. G. (2013).
- 592 Nutrient cycling in early coral life stages: *Pocillopora damicornis* larvae provide their
- 593 algal symbiont (*Symbiodinium*) with nitrogen acquired from bacterial associates.
- 594 *Ecology and Evolution*, 3(8), 2393–2400. <https://doi.org/10.1002/ece3.642>
- 595
- 596 Ceh, J., van Keulen, M., & Bourne, D. G. (2011). Coral-associated bacterial communities on
- 597 Ningaloo Reef, Western Australia: Coral bacterial communities, Ningaloo Reef. *FEMS*
- 598 *Microbiology Ecology*, 75(1), 134–144. [https://doi.org/10.1111/j.1574-](https://doi.org/10.1111/j.1574-6941.2010.00986.x)
- 599 [6941.2010.00986.x](https://doi.org/10.1111/j.1574-6941.2010.00986.x)
- 600
- 601 Chen, C.-P., Tseng, C.-H., Chen, C. A., & Tang, S.-L. (2011). The dynamics of microbial
- 602 partnerships in the coral *Isopora palifera*. *The ISME Journal*, 5, 728–740.
- 603
- 604 Chen, H., & Boutros, P. C. (2011). *VennDiagram: A package for the generation of highly-*
- 605 *customizable Venn and Euler diagrams in R*. 12(1), 35.
- 606

- Commonwealth of Australia. (2019). <http://www.bom.gov.au/?ref=logo>
- Cooper, T. F., Berkelmans, R., Ulstrup, K. E., Weeks, S., Radford, B., Jones, A. M., Doyle, J., Canto, M., O’Leary, R. A., & van Oppen, M. J. H. (2011). Environmental Factors Controlling the Distribution of Symbiodinium Harboured by the Coral *Acropora millepora* on the Great Barrier Reef. *PLoS ONE*, 6(10), e25536. <https://doi.org/10.1371/journal.pone.0025536>
- Cunning, R., & Baker, A. C. (2014). Not just who, but how many: the importance of partner abundance in reef coral symbioses. *Frontiers in microbiology*, 5, 400.
- Da-Anoy, J. P., Cabaitan, P. C., & Conaco, C. (2019). Species variability in the response to elevated temperature of select corals in north-western Philippines. *Journal of the Marine Biological Association of the United Kingdom*, 99(6), 1273-1279.
- Dana, J. D. (1846). *United States Exploring Expedition during the years 1838-1842*. 7, 1–740.
- Devlin M.J. & Lourey M.J. (2000) Water quality - field and analytical procedures. In: L–tMotGB R, editor. Standard Operational Procedure, vol. 6. Townsville: Australian Institute of Marine Science.
- Dormann, C. F., Elith, J., Bacher, S., Buchmann, C., Carl, G., Carré, G., Marquéz, J. R. G., Gruber, B., Lafourcade, B., Leitão, P. J., Münkemüller, T., McClean, C., Osborne, P. E., Reineking, B., Schröder, B., Skidmore, A. K., Zurell, D., & Lautenbach, S. (2013). Collinearity: A review of methods to deal with it and a simulation study evaluating their performance. *Ecography*, 36(1), 27–46. <https://doi.org/10.1111/j.1600-0587.2012.07348.x>
- Edgar, R. C. (2010). Search and clustering orders of magnitude faster than BLAST. *Bioinformatics*, 26(19), 2460–2461. <https://doi.org/10.1093/bioinformatics/btq461>

- Ehrenberg, C. (1834). *Contributions to the physiological knowledge of the coral animals in general, and especially of the Red Sea, together with an experiment on the physiological systematics of the same. I*, 225–380.
- Engelen, A. H., Aires, T., Vermeij, M. J. A., Herndl, G. J., Serrão, E. A., & Frade, P. R. (2018). Host Differentiation and Compartmentalization of Microbial Communities in the Azooxanthellate Cupcorals *Tubastrea coccinea* and *Rhizopsammia goesi* in the Caribbean. *Frontiers in Marine Science*, 5, 391. <https://doi.org/10.3389/fmars.2018.00391>
- Enriquez, S., Mendez, E. R., & Iglesias-Prieto, R. I. (2005). Multiple scattering on coral skeletons enhances light absorption by symbiotic algae. *Limnology and Oceanography*, 50(4), 1025–1032. <https://doi.org/10.4319/lo.2005.50.4.1025>
- Fabricius, K. E. (2005). Effects of terrestrial runoff on the ecology of corals and coral reefs: Review and synthesis. *Marine Pollution Bulletin*, 50(2), 125–146. <https://doi.org/10.1016/j.marpolbul.2004.11.028>
- Frade, P. R., Bongaerts, P., Winkelhagen, A. J. S., Tonk, L., Bak, R. P. M. (2008). In situ photobiology of corals over large depth ranges: A multivariate analysis on the roles of environment, host, and algal symbiont. *Limnology and Oceanography*, 53(6), 2711-2723. <https://doi.org/10.4319/LO.2008.53.6.2711>
- Frade, P. R., Roll, K., Bergauer, K., & Herndl, G. J. (2016a). Archaeal and bacterial communities associated with the surface mucus of Caribbean corals differ in their degree of host specificity and community turnover over reefs. *PLOS ONE*, 11(1), e0144702. <https://doi.org/10.1371/journal.pone.0144702>
- Frade, P. R., Schwaninger, V., Glasl, B., Sintes, E., Hill, R. W., Simó, R., & Herndl, G. J. (2016b). Dimethylsulfoniopropionate in corals and its interrelations with bacterial

- assemblages in coral surface mucus. *Environmental Chemistry*, 13(2), 252.
<https://doi.org/10.1071/EN15023>
- Frias-Lopez, J., L. Zerkle, A., T. Bonheyo, G., & W. Fouke, B. (2002). Partitioning of bacterial communities between seawater and healthy, black band diseased, and dead coral surfaces. *Applied and Environmental Microbiology*, 68(5), 2214–2228.
<https://doi.org/10.1128/AEM.68.5.2214-2228.2002>
- Garren, M., & Azam, F. (2012). Corals shed bacteria as a potential mechanism of resilience to organic matter enrichment. *The ISME Journal*, 6(6), 1159–1165.
<https://doi.org/10.1038/ismej.2011.180>
- Glasl, B., Bongaerts, P., Elisabeth, N. H., Hoegh-Guldberg, O., Herndl, G. J., & Frade, P. R. (2017). Microbiome variation in corals with distinct depth distribution ranges across a shallow–mesophotic gradient (15–85 m). *Coral Reefs*, 36(2), 447–452.
<https://doi.org/10.1007/s00338-016-1517-x>
- Glasl, B., Bourne, D. G., Frade, P. R., Thomas, T., Schaffelke, B., & Webster, N. S. (2019a). Microbial indicators of environmental perturbations in coral reef ecosystems. *Microbiome*, 7(1), 94. <https://doi.org/10.1186/s40168-019-0705-7>
- Glasl, B., Herndl, G. J., & Frade, P. R. (2016). The microbiome of coral surface mucus has a key role in mediating holobiont health and survival upon disturbance. *The ISME Journal*, 10(9), 2280–2292. <https://doi.org/10.1038/ismej.2016.9>
- Glasl, B., Smith, C. E., Bourne, D. G., & Webster, N. S. (2019b). Disentangling the effect of host-genotype and environment on the microbiome of the coral *Acropora tenuis*. *PeerJ*, 7, e6377. <https://doi.org/10.7717/peerj.6377>
- Grottoli, A. G., Dalcin Martins, P., Wilkins, M. J., Johnston, M. D., Warner, M. E., Cai, W.-J., Melman, T. F., Hoadley, K. D., Pettay, D. T., Levas, S., & Schoepf, V. (2018). Coral

- physiology and microbiome dynamics under combined warming and ocean acidification.
PLOS ONE, 13(1), e0191156. <https://doi.org/10.1371/journal.pone.0191156>
- Guppy, R., & Bythell, J. (2006). Environmental effects on bacterial diversity in the surface mucus layer of the reef coral *Montastraea faveolata*. *Marine Ecology Progress Series*, 328, 133–142. <https://doi.org/10.3354/meps328133>
- Hong, M.-J., Yu, Y.-T., Chen, C. A., Chiang, P.-W., & Tang, S.-L. (2009). Influence of species specificity and other factors on bacteria associated with the coral *Stylophora pistillata* in Taiwan. *Applied and Environmental Microbiology*, 75(24), 7797–7806. <https://doi.org/10.1128/AEM.01418-09>
- Howells, E. J., Beltran, V. H., Larsen, N. W., Bay, L. K., Willis, B. L., & van Oppen, M. J. H. (2012). Coral thermal tolerance shaped by local adaptation of photosymbionts. *Nature Climate Change*, 2(2), 116–120. <https://doi.org/10.1038/nclimate1330>
- Iglesias-Prieto, R. & Trench R. K. (1997) Acclimation and adaptation to irradiance in symbiotic dinoflagellates. II. Response of chlorophyll-protein complexes to different photon-flux densities. *Marine Biology*, 130(1), 23-33.
- Johnson, J. B., & Omland, K. S. (2004). Model selection in ecology and evolution. *Trends in Ecology & Evolution*, 19(2), 101–108. <https://doi.org/10.1016/j.tree.2003.10.013>
- Kelly, L. W., Williams, G. J., Barott, K. L., Carlson, C. A., Dinsdale, E. A., Edwards, R. A., Haas, A. F., Haynes, M., Lim, Y. W., McDole, T., Nelson, C. E., Sala, E., Sandin, S. A., Smith, J. E., Vermeij, M. J. A., Youle, M., & Rohwer, F. (2014). Local genomic adaptation of coral reef-associated microbiomes to gradients of natural variability and anthropogenic stressors. *Proceedings of the National Academy of Sciences*, 111(28), 10227–10232. <https://doi.org/10.1073/pnas.1403319111>

- Klaus, J. S., Janse, I., Heikoop, J. M., Sanford, R. A., & Fouke, B. W. (2007). Coral microbial communities, zooxanthellae and mucus along gradients of seawater depth and coastal pollution. *Environmental Microbiology*, 9(5), 1291–1305. <https://doi.org/10.1111/j.1462-2920.2007.01249.x>
- Koren, O., & Rosenberg, E. (2006). Bacteria Associated with Mucus and Tissues of the Coral *Oculina patagonica* in Summer and Winter. *Applied and Environmental Microbiology*, 72(8), 5254–5259. <https://doi.org/10.1128/AEM.00554-06>
- LaJeunesse, T. C., Parkinson, J. E., Gabrielson, P. W., Jeong, H. J., Reimer, J. D., Voolstra, C. R., & Santos, S. R. (2018). Systematic revision of Symbiodiniaceae highlights the antiquity and diversity of coral endosymbionts. *Current Biology*, 28(16), 2570-2580.
- Lane, D. J. (1991). *16S/23S rRNA sequencing*. 115–175.
- Larsson, J. (2020). eulerr: Area-proportional euler and venn diagrams with ellipses. R package version 6.1.0, <https://cran.r-project.org/package=eulerr>.
- Lee, S. T. M., Davy, S. K., Tang, S.-L., Fan, T.-Y., & Kench, P. S. (2015). Successive shifts in the microbial community of the surface mucus layer and tissues of the coral *Acropora muricata* under thermal stress. *FEMS Microbiology Ecology*, 91(12), fiv142. <https://doi.org/10.1093/femsec/fiv142>
- Legendre, P., & Anderson, M. J. (1999). Distance-based redundancy analysis: testing multispecies responses in multifactorial ecological experiments. *Ecological Monographs*, 69(1), 1–24. [https://doi.org/10.1890/0012-9615\(1999\)069\[0001:DBRATM\]2.0.CO;2](https://doi.org/10.1890/0012-9615(1999)069[0001:DBRATM]2.0.CO;2)
- Lema, K. A., Willis, B. L., & Bourne, D. G. (2012). Corals form characteristic associations with symbiotic nitrogen-fixing bacteria. *Applied and Environmental Microbiology*, 78(9), 3136–3144. <https://doi.org/10.1128/AEM.07800-11>

- Lesser M.P., Mazel C.H., Gorbunov M.Y., Falkowski P.G. (2004). Discovery of symbiotic nitrogen-fixing cyanobacteria in corals. *Science*, 305, 997–1000
- Li, J., Chen, Q., Long, L.-J., Dong, J.-D., Yang, J., & Zhang, S. (2015). Bacterial dynamics within the mucus, tissue and skeleton of the coral *Porites lutea* during different seasons. *Scientific Reports*, 4(1), 7320. <https://doi.org/10.1038/srep07320>
- Little, A. F. (2004). Flexibility in algal endosymbioses shapes growth in reef corals. *Science*, 304(5676), 1492–1494. <https://doi.org/10.1126/science.1095733>
- Littman, R. A., Bourne, D. G., & Willis, B. L. (2010). Responses of coral-associated bacterial communities to heat stress differ with *Symbiodinium* type on the same coral host. *Molecular Ecology*, 19(9), 1978–1990. <https://doi.org/10.1111/j.1365-294X.2010.04620.x>
- Littman, R. A., Willis, B. L., Pfeffer, C., & Bourne, D. G. (2009a). Diversities of coral-associated bacteria differ with location, but not species, for three acroporid corals on the Great Barrier Reef: Diversity of coral-associated bacteria. *FEMS Microbiology Ecology*, 68(2), 152–163. <https://doi.org/10.1111/j.1574-6941.2009.00666.x>
- Littman, R., Willis, B., & Bourne, D. G. (2009b). Bacterial communities of juvenile corals infected with different *Symbiodinium* (dinoflagellate) clades. *Marine Ecology Progress Series*, 389, 45–59. <https://doi.org/10.3354/meps08180>
- Magoc, T., & Salzberg, S. L. (2011). FLASH: Fast length adjustment of short reads to improve genome assemblies. *Bioinformatics*, 27(21), 2957–2963. <https://doi.org/10.1093/bioinformatics/btr507>
- McDevitt-Irwin, J. M., Baum, J. K., Garren, M., & Vega Thurber, R. L. (2017). Responses of coral-associated bacterial communities to local and global stressors. *Frontiers in Marine Science*, 4, 262. <https://doi.org/10.3389/fmars.2017.00262>

- McMurdie, P. J., & Holmes, S. (2013). phyloseq: An R package for reproducible interactive analysis and graphics of microbiome census data. *PLoS ONE*, 8(4), e61217. <https://doi.org/10.1371/journal.pone.0061217>
- Morris, J. J., Johnson, Z. I., Szul, M. J., Keller, M., & Zinser, E. R. (2011). Dependence of the cyanobacterium *Prochlorococcus* on hydrogen peroxide scavenging microbes for growth at the ocean's surface. *PLoS ONE*, 6(2), e16805. <https://doi.org/10.1371/journal.pone.0016805>
- Muller-Parker, G., D'Elia, C. F., & Cook, C. B. (2015). Interactions between corals and their symbiotic algae. In C. Birkeland (Ed.), *Coral Reefs in the Anthropocene* (pp. 99–116). Springer Netherlands. https://doi.org/10.1007/978-94-017-7249-5_5
- Neave, M. J., Apprill, A., Ferrier-Pagès, C., & Voolstra, C. R. (2016). Diversity and function of prevalent symbiotic marine bacteria in the genus *Endozoicomonas*. *Applied Microbiology and Biotechnology*, 100(19), 8315–8324. <https://doi.org/10.1007/s00253-016-7777-0>
- Neave, M. J., Rachmawati, R., Xun, L., Michell, C. T., Bourne, D. G., Apprill, A., & Voolstra, C. R. (2017). Differential specificity between closely related corals and abundant *Endozoicomonas* endosymbionts across global scales. *The ISME Journal*, 11(1), 186–200. <https://doi.org/10.1038/ismej.2016.95>
- Oksanen J., Blanchet F.G., Kindt R., Legendre P., Minchin P.R., O'Hara R.B., Simpson G.L., Solymos P., Stevens M.H.H., Wagner H. (2013). *vegan: community ecology package*. R package version 2.0-9. Available at <http://CRAN.R-project.org/package=vegan>.
- Pernice, M., Meibom, A., van den Heuvel, A., Kopp, C., Domart-Coulon, I., Hoegh-Guldberg, O., & Dove, S. (2012). A single-cell view of ammonium assimilation in coral–dinoflagellate symbiosis. *The ISME Journal*, 6(7), 1314–1324. <https://doi.org/10.1038/ismej.2011.196>

- 822
- 823 Pernice, M., Dunn, S.R., Tonk, L., Dove, S., Domart-Coulon, I., Hoppe, P., Schintlmeister, A.,
824 Wagner, M., & Meibom, A. (2015) A nanoscale secondary ion mass spectrometry study
825 of dinoflagellate functional diversity in reef-building corals. *Environmental Microbiology*
826 17(10), 3570–3580. doi:10.1111/1462-2920.12518
- 827
- 828 Pogoreutz C, Radecker N, Cardenas A, Gardes A, Wild C, and Voolstra CR. 2018. Dominance
829 of Endozoicomonas bacteria throughout coral bleaching and mortality suggests structural
830 inflexibility of the Pocillopora verrucosa microbiome. *Ecology and Evolution* 8:2240-
831 2252. 10.1002/ece3.3830
- 832
- 833 Pollock, F. J., Lamb, J. B., Field, S. N., Heron, S. F., Schaffelke, B., Shedrawi, G., Bourne, D.
834 G., & Willis, B. L. (2014). Sediment and turbidity associated with offshore dredging
835 increase coral disease prevalence on nearby reefs. *PLoS ONE*, 9(7), e102498.
836 <https://doi.org/10.1371/journal.pone.0102498>
- 837
- 838 Pollock, F. J., McMinds, R., Smith, S., Bourne, D. G., Willis, B. L., Medina, M., Thurber, R. V.,
839 & Zaneveld, J. R. (2018). Coral-associated bacteria demonstrate phylosymbiosis and
840 cophylogeny. *Nature Communications*, 9(1), 4921. [https://doi.org/10.1038/s41467-018-](https://doi.org/10.1038/s41467-018-07275-x)
841 [07275-x](https://doi.org/10.1038/s41467-018-07275-x)
- 842
- 843 Quigley, K. M., Alvarez-Roa, C., Torda, G., Bourne, D. G., & Willis, B. L. (2019). Co-dynamics
844 of Symbiodiniaceae and bacterial populations during the first year of symbiosis with
845 *Acropora tenuis* juveniles. *MicrobiologyOpen*. <https://doi.org/10.1002/mbo3.959>
- 846
- 847 R Core Team. (2015). *R: A language and environment for statistical computing*. R foundation
848 for statistical computing, Vienna, Austria. Available at <http://www.R-project.org/>.
- 849
- 850 Radecker, N., Pogoreutz, C., Voolstra, C. R., Wiedenmann, J., & Wild, C. (2015). Nitrogen
851 cycling in corals: The key to understanding holobiont functioning? *Trends in*
852 *Microbiology*, 23(8), 490–497. <https://doi.org/10.1016/j.tim.2015.03.008>

- 853
- 854 Rocker, M. M., Willis, B. L., & Bay, L. K. (2012). Thermal stress-related gene expression in
- 855 corals with different Symbiodinium types. *In Proc 12th Int Coral Reef Symp* 9, 1-5).
- 856
- 857 Rosado, P. M., Leite, D. C. A., Duarte, G. A. S., Chaloub, R. M., Jospin, G., Nunes da Rocha,
- 858 U., P. Saraiva, J., Dini-Andreote, F., Eisen, J. A., Bourne, D. G., & Peixoto, R. S. (2019).
- 859 Marine probiotics: Increasing coral resistance to bleaching through microbiome
- 860 manipulation. *The ISME Journal*, 13(4), 921–936. [https://doi.org/10.1038/s41396-018-](https://doi.org/10.1038/s41396-018-0323-6)
- 861 [0323-6](https://doi.org/10.1038/s41396-018-0323-6)
- 862
- 863 Rohwer, F., Seguritan, V., Azam, F., & Knowlton, N. (2002). Diversity and distribution of coral-
- 864 associated bacteria. *Marine Ecology Progress Series*, 243, 1-10.
- 865
- 866 Schloss, P. D., Westcott, S. L., Ryabin, T., Hall, J. R., Hartmann, M., Hollister, E. B.,
- 867 Lesniewski, R. A., Oakley, B. B., Parks, D. H., Robinson, C. J., Sahl, J. W., Stres, B.,
- 868 Thallinger, G. G., van Horn, D. J., & Weber, C. F. (2009). Introducing mothur: Open-
- 869 Source, Platform-Independent, Community-Supported Software for Describing and
- 870 Comparing Microbial Communities. *Applied and Environmental Microbiology*, 75(23),
- 871 7537–7541. <https://doi.org/10.1128/AEM.01541-09>
- 872
- 873 Shashar, N., Cohen, Y., & Loya, Y. (1993). Extreme diel fluctuations of oxygen in diffusive
- 874 boundary layers surrounding stony corals. *The Biological Bulletin*, 185(3), 455–461.
- 875 <https://doi.org/10.2307/1542485>
- 876
- 877 Shaver, E. C., Shantz, A. A., McMinds, R., Burkepile, D. E., Vega Thurber, R. L., & Silliman, B.
- 878 R. (2017). Effects of predation and nutrient enrichment on the success and microbiome of
- 879 a foundational coral. *Ecology*, 98(3), 830–839. <https://doi.org/10.1002/ecy.1709>
- 880
- 881 Siboni, N., Ben-Dov, E., Sivan, A., & Kushmaro, A. (2008). Global distribution and diversity of
- 882 coral-associated *Archaea* and their possible role in the coral holobiont nitrogen cycle.

- Environmental Microbiology*, 10(11), 2979–2990. <https://doi.org/10.1111/j.1462-2920.2008.01718.x>
- Ssekagiri, A., Sloan, W. T., & Ijaz, U. Z. (2017). microbiomeSeq: An R package for analysis of microbial communities in an environmental context. <https://doi.org/10.13140/rg.2.2.17108.71047>
- Sweet, M. J., Croquer, A., & Bythell, J. C. (2011). Bacterial assemblages differ between compartments within the coral holobiont. *Coral Reefs*, 30(1), 39–52. <https://doi.org/10.1007/s00338-010-0695-1>
- Tanaka, Y., Ogawa, H., & Miyajima, T. (2010). Effects of nutrient enrichment on the release of dissolved organic carbon and nitrogen by the scleractinian coral *Montipora digitata*. *Coral Reefs*, 29(3), 675–682. <https://doi.org/10.1007/s00338-010-0639-9>
- Thompson, J. R., Rivera, H. E., Closek, C. J., & Medina, M. (2015). Microbes in the coral holobiont: Partners through evolution, development, and ecological interactions. *Frontiers in Cellular and Infection Microbiology*, 4. <https://doi.org/10.3389/fcimb.2014.00176>
- Thurber, R. V., Willner-Hall, D., Rodriguez-Mueller, B., Desnues, C., Edwards, R. A., Angly, F., Dinsdale, E., Kelly, L., & Rohwer, F. (2009). Metagenomic analysis of stressed coral holobionts. *Environmental Microbiology*, 11(8), 2148–2163. <https://doi.org/10.1111/j.1462-2920.2009.01935.x>
- Tremblay, P., Weinbauer, M. G., Rottier, C., Guérardel, Y., Nozais, C., & Ferrier-Pagès, C. (2011). Mucus composition and bacterial communities associated with the tissue and skeleton of three scleractinian corals maintained under culture conditions. *Journal of the Marine Biological Association of the United Kingdom*, 91(3), 649–657. <https://doi.org/10.1017/S002531541000130X>

- Turner, S., Pryer, K. M., Miao, V. P., & Palmer, J. D. (1999). Investigating deep phylogenetic relationships among cyanobacteria and plastids by small subunit rRNA sequence analysis 1. *Journal of Eukaryotic Microbiology*, 46(4), 327-338.
- Ulstrup, K. E., & Van Oppen, M. J. H. (2003). Geographic and habitat partitioning of genetically distinct zooxanthellae (Symbiodinium) in Acropora corals on the Great Barrier Reef. *Molecular Ecology*, 12(12), 3477-3484.
- Uysal, Z., & Köksalan, İ. (2006). The annual cycle of Synechococcus (cyanobacteria) in the northern Levantine Basin shelf waters (Eastern Mediterranean). *Marine Ecology*, 27(3), 187-197.
- van Oppen, M. J., Palstra, F. P., Piquet, A. M. T., & Miller, D. J. (2001). Patterns of coral–dinoflagellate associations in Acropora: significance of local availability and physiology of Symbiodinium strains and host–symbiont selectivity. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 268(1478), 1759-1767.
- Wada, N., Ishimochi, M., Matsui, T., Pollock, F. J., Tang, S. L., Ainsworth, T. D., Willis, B. L., Mano, N., & Bourne, D. G. (2019). Characterization of coral-associated microbial aggregates (CAMAs) within tissues of the coral *Acropora hyacinthus*. *Scientific Reports*, 9(1), 14662. doi:10.1038/s41598-019-49651-7
- Wang, Q., Garrity, G. M., Tiedje, J. M., & Cole, J. R. (2007). Naive Bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy. *Applied and Environmental Microbiology*, 73(16), 5261–5267. <https://doi.org/10.1128/AEM.00062-07>
- Weber, M., de Beer, D., Lott, C., Polerecky, L., Kohls, K., Abed, R. M. M., Ferdelman, T. G., & Fabricius, K. E. (2012). Mechanisms of damage to corals exposed to sedimentation. *Proceedings of the National Academy of Sciences*, 109(24), E1558–E1567. <https://doi.org/10.1073/pnas.1100715109>

- Work, T., & Aeby, G. (2014). Microbial aggregates within tissues infect a diversity of corals throughout the Indo-Pacific. *Marine Ecology Progress Series* 500, 1-9.
- Yang, S., Sun, W., Zhang, F., & Li, Z. (2013). Phylogenetically diverse denitrifying and ammonia-oxidizing bacteria in corals *Alcyonium gracillimum* and *Tubastraea coccinea*. *Marine Biotechnology*, 15(5), 540–551. <https://doi.org/10.1007/s10126-013-9503-6>
- Yellowlees, D., Rees, T. A. V., & Leggat, W. (2008). Metabolic interactions between algal symbionts and invertebrate hosts. *Plant, Cell & Environment*, 31(5), 679–694. <https://doi.org/10.1111/j.1365-3040.2008.01802.x>
- Yilmaz, P., Parfrey, L. W., Yarza, P., Gerken, J., Priesse, E., Quast, C., Schweer, T., Peplies, J., Ludwig, W., & Glöckner, F. O. (2014). The SILVA and “All-species Living Tree Project (LTP)” taxonomic frameworks. *Nucleic Acids Research*, 42(D1), D643–D648. <https://doi.org/10.1093/nar/gkt1209>

Figure 1

Compartment-specific microbiome composition of *Acropora tenuis* and *Acropora millepora*.

Microbial community composition (size indicates mean relative abundance and color represents standard error) resolved for the surface mucus layer and tissue of two *Acropora* coral species (*A. tenuis* and *A. millepora*), and surrounding seawater, based on partial 16S rRNA gene amplicon sequencing. Only the 25 most abundant families across all samples are represented.




Figure 2

Alpha diversity measures of coral mucus, coral tissue, and seawater microbiomes.

(A) Venn diagram displaying the number of shared, unique, and ubiquitous zOTUs among mucus, tissue and seawater microbiomes. Two *Acropora* species (*A. tenuis* and *A. millepora*) are pooled for the tissue and mucus microbiomes. (B) zOTU richness and (C) Shannon diversity index of microbiomes associated with tissue and mucus of *A. millepora* and *A. tenuis*, as well as with seawater.




Figure 3

Compositional variability of microbiomes associated with distinct coral-compartment and the ambient seawater.

Two-dimensional non-metric multidimensional scaling (nMDS) ordination depicting variation in microbial community structure between coral compartments (mucus and tissue) of *Acropora tenuis* and *Acropora millepora*, and seawater samples. “k” is the number of dimensions.




Figure 4

Environmental and physiological drivers of the *Acropora tenuis* and *Acropora millepora* microbiomes.

Distance-based redundancy analysis (db-RDA) of the relationship between environmental/physiological variables and the relative abundance of bacteria in (A) mucus and (B) tissue microbiome of *A. millepora*, and (C) mucus and (D) tissue microbiome of *A. tenuis*. Arrow length indicates the strength of the correlation between the variables and the samples (note only significant variables are shown). The selected variables explain a total of (A) 14.98%, (B) 16.44%, (C) 10.63% and (D) 10.97% of the observed variance, respectively. Environmental/physiological variables represented are the sum of nitrite and nitrate concentrations (NO₂/NO₃), particulate organic carbon (POC), total suspended solids (TSS), ammonium concentration (NH₄), chlorophyll *a* concentration (Chl*a*) in seawater and Symbiodiniaceae density normalized to protein contents (Zoox/Pro) of coral tissue.




Figure 5

Bacterial taxa significantly correlated with environmental and physiological variables.

Pearson's coefficient based correlation matrix between the 20 most abundant bacterial families and environmental/physiological variables having a significant effect on the microbiome associated to tissue and surface mucus of (A) *Acropora millepora* ($n_{\text{tissue}} = 24$, $n_{\text{mucus}} = 24$) and (B) *Acropora tenuis* ($n_{\text{tissue}} = 30$, $n_{\text{mucus}} = 28$). Significant correlations indicated by asterisks at different levels of significance (* for $p < 0.05$, ** for $p < 0.01$, *** for $p < 0.001$) after correction for multiple comparisons (using Benjamini-Hochberg correction).

Environmental/physiological variables represented are the sum of nitrite and nitrate concentrations (NO_2/NO_3), particulate organic carbon (POC), total suspended solids (TSS), ammonium concentration (NH_4), chlorophyll a concentration (Chla) in seawater and Symbiodiniaceae density normalized to protein contents (Zoox/Pro) of coral tissue.

