

The evolution of feeding within Euchelicerata: data from the fossil groups Eurypterida and Trigonotarbida illustrate possible evolutionary pathways

Carolin Haug^{1,2}

¹ Department of Biology II, Ludwig-Maximilians-Universität München, Planegg-Martinsried, Germany

² GeoBio-Center, Ludwig-Maximilians-Universität München, Munich, Germany

ABSTRACT

When the evolution of Euarthropoda is discussed, often the lineage of Chelicerata s. str. is assumed to be the more 'primitive' or 'basal' part of the tree, especially when compared to the other major lineage, Mandibulata. This claimed primitiveness is (at least partly) based on the assumption that different morphological structures are still in an ancestral state and did not evolve any further. One of these sets of structures is the feeding apparatus, which has been stated to be highly advanced in Mandibulata, but not 'properly' developed, or at least not to such a high degree, within Chelicerata s. str. In this study, I reinvestigate the feeding apparatus of different ingroups of Euchelicerata, with a focus on assumed 'primitive' groups such as Eurypterida and Trigonotarbida. The basis of this study is a large amount of material from different museum collections, with fossils with the entire feeding apparatuses being exceptionally well preserved. Based on high-resolution micro-photography and three-dimensional imaging, it is possible to resolve fine details of the feeding apparatuses. The results make clear that the feeding apparatuses of different ingroups of Euchelicerata are highly specialised and often possess morphological structures comparable to those of the feeding apparatuses of representatives of Mandibulata, apparently convergently evolved. Though the reconstruction of the evolution of the feeding apparatus within Euchelicerata is to a certain degree hampered by unclear phylogenetic relationships, there was clearly a shortening of the feeding apparatus from posterior (i.e. only the anterior appendages being involved in the feeding apparatus), probably linked to the colonisation of land in Arachnida.

Submitted 2 October 2019

Accepted 20 July 2020

Published 13 August 2020

Corresponding author

Carolin Haug, carolin.haug@palaeo-evo-devo.info

Academic editor

Bruce Lieberman

Additional Information and Declarations can be found on page 22

DOI 10.7717/peerj.9696

© Copyright 2020 Haug

Distributed under Creative Commons CC-BY 4.0

OPEN ACCESS

Subjects Evolutionary Studies, Paleontology, Zoology

Keywords Eurypterida, Trigonotarbida, Arachnida, Character evolution, Mouthparts, Endites, Rhynie chert

INTRODUCTION

The origin of Euarthropoda, including the extant groups Myriapoda, Insecta, Eucrustacea (sensu [Walossek, 1999](#)) and Chelicerata s. str. (sensu [Maas et al., 2004](#)) and different fossil representatives, lies more than half a billion years ago (e.g. [Daley et al., 2018](#) and references therein). In this long time, an enormous species richness and morphological

diversity evolved in the different ingroups. Within Euarthropoda, the lineage of Chelicerata s. str. is often somehow treated as the more ‘basal’ or ‘primitive’ side of the tree (e.g. [Damen, 2002](#); [Dunlop, 2011](#); see the discussion in [Krell & Cranston \(2004\)](#) or [Omland, Cook & Crisp \(2008\)](#) on the topic why there is nothing like a more ‘basal’ side of the tree) and is thought to be more ancestral, especially in comparison to Mandibulata, the other major lineage within Euarthropoda ([Damen, 2002](#)). This view is most extremely applied to Xiphosurida (horseshoe ‘crabs,’ for example [Sekiguchi, 1988](#); [Malakhov, 2010](#); [Williams, 2019](#)), but also to now extinct groups such as Eurypterida (sea scorpions) or the spider-like group of Trigonotarbita ([Resh & Cardé, 2003](#)).

Especially horseshoe ‘crabs’ (though the old fashioned ‘sword tails’ would be less ambiguous) have often been treated as ‘living fossils’ (a very unscientific term, see also discussion in [Wagner et al. \(2017a\)](#); for attempts of more scientific approaches, see [Kin & Błażejowski \(2014\)](#) and [Bennett, Sutton & Turvey \(2018\)](#)). They have also been assumed to be a kind of proxy for the early terrestrialisation within Euchelicerata ([Martin, 2017](#)). Yet, this interpretation is most likely incorrect. It is unlikely that the stem species of Euchelicerata was already amphibious, hence the terrestrial behaviour of modern representatives of Xiphosurida has most probably evolved independently (see [Lamsdell \(2016\)](#) on independent evolution of non-marine life habits; see also recent discussion about the phylogenetic position of Xiphosurida and the consequences on the evolution of terrestrialisation: [Ballesteros & Sharma, 2019](#); [Giribet & Edgecombe, 2019](#); [Lozano-Fernandez et al., 2019](#)). It needs to be emphasised that modern representatives of Xiphosurida are not direct proxies for the ancestor of Euchelicerata, but possess their own specialisations (as all living groups do).

One reason why the lineage of Chelicerata s. str. is assumed to be primitive might be the organisation of the feeding apparatus in most representatives. In different textbooks, the impression is given that these forms lack ‘proper’ mouthparts, while ingroups of Mandibulata have a ‘full’ set of mouthparts. [Gruner \(1993](#), as an example for an important German text book) states that representatives of Chelicerata s. str. bear structures for acting in feeding but would lack true antagonistic jaws. He also states that mostly only the second pair of appendages and rarely the third and fourth one is incorporated in the feeding apparatus in most forms. This statement most likely refers to the state in scorpions and other arachnids, but appears to ignore the state in Xiphosurida where appendage pairs 1–7 are involved in the feeding apparatus.

Also in other cases the description of the feeding apparatuses appears to refer to only certain ingroups. For example the statement that there are always three pairs of mouthparts in Mandibulata ([Gruner, 1993](#)) is clearly correct for Insecta, but ignores maxillipeds in Chilopoda or Decapoda, or complex thoracic feeding apparatuses as for example in Branchiopoda (e.g., in fairy shrimps).

Hence, such textbook statements are at best oversimplified and tend to be based more on general assumptions than on direct observations. To improve this situation, I provide here functional morphological interpretations of the feeding apparatuses (= all external structures involved in feeding) of two supposedly ‘primitive’ chelicerate groups, namely of Eurypterida and of Trigonotarbita and present them in the context of evolution in

Euchelicerata. With this, I aim at providing a framework to evaluate the presumed ‘primitiveness’ of the feeding apparatuses of modern representatives of Euchelicerata.

MATERIALS AND METHODS

Material

For this study, fossil material from different museum collections was investigated, including the invertebrate palaeontology collection of the Yale Peabody Museum of Natural History, New Haven (YPM IP), the Natural History Museum, London (NHM), the Naturhistoriska riksmuseet, Stockholm (NRM), the Harvard Museum of Comparative Zoology, Cambridge (MCZ) and the Royal Ontario Museum, Toronto (ROM). Every specimen of Eurypterida and Trigonotarbida in these collections was briefly inspected, and those preserving morphological details of interest for this study were documented (see “Methods” below).

The specimens of Eurypterida in the YPM IP collections had been collected by a private collector, Samuel J. Czurca, from late Silurian deposits in New York State, USA, and southern Ontario, Canada, and donated by him to the YPM; it is by far the largest collection of sea scorpions worldwide. The sea scorpions in the NHM collections originate from different late Silurian deposits: few specimens come from New York State and from Scotland. Also one sea scorpion from the MCZ collections comes from the Silurian of Scotland. Most specimens in the NHM collections come from Estonia, more precisely from the village Rootsiküla on the island of Saaremaa (also called Ösel). From the same Estonian locality, also many specimens of Eurypterida in the NRM collections and most in the MCZ collections originate, as well as from the island of Gotland, Sweden. The specimens from Saaremaa/Ösel and Gotland are insofar exceptional as of most of these the surrounding limestone matrix had been dissolved already in the 19th century by [Holm \(1896, 1898\)](#), resulting in isolated specimens with only their cuticle being preserved. Subsequently, the specimens had either been mounted between two large cover slips, or between a microscopic slide and a cover slip (dry or with Canada balsam), or they have been fully embedded in resin. All these methods allow to access dorsal as well as ventral structures ([Holm, 1898](#); [Selden, 1981](#)).

The specimens of Trigonotarbida in the ROM collections come from the Upper Carboniferous Mazon Creek, Carbondale Formation, IL, USA. The NHM collections houses specimens of Trigonotarbida preserved in Rhynie Chert, Lower Devonian of Scotland, the rest of the material stems from the British Coal Measures, Upper Carboniferous.

Comparative material of extant species came from the former teaching collection of the University of Ulm (now at the University of Rostock) and the teaching collection of LMU Munich.

Methods

For the documentation of the specimens different macro- and microphotographic methods were used. All specimens were documented in their corresponding museum collections.

The Rhynie Chert specimens were documented on a Nikon Eclipse microscope at the Imaging and Analysis Centre of the NHM. Of each specimen an image stack with shifting focus was recorded to overcome the limitations in depth of field. For larger specimens several image stacks of adjacent and partly overlapping areas of the specimens were recorded to overcome limitations in field of view. From the image stacks, stereo images were recorded according to the protocol of [Haug et al. \(2012a\)](#).

The other fossil specimens were documented with a macro-photographic setup with a Canon Rebel T3i and an EF-S 18–55 mm lens or an MP-E 65 mm macro lens, depending on the size of the structures of interest. For illumination, a Canon MT 24-EX Macro Twin Flash or two Yongnuo Digital Speedlite YN560 flash lights were used. To enhance the contrast between fossil and matrix and reduce reflections, a polarisation filter was placed in front of the lens, and perpendicular filter foils were placed in front of the flashlights, resulting in cross-polarised light ([Haug et al., 2011a](#); [Kerp & Bomfleur, 2011](#)). Specimens mounted on cover slips or microscopic slides were placed at some distance to a white background to avoid shadows. Also of the non-Rhynie Chert specimens image stacks were recorded; of these, high-resolution, entirely sharp images were calculated with Combine ZM/ZP and Photoshop CS3 (for details, see [Haug, Haug & Ehrlich, 2008](#); [Haug et al., 2011a](#)). Specimens with high relief were additionally documented from different angles by moving the camera around the specimen; from matching image pairs, stereo images were produced.

Of some images, additional colour-marked versions were produced. For this purpose, the images were desaturated and the histogram was optimised before the important structures were colour-marked. For the Rhynie Chert specimens, the red channel of the stereo images was deleted; afterwards, the images were processed in the same way as those from non-Rhynie Chert specimens.

The extant specimens were documented with different macro- and micro-photographic methods, including photography under cross-polarised light and under fluorescence conditions.

RESULTS

In the following, the feeding apparatuses of different representatives of Eurypterida and Trigonotarbidia are described. The aim of these descriptions is not to provide an in-depth species-diagnostic description, but instead to present the general characteristics of the feeding apparatuses of each group.

Feeding apparatus of representatives of Eurypterida

The appendages of seven segments clearly contribute to the feeding apparatus in Eurypterida: chelicerae, five subsequent pairs of legs (often called walking legs, but see discussion), and the plate-like appendage pair of the seventh post-ocular segment, the metastoma ([Figs. 1A, 2A–2C and 3A](#); for the origin of the metastoma, see also [Holm, 1898](#); [Dunlop, 1998](#), his fig. 4; [Dunlop & Selden, 1998](#); [Jeram, 1998](#)). The appendages are arranged in a circle around the central area of the feeding apparatus, with their median parts being very close together ([Figs. 2B, 2C, 3B and 4A](#)). The basipods (= coxae in chelicerate




Figure 1 Feeding apparatus of YPM_IP_216689, *Eurypterus lacustris* (Eurypteridae, Eurypterida), Ontario, Welland County; images colour-inverted to enhance contrast. (A) Overview. (B)–(G) Close-ups. (B) Stout spine on appendage 2. (C) Spines and setae on appendages 3 and 4. (D) and (E) Surface ornamentation on basipods. (F) Setae on appendages 2 and 3. (G) Spines and setae on (presumably) appendages 4 and 5.

Full-size [DOI: 10.7717/peerj.9696/fig-1](https://doi.org/10.7717/peerj.9696/fig-1)

terminology) of the appendages of post-ocular segment 6 (the leg pair right in front of the metastoma) are much larger than those of the other appendages, forming roughly a rectangle, but additionally with a pronounced endite (Fig. 2A). The metastoma covers part of these basipods as well as the median gap between the left and right appendage (Figs. 2B and 2C).

Post-ocular appendages 2–6, that is all post-cheliceral legs, bear a more or less pronounced armature on the endites protruding from the median edges of their basipods (Figs. 1B, 1C, 1F, 1G, 2C, 5A, 5B and 5C). The armature differs between the appendages. The spines on some of the basipods (probably of the further anterior ones, but there appear to be species-specific differences) are thinner than on others (Figs. 5A and 5B). The basipods of post-ocular appendage 6 bear the stoutest spines, the entire basipodal median edge appearing strongly sclerotised, recognisable from its very dark colour (Figs. 2A, 3B, 4B, 5D and 5E).

Also the armature on the same basipod is differentiated (also here with species-specific differences). The spines closer to the anterior edge of the basipod (in relation to the orientation of the body of the animal) are stouter and partly also longer, while those closer to the posterior edge of the basipod are thinner and smaller (Figs. 5B, 5C, 6A and 6B). This differentiation can be very pronounced, with very stout and large teeth, elongate and




Figure 2 Feeding apparatus of NHM I3406_2, *Eurypterus fischeri* (Eurypteridae, Eurypterida), Rootsiküla, Saaremaa, Estonia. (A) Overview. (B) Stereo image of central area, colour-inverted; use red-cyan glasses to view. (C) Colour-marked version of one half image of B; pink = chelicerae and setae; blue and green = basipods of appendages 2-6; cyan = metastoma (appendage 7); orange = spines. Abbreviations: a3-a7 = post-ocular appendages 3-7. [Full-size DOI: 10.7717/peerj.9696/fig-2](https://doi.org/10.7717/peerj.9696/fig-2)

pointed spines, and thin setae on the same basipod (Figs. 3C, 4C and 4D). The armature on the median edge of the basipods can occur in two (or possibly three?) rows (Figs. 6B and 6C). The remaining surface of the basipods can be covered with setae of other surface structures (Figs. 1D, 1E and 5F).

Feeding apparatus of representatives of Trigonotarbida

In Trigonotarbida, the appendages of three segments contribute to the feeding apparatus: chelicerae, pedipalps and first pair of walking legs (Figs. 7A-7D, 8A, 8B, 8F-8I and 9A-9D). The proximal parts of these appendages sit close together. The pedipalps are proximally armed with a row of short spines (Fig. 7D). The basipods of the first pair of walking appendages bear a prominent endite medially, which again bear setae (Figs. 7D, 8A, 8B, 8E, 8H, 8I, 9C, 9D, 10A and 10B). The endites are in some fossils positioned very closely together (Figs. 8C, 8D, 8H and 8I), in others further apart (Figs. 9C, 9D, 10A and 10B), which may point to a high movability during life. Additionally, there are two distinct oval fields of densely arranged short setae next to each other in the area between chelicerae and pedipalps (Figs. 10A-10D). Another less distinct field of similar setae appears to be positioned slightly posteriorly to the two oval fields (Figs. 10C and 10D). Based on the three-dimensional position information, these fields are probably situated on the hypostome (though often termed 'labrum' by different authors).




Figure 3 Feeding apparatus of NRM Ar 35344, *Eurypterus fischeri* (Eurypteridae, Eurypterida), Rootsiküla, Saaremaa, Estonia. (A) Overview. (B) Close-up of median edges of basipods. (C) Further close-up of the differentiated armature of the basipods. [Full-size](#) DOI: 10.7717/peerj.9696/fig-3

DISCUSSION

While this study focuses on the feeding apparatus of different groups of Euchelicerata, a proper character polarisation demands an outgroup comparison (see overview in [Fig. 11](#)). For this purpose it is especially important to include now extinct groups as those may exhibit character states no longer present in the extant fauna ([Donoghue et al., 1989](#); [Rust, 2007](#); [Edgecombe, 2010](#)).

Reconstructing the original chelicerate feeding apparatus

The supposed sister group to Euchelicerata is Pycnogonida ([Waloszek & Dunlop, 2002](#); [Dunlop, 2010](#); [Haug et al., 2012b](#); [Sharma et al., 2014](#) and references therein), together forming Chelicerata s. str.. While being quite speciose today, the general body organisation is the same in all extant species of Pycnogonida, with a highly reduced posterior body area (the terms pro- and opisthosoma cannot be applied here yet; see discussion in [Haug, Wagner & Haug \(2019\)](#)) and a strongly modified feeding apparatus adapted to suction feeding ([Fahrenbach & Arango, 2007](#); [Wagner et al., 2017b](#)). The fossil record of




Figure 4 Feeding apparatus of *Eurypterus fischeri* (Eurypteridae, Eurypterida), Rootsiküla, Saaremaa, Estonia. (A) and (B) NRM Ar35343. (A) Overview over central area. (B) Close-up of median edges of basipods with differentiated armature and very strongly sclerotised edges of post-ocular appendage 6. (C) and (D) NRM Ar48883. (C) Overview over anterior part of feeding apparatus; posterior part not preserved. (D) Close-up of differentiated armature on basipods, including broader and blunt teeth, pointed spines of different sizes and thin setae. [Full-size](#) DOI: 10.7717/peerj.9696/fig-4

Pycnogonida dates back to the Cambrian (Waloszek & Dunlop, 2002), so more than half a billion years ago. However, as already at this time the morphology appears derived (and partly also as the Cambrian representatives of Pycnogonida are exclusively larvae), it does not provide relevant information about the evolution of the feeding apparatus in the lineage towards Euchelicerata. It is necessary to take a look at the feeding apparatus in supposed early representatives of Megacheira, the so-called ‘short great-appendage arthropods,’ among which the sister group to Chelicerata s. str. is assumed by some authors (Chen, Waloszek & Maas, 2004; Maas et al., 2004; Haug et al., 2012b; Tanaka et al., 2013; Edgecombe & Legg, 2014; see also recent review by Dunlop & Lamsdell (2017)).




Figure 5 Parts of feeding apparatuses of Eurypterida. (A)–(C) MCZ PALI 131326, *Erettopterus bilobus* (Pterygotidae, Eurypterida), Lesmahagow, Lanarkshire, Scotland. (A) Overview of a series of basipods with pronounced armature and metastoma. (B) Colour-marked version of (A). (C) Close-up of basipod armature with differentiation between different basipods; colour-inverted to enhance contrast. (D)–(F) Isolated leg of post-ocular segment 6 of MCZ PALI 185687, *Erettopterus osiliensis* (Pterygotidae, Eurypterida), Saaremaa, Estonia. (D) Overview; note pronounced endite with spines. (E) Close-up of spines. (F) Close-up of surface structure of basipod.

[Full-size !\[\]\(666e09182d4cd268646ea700ea60dcdf_img.jpg\) DOI: 10.7717/peerj.9696/fig-5](https://doi.org/10.7717/peerj.9696/fig-5)

Recent reinvestigations of different of these species as well as older publications provide a sound basis for the reconstruction of the feeding apparatus in the ground pattern of Megacheira (*Bruton & Whittington, 1983; Chen, Waloszek & Maas, 2004; Haug et al., 2012b; Haug, Briggs & Haug, 2012; Yang et al., 2013*). At this evolutionary stage, the entire set of appendages was included into the feeding apparatus (possibly including the hypostome; *Fig. 12A*). While the first pair of appendages was specialised for grasping the prey, the second to last pair of appendages all exhibit the same general morphology,




Figure 6 Isolated basipod, presumably of leg of post-ocular segment 6, of NRM Ar31829, undetermined representative of Eurypterida, Visby, Gotland, Sweden. (A) Overview. (B) Close-up of spines; note different spine sizes and arrangement in different rows. (C) Further close-up of smaller spines with apparent row arrangement.

Full-size [DOI: 10.7717/peerj.9696/fig-6](https://doi.org/10.7717/peerj.9696/fig-6)

serving for swimming and feeding at the same time. While slight modifications in the size of the second appendage pair of *Leancoilia superlata* might indicate a starting specialisation to a mouthpart (Haug, Briggs & Haug, 2012), the ‘whole-body-feeding’ method represents the original condition in Megacheira and presumably in Chelicerata s. str.

This also holds true if assuming that short great-appendage arthropods branched off earlier along the evolutionary lineage (as suggested, for example by Legg (2013)). The polarisation of characters remains the same. Also several other early representatives of Euarthropoda possess similar characters in their feeding apparatuses, most prominently opposing basipods with strong spines (Legg, 2014; Bicknell et al., 2018a; Aria & Caron, 2017, 2019), but will not be discussed in detail here.

Shortening of the feeding apparatus in Euchelicerata

As already mentioned above, the feeding apparatus of Pycnogonida is adapted to suction feeding and highly derived already in the fossil forms. This specialised feeding apparatus is an autapomorphy of Pycnogonida (see also Waloszek & Dunlop, 2002 for character evolution in the lineage towards and within Pycnogonida).




Figure 7 Feeding apparatus of NHM In24671, *Palaeocharinus* sp. (Trigonotarbida), Rhynie Chert, Scotland. (A)–(C) Stereo images, use red-cyan glasses to view. (A) Overview of entire specimen. (B) Same as A, but with background virtually removed. (C) Close-up of feeding apparatus. (D) Colour-marked version of one half image of (C). Abbreviations: a3, appendage of post-ocular segment 3; ch, chelicera; ed, endite; pp, pedipalp; st, setae. [Full-size !\[\]\(fcc3264021d438d9732560e78099f674_img.jpg\) DOI: 10.7717/peerj.9696/fig-7](https://doi.org/10.7717/peerj.9696/fig-7)

In Euchelicerata the systematic affinities are still in a certain state of flux and the feeding apparatus cannot be properly reconstructed for all early fossil representatives. The feeding apparatus and in general the tagmosis pattern can be reconstructed to a certain extent for the species *Offacolus kingi* (Sutton et al., 2002), *Dibasterium durgae* (Briggs et al., 2012) and *Weinbergina opitzi* (Stürmer & Bergström, 1981; Moore, Briggs & Bartels, 2005), successively splitting off the evolutionary lineage towards the remaining representatives of Euchelicerata (= Prosomapoda sensu Lamsdell, 2013, though unclear if also including *W. opitzi* depending on the presence of exopods on the walking appendages; see also Selden, Lamsdell & Qi, 2015; Lamsdell, 2016; Dunlop & Lamsdell, 2017).

At the level of Euchelicerata, a characteristic division into an anterior and a posterior tagma usually termed prosoma and opisthosoma appears for the first time, supposedly as an autapomorphy for this group (for problems with the correspondence of prosoma and opisthosoma in different groups of Euchelicerata, see Haug, Wagner & Haug, 2019). The subdivision into these two distinct tagmata is characterised by dorsal and ventral




Figure 8 Parts of feeding apparatuses of Trigonotarbida. (A)–(E) *Palaeocharinus* sp., Rhynie Chert, Scotland. (A) and (B) NHM RC_019; ventro-lateral view on feeding apparatus. (B) Colour-marked version of one half image of (A). (C) and (D) NHM In27357; cross section through body at position of first walking appendages (appendages of post-ocular segment 3) with prominent endites. (D) Close-up of endites of (C). (E) NHM In24687; close-up of endite of first walking appendages. (F)–(I) NHM In31241, *Trigonotarbus johnsoni*, Coal Measures, UK. (F) and (G) Overview of entire specimen. (H) Stereo image of feeding apparatus. (I) Colour-marked version of one half image of H. Images (A), (C)–(E), (G) and (H) are stereo images, use red-cyan glasses to view. Abbreviations: a3, appendage of post-ocular segment 3; ch, chelicera; ed, endite; pp, pedipalp.

Full-size [DOI: 10.7717/peerj.9696/fig-8](https://doi.org/10.7717/peerj.9696/fig-8)

structural changes (see [Haug et al. \(2012c\)](#) and [Haug, Wagner & Haug \(2019\)](#) for characteristics of tagmata).

Dorsally, the segments of the anterior tagma form a uniform shield without any visible segmental borders. The segments of the posterior tagma possess separate tergites in the ground pattern of Euchelicerata. Ventrally, the tagmatisation is characterised by a ‘division of labour’ between the different appendages. While in short great-appendage arthropods all appendages were still involved in feeding, in the ground pattern of Euchelicerata the feeding apparatus becomes shorter; only the appendages of the anterior tagma contribute to the feeding apparatus, in addition to their locomotory (walking) function ([Fig. 12B](#)).




Figure 9 Feeding apparatus of ROMIP45532, undetermined trigontarbidan, Mazon Creek, USA. (A) and (B) Overview of entire specimen. (B) and (C) Stereo images, use red-cyan glasses to view. (C) and (D) Close-up of feeding apparatus and further walking appendages. (D) Colour-marked version of one half image of (C). Abbreviations: a3–6, appendages of post-ocular segments 3–6; ch, chelicera; ed, endite; pp, pedipalp; ste, sternum. [Full-size !\[\]\(fcc3264021d438d9732560e78099f674_img.jpg\) DOI: 10.7717/peerj.9696/fig-9](https://doi.org/10.7717/peerj.9696/fig-9)

The more posterior appendages serve for swimming; if they also possessed structures for oxygen exchange remains currently unclear (Sutton *et al.*, 2002; Dunlop & Lamsdell, 2017).

The appendage of post-ocular segment 7 (the pre-genital segment) differs morphologically in *O. kingi*, *D. durgae* and *W. opitzi*, being significantly smaller in the first two species (Sutton *et al.*, 2002; Briggs *et al.*, 2012). Yet, as *W. opitzi* possesses an appendage on this segment similar to the preceding ones with apparently locomotory function (Lehmann, 1956; Stürmer & Bergström, 1981; Moore, Briggs & Bartels, 2005), this was presumably the ground pattern condition for Euchelicerata. Hence, the feeding apparatus in the ground pattern of Euchelicerata most likely consists of (possibly the hypostome and) chelicerae and six pairs of walking appendages.




Figure 10 Feeding apparatus of NHM In24702, *Palaeocharinus* sp. (Trigonotarbita), Rhynie Chert, Scotland. (A) and (B) Overview. (A) Stereo image. (B) Colour-marked version of one half image of (A). (C) and (D) Close-up of fields with setae. (C) Stereo image. (D) Colour-marked version of one half image of (C); red = chelicerae; green = pedipalps; blue = first walking appendages; yellow = fields with setae.

Full-size [DOI: 10.7717/peerj.9696/fig-10](https://doi.org/10.7717/peerj.9696/fig-10)

Further modification of the feeding apparatus in Neochelicerata

Neochelicerata is an ingroup of Euchelicerata, the ‘crown group’ (most inclusive group with extant representatives). It has been characterised by [Haug, Wagner & Haug \(2019\)](#) as certain aspects of body organisation have not evolved in Euchelicerata yet, but are present in the ground pattern of Neochelicerata. The ground pattern condition of Neochelicerata is mainly reconstructed based on the morphology of Xiphosurida (sensu [Lamsdell, 2016](#)).

The body organisation in general and the feeding apparatus in particular are very similar in the ground pattern of Neochelicerata to that of Euchelicerata. Also here the anterior tagma dorsally bears a uniform shield. The tergites of the posterior tagma are fused into an entire dorsal shield, the thoracetrone, in Xiphosurida ([Anderson & Selden, 1997](#); [Lamsdell, 2016](#)). However, this condition appears to be an autapomorphy of Xiphosurida, while in the ground pattern of Neochelicerata the tergites were most probably still separate.




Figure 11 Coarse phylogenetic overview of the groups discussed in this article. For more details, see Haug, Wagner & Haug (2019). Full-size [DOI: 10.7717/peerj.9696/fig-11](https://doi.org/10.7717/peerj.9696/fig-11)

Also ventrally, the ground pattern condition of Neochelicerata is largely the same as that of Euchelicerata (Fig. 12C). The appendages of the anterior tagma, (possibly the hypostome and) chelicerae and the following six pairs of appendages, are incorporated into the feeding apparatus. However, the last of these appendage pairs, the chilidia (appendages of the pre-genital segment or post-ocular segment 7), consists of only the shovel-shaped most proximal part (basipod in neutral euarthropodan terminology, usually called coxa in chelicerate terminology); the walking part (endopod) is lacking. With this, the appendages of post-ocular segment 7 no longer perform a combined feeding-and-walking function, but instead close the feeding apparatus from its posterior end to reduce the loss of food in Xiphosurida (e.g. Patten, 1894; see Manton, 1964 for a detailed functional explanation of the feeding process). This condition is possibly already present earlier as it looks very similar in a species splitting off the evolutionary lineage before the node of Neochelicerata, *Venustulus waukashensis* (Moore et al., 2005, their fig. 3.3 + 3.4). Therefore, in the ground pattern of Neochelicerata (or slightly earlier) the feeding apparatus became further specialised, consisting of (possibly the hypostome,) chelicerae, five pairs of walking appendages and chilidia.

The specialised feeding apparatus of Eurypterida

The exact relationships of Eurypterida, Arachnida, Xiphosurida, and other exclusively fossil groups such as Chasmataspidida are still not entirely resolved (e.g. Dunlop, 2010; Garwood & Dunlop, 2014a and references therein). The feeding apparatus of Eurypterida (and possibly already of Metastomata if this is a natural group; see Haug, Wagner &




Figure 12 Schemes of feeding apparatuses and general body organisation in the ground pattern of different evolutionary levels. (A) Megacheira. (B) Euchelicerata. (C) Neochelicerata. Grey background shadings mark different tagmata. Colour coding: black = appendages of first post-ocular segment (great appendages resp. chelicerae) and hypostome ('labrum'); dark grey = basipod; light grey = endopod; white = exopod and (possibly) limbless segments. Presence of specific respiratory structures not known for the ground patterns of Megacheira and Euchelicerata. [Full-size !\[\]\(1663bb69f307a960345edb0e712f8c02_img.jpg\) DOI: 10.7717/peerj.9696/fig-12](https://doi.org/10.7717/peerj.9696/fig-12)

Haug, 2019) shows a stronger specialisation than that in the ground pattern of Neochelicerata, but is still composed of (possibly the hypostome and) the appendages of the same segments (chelicerae, following five pairs of appendages, metastoma, see below; Fig. 13A). The further posterior segments are also here not involved in feeding or locomotion; the corresponding appendages most probably became (partly) internalised and fulfilled respiratory function (Waterston, 1975; Braddy et al., 1999).

Also in sea scorpions, feeding and locomotion appears to be performed by the same appendages. However, Selden (1981), who described the functional morphology of *Eurypterus tetragonophthalmus* in great detail, assumed that only the posterior appendages were used for locomotion to avoid coordination problems between the differently




Figure 13 Schemes of feeding apparatuses and general body organisation in the ground pattern of different evolutionary levels, continued. (A) Metastomata. (B) Scorpionida. (C) Trigonotarbida. Grey background shadings mark different tagmata. Colour coding: black = appendages of first post-ocular segment (chelicerae) and hypostome ('labrum'); dark grey = basipod; light grey = endopod; white = exopod and (possibly) limbless segments. Question marks indicate uncertainties about origin of sternum and transition area. [Full-size DOI: 10.7717/peerj.9696/fig-13](https://doi.org/10.7717/peerj.9696/fig-13)

long appendages (though this assumption does not have to account for all species of Eurypterida due to their morphological differences). He also assumed that there is a task differentiation in food handling between the anterior and posterior legs, the anterior ones gathering food, while especially the last pair crushed hard food particles (Selden, 1981). This differentiation is corroborated by the specimens investigated in this study, as the basipods of the different appendages are equipped with different types of teeth on their median edges (see Bicknell et al. (2018a) for microstructure of these teeth, and Bicknell et al. (2018b) for similar studies on *Limulus polyphemus*), best visible when (almost) the entire feeding apparatus is preserved in situ (Figs. 5A and 5B). In some specimens the teeth on the basipods of the further anterior appendages are thinner and appear less robust than those on the basipods of the fifth (last) pair of walking legs (appendage pair of

post-ocular segment 6). The antero-median edges of the latter are equipped with strong teeth and reach under the metastoma (Figs. 2A and 3A; [Holm, 1898](#); [Selden, 1981](#)). They are also significantly elongated in anterior-posterior axis (Figs. 1A and 2A; based on its position in the fossil, not on its evolutionary origin), probably to achieve a larger biting force. With this, sea scorpions possessed fully functional antagonistic jaws (in contrast to earlier statements; for example [Gruner, 1993](#)) comparable to the condition in many crustaceans ([Manton, 1964](#)), but convergently evolved.

In addition to the differentiation of armature on the basipods of different appendages, on the same basipod the armature is also differentiated. The teeth further anterior on the median edge of the basipod appear stronger than the further posterior ones on the same basipod. [Selden \(1981\)](#) describes this differentiation for *E. tetragonophthalmus*, and discusses that some of the teeth or spines would have been movable while others were not. The presence of movable teeth may be species specific based on the observations in this study, but the general pattern of differentiated armature on the same basipod apparently occurs in different species of Eurypterida. This differentiation appears very similar to the situation in Mandibulata, in which the mandibles bear the pars incisivus and the pars molaris, that is two regions with rather different armature ([Edgecombe, Richter & Wilson, 2003](#)). Apparently, also this specialisation evolved convergently.

The metastoma (appendage of pre-genital segment or post-ocular segment 7) in representatives of Eurypterida basically fulfils the same function as the chilaria in representatives of Xiphosurida, it closes the feeding apparatus from posteriorly ([Selden, 1981](#)). Yet, in sea scorpions this is only a single plate (but which can have various shapes; for example [Tollerton, 1989](#)), so most probably the conjoined basipods of formerly free appendages (see [Holm, 1898](#); [Dunlop, 1998](#), his fig. 4; [Dunlop & Selden, 1998](#); [Jeram, 1998](#)). Some species show a notch in the anterior area of the metastoma ([Holm, 1896](#)), which may represent a remnant of the not completed fusion process. Unfortunately, no extremely early ontogenetic stages are preserved (late embryos to hatchlings; but see [Lamsdell & Selden \(2013\)](#) for fairly small stages), at least not well enough to allow investigation of the development of the metastoma (see below for development of the corresponding structure in scorpions).

In addition to the closing of the feeding apparatus, the metastoma appears to provide a kind of guide rail for the movements of the basipods of the appendage pair right in front of it ([Selden, 1981](#)). This function is comparable to that of the paragnaths in eucrustaceans, which are elevations of the sternite of the mandible segment and guide the movement of the mandibles ([Haug et al., 2011b](#); [Rötzer & Haug, 2015](#)). Again, this morphological similarity evolved most likely convergently.

The condition of the metastoma as a single plate and its covering of the proximal area of the posterior appendages leads to a more tightly closed feeding apparatus in comparison to that in Xiphosurida and the ground pattern condition of Euchelicerata. Considering that representatives of Eurypterida may have had an amphibious lifestyle ([Lamsdell & Braddy, 2009](#)), a more closed feeding apparatus could have been a predisposition for going on land, as food may get lost more easily on land than in the water, where it sinks slower when it is not grabbed tightly enough. The specialised feeding apparatus of sea

scorpions in its highly differentiated morphology is probably best interpreted as an autapomorphy of Eurypterida or even of an ingroup.

Further shortening of the feeding apparatus in Arachnida

In the following, scorpions are taken as a first example for Arachnida. Scorpions have been assumed to be the sister group to the remaining groups of Arachnida ([Weygoldt & Paulus, 1979](#)). Yet, in recent studies, scorpions resulted as sister group to Megopericulata ([Sharma et al., 2014](#); [Klußmann-Fricke & Wirkner, 2016](#); [Lozano-Fernandez et al., 2019](#)); in these analyses sea scorpions were not included (which was mostly not possible due to the type of analysed characters). This deep ingroup position of scorpions may be an artefact resulting from the lack of proper character polarisation due to the absence of Eurypterida, but this problem cannot be further discussed here.

In Arachnida, the posterior border of the feeding apparatus lies further anteriorly than in the previously discussed groups. In modern scorpions (Scorpiones), only the first four appendage-bearing segments (and possibly the hypostome) are involved in the feeding apparatus: chelicerae, pedipalps, and two pairs of walking appendages. The basipods of these two pairs of walking appendages are antero-medially elongated into a pronounced endite, which reaches far anteriorly, closing the feeding apparatus from the posterior end ([Snodgrass, 1948](#)). The median areas of the basipods of walking appendage pairs 3 and 4 are oriented almost anteriorly, but apparently not included into the feeding apparatus. Also the sternum, most likely the embryonically fused appendages of the seventh post-ocular (pre-genital) segment (see [Dunlop & Webster \(1999\)](#); [Farley \(2005\)](#); discussion in [Haug, Wagner & Haug \(2019\)](#)), is oriented anteriorly, but not involved in the feeding apparatus. Dorsally, the shield extends further posteriorly, including the segments bearing chelicerae, pedipalps, all walking appendages and possibly also the sternum-bearing segment (see discussion in [Haug, Wagner & Haug \(2019\)](#)). Hence, the posterior border of the feeding apparatus in Scorpiones no longer corresponds to the posterior border of the dorsal shield, in contrast to the presumed ground pattern condition in Euchelicerata, Neochelicerata, and Metastomata.

However, the condition of the feeding apparatus in modern scorpions differs from that in early fossil scorpions. In early scorpions, the feeding apparatus extends posteriorly only to the pedipalps ([Kjellesvig-Waering, 1986](#); [Waddington, Rudkin & Dunlop, 2015](#)). The walking appendages do not bear enditic protrusions, hence do not appear to have been involved in the feeding process ([Fig. 13B](#)). If this condition is the ground pattern condition of Scorpionida, the group including Scorpiones and different fossil scorpions, the feeding apparatus in modern scorpions would be an autapomorphy of Scorpiones.

In Trigonotarbida, the chelicerae, pedipalps and first pair of walking appendages contribute to the feeding apparatus ([Fig. 13C](#)). The basipods of the pedipalps and especially of the first pair of walking appendages possess endites medially, which were probably used for food manipulation (e.g. [Garwood, Dunlop & Sutton, 2009](#); [Garwood & Dunlop, 2014b](#), their fig. 1.5; [Dunlop & Garwood, 2017](#)). This condition is unknown from other arachnids.

The phylogenetic position of Trigonotarbida is still unclear ([Garwood & Dunlop, 2014a](#)). In different analyses, they have already resolved, for example, as sister group to




Figure 14 Feeding apparatuses of different extant representatives of Arachnida. (A) Araneae. (B) Amblypygi. (C) Schizomida. (D) Thelyphonida. (E) Phalangida (Opiliones). (F) Cyphophthalmi (Opiliones). (G) Solifugae. (H) Pseudoscorpiones. (I) Palpigradi. (J) Ricinulei. (K) Mesostigmata (Acari). (L) Ixodoidea (Acari). [Full-size !\[\]\(1663bb69f307a960345edb0e712f8c02_img.jpg\) DOI: 10.7717/peerj.9696/fig-14](https://doi.org/10.7717/peerj.9696/fig-14)

Ricinulei (Dunlop, 1996; Dunlop, Kamenz & Talarico, 2009) or as closely related to Megopericulata (Shear & Selden, 1986; Selden, Shear & Bonamo, 1991). Similarities to the one respectively the other group occur, for example, in the filtering structures on the mouthparts (Dunlop, 1994; Haug, 2017), details on the pedipalps (Dunlop, Kamenz & Talarico, 2009), or the general body organisation.

The feeding apparatus does not provide clear arguments for placing Trigonotarvida near the one or the other group of Arachnida. In most representatives of Arachnida, the feeding apparatus only consists of (possibly a hypostome,) chelicerae and pedipalps (Figs. 14B–14D, 14G, 14H and 14J). In some of them, the basipods of the pedipalps are more or less closely connected medially (Figs. 14D and 14H). The basipods of the pedipalps are in many web spiders involved in the feeding process, using median projections for mastication (sometimes referred to as gnathocoxae or maxillae; for example Williams, 1979;

Sierwald, 1988; Žabka, 2001; J. Dunlop, 2020, personal observations). In certain cases, a lower lip contributes to the feeding apparatus from posteriorly (Figs. 14A and 14I). In some groups, all these structures form a single, tightly connected unit (Figs. 14K and 14L). If this short condition of the feeding apparatus would be present in all representatives of Arachnida besides Trigonotarbida, two options for the ground pattern of Arachnida would be possible: either the feeding apparatus of Trigonotarbida would represent the ground pattern condition of Arachnida, or the very short condition of the other representatives of Arachnida would be the ground pattern condition and Trigonotarbida autapomorphically elongated the feeding apparatus.

However, the entire situation is more complicated due to the feeding apparatus of Opiliones (harvestmen). In harvestmen, the feeding apparatus includes also the first pair of walking legs, which bears endites on the basipods (Fig. 14E). Also the second pair of walking legs bears endites, at least in certain harvestmen, which may have a supporting function in the feeding process (*Shultz & Pinta-da-Rocha, 2014*). The presence of a preoral chamber, stomotheca, formed by all these endites in Opiliones and Scorpionida led *Shultz (2000, 2007)* to suggest a sister group relationship between these two groups (together forming Stomothecata). The endites of the first pair of walking appendages in harvestmen look rather similar to those in Trigonotarbida, and in general also the composition of the feeding apparatus would be similar between the two groups (*Garwood, Dunlop & Sutton, 2009*). However, the morphology in harvestmen is highly variable (Figs. 14E and 14F), and it is unclear how the feeding apparatus in the ground pattern of Opiliones looks. This together with the still unresolved phylogenetic position of Opiliones (*Garwood & Dunlop, 2014a; Lozano-Fernandez et al., 2019*) does not allow to make a reliable assumption about the feeding apparatus in the ground pattern of Arachnida.

CONCLUSIONS

- (1) The feeding apparatus in different groups of Euchelicerata is far from primitive, but is in fact a highly specialised system in each group.
- (2) During evolution, the feeding apparatus became progressively shorter in Euchelicerata, though the evolution within Arachnida remains still unclear due to unresolved phylogenetic relationships. The shortness of the feeding apparatus is not an ancestral character, but in fact a highly derived one, probably evolved in adaptation to new requirements resulting from habitat changes such as terrestrialisation.
- (3) Sea scorpions possess true antagonistic mouthparts with differentiated armature and a guide rail system. These characters are all similar to the condition in the mandibles of Mandibulata, apparently as result of convergent evolution.
- (4) Representatives of Trigonotarbida show similar specialisations concerning their feeding apparatuses to harvestmen.
- (5) In conclusion, the supposedly ‘primitive’ groups Eurypterida and Trigonotarbida are astonishingly specialised.

ACKNOWLEDGEMENTS

I thank Jason Dunlop (Berlin), Greg Edgecombe (London) and an anonymous reviewer for their helpful comments that improved the manuscript. Bruce Lieberman (Lawrence, Kansas) is thanked for handling the manuscript. My thanks go to Derek Briggs, Susan Butts, Jessica Utrup (all New Haven), Jean-Bernard Caron, Janet Waddington, Peter Fenton, Maryam Akrami (all Toronto), Claire Mellish (London), Vladimir Blagoderov (formerly London, now Edinburgh), Jessica Cundiff, Philip Perkins, Charles (Whit) Farnum (all Harvard), Christian Skovsted, Steffen Kiel and Cecilia Larsson (all Stockholm) for providing access to the specimens and technical assistance. I am grateful to Joachim T. Haug and J. Matthias Starck, Munich, for inspiring discussions. Joachim T. Haug, Mario Schädel and Philipp Wagner, all Munich, assisted with the imaging.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

The research visit to the NHM London was made possible by a grant from the European Commission's (FP 6) Integrated Infrastructure Initiative programme SYNTHESYS (GB-TAF-4057). The research was funded via a DAAD incoming fellowship and a Bavarian Equal Opportunities Sponsorship (BGF) of the LMU Munich. This study is part of a project funded by the German Research Foundation (DFG) under HA 7066/3-1. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:
European Commission's (FP 6) Integrated Infrastructure Initiative programme SYNTHESYS: GB-TAF-4057.
DAAD.
LMU Munich.
German Research Foundation (DFG): HA 7066/3-1.

Competing Interests

The author declares that she has no competing interests.

Author Contributions

- Carolin Haug conceived and designed the experiments, performed the experiments, analysed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.

Data Availability

The following information was supplied regarding data availability:

The phylogenetic relationships were drawn from the text of Haug C, Wagner P, Haug JT. 2019. The evolutionary history of body organisation in the lineage towards modern scorpions. *Bulletin of Geosciences* 94 (4), 389–408.

The specimens are deposited as follows: YPM_IP_216689 (Yale Peabody Museum of Natural History, New Haven); NHM I3406_2, NHM In24671, NHM In24687, NHM In24702, NHM In27357, NHM In31241, NHM RC_019 (Natural History Museum, London); NRM Ar31829, NRM Ar35343, NRM Ar35344, NRM Ar48883 (Naturhistoriska riksmuseet, Stockholm); MCZ PALI 131326, MCZ PALI 185687 (Harvard Museum of Comparative Zoology, Cambridge); ROMIP45532 (Royal Ontario Museum, Toronto).

REFERENCES

- Anderson LI, Selden PA. 1997.** Opisthosomal fusion and phylogeny of Palaeozoic Xiphosura. *Lethaia* **30**(1):19–31 DOI [10.1111/j.1502-3931.1997.tb00440.x](https://doi.org/10.1111/j.1502-3931.1997.tb00440.x).
- Aria C, Caron JB. 2017.** Mandibulate convergence in an armoured Cambrian stem chelicerate. *BMC Evolutionary Biology* **17**(1):261 DOI [10.1186/s12862-017-1088-7](https://doi.org/10.1186/s12862-017-1088-7).
- Aria C, Caron JB. 2019.** A middle Cambrian arthropod with chelicerae and proto-book gills. *Nature* **573**(7775):586–589 DOI [10.1038/s41586-019-1525-4](https://doi.org/10.1038/s41586-019-1525-4).
- Ballesteros JA, Sharma PP. 2019.** A critical appraisal of the placement of Xiphosura (Chelicerata) with account of known sources of phylogenetic error. *Systematic Biology* **68**(6):896–917 DOI [10.1093/sysbio/syz011](https://doi.org/10.1093/sysbio/syz011).
- Bennett DJ, Sutton MD, Turvey ST. 2018.** Quantifying the living fossil concept. *Palaeontologia Electronica* **21**(1):14A DOI [10.26879/750](https://doi.org/10.26879/750).
- Bicknell RD, Ledogar JA, Wroe S, Gutzler BC, Watson WH, Paterson JR. 2018b.** Computational biomechanical analyses demonstrate similar shell-crushing abilities in modern and ancient arthropods. *Proceedings of the Royal Society B* **285**(1889):20181935 DOI [10.1098/rspb.2018.1935](https://doi.org/10.1098/rspb.2018.1935).
- Bicknell RD, Paterson JR, Caron JB, Skovsted CB. 2018a.** The gnathobasic spine microstructure of recent and Silurian chelicerates and the Cambrian artiopodan *Sidneyia*: functional and evolutionary implications. *Arthropod Structure & Development* **47**(1):12–24 DOI [10.1016/j.asd.2017.12.001](https://doi.org/10.1016/j.asd.2017.12.001).
- Braddy SJ, Aldridge RJ, Gabbott SE, Theron JN. 1999.** Lamellate book-gills in a late Ordovician eurypterid from the Soom Shale, South Africa: support for a eurypterid-scorpion clade. *Lethaia* **32**(1):72–74 DOI [10.1111/j.1502-3931.1999.tb00582.x](https://doi.org/10.1111/j.1502-3931.1999.tb00582.x).
- Briggs DEG, Siveter DJ, Siveter DJ, Sutton MD, Garwood RJ, Legg D. 2012.** Silurian horseshoe crab illuminates the evolution of arthropod limbs. *Proceedings of the National Academy of Sciences of the United States of America* **109**(39):15702–15705 DOI [10.1073/pnas.1205875109](https://doi.org/10.1073/pnas.1205875109).
- Bruton DL, Whittington HB. 1997.** *Emeraldella* and *Leancoilia*, two arthropods from the Burgess Shale, Middle Cambrian, British Columbia. *Philosophical Transactions of the Royal Society of London B, Biological Sciences* **300**(1102):553–582 DOI [10.1098/rstb.1983.0020](https://doi.org/10.1098/rstb.1983.0020).
- Chen J, Waloszek D, Maas A. 2004.** A new ‘great-appendage’ arthropod from the Lower Cambrian of China and homology of chelicerate chelicerae and raptorial antero-ventral appendages. *Lethaia* **15**:3–20.
- Daley AC, Antcliffe JB, Drage HB, Pates S. 2018.** Early fossil record of Euarthropoda and the Cambrian Explosion. *Proceedings of the National Academy of Sciences of the United States of America* **115**(21):5323–5331 DOI [10.1073/pnas.1719962115](https://doi.org/10.1073/pnas.1719962115).
- Damen WG. 2002.** Parasegmental organization of the spider embryo implies that the parasegment is an evolutionary conserved entity in arthropod embryogenesis. *Development* **129**(5):1239–1250.

- Donoghue MJ, Doyle JA, Gauthier J, Kluge AG, Rowe T. 1989. The importance of fossils in phylogeny reconstruction. *Annual Review of Ecology and Systematics* 20(1):431–460 DOI 10.1146/annurev.es.20.110189.002243.
- Dunlop JA. 1994. Filtration mechanisms in the mouthparts of tetrapulmonate arachnids (Trigonotarbida, Araneae, Amblypygi, Uropygi, Schizomida). *Bulletin of the British Arachnological Society* 9:267–273.
- Dunlop JA. 1996. Evidence for a sister group relationship between Ricinulei and Trigonotarbida. *Bulletin of the British Arachnological Society* 10:193–204.
- Dunlop JA. 1998. The origins of tetrapulmonate book lungs and their significance for chelicerate phylogeny. In: *Proceedings of the 17th European Colloquium of Arachnology, Edinburgh*. 9–16.
- Dunlop JA. 2010. Geological history and phylogeny of Chelicerata. *Arthropod Structure & Development* 39(2–3):124–142 DOI 10.1016/j.asd.2010.01.003.
- Dunlop JA. 2011. Fossil Focus: Chelicerata. *Palaeontology Online* 1(1):1–8.
- Dunlop JA, Garwood RJ. 2017. Terrestrial invertebrates in the Rhynie chert ecosystem. *Philosophical Transactions of the Royal Society B, Biological Sciences* 373(1739):20160493 DOI 10.1098/rstb.2016.0493.
- Dunlop JA, Kamenz C, Talarico G. 2009. A fossil trigonotarbid arachnid with a ricinuleid-like pedipalpal claw. *Zoomorphology* 128(4):305–313 DOI 10.1007/s00435-009-0090-z.
- Dunlop JA, Lamsdell JC. 2017. Segmentation and tagmosis in Chelicerata. *Arthropod Structure & Development* 46(3):395–418 DOI 10.1016/j.asd.2016.05.002.
- Dunlop JA, Selden PA. 1998. The early history and phylogeny of the chelicerates. In: Fortey RA, Thomas RH, eds. *Arthropod Relationships, Systematics Association Special Volume Series* 55. London: Chapman & Hall, 221–235.
- Dunlop JA, Webster M. 1999. Fossil evidence, terrestrialization and arachnid phylogeny. *Journal of Arachnology* 27:86–93.
- Edgecombe GD. 2010. Palaeomorphology: fossils and the inference of cladistic relationships. *Acta Zoologica* 91(1):72–80 DOI 10.1111/j.1463-6395.2009.00426.x.
- Edgecombe GD, Legg DA. 2014. Origins and early evolution of arthropods. *Palaeontology* 57(3):457–468 DOI 10.1111/pala.12105.
- Edgecombe GD, Richter S, Wilson GD. 2003. The mandibular gnathal edges: homologous structures throughout Mandibulata? *African Invertebrates* 44(1):115–135.
- Fahrenbach WH, Arango CP. 2007. Microscopic anatomy of Pycnogonida: II. Digestive system. III. Excretory system. *Journal of Morphology* 268(11):917–935.
- Farley RD. 2005. Developmental changes in the embryo, pronymph, and first molt of the scorpion *Centruroides vittatus* (Scorpiones: Buthidae). *Journal of Morphology* 265(1):1–27 DOI 10.1002/jmor.10227.
- Garwood RJ, Dunlop J. 2014a. Three-dimensional reconstruction and the phylogeny of extinct chelicerate orders. *PeerJ* 2(2):e641 DOI 10.7717/peerj.641.
- Garwood R, Dunlop J. 2014b. The walking dead: blender as a tool for paleontologists with a case study on extinct arachnids. *Journal of Paleontology* 88(4):735–746 DOI 10.1666/13-088.
- Garwood R, Dunlop JA, Sutton MD. 2009. High-fidelity X-ray micro-tomography reconstruction of siderite-hosted Carboniferous arachnids. *Biology Letters* 5(6):841–844 DOI 10.1098/rsbl.2009.0464.
- Giribet G, Edgecombe GD. 2019. The phylogeny and evolutionary history of arthropods. *Current Biology* 29(12):R592–R602 DOI 10.1016/j.cub.2019.04.057.

- Gruner HE. 1993.** Teil 4: Arthropoda (ohne Insecta). Kaestner A (Begr.), *Lehrbuch der speziellen Zoologie*. Jena: Gustav Fischer Verlag.
- Haug C. 2017.** Feeding strategies in arthropods from the Rhynie and Windyfield cherts: ecological diversification in an early non-marine biota. *Philosophical Transactions of the Royal Society B: Biological Sciences* **373**(1739):20160492 DOI [10.1098/rstb.2016.0492](https://doi.org/10.1098/rstb.2016.0492).
- Haug JT, Briggs DEG, Haug C. 2012.** Morphology and function in the Cambrian Burgess Shale megacheiran arthropod *Leancoilia superlata* and the application of a descriptive matrix. *BMC Evolutionary Biology* **12**:162.
- Haug JT, Haug C, Ehrlich M. 2008.** First fossil stomatopod larva (Arthropoda: Crustacea) and a new way of documenting Solnhofen fossils (Upper Jurassic, Southern Germany). *Palaeodiversity* **1**:103–109.
- Haug C, Haug JT, Fayers SR, Trewin NH, Castellani C, Waloszek D, Maas A. 2012a.** Exceptionally preserved nauplius larvae from the Devonian Windyfield chert, Rhynie, Aberdeenshire, Scotland. *Palaeontologia Electronica* **15**(2):24A.
- Haug JT, Haug C, Kutschera V, Mayer G, Maas A, Liebau S, Castellani C, Wolfram U, Clarkson ENK, Waloszek D. 2011a.** Autofluorescence imaging, an excellent tool for comparative morphology. *Journal of Microscopy* **244**(3):259–272 DOI [10.1111/j.1365-2818.2011.03534.x](https://doi.org/10.1111/j.1365-2818.2011.03534.x).
- Haug JT, Olesen J, Maas A, Waloszek D. 2011b.** External morphology and post-embryonic development of *Derocheilocaris remanei* (Crustacea: Mystacocarida) revisited, with a comparison to the Cambrian taxon *Skara*. *Journal of Crustacean Biology* **31**(4):668–692 DOI [10.1651/11-3481.1](https://doi.org/10.1651/11-3481.1).
- Haug C, Sallam WS, Maas A, Waloszek D, Kutschera V, Haug JT. 2012c.** Tagmatization in Stomatopoda—reconsidering functional units of modern-day mantis shrimps (Verunipeltata, Hoplocarida) and implications for the interpretation of fossils. *Frontiers in Zoology* **9**(1):31 DOI [10.1186/1742-9994-9-31](https://doi.org/10.1186/1742-9994-9-31).
- Haug C, Wagner P, Haug JT. 2019.** The evolutionary history of body organisation in the lineage towards modern scorpions. *Bulletin of Geosciences* **94**:389–408 DOI [10.3140/bull.geosci.1750](https://doi.org/10.3140/bull.geosci.1750).
- Haug JT, Waloszek D, Maas A, Liu Y, Haug C. 2012b.** Functional morphology, ontogeny and evolution of mantis shrimp-like predators in the Cambrian. *Palaeontology* **55**(2):369–399 DOI [10.1111/j.1475-4983.2011.01124.x](https://doi.org/10.1111/j.1475-4983.2011.01124.x).
- Holm G. 1896.** Über eine neue Bearbeitung des *Eurypterus fischeri* Eichw. *Bulletin de l'Académie Impériale des Sciences de St.-Petersbourg* **4**:369–372.
- Holm G. 1898.** Über die Organisation des *Eurypterus fischeri* Eichw. *Mémoires de l'Académie Impériale des Sciences de St.-Petersbourg* **8**:1–57.
- Jeram AJ. 1998.** Phylogeny, classification and evolution of Silurian and Devonian scorpions. In: *Proceedings of the 17th European Colloquium of Arachnology*, Edinburgh. 17–31.
- Kerp H, Bomfleur B. 2011.** Photography of plant fossils—new techniques, old tricks. *Review of Palaeobotany and Palynology* **166**(3–4):117–151 DOI [10.1016/j.revpalbo.2011.05.001](https://doi.org/10.1016/j.revpalbo.2011.05.001).
- Kin A, Błażejowski B. 2014.** The horseshoe crab of the genus *Limulus*: living fossil or stabilomorph? *PLOS ONE* **9**(10):e108036 DOI [10.1371/journal.pone.0108036](https://doi.org/10.1371/journal.pone.0108036).
- Kjellesvig-Waering EN. 1986.** A restudy of the fossil Scorpionida of the world. *Palaeontographica Americana* **55**:1–287.
- Klußmann-Fricke BJ, Wirkner CS. 2016.** Comparative morphology of the hemolymph vascular system in Uropygi and Amblypygi (Arachnida): complex correspondences support Arachnopulmonata. *Journal of Morphology* **277**(8):1084–1103 DOI [10.1002/jmor.20559](https://doi.org/10.1002/jmor.20559).

- Krell FT, Cranston PS. 2004. Which side of the tree is more basal? *Systematic Entomology* 29(3):279–281 DOI 10.1111/j.0307-6970.2004.00262.x.
- Lamsdell JC. 2013. Revised systematics of Palaeozoic ‘horseshoe crabs’ and the myth of monophyletic Xiphosura. *Zoological Journal of the Linnean Society* 167(1):1–27 DOI 10.1111/j.1096-3642.2012.00874.x.
- Lamsdell JC. 2016. Horseshoe crab phylogeny and independent colonizations of fresh water: ecological invasion as a driver for morphological innovation. *Palaeontology* 59(2):181–194 DOI 10.1111/pala.12220.
- Lamsdell JC, Braddy SJ. 2009. Cope’s rule and romer’s theory: patterns of diversity and gigantism in eurypterids and Palaeozoic vertebrates. *Biology Letters* 6(2):265–269 DOI 10.1098/rsbl.2009.0700.
- Lamsdell JC, Selden PA. 2013. Babes in the wood—a unique window into sea scorpion ontogeny. *BMC Evolutionary Biology* 13(1):98 DOI 10.1186/1471-2148-13-98.
- Legg D. 2013. Multi-segmented arthropods from the Middle Cambrian of British Columbia (Canada). *Journal of Paleontology* 87(3):493–501 DOI 10.1666/12-112.1.
- Legg DA. 2014. *Sanctacaris uncata*: the oldest chelicerate (Arthropoda). *Naturwissenschaften* 101(12):1065–1073 DOI 10.1007/s00114-014-1245-4.
- Lehmann WM. 1956. Beobachtungen an *Weinbergina opitzi* (Merost., Devon). *Senckenbergiana Lethaia* 37:67–77.
- Lozano-Fernandez J, Tanner AR, Giacomelli M, Carton R, Vinther J, Edgecombe GD, Pisani D. 2019. Increasing species sampling in chelicerate genomic-scale datasets provides support for monophyly of Acari and Arachnida. *Nature Communications* 10(1):2295 DOI 10.1038/s41467-019-10244-7.
- Maas A, Waloszek D, Chen J, Braun A, Wang X, Huang D. 2004. Phylogeny and life habits of early arthropods—predation in the Early Cambrian sea. *Progress in Natural Science* 14(2):158–166 DOI 10.1080/10020070412331343301.
- Malakhov VV. 2010. A new system of Bilateria. *Herald of the Russian Academy of Sciences* 80(1):29–41 DOI 10.1134/S1019331610010041.
- Manton SM. 1964. Mandibular mechanisms and evolution of arthropods. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences* 247(737):1–183 DOI 10.1098/rstb.1964.0001.
- Martin AJ. 2017. *The evolution underground: burrows, bunkers, and the marvelous subterranean world beneath our feet*. New York: Pegasus Books.
- Moore RA, Briggs DE, Bartels C. 2005. A new specimen of *Weinbergina opitzi* (Chelicerata: Xiphosura) from the Lower Devonian Hunsrück Slate, Germany. *Paläontologische Zeitschrift* 79(3):399–408.
- Moore RA, Briggs DE, Braddy SJ, Anderson LI, Mikulic DG, Kluessendorf J. 2005. A new synziphosurine (Chelicerata: Xiphosura) from the late Llandovery (Silurian) Waukesha lagerstätte, Wisconsin, USA. *Journal of Paleontology* 79(2):242–250 DOI 10.1666/0022-3360(2005)079<0242:ANSCXF>2.0.CO;2.
- Omland KE, Cook LG, Crisp MD. 2008. Tree thinking for all biology: the problem with reading phylogenies as ladders of progress. *BioEssays* 30(9):854–867 DOI 10.1002/bies.20794.
- Patten W. 1894. On the morphology and physiology of the brain and sense organs of *Limulus*. *Quarterly Journal of Microscopical Science* 35:1–96.
- Resh VH, Cardé RT. 2003. *Encyclopedia of Insects*. San Diego: Academic Press.

- Rust J. 2007. Die Bedeutung von Fossilien für phylogenetische Rekonstruktionen. *Species, Phylogeny and Evolution* 1:73–87.
- Rötzer MAIN, Haug JT. 2015. Larval development of the European lobster and how small heterochronic shifts lead to a more pronounced metamorphosis. *International Journal of Zoology* 2015:345172.
- Sekiguchi K. 1988. *Biology of horseshoe crabs*. Tokyo: Science House.
- Selden PA. 1981. Functional morphology of the prosoma of *Baltoeurypterus tetragonophthalmus* (Fischer) (Chelicerata: Eurypterida). *Earth and Environmental Science Transactions of the Royal Society of Edinburgh* 72(1):9–48 DOI 10.1017/S0263593300003217.
- Selden PA, Lamsdell JC, Qi L. 2015. An unusual euchelicerate linking horseshoe crabs and eurypterids, from the Lower Devonian (Lochkovian) of Yunnan, China. *Zoologica Scripta* 44(6):645–652 DOI 10.1111/zsc.12124.
- Selden PA, Shear WA, Bonamo PM. 1991. A spider and other arachnids from the Devonian of New York, and reinterpretations of Devonian Araneae. *Palaeontology* 34:241–281.
- Sharma PP, Kaluziak ST, Perez-Porro AR, Gonzalez VL, Hormiga G, Wheeler WC, Giribet G. 2014. Phylogenomic interrogation of Arachnida reveals systemic conflicts in phylogenetic signal. *Molecular Biology and Evolution* 31(11):2963–2984 DOI 10.1093/molbev/msu235.
- Shear WA, Selden PA. 1986. Phylogenetic relationships of the Trigonotarbitida, an extinct order of arachnids. *Actas X Congr. Int. Aracnol. Jaca/España* 1:393–397.
- Shultz JW. 2000. Skeletomuscular anatomy of the harvestman *Leiobunum aldrichi* (Weed, 1893) (Arachnida: Opiliones: Palpatores) and its evolutionary significance. *Zoological Journal of the Linnean Society* 128(4):401–438 DOI 10.1111/j.1096-3642.2000.tb01522.x.
- Shultz JW. 2007. A phylogenetic analysis of the arachnid orders based on morphological characters. *Zoological Journal of the Linnean Society* 150(2):221–265 DOI 10.1111/j.1096-3642.2007.00284.x.
- Shultz JW, Pinta-da-Rocha R. 2014. Morphology and functional anatomy. In: Pinta-da-Rocha R, Machado G, Giribet G, eds. *Harvestmen—The Biology of Opiliones*. Harvard: Harvard University Press, 14–61.
- Sierwald P. 1988. Notes on the behavior of *Thalassius spinosissimus* (Arachnida: Araneae: Pisauridae). *Psyche: A Journal of Entomology* 95(3–4):243–252 DOI 10.1155/1988/91410.
- Snodgrass RE. 1948. The feeding organs of Arachnida, including mites and ticks. *Smithsonian Miscellaneous Collections* 110:1–93 DOI 10.5962/bhl.part.9028.
- Stürmer W, Bergström J. 1981. *Weinbergina*, a xiphosuran arthropod from the Devonian Hunsrück Slate. *Paläontologische Zeitschrift* 55(3–4):237–255.
- Sutton MD, Briggs DE, Siveter DJ, Siveter DJ, Orr PJ. 2002. The arthropod *Offacolus kingi* (Chelicerata) from the Silurian of Herefordshire, England: computer based morphological reconstructions and phylogenetic affinities. *Proceedings of the Royal Society of London Series B: Biological Sciences* 269(1497):1195–1203 DOI 10.1098/rspb.2002.1986.
- Tanaka G, Hou X, Ma X, Edgecombe GD, Strausfeld NJ. 2013. Chelicerate neural ground pattern in a Cambrian great appendage arthropod. *Nature* 502(7471):364–367 DOI 10.1038/nature12520.
- Tollerton VP. 1989. Morphology, taxonomy, and classification of the order Eurypterida Burmeister, 1843. *Journal of Paleontology* 63(5):642–657 DOI 10.1017/S0022336000041275.
- Waddington J, Rudkin DM, Dunlop JA. 2015. A new mid-Silurian aquatic scorpion—one step closer to land? *Biology Letters* 11(1):20140815 DOI 10.1098/rsbl.2014.0815.

- Wagner P, Dömel JS, Hofmann M, Hübner J, Leese F, Melzer RR. 2017b. Comparative study of bisected proboscides of Pycnogonida. *Organisms Diversity & Evolution* 17(1):121–135 DOI 10.1007/s13127-016-0310-6.
- Wagner P, Haug JT, Sell J, Haug C. 2017a. Ontogenetic sequence comparison of extant and fossil tadpole shrimps: no support for the “living fossil” concept. *PalZ* 91(4):463–472 DOI 10.1007/s12542-017-0370-8.
- Walossek D. 1999. On the Cambrian diversity of Crustacea. In: Schram FR, Von Vaupel Klein JC, eds. *Crustaceans and the Biodiversity Crisis, Vol. 1: Proceedings of the Fourth International Crustacean Congress, Amsterdam, the Netherlands, July 20–24, 1998*. Leiden: Brill Academic Publishers, 3–27.
- Waloszek D, Dunlop JA. 2002. A larval sea spider (Arthropoda: Pycnogonida) from the upper Cambrian ‘Orsten’ of Sweden, and the phylogenetic position of pycnogonids. *Palaeontology* 45(3):421–446 DOI 10.1111/1475-4983.00244.
- Waterston CD. 1975. Gill structure in the Lower Devonian eurypterid *Tarsopterella scotica*. *Fossils and Strata* 4:241–254.
- Weygoldt P, Paulus HF. 1979. Untersuchungen zur Morphologie, Taxonomie und Phylogenie der Chelicerata II: Cladogramme und die Entfaltung der Chelicerata. *Journal of Zoological Systematics and Evolutionary Research* 17(3):177–200 DOI 10.1111/j.1439-0469.1979.tb00699.x.
- Williams DS. 1979. The feeding behaviour of New Zealand *Dolomedes* species (Araneae: Pisauridae). *New Zealand Journal of Zoology* 6(1):95–105 DOI 10.1080/03014223.1979.10428352.
- Williams KL. 2019. *Limulus* as a model organism. In: Williams KL, ed. *Endotoxin Detection and Control in Pharma, Limulus, and Mammalian Systems*. Cham: Springer, 597–629.
- Yang J, Ortega-Hernández J, Butterfield NJ, Zhang XG. 2013. Specialized appendages in fuxianhuiids and the head organization of early euarthropods. *Nature* 494(7438):468–471 DOI 10.1038/nature11874.
- Žabka M. 2001. Salticidae (Arachnida: Araneae) from the Oriental, Australian and Pacific regions, XIV: the genus *Adoxotoma* Simon. *Records of the Western Australian Museum* 20:323–332.