

Intergenerational implications of alcohol intake: metabolic disorders in alcohol-naïve rat offspring

Paweł Mierzejewski¹, Alicja Zakrzewska¹, Julita Kuczyńska¹, Edyta Wyszogrodzka¹, Monika Dominiak^{Corresp. 1}

¹ Department of Pharmacology, Institute of Psychiatry and Neurology, Warsaw, Poland

Corresponding Author: Monika Dominiak
Email address: mdominia@wp.pl

Alcohol drinking may be associated with an increased risk of various metabolic diseases. Rat lines selectively bred for alcohol preference and alcohol avoidance constitute an interesting model to study inherited factors related to alcohol drinking and metabolic disorders. The aim of the present study was to compare the levels of selected laboratory biomarkers of metabolic disorders in blood samples from naïve offspring of Warsaw alcohol high-preferring (WHP), Warsaw alcohol low-preferring (WLP), and wild Wistar rats. Blood samples were collected from 3-month old (300-350 g) alcohol-naïve, male offspring of WHP (n=8) and WLP rats (n=8), as well as alcohol-naïve, male, wild Wistar rats. Markers of metabolic, hepatic, and pancreatic disorders were analysed (levels of homocysteine, glucose, total cholesterol, triglycerides and γ -glutamyl transferase (GGT), aspartate (AST), alanine aminotransferase (ALT), and amylase serum activities). Alcohol-naïve offspring of WHP, WLP, and wild Wistar rats differed significantly in the levels of triglycerides, total cholesterol, homocysteine, as well as in the activity of GGT, ALT, AST, and amylase enzymes. Most markers in the alcohol-naïve offspring of WHP rats were altered even though they were never exposed to alcohol pre- or postnatally. This may suggest that parental alcohol abuse can have a detrimental influence on offspring vulnerability to metabolic disorders.

Intergenerational implications of alcohol intake: metabolic disorders in alcohol-naïve rat offspring

Paweł Mierzejewski¹, Alicja Zakrzewska¹, Julita Kuczyńska¹, Edyta Wyszogrodzka¹, Monika Dominiak^{1*}

¹*Department of Pharmacology, Institute of Psychiatry and Neurology, Warsaw, Poland*

***Corresponding author:**

Monika Dominiak

ul. Sobieskiego 9, 02-957 Warsaw, Poland

email: mdominia@wp.pl

Abstract

Alcohol drinking may be associated with an increased risk of various metabolic diseases. Rat lines selectively bred for alcohol preference and alcohol avoidance constitute an interesting model to study inherited factors related to alcohol drinking and metabolic disorders. The aim of the present study was to compare the levels of selected laboratory biomarkers of metabolic disorders in blood samples from naïve offspring of Warsaw alcohol high-preferring (WHP), Warsaw alcohol low-preferring (WLP), and wild Wistar rats. Blood samples were collected from 3-month old (300-350 g) alcohol-naïve, male offspring of WHP (n=8) and WLP rats (n=8), as well as alcohol-naïve, male, wild Wistar rats. Markers of metabolic, hepatic, and pancreatic disorders were analysed (levels of homocysteine, glucose, total cholesterol, triglycerides and γ -glutamyl transferase (GGT), aspartate (AST), alanine aminotransferase (ALT), and amylase serum activities). Alcohol-naïve offspring of WHP, WLP, and wild Wistar rats differed significantly in the levels of triglycerides, total cholesterol, homocysteine, as well as in the activity of GGT, ALT, AST, and amylase enzymes. Most markers in the alcohol-naïve offspring of WHP rats were altered even though they were never exposed to alcohol pre- or postnatally. This may suggest that parental alcohol abuse can have a detrimental influence on offspring vulnerability to metabolic disorders.

Introduction

Heavy alcohol drinking can lead to serious health problems, including cardiovascular diseases (CVD) and liver cirrhosis [1-3]. In numerous studies, heavy alcohol drinking was linked to both an increased risk of CVD and metabolic syndrome, leading to the hypothesis that alcohol-induced metabolic disorders could stand behind the relationship between excessive alcohol consumption and CVD [4-6]. The above-mentioned associations may be related to environmental as well as genetic factors. Twin, family, and adoption studies have consistently shown that genetic factors play an important role in the development of alcohol dependence [7-9]. Genetic loading is also an important risk factor for dyslipidaemia, diabetes, and metabolic syndrome [10-12]. Hence, one may hypothesise that excessive alcohol consumption and metabolic abnormalities are related to the same genetic mechanism(s).

Prenatal alcohol consumption has been shown to interfere with the development of multiple organs and systems, resulting in Foetal Alcohol Spectrum Disorders (FASD) [13]. Epidemiological studies have revealed that alcohol is the most common teratogen to which humans are exposed. It was estimated that 8 children per 1,000 live births are diagnosed with some degree of FASD [14]. This condition is characterised by a spectrum of structural defects, central nervous system disorders, as well as growth deficits, and metabolic disorders [15,16]. Although it is perceived as a childhood disorder, the majority of health problems, including growth restriction, hyperinsulinemia, and other metabolic disruptions are lifelong, persisting into adulthood [17,18].

Apart from this well-known unfavourable effect on the foetus, recent studies on rats have highlighted the possible negative impact on offspring of maternal alcohol exposure also around the time of conception. Several lines of evidence have suggested that periconceptional alcohol

use may have a negative impact on adult offspring, including increased obesity risk, altered plasma lipids, leptin profiles, and liver steatosis [19]. Furthermore, the study of Dorey et al. (2018) [20] revealed a greater risk of obesity in the offspring of females exposed to alcohol around the time of conception. It was associated with the altered expression of key components of the mesolimbic reward pathway and an increased preference for palatable foods in offspring.

However, apart from reports on maternal alcohol use during pregnancy or in the periconceptional period, studies on paternal preconception alcohol exposure on offspring also provide interesting findings. Paternal alcohol intake during the preconception period induces a spectrum of health problems and alters gene expression in rodent offspring [21-24]. The results of the study of Kim et al. (2014) [25] suggest that the preconceptional exposure to ethanol of male mice before mating induces ADHD-like behaviours in offspring, possibly via epigenetic changes in gene expression. Asimes et al. (2017) [26] demonstrated that adolescent binge ethanol exposure altered DNA methylation patterns in the hypothalamus of alcohol-naïve offspring. In another study examining parental binge alcohol abuse, differences in the expression of genes involved in neurogenesis, reproductive function, and regulation of obesity were observed in offspring [27]. Also, the studies of Chang et al. (2019a, 2019b) [28,29] using a mouse model of paternal alcohol exposure have identified the markers of hepatic fibrosis and abnormalities in both lipid production and insulin signaling in the offspring of alcohol-exposed sires. The aforementioned studies raise the possibility that alcohol affects naïve offspring even in the absence of direct foetal alcohol exposure, and epigenetic inheritance may be the mechanism responsible for this phenomenon.

In light of the metabolic changes associated with alcohol consumption, whether common metabolic markers differ in alcohol-naïve offspring of high- and low-preferring rats is of interest.

Rat lines selectively bred for alcohol preference and alcohol avoidance constitute an interesting model to study the inherited factors related to alcohol drinking and metabolic disorders. The breeding of rats was carried out as previously described by Rok-Bujko et al. (2006) [30]. Specifically, Warsaw high-preferring (WHP) and Warsaw low-preferring (WLP) rats were bred from Wistar rats for opposite ethanol preference [30,31]. When given a choice between water and 10% ethanol, WHP rats voluntarily drink excessive amounts of ethanol (>5.0 g/kg/24 h), while WLP rats drink marginal quantities of ethanol [31].

The aim of the present study was to compare the common markers reflecting cardiovascular, liver, and pancreatic function in alcohol-naïve offspring of WHP, WLP, and Wistar rats: the levels of homocysteine, glucose, total cholesterol, triglycerides, γ -glutamyl transferase (GGT), aspartate aminotransferase (AST), alanine aminotransferase (ALT), and amylase serum activities. We hypothesise that markers of cardiovascular, liver, and pancreatic function differ between alcohol-naïve offspring of WHP, WLP, and Wistar rats.

Materials and methods

Rats and experimental design

Twenty-four (8 per group), twelve-week-old, alcohol-naïve Warsaw high-preferring (WHP), alcohol-naïve Warsaw low-preferring (WLP) rats, and male albino Wistar rats (Charles River, Sulzfeld, Germany) were used for the study. The WLP and WHP rats were obtained from

the Department of Pharmacology and Physiology of the Nervous System at the Institute of Psychiatry and Neurology in Warsaw, Poland.

During the selection, which is a standardized procedure repeated in every generation, WHP and WLP rats were exposed to alcohol for 30 days, after reaching a weight of 250 g. In males, this occurs at around 60 days of age. When given a choice between water and 10% ethanol, WHP rats voluntarily drank excessive amounts of ethanol (>5.0 g/kg/24 h), while WLP rats consumed negligible quantities of ethanol [31]. Periods of binge drinking are observed only in WHP rats. In this study, all parents of examined offspring were previously subject to a standardized selection procedure. After a 30-day period of alcohol exposure, they were mated 24 hours after the last ethanol consumption and subsequent offspring were produced. The parents were not intoxicated at the time of mating and were not exposed to alcohol at any time during pregnancy, thus, the analysed offspring were never directly exposed to alcohol. Two rats per litter were used (4 litters in total for each group).

All the rats were housed in the same environmental conditions and handled by the same experimenters for at least 30 days before blood sampling. The environmental conditions were as previously described in the study of Zakrzewska et al. (2019) [32]. Specifically, the rats were housed in standard plastic cages, in groups of three, with wood chip bedding on the floor, in fully controlled environmental conditions: temperature 22 ± 2 °C, humidity $50\% \pm 5$, and 12-hour light/12-hour dark cycles with lights on at 8.00 a.m. All the rats had unlimited access to tap water and standard laboratory chow (Labofeed H, WPIK, Kcynia, Poland) throughout the duration of the study.

The offspring of all rat lines were weighed at 21, 60, and 90 days of age. There were no differences in body weight between the tested rat lines at any stage of their development or their

food intake (both $F < 1.0$, $p > 0.05$). In the last measurement at 90 days of age (at the time of sampling) the offspring of WHP, WLP, and Wistar rats weighed 345 g (± 27.6), 338 g (± 30.5) and 341 g (± 25.8), respectively.

Blood samples were obtained from 3-month-old (average of 300-350 g) drug- and alcohol-naïve male offspring of WHP ($n=8$) and WLP rats ($n=8$). Non-selected 3-month-old, drug- and alcohol-naïve male Wistar rats (Charles River, Sulzfeld, Germany; 300-350 g; $n=8$) were used as additional controls. The rats were deprived of food for 12 h overnight and were sacrificed by rapid decapitation with a guillotine designed for rodents. Trunk blood was collected into a tube immediately after decapitation. Serum was obtained after centrifugation at 1600 g for 15 minutes. The serum was collected into a plain tube and stored at -80°C until the analysis. No haemolysis was observed in any collected sample.

The Ethics Committee for Animal Care provided its consent for the study (Agreement no. 10/2010). The study was performed in full compliance with respective Polish and European ethical regulations (Directive No. 86/609/EEC).

Biochemical analysis

The activities of AST, ALT, GGT, and amylase, as well as the levels of glucose, total cholesterol, and triglycerides were measured in the serum using automated enzymatic colorimetric methods (Cobas Integra, Roche Diagnostics, Rotkreuz, Switzerland). Serum homocysteine determination was carried out using fully automated chemiluminescence assay (ADVIA Centaur, Siemens Healthcare Diagnostics, Erlangen, Germany). The assays were performed in accordance with the manufacturers' instructions in the analytical laboratory of the Institute of Psychiatry and Neurology in Warsaw.

Statistical analysis

A one-way ANOVA was used to evaluate between-group differences in the biochemical measures. The Newman-Keuls test was applied for post hoc comparisons. The normality of distribution of the tested variables was verified using the Shapiro-Wilk test. For two tested variables: ALT and triglycerides, the distribution could not be considered as normal. For the analysis of these parameters, the Kruskal-Wallis test was applied, while Dunn's multiple comparisons test was used for post hoc comparisons. P-values of less than 0.05 were considered significant. The study was exploratory in nature, thus, no correction for multiple comparisons was applied. The Statistica 12.0 software package (StatSoft, Inc., Tulsa, OK, USA) was used to analyse all the data.

Results

The one-way ANOVA revealed that the offspring of WHP, WLP, and Wistar rats differed significantly in the following laboratory parameters: homocysteine ($F(2, 21)=9.65$, $p=0.001$), total cholesterol ($F(2, 21)=44.63$, $p<0.001$), AST ($F(2, 21)=6.74$, $p=0.005$), amylase ($F(2, 21)=14.30$, $p<0.001$), and GGT ($F(2, 21)=135.87$, $p<0.001$) (Table 1). There were no differences in blood glucose levels ($F(2, 21) = 0.86$, $p=0.437$).

[Table 1]

The post hoc Newman-Keuls comparisons indicated that the alcohol-naïve offspring of WHP rats had significantly higher levels of total cholesterol and homocysteine, as well as significantly higher AST and amylase activity as compared to the alcohol-naïve WLP and the Wistar controls. The offspring of WHP rats had a significantly higher level of homocysteine as compared to the offspring of WLP rats, but not the Wistar rats. Surprisingly, the offspring of WHP subjects showed significantly lower GGT activity as compared to the WLP and the Wistar rats.

Regarding ALT as well as triglycerides, the Kruskal-Wallis test revealed a highly significant difference between the tested groups: ($H(2)=17.6$, $p<0.001$, ($H(2)=20.2$, $p<0.001$, respectively. Post hoc comparisons for these markers revealed significant differences between WHP and Wistar rats (both $p<0.001$) for both markers, as well as between WHP and WLP rats ($p=0.04$, $p<0.001$, for ALT and triglycerides, respectively).

Discussion

This is the first study to show changes in selected biochemical markers of metabolic disorders in naïve Wistar rats selected for high alcohol preference (WHP rats). A major finding of the present study is that the alcohol-naïve offspring of WHP rats have significantly higher blood levels of total cholesterol, triglycerides, and homocysteine than their WLP counterparts and wild Wistar rats. Some differences were also observed in the markers of hepatic and pancreatic function between alcohol-naïve offspring of WHP, WLP, and Wistar rats (ALT, AST, and amylase were increased, whereas GGT activity was decreased in WHPs).

Interestingly, our previous study showed an increased level of leptin—a risk factor for obesity and hypertension— in alcohol-naïve WHP rats as compared to alcohol-naïve WLP rats [33]. Although leptin is involved in lipid metabolism [34], the exact mechanism of the above observation in alcohol-naïve WHP rats remains unclear. In other studies, chronic alcohol exposure in rodents was shown to reduce the serum level of leptin [35]. Furthermore, leptin deficiency was associated with the development of fatty liver in mice [36].

A high level of total cholesterol or triglycerides is associated with an increased risk of atherosclerosis, myocardial infarction, and stroke [37-39]. Elevated levels of homocysteine are thought to facilitate the development of atherosclerosis and are considered a risk factor for coronary disease and stroke [40-41]. Several studies have shown a positive correlation between the amount of alcohol consumption and the blood homocysteine level [42-43]. It has also been shown that active drinking and early abstinent alcoholics have a higher level of homocysteine in the serum as compared to the general population [44-45]. There are also some epidemiological analyses showing a link between familial alcoholism and obesity among men and women [46].

There are many hypotheses aimed at explaining the increased risk of metabolic diseases in alcoholics. Our study indicates that some inherited factors (genetic, epigenetic) may play a role. These factors transmitted from alcoholic parents to their offspring could, at least partially, be responsible for the increased risk of metabolic disorders. A few studies have attempted to identify the genes responsible for the differences in alcohol preference among the naïve offspring of high- and low-preferring rats [47-49]. One such study indicated that alcohol preference may share some genetic factors with the cardiovascular system and metabolism [49], however, this study failed to identify any gene differences between high- and low-preferring rats. Thus, it does not support the hypothesis that the observed phenotypic differences are gene related. One can

hypothesise that other factors, for example, the epigenetic mechanism, may have a dominant role in the observed phenomenon. Many reports in the literature have indicated that environmental factors can have long-term effects on gene expression through epigenetic mechanisms [50]. Epigenetics refers to mechanisms that modify gene expression without any changes in the DNA sequence. Recent evidence suggests that epigenetic modifications can be carried to the offspring via the germline and are responsible for the transmission of alcohol-related disorders across generations [51-52]. Interestingly, there are studies showing the association of parental alcohol intake with subsequent changes in gene expression in naïve offspring [24-26], as well as with changes in lipid metabolism [27]. Moreover, Pennington et al. (2002) [53] showed that maternal preconceptional alcohol intake may be associated with hypertriglyceridemia in adult offspring who have never been exposed to alcohol. Paternally inherited alterations in epigenetic programming were also shown to be related to metabolic defects observed in foetal alcohol spectrum disorders. In rodents, chronic paternal alcohol use affected insulin signaling and lipid homeostasis in the offspring through paternally inherited alterations in liver x receptor activity [28]. Similar findings were reported by Chang et al. (2019b) [29], assessing long-term impacts of chronic preconception paternal alcohol use in mice. Evidence of metabolic programming in offspring included suppressed cytokine profiles within the liver and pancreas, consistent with findings from this study.

In our study, most of the tested hepatic biomarkers in the alcohol-naïve offspring of WHP rats had changed similarly to the heavy alcohol abusers. Surprisingly, the serum activity of the most specific alcohol-related biomarker, GGT, was decreased as compared to the WLP counterparts and the Wistar controls. These contrary results are difficult to explain. GGT, unlike AST and ALT, is located on the external surface of cellular membranes. Chronic alcohol use is

considered to induce membrane GGT solubilization and increase the release of the enzyme, thus, the mechanism of release of given liver enzymes is related to the degree of cell damage. One can also hypothesise that different epigenetic mechanisms regulate the activity of these enzymes [54] and one of the possible mechanisms could be linked to the changes in cellular membrane properties. Nevertheless, the mechanism responsible for this is yet to be discovered.

The results of this study should be interpreted in light of certain limitations. WHP and WLP line breeding involves the process of selecting rats in each generation through exposure to alcohol. Therefore, the design of the study could not include a control group of the offspring of WHP unexposed rats and, similarly, for this reason distinguishing between the paternal and maternal effects was also unfeasible as both males and females were required to be exposed to alcohol. Also, due to ethical concerns, we were only able to include a limited number of rats per group, thus, only male offspring were examined. However, there are sex-specific effects on offspring growth and long-term metabolic programming [29], thus, comparable studies on female offspring are warranted.

It should be emphasized that although the offspring were not directly exposed to alcohol, the last estrogenic cycle for females, as well as spermatogenesis in males, were exposed to alcohol. Thus, it cannot be excluded that periconceptional exposure could have an impact on gametes and affect, for example, oocyte maturation [55-57]. Similarly, this may influence the perinatal behaviours of parents and food intake in such a way that it may have indirect effects on metabolism. It has been shown, for example, that exposure to alcohol three weeks before conception affects the birth weight and body growth of offspring [23]. In our study, however, we failed to observe any differences in weight gain between the examined rat lines.

Conclusions

In conclusion, the present study revealed that the alcohol-naïve offspring of high alcohol preference rats differed significantly from their low alcohol preference counterparts and wild Wistar rats in terms of common markers reflecting cardiovascular, liver, and pancreatic disorders. To our knowledge, this is the first study showing the differences in selected biochemical markers of metabolic disorders in high- and low-preferring rats. One can hypothesise that these changes are associated with epigenetic changes at the level of the gametes and/or with the genetically inherited phenotype. Either way, this study suggests that parental alcohol abuse can have a detrimental influence on offspring vulnerability to metabolic disorders.

Given the detrimental influence of excessive alcohol consumption and metabolic disorders on individuals and societies, the present findings should be verified in further preclinical and clinical studies.

Acknowledgments

None.

References

1. Corrao G, Bagnardi V, Zambon A, La Vecchia C. A meta-analysis of alcohol consumption and the risk of 15 diseases. *Prev Med.* 2004;38:613-619.
2. Ronksley PE, Brien SE, Turner BJ, Mukamal KJ, Ghali WA. Association of alcohol consumption with selected cardiovascular disease outcomes: a systematic review and meta-analysis. *Br Med J.* 2011;342:d671.

3. Yang AL, Vadhavkar S, Singh G, Omary MB. Epidemiology of alcohol-related liver and pancreatic disease in the United States. *Arch Intern Med.* 2008;168:649-656.
4. Athyros VG, Liberopoulos EN, Mikhailidis DP, Papageorgiou AA, Ganotakis ES, Tziomalos K, Kakafika AI, Karagiannis A, Lambropoulos S, Elisaf M. Association of drinking pattern and alcohol beverage type with the prevalence of metabolic syndrome, diabetes, coronary heart disease, stroke, and peripheral arterial disease in a Mediterranean cohort. *Angiology.* 2007;58:689-697.
5. Alkerwi A, Boutsen M, Vaillant M, Barre J, Lair ML, Albert A, Guillaume M, Dramaix M. Alcohol consumption and the prevalence of metabolic syndrome: a meta-analysis of observational studies. *Atherosclerosis.* 2009;204:624-625.
6. Kahl KG, Hillemacher T. The metabolic syndrome in patients with alcohol dependency: Current research and clinical implications. *Prog Neuro-Psychopharmacol Biol Psychiatry.* 2016;70:49-56.
7. Kendler KS, Heath AC, Neale MC, Kessler RC, Eaves LJ. A population-based twin study of alcoholism in women. *J Am Med Assoc.* 1992;268:1877-1882.
8. Kendler KS, Davis CG, Kessler RC. The familial aggregation of common psychiatric and substance use disorders in the National Comorbidity Survey: a family history study. *Br J Psychiatry.* 1997;170:541-548.
9. Heath AC, Bucholz KK, Madden PA, Dinwiddie SH, Slutske WS, Bierut LJ, Statham DJ, Dunne MP, Whitfield JB, Martin NG. Genetic and environmental contributions to alcohol dependence risk in a national twin sample: consistency of findings in women and men. *Psychol Med.* 1997;27:1381-1396.

10. Anjos S, Polychronakos C. Mechanisms of genetic susceptibility to type I diabetes: beyond HLA. *Mol Genet Metab.* 2004;81:187-195.
11. Joy T, Lahiry P, Pollex RL, Hegele RA. Genetics of metabolic syndrome. *Curr Diabetes Rep.* 2008;8:141-148.
12. Blakemore AI, Buxton JL. Obesity, genetic risk, and environment. *Br Med J.* 2004;348:g1900.
13. Jones KL. The effects of alcohol on fetal development. *Birth Defects Res, Part C.* 2011;93:3-11.
14. Lange S, Probst C, Gmel G, Rehm J, Burd L, Popova S. Global Prevalence of Fetal Alcohol Spectrum Disorder Among Children and Youth: A Systematic Review and Meta-analysis. *JAMA Pediatr.* 2017;17:948-956.
15. Roozen S, Peters GJ, Kok G, Townend D, Nijhuis J, Curfs L. Worldwide prevalence of fetal alcohol spectrum disorders: A systematic literature review including meta-analysis. *Alcohol Clin Exp Res.* 2016;40:18-32.
16. Carter RC, Jacobson JL, Molteno CD, Dodge NC, Meintjes EM, Jacobson SW. Fetal alcohol growth restriction and cognitive impairment. *Pediatrics.* 2016;138:1–9.
17. Carter RC, Jacobson JL, Sokol RJ, Avison MJ, Jacobson SW. Fetal alcohol-related growth restriction from birth through young adulthood and moderating effects of maternal prepregnancy weight. *Alcohol Clin Exp Res.* 2013;37:452–62.
18. Moore EM, Riley EP. What happens when children with fetal alcohol spectrum disorders become adults? *Curr Dev Disord Rep.* 2015;2:219–27.
19. Gardejber EM, Cuffe JSM, Ward LC, Steane S, Anderson ST, Dorey ES, Kalish-Smith JJ, Pantalein M, Chong S, Yamada L, Wlodek ME, Bielefeldt-Ohmann H, Moritz Km.

- Effects of periconceptional maternal alcohol intake and a postnatal high-fat diet on obesity and liver disease in male and female rat offspring. *Am J Physiol: Endocrinol Metab.* 2018;315:E694-E704.
20. Dorey ES, Cullen CL, Lucia D, Mah KM, Manchadi MR, Muhlhausler BS, Moritz KM. The impact of periconceptional alcohol exposure on fat preference and gene expression in the mesolimbic reward pathway in adult rat offspring. *J Dev Origins Health Dis.* 2018;9:223-231.
21. Knezovich JG, Ramsay M. The effect of preconception paternal alcohol exposure on epigenetic remodeling of the h19 and rasgrfl imprinting control regions in mouse offspring. *Front Genet.* 2012;3:10.
22. Finegersh A, Rompala GR, Martin DIK, Homanics GE. Drinking beyond a lifetime: New and emerging insights into paternal alcohol exposure on subsequent generations. *Alcohol.* 2015;49:461-470.
23. Jabbar S, Chastain LG, Gangisetty O, Cabrera MA, Sochacki K, Sarkar DK. Preconception Alcohol Increases Offspring Vulnerability to Stress. *Neuropsychopharmacology.* 2016;41:2782-2793.
24. Rompala GR, Homanics GE. Intergenerational Effects of Alcohol: A Review of Paternal Preconception Ethanol Exposure Studies and Epigenetic Mechanisms in the Male Germline. *Alcohol: Clin Exp Res.* 2019;43:1032-1045.
25. Kim P, Choi CS, Park JH, Joo SH, Kim HM, Ko HM, Kim KC, Jeon SJ, Park SH, Han SH, Ryu JH, Cheong JH, Han JY, Ko KN, Shin CY. Chronic exposure to ethanol of male mice before mating produces attention deficit hyperactivity disorder-like

- phenotype along with epigenetic dysregulation of dopamine transporter expression in mouse offspring. *J Neurosci Res.* 2014;92:658-670.
26. Asimes A, Torasco A, Pinceti E, Kim CK, Zeleznik-Le NJ, Pak TR. Adolescent binge-pattern alcohol exposure alters genome-wide DNA methylation patterns in the hypothalamus of alcohol-naïve male offspring. *Alcohol.* 2017;60:179-189.
27. Przybycien-Szymanska MM, Rao YS, Prins SA, Pak TR. Parental Binge Alcohol Abuse Alters F1 Generation Hypothalamic Gene Expression in the Absence of Direct Fetal Alcohol Exposure. *PloS One.* 2014;9:e89320.
28. Chang RC, Thomas KN, Bedi YS, Golding MC. Programmed increases in LXR α induced by paternal alcohol use enhance offspring metabolic adaptation to high-fat diet induced obesity. *Molecular metabolism*, 2019a. 30:161-172.
29. Chang RC, Wang H, Bedi Y, Golding MC. Preconception paternal alcohol exposure exerts sex-specific effects on offspring growth and long-term metabolic programming. *Epigenetics Chromatin* 2019b, 12(1):9.
30. Rok-Bujko P, Dyr W, Kostowski W. Operant self-administration of ethanol in Warsaw High-Preferring (WHP) and Warsaw Low-Preferring (WLP) lines of rats. *Pharmacological Reports*; 2006;58:931-935.
31. Dyr W, Kostowski W. Warsaw high-preferring (WHP) and Warsaw low-preferring (WLP) lines of rats selectively bred for high and low voluntary ethanol intake: preliminary phenotypic characterization. *Alcohol.* 2008;42:161-170.
32. Zakrzewska A, Kuczyńska J, Dermanowski M, Wyszogrodzka E, Dominiak M, Mierzejewski P. Levetiracetam concentration in hair is associated with the time schedule of administration: Study on rats. *Basic Clin Pharmacol Toxicol.* 2020;126:51–58.

33. Mikołajczak P, Okulicz-Kozaryn I, Kamińska E, Wiktorowicz K, Leśniewska K, Dyr W, Kostowski W. Effect of subchronic ethanol treatment on plasma and cerebrospinal fluid leptin levels in rats selectively bred for high and low alcohol preference. *Pol J Pharmacol.* 2002;54:127-132.
34. Hynes GR, Jones PJ. Leptin and its role in lipid metabolism. *Curr Opin Lipidol.* 2001;12(3):321-327.
35. Otaka M, Konishi N, Odashima M, Jin M, Wada I, Matsuhashi T, Ohba R, Watanabe S. Effect of alcohol consumption on leptin level in serum, adipose tissue, and gastric mucosa. *Dig Dis Sci*,52 (2007)3066-3069.
36. Tan X, Sun X, Li Q, Zhao Y, Zhong W, Sun X, Jia W, McClain CJ, Zhou Z. Leptin Deficiency Contributes to the Pathogenesis of Alcoholic Fatty Liver Disease in Mice. *The American Journal of Pathology*,181, 2012:1279-1286.
37. Lindenstrøm E, Boysen G, Nyboe J. Influence of total cholesterol, high density lipoprotein cholesterol, and triglycerides on risk of cerebrovascular disease: the Copenhagen City Heart Study. *Br Med J.* 1994;309:11-15.
38. Langsted A, Freiberg JJ, Tybjaerg-Hansen A, Schnohr P, Jensen GB, Nordestgaard BG. Nonfasting cholesterol and triglycerides and association with risk of myocardial infarction and total mortality: the Copenhagen City Heart Study with 31 years of follow-up. *J Intern Med.* 2010;270:65-75.
39. Triglyceride Coronary Disease Genetics Consortium and Emerging Risk Factors Collaboration, Sarwar N, Sandhu MS, Ricketts SL, Butterworth AS, Di Angelantonio E, Boekholdt SM, Ouwehand W, Watkins H, Samani NJ, Saleheen D, Lawlor D, Reilly MP,

- Hingorani AD, Talmud PJ, Danesh J. Triglyceride-mediated pathways and coronary disease: collaborative analysis of 101 studies. *Lancet*. 2010;375:1634-1639.
40. Lawrence de Koning AB, Werstuck GH, Zhou J, Austin RC. Hyperhomocysteinemia and its role in the development of atherosclerosis. *Clin Bioch*. 2003;36:431-441.
41. Guthikonda S, Haynes WG. Homocysteine: role and implications in atherosclerosis. *Curr Artheroscler Rep*. 2006;8:100-106.
42. Van der Gaag MS, Ubbink JB, Sillanaukee P, Nikkari S, Hendriks HF. Effect of consumption of red wine, spirits, and beer on serum homocysteine. *Lancet*. 2000; 355:1522.
43. Gibson A, Woodside JV, Young IS, Sharpe PC, Mercer C, Patterson CC, McKinley MC, Kluijtmans LA, Whitehead AS, Evans A. Alcohol increases homocysteine and reduces B vitamin concentration in healthy male volunteers - a randomized, crossover intervention study. *QJM*. 2008;101:881-887.
44. Bleich S, Degner D, Wiltfang J, Maler JM, Niedmann P, Cohrs S, Mangholz A, Porzig J, Sprung R, R  ther E, Kornhuber J. Elevated homocysteine levels in alcohol withdrawal. *Alcohol*. 2000;35:351-354.
45. Bleich S, Carl M, Bayerlein K, Reulbach U, Biermann T, Hillemacher T, B  nsch D, Kornhuber J. Evidence of increased homocysteine levels in alcoholism: the Franconian alcoholism research studies (FARS). *Alcohol: Clin Exp Res*. 2005;29:334-336.
46. Grucza RA, Krueger RF, Racette SB, Norberg KE, Hipp PR, Bierut LJ. The emerging link between alcoholism risk and obesity in the United States. *Arch Gen Psychiatry*. 2010;67:1301-1308.

47. Carr LG, Kimpel MW, Liang T, McClintick JN, McCall K, Morse M, Edenberg HJ. Identification of candidate genes for alcohol preference by expression profiling of congenic rat strains. *Alcohol: Clin Exp Res.* 2007;31:1089-1098.
48. Kimpel MW, Strother WN, McClintick JN, Carr LG, Liang T, Edenberg HJ, McBride WJ. Functional gene expression differences between inbred alcohol-preferring and non-preferring rats in five brain regions. *Alcohol.* 2007;41:95-132
49. Stankiewicz AM, Goscik J, Dyr W, Juszczak GR, Ryglewicz D, Swiergiel AH, Wiczorek M, Stefanski R. Novel candidate genes for alcoholism - transcriptomic analysis of prefrontal medial cortex, hippocampus and nucleus accumbens of Warsaw alcohol-preferring and non-preferring rats. *Pharmacol Biochem Behav.* 2015;139:27-38.
50. Shukla SD, Velazquez J, French SW, Lu SC, Ticku MK, Zakhari S. Emerging Role of Epigenetics in the Actions of Alcohol. *Alcohol: Clin Exp Res.* 2008;32:1525–1534.
51. Mahnke AH, Miranda RC, Homanics GE. Epigenetic Mediators and Consequences of Excessive Alcohol Consumption. *Alcohol.* 2017;60:1-6.
52. Chastain LG, Sarkar DK. Alcohol effects on the epigenome in the germline: role in the inheritance of alcohol-related pathology. *Alcohol.* 2017;60:53-66.
53. Pennington JS, Shuvaeva T, Pennington SN. Maternal dietary ethanol consumption is associated with hypertriglyceridemia in adult rat offspring. *Alcohol: Clin Exp Res.* 2002;26:848-55.
54. Teschke R. Alcoholic Liver Disease: Alcohol Metabolism, Cascade of Molecular Mechanisms, Cellular Targets, and Clinical Aspects. *Biomedicines.* 2018;6:106.
55. Fleming TP, Kwong WY, Porter R, Ursell E, Fesenko I, Wilkins A, Miller DJ, Watkins AJ, Eckert JJ. The embryo and its future. *Biol Reprod.* 2004;71(4):1046-1054.

- 451 56. Bielawski DM, Zaher FM, Svinarich DM, Abel EL. Paternal alcohol exposure affects sperm
452 cytosine methyltransferase messenger RNA levels. *Alcohol Clin Exp Res.* 2002;26:347-
453 351.
- 454 57. Kalisch-Smith J, Steane SE, Simmons DG, Pantaleon M, Anderson ST, Akison LK, Wlodek
455 ME, Moritz KM. Periconceptional alcohol exposure causes female-specific perturbations t
456 o trophoblast differentiation and placental formation in the rat. *Development*
457 2019;146:dev172205.

Table 1(on next page)

Biochemical characteristics of the alcohol-naïve offspring of WHP, WLP and alcohol-naïve wild Wistar rats.

*offspring of WHP rats vs offspring of WLP rats; # offspring of WHP rats vs Wistar rats;

†offspring of WLP rats vs Wistar rats; values represent mean (SD); p-value from Newman-Keuls test or Dunn's test (for ALT and triglycerides)

Table 1. Biochemical characteristics of the alcohol-naïve offspring of WHP, WLP and alcohol-naïve wild Wistar rats.

Parameters	WHP (n=8)	WLP (n=8)	Wistar (n=8)	P-value
Homocysteine (μmol/L)	7.48 (1.37)*#	4.77 (1.55)	6.57 (0.68)	*p<0.001; #p=0.009
Glucose (mg/dL)	143.5 (13.55)	149.83 (12.11)	143.16 (8.03)	all p>0.05
Total cholesterol (mg/dL)	79.37 (16.30)*#	34.87 (3.87)†	46.00 (2.72)	*p<0.001; #p<0.001; †p=0.033
Triglycerides (mg/dL)	74.28 (22.94)*#	25.06 (4.63)	37.91 (2.47)	*p<0.001; #p<0.001
ALT (U/L)	97.75 (13.18)*#	74.25 (7.95)†	63.81 (5.18)	*p=0.04; #p<0.001; †p=0.02
AST (U/L)	200.5 (22.78)#	177.66 (29.34)	152.53 (25.87)	#p=0.004
GGT (U/L)	4.95 (0.74)*#	9.85 (0.54)	9.67 (0.72)	*p<0.001; #p<0.001
Amylase (U/L)	2174.37 (359.72)#	2073.25 (189.26)†	1559.37 (131.57)	#p<0.001; †p<0.001

*offspring of WHP rats vs offspring of WLP rats; # offspring of WHP rats vs Wistar rats;
 †offspring of WLP rats vs Wistar rats;
 values represent mean (SD); p-value from Newman-Keuls test or Dunn's test (for ALT and triglycerides)