

Genetic diversity and structure of the critically endangered *Artocarpus annulatus*, a crop wild relative of jackfruit (*A. heterophyllus*)

Leta Dickinson¹, Hilary Noble², Elliot Gardner^{3, 4, 5}, Aida Shafreena Ahmad Puad⁶, Wan Nuur Fatiha Wan Zakaria⁶, Nyree JC Zerega^{1, 2}

¹ Plant Biology and Conservation, Northwestern University, Evanston, IL, United States

² The Negaunee Institute for Plant Conservation Science and Action, Chicago Botanic Garden, Glencoe, IL, United States

³ Department of Biology, Case Western Reserve University, Cleveland, OH, United States

⁴ The Morton Arboretum, Lisle, IL, United States

⁵ Singapore Botanic Gardens, Singapore, Singapore

⁶ Faculty of Resource Science and Technology, Universiti Malaysia Sarawak, Kota Samarahan, Sarawak, Malaysia

Corresponding Author: Nyree JC Zerega
Email address: n-zerega@northwestern.edu

Limestone karsts of Southeast Asia can harbor high levels of endemism, but are highly fragmented, increasingly threatened, and their biodiversity is often poorly studied. This is true of the Padawan Limestone Area of Sarawak, Malaysia, home to the endemic *Artocarpus annulatus*, the closest known wild relative of two important and underutilized fruit tree crops, jackfruit (*A. heterophyllus*) and cempedak (*A. integer*). Identifying and conserving crop wild relatives is critical for the conservation of crop genetic diversity and breeding. In 2016 and 2017, five *A. annulatus* populations were located, and leaf material, locality information, and demographic data were collected. Microsatellite markers were used to assess genetic diversity and structure among populations, and to compare levels of genetic diversity to closely related congeneric species. Results indicate no evidence of inbreeding in *A. annulatus*, and there is no genetic structure among the five populations. However, diversity measures trended lower in seedlings compared to mature trees, suggesting allelic diversity may be under threat in the youngest generation of plants. Also, genetic diversity is lower in *A. annulatus* compared to closely related congeners. The present study provides a baseline estimate of *A. annulatus* genetic diversity that can be used for comparison in future studies and to other species in the unique limestone karst ecosystems. Considerations for *in situ* and *ex situ* conservation approaches are discussed.

Genetic diversity and structure of the critically endangered *Artocarpus annulatus*, a crop wild relative of jackfruit (*A. heterophyllus*)

Leta Dickinson¹, Equal first author, Hilary Noble², Elliot M. Gardner^{3,4,5}, Aida S.A. Puad⁶, W.N.F. Wan Zakaria⁶, Nyree J.C. Zerega^{1,2}, Equal first author

¹ Northwestern University, Plant Biology and Conservation, Evanston, IL, USA

² Chicago Botanic Garden, The Negaunee Institute for Plant Conservation Science and Action, Glencoe, IL, USA

³ The Morton Arboretum, Lisle, Illinois, USA

⁴ Case Western Reserve University, Department of Biology, Cleveland, Ohio, USA

⁵ Singapore Botanic Gardens, Singapore, Singapore

⁶ Faculty of Resource Science and Technology, Universiti Malaysia Sarawak, Kota Samarahan, Sarawak, Malaysia

Corresponding Author:

Nyree J.C. Zerega

Email address: n-zerega@northwestern.edu

Abstract

Limestone karsts of Southeast Asia can harbor high levels of endemism, but are highly fragmented, increasingly threatened, and their biodiversity is often poorly studied. This is true of the Padawan Limestone Area of Sarawak, Malaysia, home to the endemic *Artocarpus annulatus*, the closest known wild relative of two important and underutilized fruit tree crops, jackfruit (*A. heterophyllus*) and cempedak (*A. integer*). Identifying and conserving crop wild relatives is critical for the conservation of crop genetic diversity and breeding. In 2016 and 2017, five *A. annulatus* populations were located, and leaf material, locality information, and demographic data were collected. Microsatellite markers were used to assess genetic diversity and structure among populations, and to compare levels of genetic diversity to closely related congeneric species. Results indicate no evidence of inbreeding in *A. annulatus*, and there is no genetic structure among the five populations. However, diversity measures trended lower in seedlings

compared to mature trees, suggesting allelic diversity may be under threat in the youngest generation of plants. Also, genetic diversity is lower in *A. annulatus* compared to closely related congeners. The present study provides a baseline estimate of *A. annulatus* genetic diversity that can be used for comparison in future studies and to other species in the unique limestone karst ecosystems. Considerations for *in situ* and *ex situ* conservation approaches are discussed.

Introduction

The karst limestone formations of Southeast Asia are dramatic, rugged landscapes that are home to some of the most unique and understudied flora and fauna in the world (Tuyet 2001; Clements et al. 2006; Chung et al. 2014; Hughes 2017). There are approximately 800,000 km² of karst ecosystems in tropical Southeast Asia and southern China, and they include several UNESCO World Heritage sites (Day and Ulrich 2000; Williams 2008). Most karsts were formed millions of years ago by calcium-secreting marine organisms, and the current landscapes are the result of geomorphological processes that have formed complex terrains such as fissured and precipitous cliffs, jagged towers, sinkholes, extensive cave systems, and subterranean rivers, harboring a range of ecological niches (Clements et al. 2006; Williams 2008). The resulting landscapes are naturally fragmented mosaics of karst and ordinary terrain, promoting edaphic isolation and high levels of endemism (Kruckeberg and Rabinowitz 1985; Farjon et al. 2002; Schilthuizen 2004; Musser et al. 2005; Williams 2008). Healthy karst habitats provide tangible benefits to humans, such as water from below-ground aquifers, as well as a range of ecosystem services (Beynan 2011), some of which are due to their unique biodiversity (Schilthuizen 2004; Clements et al. 2006). However, due to human practices, the karst ecosystems and the species inhabiting them are themselves increasingly at risk of further fragmentation and degradation (Clements et al. 2006, Clements et al. 2008, Struebig et al. 2009, Latinne et al. 2011, Hughes et al. 2017). The

most immediate and direct threat is overexploitation of resource extraction, especially limestone and mineral mining (Kiew et al. 2017; Beynan 2011, Hughes et al. 2017). These extremely lucrative but destructive extractive practices can have cascading effects of severe fragmentation and habitat degradation, leading to serious negative impacts on flora, fauna, and humans.

In Sarawak, Malaysia, the Padawan limestone formation (in the southwest part of Kuching Division) is the largest and one of the oldest (dating from about 163 to 100 million years ago) outcrops among the six limestone biodistricts there (Lim 2008) (Cranbrook 2004). One of the species endemic to this habitat is the critically endangered tree, *Artocarpus annulatus* Jarrett, known only from the Padawan-Serian-Tebedu karst forests (Fig. 1). This species is a member of the mulberry (Moraceae) family and the closest wild relative of the important tropical food crops jackfruit (*A. heterophyllus* Lam.) and cempedak (*A. integer* (Thunb.) Merr.) (Zerega et al. 2010; Williams et al. 2017, Zerega et al. 2019). *Artocarpus annulatus* was first described nearly 45 years ago (Jarrett 1975) but remains poorly understood in terms of its biology and distribution. As of 2015, the species was known from only nine herbarium specimens. They were all collected between 1960-1999 and came from just four limestone hill forest outcrops within the Padawan-Serian-Tebedu limestone karst formation in Kuching (Gunung Mentawa, Gunung Gayu, and Gunung Teng Bukap) and Serian (Gunung Payang) Divisions (Fig. 2). According to Zerega et al. (2019), there are thought to be fewer than 50 mature individuals in an area of occupancy (AOO) of 32 km².

The importance of studying and conserving *A. annulatus* has broad implications beyond karst systems, as the genetic diversity of crop wild relatives can prove a useful source of novel traits (Lobell et al. 2008). The economic and food value of its cultivated close relatives, cempedak and jackfruit, are likely to increase, as they grow in tropical regions where food insecurity is high,

and provide greater local autonomy in food production systems (Ford-Lloyd 2011; Jones et al. 2011, Wang et al. 2018, Witehrup et al. 2019). As agriculture adapts to climate change, long-lived tree crops may play an increasingly important role in the global diet. It is known that jackfruit and cempedak are interfertile, because hybrid cultivars exist (Wang et al. 2018). Given that *A. annulatus* is the only other member of the lineage including jackfruit and cempedak (Williams et al. 2017), it is possible that they could all be interfertile and traditional plant breeding could incorporate desirable traits, such as expanding the edaphic conditions of the crop species.

Depending on what we know about the genetic diversity and structure of endangered species, different approaches to conservation may be warranted (Widener and Fant 2018, Münzbergová 2005, Crozier 1997). Due to the unique karst environment, it is possible that *A. annulatus* populations have long been isolated and small. However, human-caused fragmentation may have further reduced genetic diversity, making it harder for the species to rebound and recover (Leimu et al. 2006). While there are limited studies examining population genetics of endangered flora in limestone karst systems of Southeast Asia, the unique habitat and fragmented nature of karst systems have made them of interest for understanding species diversification through isolation. Recent studies of common karst endemic plant genera (*Begonia* and *Primulina*) found substantial differentiation and significant isolation by distance across populations (Gao et al. 2015, Tseng et al. 2019). Understanding genetic patterns and diversity in *A. annulatus* is an important step towards developing conservation strategies for this crop wild relative. This study uses microsatellite and demographic data to assess the genetic diversity and structure of *A. annulatus*, an endemic tree of the Padawan-Serian-Tebedu karst forests of Sarawak, Malaysia. Comparisons of genetic diversity are also made to its two closest relatives, which are commonly occurring

underutilized crops. Specifically, we test the following hypotheses: Given the relatively recent (in decades) destruction of *A. annulatus* habitat, the known populations of *A. annulatus* will show recent evidence of gene flow and as there will not be population specific alleles; due to pressures of increased fragmentation and habitat loss, recruitment of younger individuals may be more tenuous and lead to decreased genetic diversity in younger individuals compared to mature trees; and as a critically endangered species with restricted distribution, *A. annulatus* will harbor less genetic diversity than its widely distributed congeners. It is hoped that the results can inform management plans for *in situ* and *ex situ* conservation efforts and provide much needed information about the unique biota of the Southeast Asian karst formations. Understanding how diversity is distributed among populations will provide insight into the importance of whether *in situ* or *ex situ* conservation efforts should be targeted toward all or some populations to capture the existing diversity of this critically endangered species.

Materials & Methods

Species information

Artocarpus annulatus is a small to mid-sized evergreen tree, growing up to 10 m tall on karst outcrops in Sarawak, Malaysia. It is monoecious, with separate male and female inflorescences occurring on the same individual tree. The male inflorescences have rings around them, for which the specific epithet “*annulatus*” is named (Fig. 1). Once fertilized, the female inflorescence develops into a multiple fruit structure (called a syncarp) that is cauliflorous (i.e. grows directly from the tree trunk) (Fig. 1). Like all members of the genus *Artocarpus*, it produces copious white exudate from all parts when cut. The closest relatives of *A. annulatus* are two important crop species: *A. heterophyllus* (jackfruit) and *A. integer* (cempedak). They also

bear cauliflorous fruit structures, which are edible and much larger than *A. annulatus* (reaching up to 45 kg in *A. heterophyllus*). Jackfruit is thought to be native to the western Ghats of India (with a possible secondary center of diversity in Bangladesh) (Witherup et al. 2019). It is grown throughout much of the tropics today. Cempedak (*A. integer* var. *integer*) is thought to be native to peninsular Malaysia and possibly Borneo; bangkong (*A. integer* var. *silvestris*) is a close wild relative native to peninsular Malaysia (Wang et al. 2018).

Site information

Artocarpus annulatus is known only from the Padawan limestone area in Kuching District, Sarawak, Malaysia (Fig. 2). This area consists of small mountains with tower karst formations. The landscape is dominated by very steep, or even vertical sided limestone “towers,” known as tower karst formations. It is the result of once continuous limestone rock being eroded (Brenda et al, 2004), The towers are sculptured by chemical and mechanical processes forming a terrain covered by jagged pinnacles and deep pits and crevices. *Artocarpus annulatus* can be found at the base of the karst forest mountains on relatively flat ground, as well as at higher elevations among the karst towers and growing sideways from outcrops, with roots clinging to the limestone rock (Fig. 1).

Sampling

In 2016 and 2017, fieldwork was undertaken in the Padawan-Serian-Tebedu karst limestone forests in Sarawak, Malaysia for collection and observation of *A. annulatus* (Zerega et al. 2019). Research permits were approved by the Ibu Pejabat Jabatan Hutan Wisma Sumber Alam (National Parks and Nature Reserves, Forest Department), Sarawak for 2016 (Permit no. NCCD.907.4.4(JLD.13)-195) and renewed for 2017 (Permit no. NPW.907.4.4(JLD.14)-137). All known sites from which *A. annulatus* had previously been recorded (Gunung Gayu, Gunung

Mentawa, Gunung Teng Bukap, and Gunung Payang), as well as four additional sites in surrounding areas of limestone hill forest (Gunung Manok, Gunung Bedoh, Gunung Jambusan, and Fairy Caves) were visited to search for *A. annulatus* individuals (Zerega et al. 2019). Individuals of *A. annulatus* were found in five sites (Gunung Mentawa, Gunung Gayu, Gunung Teng Bukap, Gunung Manok, and Gunung Bedoh) (Fig. 2). Representative collections were made at each site, and the number and age class of individuals from each population varied (Table S1). At one of the sites where collections had previously been made in 1999 (Gunung Payang), the area was found in 2017 to have a small area of limestone hill forest remaining, and a nearby rubber plantation. However, no *A. annulatus* was found (pers. obs. Zakaria). For all sites where *A. annulatus* was found, GPS coordinates, images, and herbarium collections were taken for representative samples. In addition, leaf tissue samples (dried in silica), for 131 *A. annulatus* trees were taken across the five sites (Table S1). For most of these 131 trees, diameter at breast height (dbh) measurements, and notes about the age class of the samples were also taken. The trees sampled were of varying maturity (seedling, sapling, and reproductively mature). Seedlings were defined as having a height less than 1 meter, while mature plants were defined as having a dbh of 5.5 cm or more, because plants that met this criterion showed signs of past reproduction. Anything taller than 1 meter and with a dbh less than 5.5 cm was considered a sapling. The dried leaf material was shipped to the Chicago Botanic Garden where it was stored at -20°C. Additional data for comparison to closely related congeners of *A. annulatus* came from jackfruit (*A. heterophyllus*, Witherup et al. 2019, n=373 from Bangladesh; and Melhem 2015, n=373 from India), cempedak (*A. integer* var. *integer*, n=344), and *A. integer* var. *silvestris* (n=187, bangkong, a wild relative of cempedak) (Wang et al. 2018).

Genetic data collection

Genomic DNA was extracted from 1 cm² dried leaf using either the Qiagen DNeasy kits (Hilden, Germany) following standard protocol or a modified cetyltrimethylammonium bromide (CTAB) method (Doyle and Doyle 1987). Nine different microsatellite primers developed for *A. altilis* (Witherup et al. 2013) were used following the methods of Witherup et al. (2013). These loci previously revealed diversity in the two most closely relatives of *A. annulatus*: jackfruit (*A. heterophyllus*) and cempedak (*A. integer*) (Wang et al. 2018, Witherup et al. 2019). The microsatellite regions amplified in this study were all dinucleotide repeats: MAA26 (black D2 dye), MAA54 (blue D4 dye), MAA105 (black D2 dye), MAA122 (green D3 dye), MAA140 (blue D4 dye), MAA156 (green D3 dye), MAA182 (blue D4 dye), MAA178 (green D3 dye), and MAA196 (black D2 dye). The forward primers were labelled with WellRed fluorescent dyes, shown in parentheses above.

The PCR reactions contained 5 µL of 2x myTaq Mix (Bioline, Taunton, Massachusetts, USA), 0.15 µL 20 mg/ml BSA (bovine serum albumin), 0.25 µL each of labelled forward and reverse primers (10 µM concentration), 3.35 µL of DNA-free water, and 1 µL of DNA (with concentrations between 10 and 80 ng/µL). The PCR cycles were: 2 minutes at 94°C for initial denaturization, followed by 34 cycles of 94°C for 30 seconds of denaturation, 52°C for 1 minute of annealing, then 72°C for 2 minutes of extension; the 34 cycles were then followed by 72°C for 10 minutes of final extension (Wang et al. 2018).

The PCR products were run on a Beckman Coulter CEQ 8000 Genetic Analysis System (Beckman Coulter, Brea, CA) following the methods of Witherup et al. (2013) and multiplexed via the same groupings of Wang et al. (2018). The samples were scored alongside a GenomeLab 400 bp internal size standard ladder that was added to HiDi formamide (20 µL ladder per 1.5 mL of HiDi) (Azco Biotech., San Diego, CA, USA). In each well, 30 µL of the HiDi /ladder mixture

were added to between 1.0 and 2.0 μ L of PCR product, depending on the strength of the fluorescence color. These concentrations were determined by running a few different combinations and dilutions of primers and determining which resulted in the clearest peaks.

The resulting peaks were scored manually using Beckman Coulter CEQ 8000 Software 9.0. Allele sizes are recorded based on the size of the ladder and electrophoresis rates as described in Wang et al. (2018). *Artocarpus annulatus* is thought to be functionally diploid, and as such each region should display either one (homozygous) or two (heterozygous) alleles. Previous studies have revealed that some primers (including primers MAA196 and MAA156) may amplify two regions, and up to four alleles can be present (Wang et al. 2018; Witherup et al. 2013; Witherup et al. 2019). This was rarely observed in *A. annulatus*. In a few samples when more than three strong peaks were observed, the rarest allele was omitted (allele size 268 for primer MAA196; allele size 170 for primer MAA54) (Wang et al. 2018).

Analysis of genetic data

Microchecker analysis (Van Oosterhout et al. 2004), using the Oosterhout method was run to check for null alleles. The microsatellite data were analyzed using GenAlEx v.6.5 (Peakall and Smouse 2012) to calculate diversity measures, including average number of alleles per locus (N_a), number of effective alleles (N_e), observed and expected heterozygosities (H_o and H_e), number of private alleles (P_a), inbreeding coefficient (F_{IS}) and fixation index (F_{ST}). Allelic richness was calculated in the R package Hierfstat (Goudet 2005). To test if diversity measures were significantly different across populations and age classes, single factor ANOVAs were conducted in Excel v. 16.16.14.

Genetic structure among populations was visualized using STRUCTURE v.2.3.4 (Pritchard, Stephens, and Donnelly 2000). Twenty independent runs per K were carried out with a burn-in

period of 10,000 and 10,000 MCMC iterations for $K = 1 - 20$. Structure Harvester (Earl and vonHoldt 2012) was used to determine the most likely value of genetic clusters (K) implementing the Evanno method (Evanno et al. 2005). Spatial clustering of populations was assessed using Principle Coordinate Analysis (PCoA) of Nei's genetic distance in GeneAEx (Peakall and Smouse 2012). Additionally, GENEALEx v.6.5 (Peakall and Smouse 2012) was used to assess genetic differentiation among populations AMOVA. To assess whether there was any genetic differentiation among age classes, PCoA, F_{ST} , and AMOVA analyses were also run for the three age classes in GeneAEx v.6.5 (Peakall and Smouse 2012). For PCoA, the two principle coordinates were plotted with 95% confidence ellipse in R (ggplot) (Wickham 2016).

Results

Microsatellite loci

Of the nine different microsatellite primers that were tested, one (MAA178) consistently amplified two readily distinguishable regions, which were scored as separate loci and differentiated as primer MAA178A and primer MAA178B (following Wang et al. 2018). Across the resulting 10 loci, there were 44 detected alleles, however, two loci were excluded from further analyses as they were monomorphic across all samples (MAA122 and MAA140) (Table S2). Another locus (MAA54) was excluded from further analyses as it was functionally monomorphic, in that nearly all individuals were heterozygotes with the same two alleles, suggesting that this primer may amplify two fixed homozygote loci (MAA54 has been shown to amplify two loci in other *Artocarpus* species) (Wang et al. 2018). Although these three monomorphic loci were excluded from subsequent analyses, the fact that the same loci were found to be polymorphic (with 3 – 18 alleles) in the two closest relatives to *A. annulatus* (Wang et al. 2018, Witherup et al. 2019), suggests that *A. annulatus* is more depauperate in genetic

diversity. Finally, MAA105 was excluded from further analysis due to a null allele frequency value > 0.20 , which can cause significant underestimation of population differentiation (Table S2). This threshold value has been used in other studies to exclude primers that may significantly affect heterozygosity measures (Chapuis and Estoup, 2007; Minn et al., 2014; Muzzalupo et al., 2014). The following loci were included in subsequent genetic diversity and structure analyses: MAA26, MAA156, MAA182, MAA178 A, MAA178 B, MAA196.

Genetic diversity and structure

Analysis of the polymorphic loci with no evidence of null alleles revealed no significant differences in diversity measures across populations (Table 1). The Fixation Index of individuals (F_{IS}) for all populations was negative (indicating no signs of inbreeding), or in the case of Gunung Mentawa it was barely positive (0.05). The population at Gunung Mentawa had no private alleles, while the other four populations had between one (Gunung Manok) and six (Gunung Bedoh) private alleles. Pairwise comparisons of F_{ST} showed limited levels of genetic differentiation across populations (Table 2). Results of AMOVA among and within populations showed that only 8% of diversity can be accounted for among populations, and 92% is within populations (Table 3).

All samples were sorted into age classes as described in the methods. Analysis of the six microsatellite loci revealed no significant differences in diversity measures across size classes, except in the case of private alleles; seedlings had significantly fewer private alleles than saplings or mature trees (Table 4). Pairwise comparisons of F_{ST} across tree size class showed no signs of genetic differentiation across size classes (Table 2). Results of AMOVA among and within tree size classes indicated that only 6% of diversity can be accounted for among populations, and 94% is within populations (Table 3).

The first two principle coordinates of the PCoA account for 25% and 10%, respectively, of the variability in the dataset, and there are no clear clusters based on populations or tree size class (Fig. 3). The population at Gunung Mentawa occupies the smallest space, as does the mature tree size class. STRUCTURE was run on all samples with six loci, and results were analyzed using the Evanno method in Structure Harvester (Earl and vonHoldt 2012) to determine the most probable number of genetic clusters (K). The most likely number was $K = 3$. The structure plot shows that the genetic clusters are restricted to neither distinct geographic populations nor size classes of trees, with all three clusters being fairly equally present in each population and each size class (Fig. S1).

Genetic diversity comparisons with closely related species

Expected heterozygosities (H_e) and inbreeding coefficients (F_{IS}), based on the same six microsatellite loci, were calculated for *A. annulatus* and its closest relatives: *A. integer* var. *integer* (cempedak), *A. integer* var. *silvestris* (bangkong, a wild relative of cempedak) (Data from Wang et al. 2018), and *A. heterophyllus* (jackfruit) (Data from Witherup et al. 2019). A one-way ANOVA found no significant difference ($p=0.113$) across the species for H_e , though *A. annulatus* tended to have the lowest values (Table 5). A one-way ANOVA did find significant difference ($p=0.020$) across species for F_{IS} , and a posthoc Tukey HSD test then found significant differences between *A. annulatus* and the other three taxa (*A. heterophyllus*, $p<0.01$; *A. integer* var. *integer*, $p=0.013$; *A. integer* var. *silvestris*, $p=0.039$)

Discussion

Genetic diversity of closely related *Artocarpus* species

Understanding genetic diversity and its spatial and temporal structure within endangered species and populations is important in order to make sound conservation management plans, especially for imperiled species with very restricted habitats. When starting with zero information about the population genetics of a species, assessing standard measures of genetic diversity is important to establish a baseline of information, but it is also valuable to consider those measures in a broader context. Looking at closely related, more common congeners allows for an evolutionary, comparative perspective. For example, Widener and Fant (2018) found higher levels of inbreeding in an endangered edaphic specialist compared to its more common congener. In the case of *A. annulatus* and its closest relatives, when considering measures of diversity (H_e) of its most closely related congeners (*A. heterophyllus*, *A. integer* var. *integer*, and *A. integer* var. *silvestris*) using the same six microsatellite markers, *A. annulatus* tends to have lower measures, but there is no significant difference ($p = 0.597$) (Table 5). We also considered broader comparisons with other long-lived, outcrossing wild trees and closely related domesticates. Miller and Gross (2011) reviewed genetic diversity in perennial crops and their wild relatives and reported that based on neutral codominant markers (SSRs and allozymes) cultivated perennial fruit crops retain on average 95% (from 65% - 127%) of the neutral variation found in wild populations. Cultivated *Artocarpus* crops had elevated levels of genetic variation compared to the closely related *A. annulatus* (Table 5). This could be an artifact of smaller sampling size in *A. annulatus*, or it could reflect loss of diversity due to habitat destruction. Another possible explanation is that the cultivated populations may represent descendants of crosses between geographically and genetically distinct individuals, giving rise to novel diversity. Studies have demonstrated that allelic diversity in small populations are likely to be very low simply by chance (Cole 2003), which may be the case for *A. annulatus*.

Despite the lower than expected levels of diversity, *A. annulatus* does not show signs of inbreeding, as evidenced by the negative value of the F_{IS} compared to its three relatives—displaying F_{IS} values between 0.149 and 0.314 (Table 5). The results might seem counter-intuitive for such small and restricted populations. However, since most of the variation present in *A. annulatus* is distributed within populations (93%), and because *A. annulatus* is a long-lived, likely outcrossing tree (as are other members of the genus), these results are put into perspective. Genetic diversity levels could be remnant from when the population of *A. annulatus* may have been larger and may have had higher levels of diversity, and the remaining individuals still retain this because of their lifespan (Guries and Ledig 1979). Additionally, it is possible that *A. annulatus* shows no signs of inbreeding, because long-lived perennials often exhibit stronger selection than non-perennials against inbred plants. Indeed, most woody perennials exhibit high outcrossing rates to maintain greater levels of diversity (Duminil, Hardy, and Petit 2009). Meanwhile, crops like jackfruit (*A. heterophyllus*) are under human selection and are frequently vegetatively propagated (i.e. grafting) to maintain desired traits (Witherup et al. 2019).

Genetic structure within and across *Artocarpus annulatus* populations

The results of several analyses indicate that there is little to no geographical genetic structure exhibited across the five populations of *A. annulatus*. Principle Coordinate Analysis and Structure analyses indicate there is no genetic differentiation based on the geography of the populations (Fig. 3, Fig. S1). The results from the AMOVA test revealed that only 8% of diversity was found among populations (Table 3), and the F_{ST} values demonstrated low genetic differentiation among populations (Table 2). Values below 0.05 are considered to represent very low to no genetic differentiation (Frankham et al. 2002), and the values in this study ranged from 0.028 to 0.089. The five populations have an area of occupancy of approximately 32 km² (Zerega

et al. 2019). Three of the populations (Gunung Gayu, Gunung Mentawa, and Gunung Teng
Bukap) are geographically closely clustered, each being ca. 0.3 km apart. Gunung Manok is the
most isolated, situated ca. 6 km from the cluster; and Gunung Bedoh is in the middle, ca. 3 km
away from both Gunung Manok and the cluster of three (Fig. 2). Not surprisingly, Gunung
Manok is the most differentiated from the others, being the least differentiated from the closest
population (Gunung Bedoh). These results, taken in aggregate, suggest that gene flow has
occurred among the populations, but that it may be limited across greater distances. The five *A.*
annulatus populations may be part of one largely intermixing population across which genetic
material is shared via gene flow or may represent remnants of populations that were once larger
and less fragmented. Because it is a long-lived tree, the effects of fragmentation and isolation
may be slower manifested in the genetic structure.

It is of interest to consider methods of gene flow to understand what might be causing low
levels of genetic differentiation across populations in such a dramatic landscape as karst forests,
where dispersal ability of inhabitants may be limited (Hughes 2017). The two ways for plants to
spread genes across space are via pollen dispersal (pollination) and fruit/seed dispersal. While we
know virtually nothing about this in *A. annulatus*, we can turn to its closest relatives to consider
possibilities. Recent work has been conducted on elucidating pollination mechanisms in its two
closest relatives, jackfruit (*A. heterophyllus*) and cempedak (*A. integer*). Both of these
monoecious species (separate male and female inflorescences present on the same individual
tree) are thought to be pollinated via a three-way mutualistic relationship between the plant itself,
gall midges (genus: *Clinodiplosis*, Family: Cecidomyiidae, Order: Diptera), and a fungus (yet to
be identified, but possibly Genus: *Choanephora*, Family: Choanephoraceae, Order: Mucorales)
that grows on the male inflorescences (Gardner et al. 2018; Sakai 2000). The adult gall midges

are thought to be attracted to the same volatile compounds that are similarly emitted by both male and female inflorescences. The midges feed on the pollen of male flowers, and at the same time a fungus (which does not appear to infest other parts of the tree) gradually grows over the length of the male inflorescence. Gravid female midges lay their eggs on the fungus and the young larvae feed on the liquid exuded by the fungal mycelia. As the midges seek out male inflorescences for feeding and egg-laying, they sometimes mistakenly visit female flowers (which offer no apparent reward for the midges but mimic the scent of the male flowers) (Gardner et al. 2018), and in the process may transfer pollen to effect pollination. Given that its closest relatives are both pollinated by gall midges, it is hypothesized that *A. annulatus* may share a similar mechanism (Gardner et al. 2018). Gall midges are tiny insects, about 1 mm in length, and have short lifespans (adults live between a few hours to a few days), so they are likely limited in the distances they can travel to disperse pollen (Gardner et al. 2018; Skuhrava 1991). Some of the *A. annulatus* populations in this study are found within just 0.25 km of each other, while the most isolated population (Gunung Manok) is about 3 km to the closet population (Gunung Bedoh) and about 6 km to the next closet populations. Most of what is known about flight range of members of the gall midge family comes from studies of pests. For example, studies of Hessian flies (*Mayetiola destructor*, *Cecidomyiidae*) suggest that they choose when, where, and how far to move based on cues from the environment, and are also affected by wind. They can disperse from a few meters to a few kilometers and exhibit nonrandom movement directed by things like scents or certain wavelengths of color (Harris and Rose 1990, Withers et al. 1997, Anderson et al. 2012). Additionally, female midges are more likely to stay in an area where their plant of interest is located as opposed to moving to an area with less attractive plants (Withers and Harris 1996). We still need to learn more about pollination in *A. annulatus*, but it

seems possible that if it is similar to its closest relatives, gall midges do not typically travel long distances, and the more fragmented *A. annulatus* populations become over time, the more greatly genetically differentiated they could become. This may explain why the most geographically distant population (Gunung Manok) is also the most genetically differentiated from the others.

However, none of the populations are very highly differentiated from one another, and gene flow appears to be occurring (or has recently occurred) among them (Table 2, Fig. S1). This brings us to the other mechanism for gene flow: fruit (i.e. seed) dispersal. The fruits of *A. annulatus* are large (ca. 6 x 8 cm) and distinctive with bumpy or spiky skin and a strong fragrance (Jarrett 1975). Just like its crop relatives, jackfruit and cempedak, the fruit structure grows from the trunk of the tree (Fig. 1), but the wild *A. annulatus* has much smaller and less fleshy fruits than its crop relatives (Jarrett 1975). While research is limited on fruit dispersal in *Artocarpus*, there are studies that suggest fruits are spread by large mammals, and this may also be the case in *A. annulatus*. In India, the related *A. chaplasha* depends on large mammals such as the Asian elephant (*Elephas maximus*), domestic cows (*Bos primigenius*), buffalo (*Bubalus bubalis*), and rhesus macaques (*Macaca mulatta*) as seed dispersers (Sekar 2014). The study species, *A. annulatus*, shares its habitat with a few sizeable mammals, including François's leaf monkey (*Trachypithecus francoisi*) and the serow (*Capricornis sumatraensis*) (Kiew et al. 2017). While anecdotal information from fieldwork suggests that many of the seeds simply germinate under the mother tree, where the fruits drop (i.e. numerous saplings were observed growing in clusters beneath mature trees), it is possible that birds or mammals disperse some *A. annulatus* seeds, accounting for gene flow and low genetic differentiation across sites (Zerega et al. 2019). Future research could focus on better understanding the contributions of these two modes of gene

flow by studying the distribution of genetic structure of seed (maternal chloroplast loci) vs. pollen dispersed alleles.

Genetic structure across age classes of *Artocarpus annulatus*

When an endangered species occurs in habitats that are under threat, it is valuable to know if genetic diversity is being lost over time, such that older, mature individuals harbor greater or unique genetic diversity compared to young seedlings and saplings. As the highly specialized karst habitat becomes increasingly smaller and fragmented due to human pressures, it might be expected that the *A. annulatus* gene pool could be reduced due to random, stochastic loss of individuals, leading to a decreased level of diversity in younger generations and a genetic bottleneck. When *A. annulatus* individuals were analyzed by size class (seedling, sapling, or reproductively mature), diversity measures tended to be lower in seedlings compared to saplings and mature trees. However, only the number of private alleles was significantly less in seedlings (Table 4). Additionally, PCoA clustering analysis shows that the age classes do not cluster separately (Fig. 3), and AMOVA (Table 3) suggests that only 6% of diversity was found within age classes. This all suggests that there is no genetic differentiation across size classes, but that allelic diversity may be under threat in the youngest generation of plants. To better understand possible temporal changes, it would be useful to look at a much longer trajectory of time and over a larger space. Assessing the genetic diversity of older herbarium specimens would be one way to help determine if diversity has been lost. Unfortunately, there are only nine historical collections known of *A. annulatus*, the oldest only dates back to the 1960s, and due to its critically endangered status, sampling of herbarium specimens for DNA was not possible. However, the present study provides a baseline estimate of genetic diversity that can be used for comparison of future studies.

Informing conservation strategies

Understanding *A. annulatus* genetic diversity can inform conservation management, including *ex situ* conservation. In the case of *A. annulatus*, different populations were not all strongly genetically differentiated from one another, so there may be little justification for collecting germplasm from multiple individuals from all populations for *ex situ* conservation in local arboreta and conservation collections. Instead, seeds or seedlings from populations with the most individuals or the most accessible population could be targeted for conservation. For *in situ* conservation efforts, all sites should ideally be protected (Crozier 1997), however, focusing conservation efforts on the largest population would likely yield similar results in terms of conserving the breadth of *A. annulatus* genetic diversity. In many cases, *in situ* conservation may be the preferred approach. A recent meta-analysis of 32 plant species from various habitats compared how well plants grew in their local habitats versus a foreign transplant site. They found that in over 70% of the studies the local plants consistently did better in their habitats of origin, further supporting the argument for conservation of karst areas (Leimu et al. 2008). Additionally, the unique karst habitat is difficult to replicate, and due to the economic importance and biodiversity that karsts hold, *in situ* conservation is especially important as it conserves an entire ecosystem.

Conclusions

The threats to limestone karst ecosystems in Southeast Asia are immense, and we still have much to learn about them (Clements et al. 2006, Hughes et al. 2017). Laws to protect karst ecosystems are severely lacking, and policies in place are often not enforced (Clements et al. 2006). Unfortunately, the lack of information about the ecosystems and the biodiversity they

house can make it more difficult to justify their conservation. For this reason, it is important to conduct research in these ecosystems, and by combining available data across studies it can help understand patterns and prioritize regions for protection. *Artocarpus annulatus* is just one of many endemic species in an ecosystem that is as incredibly unique and diverse as it is fragile, and points to the need for protection of the karst forests of Southeast Asia. The health of *A. annulatus*, as a long-lived, fruit-bearing tree, is crucial for the complex web of plants and animals that depend upon it. Finally, given that *A. annulatus* is the closest wild relative of two important fruit tree species, it may be valuable for breeding programs to expand the growing conditions of jackfruit and cempedak to limestone soils.

Acknowledgements

We would like to thank Jeremie Fant, Fabrizio Grassi, and two anonymous reviewers for valuable and insightful comments that improved this manuscript; Lynnaun Johnson for assistance in the lab; Jegong anak Suka and Jugah anak Tagi for assistance in the field; and the following herbaria for access to collections: SAR, KEP, L, K, HUH.

References

- Anderson KM, Hillbur Y, Reber J, Hanson B, Ashley RO, Harris MO (2012) Using sex pheromone trapping to explore threats to wheat from Hessian fly (Diptera: Cecidomyiidae) in the Upper Great Plains. *J. Econ. Entomol* 105: 1988–1997
<https://doi.org/10.1603/EC12011>
- Beynan PE (2011) *Karst Management*, pp. 489. Springer. Netherlands.

- 470 Brenda RM, Gendang R, Ambun AU (2004) Geology and geochemistry of the limestone in
471 Sarawak. In: Yong HS, Ng FSP and Yen EEL (eds.) Sarawak Bau Limestone Biodiversity.
472 Sarawak Museum Journal 59, no. 80; Special Issue 6: 41–61
- 473 Chapuis MP, Estoup A (2007) Microsatellite null alleles and estimation of population
474 differentiation. *Molecular Biology and Evolution* 24: 621-631
475 <https://doi.org/10.1093/molbev/msl191>
- 476 Chung KF, Leong WC, Rubite RR, Repin R, Kiew R, Liu Y, Peng CI (2014) Phylogenetic
477 analyses of *Begonia sect. Coelocentrum* and allied limestone species of China shed light on
478 the evolution of Sino-Vietnamese karst flora. *Botanical Studies* 55: 1
479 <https://doi.org/10.1186/1999-3110-55-1>
- 480 Cole CT (2003) Genetic Variation in Rare and Common Plants. *Annual Review* 34: 213-237.
481 <https://doi.org/10.1146/annurev.ecolsys.34.030102.151717>
- 482 Clements R, Sodhi NS, Schilthuizen M, Ng PKL (2006) Limestone Karsts of Southeast Asia:
483 Imperiled Arks of Biodiversity. *BioScience* 56 (9): 733-742 [https://doi.org/10.1641/0006-3568\(2006\)56\[733:LKOSAI\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2006)56[733:LKOSAI]2.0.CO;2)
484
- 485 Clements R, Ng PK, Lu XX, Ambu S, Schilthuizen M, Bradshaw CJ (2008) Using
486 biogeographical patterns of endemic land snails to improve conservation planning for
487 limestone karsts. *Biological Conservation* 141: 2751-2764
488 <https://doi.org/10.1016/j.biocon.2008.08.011>
- 489 Cranbrook EO (2004) A history of animal diversity of the Bau Limestone Area. Sarawak Bau
490 limestone biodiversity. *The Sarawak Museum Journal* 80(6): 193–220

491 Crozier RH (1997) Preserving the Information Content of Species: Genetic Diversity,
492 Phylogeny, and Conservation Worth. *Annual Review of Ecology and Systematics* 28: 243-
493 268 <https://doi.org/10.1146/annurev.ecolsys.28.1.243>

494 Day M, Urich P (2000) An assessment of protected karst landscapes in South- east Asia. *Cave*
495 *Karst Sci* 27(2):61–70

496 Doyle J, Doyle JL (1987) Genomic plant DNA preparation from fresh tissue-CTAB method.
497 *Phytochemical Bulletin* 19: 11–15

498 Duminil J, Hardy OJ, Petit RJ (2009) Plant traits correlated with generation time directly affect
499 inbreeding depression and mating system and indirectly genetic structure. *BMC*
500 *Evolutionary Biology* <https://doi.org/10.1186/1471-2148-9-177>

501 Earl DA, vonHoldt BM (2012) STRUCTURE Harvester: a website and program for visualizing
502 STRUCTURE output and implementing the Evanno method. *Conservation Genetics*
503 *Resources* vol 4 (2): 359-361 <https://doi.org/10.1007/s12686-011-9548-7>

504 Evanno G, Regnaut S, Goudet J (2005) Detecting the number of clusters of individuals using the
505 software STRUCTURE: a simulation study. *Molecular Ecology* 8: 2611-2620
506 <https://doi.org/10.1111/j.1365-294X.2005.02553.x>

507 Farjon A, Nguyen TH, Harder DK, Phan KL, Averyanov LV (2002) A new genus and species in
508 Cupressaceae (Coniferales) from northern Vietnam, *Xanthocyparis vietnamensis*. *Novon* 12:
509 179–189

510 Ford-Lloyd BV, Schmidt M, Armstrong SJ, Barazani O, Engels J, Hadas R, Hammer K, Kell SP,
511 Kang D, Khoshbakht K, Li Y, Long C, Lu BR, Ma K, Nguyen VT, Qiu L, Ge S, Wei W,
512 Zhang Z, Maxted N (2011) Crop Wild Relatives—Undervalued, Underutilized and under
513 Threat? *BioScience* 61 (11): 559-565 <https://doi.org/10.1525/bio.2011.61.7.10>

514 Frankham R, Ballou JD, Briscoe DA (2002) Introduction to Conservation Genetics. Cambridge
 515 University Press, Cambridge

516 Gao Y, Bin A, Kong H, Kang M, Huang H (2015) Geographical pattern of isolation and
 517 diversification in karst habitat islands: a case study in the *Primulina 23burnean* complex.
 518 Journal of Biogeography 42 (11): 2131-2144 doi: [10.1111/eva.12495](https://doi.org/10.1111/eva.12495)

519 Gardner EM, Gagné RJ, Kendra PE, Montgomery WS, Raguso RA, McNeil TT, Zerega NJC
 520 (2018) A Flower in Fruit's Clothing: Pollination of Jackfruit by a New Species of Gall
 521 Midge. Chicago: Int. J. Plant Sci, University of Chicago
 522 <https://www.journals.uchicago.edu/doi/10.1086/697115>

523 Goudet, J (2005) Hierfstat, a package for R to compute and test hierarchical F-statistics.
 524 *Molecular Ecology Notes* 5: 184-186.

525 Guries RP, Ledig FT (1979) Genetic Structure of Populations and Differentiation in Forest
 526 Trees. Proceedings of the Symposium on Isozymes of North American Forest Trees and
 527 Forest Insects, p. 42.

528 Harris MO, Rose S (1990) Chemical, color, and tactile cues influencing oviposition behavior of
 529 the Hessian fly (Diptera, Cecidomyiidae). *Environ. Entomol.* 19: 303–308
 530 <https://doi.org/10.1093/ee/19.2.303>

531 Hughes AC (2017) Understanding the drivers of Southeast Asian biodiversity loss. *Ecosphere* 8:
 532 1

533 Jarrett FM (1975) Four New *Artocarpus* Species From Indo-Malesia (Moraceae). *Blumea* 22:
 534 409-410

535 Jones AMP, Ragone D, Tavana NG, Bernotas DW, Murch SJ (2011) Beyond the Bounty:
536 Breadfruit (*Artocarpus altilis*) for food security and novel foods in the 21st Century.
537 Ethnobotany Research and Applications.

538 Kiew R, Rafidah AR, Ong PT, Ummul-Nazrah AR (2017). Limestone treasures: Rare plants in
539 Peninsular Malaysia – What they are, where they grow and how to conserve them?
540 Malaysian Naturalist 71: 32–41

541 Kruckeberg AR, Rabinowitz D (1985) Biological aspects of endemism in higher plants. Annual
542 Review of Ecology and Systematics 16: 447–479
543 <https://doi.org/10.1146/annurev.es.16.110185.002311>

544 Latinne AS, Waengsothorn VH, Michaux JR (2011) Evidence of complex phylogeographic
545 structure for the threatened rodent *Leopoldamys neilli*, in Southeast Asia. Conservation
546 Genetics 12: 1495-1511 <https://doi.org/10.1007/s10592-011-0248-3>

547 Leimu R, Mutikainen P, Koricheva J, Fischer M (2006) How general are positive relationships
548 between plant population size, fitness and genetic variation? Journal of Ecology 94: 942-952
549 [10.1111/j.1365-2745.2006.01150.x](https://doi.org/10.1111/j.1365-2745.2006.01150.x)

550 Lim M Y L (2008). Diversity of limestone orchids in Central and Northern Padawan, Kuching,
551 Sarawak. MSc. Diss., Universiti Putra Malaysia

552 Lobell DB, Burke MB, Tebaldi C, Mastrandrea MD, Falcon WP, Naylor RL (2008) Prioritizing
553 Climate Change Adaptation Needs for Food Security in 2030. Science 319 no. 5863: 607-
554 610 DOI: 10.1126/science.1152339

555 Melhem, T. 2015. Origin and diversity of jackfruit (*Artocarpus heterophyllus*, Moraceae). M.Sc.
556 thesis, Northwestern University, Evanston, IL, USA.

557 Miller A, Gross BL (2011) From forest to field: Perennial fruit crop domestication. *American*
558 *Journal of Botany* 98 (9): 1389-414 doi: 10.3732/ajb.1000522

559 Münzbergová Z (2005) Determinants of Species Rarity: Population Growth Rates of Species
560 Sharing the Same Habitat. *American Journal of Botany* 92: 1887-1994 doi:
561 10.3732/ajb.92.12.1987

562 Musser GG, Smith AL, Robinson MF, Lunde DP. 2005. Description of a new genus and species
563 of rodent (Murinae, Muridae, Rodentia) from the Khammouan Limestone National
564 Biodiversity Conservation Area in Lao PDR. *American Museum Novitates* 497: 1–31
565 <http://digitallibrary.amnh.org/handle/2246/5666>

566 Nielsen SV, Oliver PM. 2017 Morphological and genetic evidence for a new karst specialist
567 lizard from New Guinea (*Cyrtodactylus*: Gekkonidae). *R. Soc. Open sci.* 4: 170781
568 <https://doi.org/10.1098/rsos.170781>

569 Peakall R, Smouse PE (2012) GenAlEx 6.5: Genetic Analysis in Excel. *Population Genetic*
570 *Software for Teaching and Research—An Update.* *Bioinformatics* 28: 2537-2539
571 <https://doi.org/10.1093/bioinformatics/bts460>

572 Pritchard JK, Stephens M, Donnelly P (2000). Inference of population structure using multilocus
573 genotype data. *Genetics* 155: 945-959

574 Schmid RB, Knutson A, Giles KL, McCornack BP (2018) Hessian Fly (Diptera: Cecidomyiidae)
575 Biology and Management in Wheat. *Journal of Integrated Pest Management* 9 (1): 14
576 <https://doi.org/10.1093/jipm/pmy008>

577 Sakai S, Kato M, Nagamasu H (2000) Artocarpus (Moraceae)-gall midge pollination mutualism
578 mediated by a male- flower parasitic fungus. *American Journal of Botany* 87 (3): 440-445
579 <https://doi.org/10.2307/2656640>

Schilthuizen M (2004) Land snail conservation in Borneo: Limestone outcrops act as arks. Journal of Conchology Special Publication 3: 149–154 [https://doi.org/10.1641/0006-3568\(2006\)56\[733:LKOSAI\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2006)56[733:LKOSAI]2.0.CO;2)

Sekar N (2014) Asian elephants are essential as seed dispersers in a disturbed tropical forest. Dissertation, Princeton University

Skuhrava M (1991) Gall Midges (Diptera: Cecidomyiidae) in forest ecosystems. In Baranchikov, Yuri N.; Mattson, William J.; Hain, Fred P.; Payne, Thomas L., eds. Forest Insect Guilds: Patterns of Interaction with Host Trees; pp 293-297. Invited keynote 1989 August 13-17; U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station.

Struebig MJ, Kingston T, Zubaid A, LeComber SC, Mohd-Adnan A, Turner A, Rossiter SJ (2009) Conservation importance of limestone karst outcrops for Palaeotropical bats in a fragmented landscape. Biological Conservation 142: 2089-2096 <https://doi.org/10.1016/j.biocon.2009.04.005>

Tuyet D (2001) Characteristics of karst ecosystems of Vietnam and their vulnerability to human impact. Acta Geologica Sinica 75: 325-329 <https://doi.org/10.1111/j.1755-6724.2001.tb00539.x>

Van Oosterhout C, Hutchinson WF, Wills D, Shipley PS (2004) Microchecker: software for identifying and correcting genotyping errors in microsatellite data. Molecular Ecology 4 (3): 535-538 <https://doi.org/10.1111/j.1471-8286.2004.00684.x>

Wang MMH, Gardner EM, Chung RCK, Chew MY, Milan AR, Pereira JT, Zerega NJC (2018) Diversity and origins of an underutilized tropical fruit tree, cempedak (*Artocarpus integer*, Moraceae), and its wild relative bangkong (*Artocarpus integer* var. *silvestris*). American Journal of Botany: 898-914 <https://doi.org/10.1002/ajb2.1094>

Wickham H (2016) ggplot2. Elegant Graphics for Data Analysis. Springer-Verlag New York

Widener L, Fant JB (2018) Genetic differentiation and diversity of two sympatric subspecies of *Castilleja affinis*; a comparison between the endangered serpentine endemic (ssp. *Neglecta*) and its widespread congener (ssp. *affinis*). Conservation Genetics 19 (2) 365-381

<https://doi.org/10.1007/s10592-017-1009-8>

Williams EW, Gardner EM, Harris R 3rd, Chaveerach A, Pereira JT, Zerega NJ (2017) Out of Borneo: biogeography, phylogeny and divergence date estimates of *Artocarpus* (Moraceae). *Annals of Botany* 119(4):611-627 <https://doi.org/10.1093/aob/mcw249>

Williams P (2008) World Heritage caves and karst: a thematic study. IUCN (International Union for Conservation of Nature) World Heritage Studies Ser. 2, pp. 34.

Withers TM, Harris MO (1996) Foraging for oviposition sites in the Hessian fly: random and non-random aspects of movement. Ecol. Entomol. 21: 382–395

Withers TM, Harris MO, Madie C (1997) Dispersal of mated female Hessian flies (Diptera: Cecidomyiidae) in field arrays of host and nonhost plants. Environ. Entomol. 26: 1247–1257

Wetherup CE, Ragone D, Wiesner-Hanks T, Irish B, Scheffler B, Simpson S, Zee F, Zuberi MI, Zerega NJC (2013) Development of Microsatellite Loci in *Artocarpus altilis* (Moraceae) and Cross-Amplification in Congeneric Species. Applications in Plant Science 1 (7): 1200423

<http://dx.doi.org/10.3732/apps.1200423>

Wetherup CE, Zuberi MI, Hossain S, Zerega NJC (2019) Genetic Diversity of Bangladeshi Jackfruit (*Artocarpus heterophyllus*) over Time and Across Seedling Sources. Journal of Economic Botany 73 (2): 233-248 <https://doi.org/10.1007/s12231-019-09452-5>

625 Tseng YH, Huang HY, Xu WB, Yang HA, Peng CI, Liu Y, Chung KF (2019) Phylogeography
 626 of *Begonia luzhaiensis* suggests both natural and anthropogenic causes for the marked
 627 population genetic structure. *Botanical Studies* 60: 20 [https://doi.org/10.1186/s40529-](https://doi.org/10.1186/s40529-019-0267-9)
 628 [019-0267-9](https://doi.org/10.1186/s40529-019-0267-9)

629 Zerega NJC, Nur Supardi MN, Motley TJ (2010) Phylogeny and recircumscription of
 630 Artocarpeae (Moraceae) with a focus on *Artocarpus*. *Systematic Botany* 35(4): 766-783

631 Zerega N, Gardner E, Wan Zakaria W, Puad ASA (2019) *Artocarpus annulatus*. The IUCN Red
 632 List of Threatened Species [https://dx.doi.org/10.2305/IUCN.UK.2019-](https://dx.doi.org/10.2305/IUCN.UK.2019-2.RLTS.T86528602A86528612.en)
 633 [2.RLTS.T86528602A86528612.en](https://dx.doi.org/10.2305/IUCN.UK.2019-2.RLTS.T86528602A86528612.en)

Figure 1

Artocarpus annulatus

a) Karst limestone habitat in Kuching Division, Sarawak, b) Roots growing on limestone outcrops, c) White exudate found throughout trees, d) Tree, e) Cauliflorous syncarp (multiple fruit structure), f) Close up of syncarp, and g) Male inflorescence with annulate rings. Photo credits: Nyree Zerega (a-c), Michael Lo (d and g) and Benedikt Kuhnhäuser (e and f)




Figure 2

Artocarpus annulatus distribution

a) Malaysia is shown in dark gray; the area in Sarawak where *A. annulatus* populations are found is indicated by the white box, b) Close up of boxed area from above panel shows extant populations of *A. annulatus* (circles), locations of historical specimens (triangles), and locations that were searched in 2016 and no *A. annulatus* were found (squares), c) Close up of boxed area from above panel shows known extant *A. annulatus* populations that were sampled in this study. Map Credit: SimpleMappr (2019), CC 0.




Figure 3

Principle Coordinate Analysis of five *A. annulatus* populations

a) Populations are displayed in different colors, tree size class is represented by different shapes, 95% confidence ellipses are drawn around populations; b) Tree size classes are displayed in different colors; populations are represented by different shapes; 95% confidence ellipses are drawn around tree size classes.




Table 1(on next page)

Genetic diversity measures of *A. annulatus* populations.

Number of individuals sampled in each population by size class. Numbers in parentheses after population names refer to number of individuals from each size class (mat = mature, sap = sapling, sdlg = seedling).

Population	N	N _a	N _e	A _R	H _o	H _e	F _{IS}	I	P _a
Gunung Gayu (5 mat, 16 sap, 1sdlg)	22	3.33	2.01	3.44	0.60	0.49	-0.23	0.85	5
Gunung Bedoh (11 mat, 12 sap, 3sdlg)	26	3.83	1.81	3.14	0.49	0.41	-0.20	0.72	6
Gunung Manok (13 mat, 18 sap)	21	2.00	1.50	2.53	0.42	0.29	-0.36	0.45	1
Gunung Mentawa (6 mat, 3 sap, 6sdlg)	15	2.17	1.56	2.96	0.31	0.29	0.05	0.48	0
Gunung Teng Bukap (6 mat, 20 sap, 21sdlg)	47	3.00	1.71	3.08	0.38	0.35	-0.06	0.63	3
Total (41 mat, 59, sap, 31 sdlg)	131								
P value	NA	0.07	0.31	0.18	0.46	0.40	0.40	0.26	0.20

N_a = no. of different alleles, N_e = no. of effective alleles, A_R = allelic richness, H_o = observed heterozygosity, H_e = expected heterozygosity, F_{IS} = Fixation Index, P_a = number of private alleles, I = Shannon's Informational Index

Table 2 (on next page)

Pairwise comparisons of F_{ST} values

Comparison of *A. annulatus* by population and tree size.

Pairwise F_{ST} Comparison by Populations					
Gunung Gayu (Pop. 1)	Gunung Bedoh (Pop. 2)	Gunung Manok (Pop. 3)	Gunung Mentawa (Pop. 4)	Gunung Teng Bukap (Pop. 5)	
0.00	-	-	-	-	1
0.067	0.000	-	-	-	2
0.118	0.032	0.000	-	-	3
0.101	0.052	0.068	0.000	-	4
0.061	0.032	0.059	0.028	0.000	5
Pairwise F_{ST} Comparison by Tree Size					
Sapling	Seedling	Mature			
0.00	-	-		Sapling	
0.021	0.000	-		Seedling	
0.020	0.014	0.000		Mature	

Table 3(on next page)

AMOVA

Results are shown among and within five *A. annulatus* populations, and among and within tree size classes across all five populations.

AMOVA across five populations					
Source	df	SS	MS	Est. Var.	%
Among populations	4	42.80	10.70	0.17	8
Within populations	257	534.27	2.08	2.08	92
Total	261	577.07	-	2.25	100
AMOVA across three tree size classes					
Among populations	2	28.15	14.08	0.14	6
Within populations	259	548.92	2.12	2.12	94
Total	261	577.07		2.26	100

df = degrees of freedom, SS = sum of squares, MS = mean square, Est. Var. = estimated variance

Table 4(on next page)

Genetic diversity measures by tree size class of *A. annulatus* across five populations.

Total values are listed for sample size (N) and private alleles (P_a). Mean values across all loci are listed for all other measures. If p-value are < 0.05 , different superscript letters indicate significant difference between groups.

Size Class	N	N _a	N _e	A _R	H _o	H _e	F _{is}	I	P _a
Seedling	28	3.00	1.77	3.46	0.41	0.36	-0.16	0.65	0 ^a
Sapling	46	4.33	1.80	4.02	0.45	0.42	-0.06	0.77	8 ^b
Mature	37	4.00	1.73	3.30	0.43	0.39	-0.10	0.70	6 ^b
P value	NA	0.26	0.97	0.12	0.95	0.89	0.79	0.84	0.03

N_a = no. of different alleles, N_e = no. of effective alleles, A_R = allelic richness, H_o = observed heterozygosity, H_e = expected heterozygosity, F_{is} = Fixation Index, P_a = number of private alleles, I = Shannon's Informational Index

Table 5 (on next page)

Comparison of genetic diversity (H_e).

Comparisons are shown between *A. annulatus* and its closest relatives, *A. heterophyllus* (jackfruit) and *A. integer* (cempedak) based on the same microsatellite loci. There is no significant difference between species for H_e , but there is for F_{IS} (indicated by superscript letters).

1

Locus	<i>A. heterophyllus</i> (N = 746)		<i>A. integer</i> var. <i>integer</i> (N=344)		<i>A. integer</i> var. <i>silvestris</i> (N=187)		<i>A. annulatus</i> (N = 131)	
	H_e	F_{IS}	H_e	F_{IS}	H_e	F_{IS}	H_e	F_{IS}
MAA26	0.774	0.106	0.348	0.341	0.480	0.154	0.427	-0.094
MAA156	0.373	-0.084	0.566	0.016	0.613	-0.088	0.287	-0.429
MAA182	0.554	0.223	0.178	0.360	0.594	0.182	0.087	-0.146
MAA178A	0.580	0.638	0.610	0.092	0.553	0.261	0.484	0.091
MAA178B	0.776	0.391	0.558	0.255	0.870	0.207	0.535	-0.105
MAA196	0.730	0.609	0.599	0.165	0.521	0.179	0.497	-0.439
Mean	0.631	0.314^a	0.477	0.205^a	0.605	0.149^a	0.386	-0.184^b

2