

Cocoa colonic phenolic metabolites are related to blood lipid lowering effects and methylxanthine metabolites and insoluble dietary fibre to anti-inflammatory and hypoglycemic effects in humans

Beatriz Sarriá^{Corresp., 1}, Miren Gomez-Juaristi¹, Sara Martínez López¹, Joaquín García Cordero¹, Laura Bravo¹, M^a Raquel Mateos Briz^{Corresp. 1}

¹ Department of Metabolism and Nutrition, Institute of Food Science, Technology and Nutrition (ICTAN-CSIC); Spanish National Research Council (CSIC), Madrid, España

Corresponding Authors: Beatriz Sarriá, M^a Raquel Mateos Briz
Email address: beasarria@ictan.csic.es, raquel.mateos@ictan.csic.es

Background. In many cocoa intervention studies, health outcomes are related to cocoa components without taking into account the bioavailability of the main bioactive components: phenolic compounds and methylxanthines.

Methods. The present work associates the results of bioavailability and randomised controlled crossover studies in humans carried out with similar cocoa products, so that the main phenol and methylxanthine metabolites observed in plasma and urine are associated to the health effects observed in the chronic studies. We outstand that doses of cocoa and consumption rate used are realistic. In the bioavailability study, a conventional (CC) and methylxanthine-polyphenol rich (MPC) cocoa product were used, whereas in the chronic study a dietary fibre-rich (DFC) and polyphenol-rich (PC) product were studied in healthy and cardiovascular risk subjects.

Results and Discussion. The main phenolic metabolites formed after CC and MPC intake, 5-(4'-hydroxyphenyl)- γ -valerolactone-3'-sulfate, 3'-methyl-epicatechin-5-sulfate, 4-hydroxy-5-(4'-hydroxyphenyl)valeric acid-sulfate, 5-phenyl- γ -valerolactone--sulfate and 5-(4'-hydroxyphenyl)- γ -valerolactone-3'-glucuronide, may contribute to the changes in cholesterol (and indirectly HDL-cholesterol) observed after the regular intake of both DFC and PC, in healthy and cardiovascular risk subjects. Whereas, 7-methylxanthine (the main cocoa methylxanthine metabolite) and theobromine, together with its content in insoluble dietary fiber, may be responsible for the decrease of IL-1 β and hypoglycemic effects observed with DFC. With both phenolic and methylxanthine metabolites a strong dose-response effect was observed.

Conclusion. After the regular consumption of both DFC and PC, positive changes were observed in volunteer's lipid profile, which may be related to the long-lasting presence of colonic phenolic metabolites in blood. In contrast, the anti-inflammatory and hypoglycemic effects were only observed with DFC, and these may be related to methylxanthine metabolites, and it is likely that insoluble dietary fibre may have also played a role.

Cocoa colonic phenolic metabolites are related to blood lipid lowering effects and methylxanthine metabolites and insoluble dietary fibre to anti-inflammatory and hypoglycemic effects in humans

Beatriz Sarriá*, Miren Gómez-Juaristi, Sara Martínez-López, Joaquín García-Cordero, Laura Bravo-Clemente and Raquel Mateos*

Department of Metabolism and Nutrition, Institute of Food Science, Technology and Nutrition (ICTAN-CSIC). Spanish National Research Council (CSIC), José Antonio Nováis 10, 28040 Madrid, Spain.

Corresponding Authors:

Beatriz Sarriá and Raquel Mateos

Department of Metabolism and Nutrition, Institute of Food Science, Technology and Nutrition (ICTAN-CSIC). Spanish National Research Council (CSIC), José Antonio Nováis 10, 28040 Madrid, Spain

Beatriz Sarriá (beasarria@ictan.csic.es); Raquel Mateos (raquel.mateos@ictan.csic.es)

Abstract

Background. In many cocoa intervention studies, health outcomes are related to cocoa components without taking into account the bioavailability of the main bioactive components: phenolic compounds and methylxanthines.

Methods. The present work associates the results of bioavailability and randomised controlled crossover studies in humans carried out with similar cocoa products, so that the main phenol and methylxanthine metabolites observed in plasma and urine are associated to the health effects observed in the chronic studies. We outstand that doses of cocoa and consumption rate used are realistic. In the bioavailability study, a conventional (CC) and methylxanthine-polyphenol rich (MPC) cocoa product were used, whereas in the chronic study a dietary fibre-rich (DFC) and polyphenol-rich (PC) product were studied in healthy and cardiovascular risk subjects.

Results and Discussion. The main phenolic metabolites formed after CC and MPC intake, 5-(4'-hydroxyphenyl)- γ -valerolactone-3'-sulfate, 3'-methyl-epicatechin-5-sulfate, 4-hydroxy-5-(4'-hydroxyphenyl)valeric acid-sulfate, 5-phenyl- γ -valerolactone--sulfate and 5-(4'-hydroxyphenyl)- γ -valerolactone-3'-glucuronide, may contribute to the changes in cholesterol (and indirectly HDL-cholesterol) observed after the regular intake of both DFC and PC, in healthy and cardiovascular risk subjects. Whereas, 7-methylxanthine (the main cocoa methylxanthine metabolite) and theobromine, together with its content in insoluble dietary fiber, may be responsible for the decrease of IL-1 β and hypoglycemic effects observed with DFC. With both phenolic and methylxanthine metabolites a strong dose–response effect was observed.

Conclusion. After the regular consumption of both DFC and PC, positive changes were observed in volunteer's lipid profile, which may be related to the long-lasting presence of colonic phenolic metabolites in blood. In contrast, the anti-inflammatory and hypoglycemic effects were only observed with DFC, and these may be related to methylxanthine metabolites, and it is likely that insoluble dietary fibre may have also played a role.

Abbreviation list

Body Mass Index (BMI)
Caffeine (CF)
Catechin (CA)
Conventional cocoa (CC)
Dietary fiber (DF)
Dietary fiber-rich cocoa (DFC)
Epicatechin (EC)
Insoluble DF (IDF)
Malondialdehyde (MDA)
Methyluric acid (MU)
Methylxanthine-polyphenol rich cocoa (MPC)
Paraxanthine (PX)
Polyphenol-rich cocoa (PC)

62 Procyanidin B1 (PB1)
 63 Procyanidin B2 (PB2)
 64 Raw cocoa (RC)
 65 Soluble DF (SDF)
 66 Theobromine (TB)
 67 Theophylline (TP)
 68 Trolox Equivalent (TE)
 69 5-(3',4'-Dihydroxyphenyl)-γ-valerolactone (DHPVL)
 70 4-Hydroxy-5-(Hydroxyphenyl)valeric acid sulfate (HHPVA-sulfate)
 71 5-(4'-Hydroxyphenyl)-γ-valerolactone-3'-sulfate (HPVL-3'-sulfate)
 72 5-(4'-Hydroxyphenyl)-γ-valerolactone-3'-glucuronide (HPVL-3'-glucuronide)
 73 5-(Phenyl)-γ-valerolactone-sulfate (PVL-sulfate)

74

75

76 **Introduction**

77 Cocoa is consumed worldwide in different population groups being Europe the world's
 78 largest cocoa consuming area with a 40% of global consumption (Food and Agricultural
 79 Organization of the United Nations, 2003). In response to consumer's demand, the food
 80 industry continues producing functional cocoa products enriched with bioactive
 81 components, such as dietary fibre, polyphenols or methylxanthines, or making products
 82 without fat, sugar or sweeteners (World Cocoa Conference, 2018).

83 A body of scientific evidence supports the beneficial cardiovascular health effects of
 84 cocoa consumption (Arranz et al., 2013; Ellam & Williamson, 2013; Gómez-Juaristi et
 85 al., 2011; Hooper et al., 2012). However, there are aspects in cocoa's health effects that
 86 need to be clarified. The aim of this work is to look into the effects of cocoa on
 87 cardiovascular health through associating the lipid lowering, anti-inflammatory and
 88 hypoglycemic effects observed in two chronic, randomized, controlled studies, with the
 89 phenolic and methylxanthine metabolites observed in bioavailability studies in humans
 90 with similar cocoa products. It is noteworthy that the cocoa products used in both the
 91 chronic and bioavailability studies, produced by the same cocoa manufacturer, had a

similar cocoa matrix and realistic amounts of cocoa were consumed by the volunteers in both studies.

In many cocoa intervention studies, health outcomes are related to cocoa component intake without taking into account the bioavailability of the main bioactive components in the test food. The present work goes one step ahead and considers the bioavailability of two of the main bioactive components in cocoa products: the phenolic compounds, mainly flavanols epicatechin, catechin and procyanidins (Gómez-Juaristi, et al., 2019) as well as methylxanthines, mainly theobromine followed by theophylline and caffeine (Martínez-López et al., 2014a). The main metabolites observed in plasma and urine are related to the health effects observed in two randomised controlled human studies carried out with cocoa products similar to that used in the bioavailability study, i.e. a functional cocoa product rich in dietary fibre (Sarriá et al., 2014) and a cocoa product rich in polyphenols (Martínez-López et al., 2014b). As said, the cocoa products used in the mentioned studies were provided by a well-known Spanish cocoa product manufacturer who prepared individual sachets containing the dose recommended on the label to prepare the chocolate drink. It is important to note that the doses used are realistic, contrary to many studies that use exaggerated cocoa conditions. Moreover, the consumption rate corresponds to a real consumption pattern, two sachets per day.

Materials & Methods

Characterisation of the cocoa products

Five cocoa powders were analyzed, an un-processed cocoa powder (cocoa RC), and four new soluble cocoa products commercially available in Spain: cocoa DFC [rich in dietary fibre (DF)], cocoa MPC (rich in methylxanthines and cocoa), cocoa CC [conventional cocoa low in sugar (2.8)], and cocoa PC [rich in cocoa, and thus phenolic compounds, and low in sugar (2.8%)].

In all the cocoa products studied, polyphenols and methylxanthines were extracted following a procedure developed by our group (Bravo & Saura-Calixto, 1998). Total polyphenols were determined using Folin-Ciocalteu reagent and gallic acid as standard and the phenolic and methylxanthine composition of the cocoa extracts was

characterized by high-performance liquid chromatography (HPLC) with diode-array detection (DAD) using an Agilent 1200 series equipment, the procedure followed is described in Gomez-Juarist et al., (2019).

The total DF of the cocoa products was analyzed in triplicate from defatted samples following the AOAC method modified in our laboratory (Saura-Calixto et al., 2000) and is described in Sarriá et al., (2012).

The antioxidant capacity of the all the cocoa products was evaluated in the soluble extracts by three different methods. The reducing power of the samples was measured using the ferric reducing/antioxidant power (FRAP) assay (Pulido, Bravo & Saura-Calixto, 2000). The capacity of samples to scavenge the stable radical ABTS was determined by the ABTS discoloration method (Re et al., 1999). And the oxygen radical absorbance capacity (ORAC) was determined according to (Huang et al, 2002). In the three parameters, Trolox was used as standard and results were expressed as μmol of Trolox Equivalent (TE) per gram of dry matter (d.m.) of the product. Additionally, the FRAP and ABTS methods were also used to determine the antioxidant capacity in serum samples obtained in the bioavailability study (see below), and results were expressed as μM TE.

Methylxanthine and polyphenol bioavailability studies

A crossover, single-blind study was carried out in healthy men and women, aged 18-45 years old, with body mass index between 18 and 25 kg/m^2 . They were also non-smoker, non-vegetarian, non-pregnant women, who were not taking any medication or nutritional supplements, not suffering from any chronic pathology or gastrointestinal disorder. Fourteen volunteers gave their written informed consent prior to participation in the bioavailability study, but only thirteen completed the study.

On two different days, separated by a 10-day period, after an overnight fast, volunteers consumed 15 g of the conventional cocoa product low in sugar (CC) and 25 g of a cocoa product enriched in methylxanthines and polyphenols (MPC) in 200 mL of semi-

skimmed milk. A nurse inserted a cannula in the cubital vein of one of their arms and blood samples were collected into EDTA-coated tubes at baseline (t=0) and 0.5, 1, 2, 3, 4, 6 and 8 h after consuming the cocoa drinks and plasma was separated by centrifugation. Analytical methods were optimized to measure by chromatography the absorption of methylxanthines and phenolics in plasma. The two days previous to each intervention, certain polyphenol-rich-foods, such as some fruits, vegetables and their derivate beverages, as well as methylxanthine-rich-foods, such as coffee, tea or chocolate products, were restricted from their diets in order to reduce inter and intraindividual differences. Volunteers were asked to complete a 24 h food intake recall the day before each intervention in order to control any possible food restriction incompliance. The bioavailability studies of polyphenols and methylxanthines in the cocoa products have already been described in detail in Gómez-Juaristi et al., (2019) and Martínez-López et al., (2014a).

Chronic cocoa studies

To evaluate the effects of regularly consuming DFC (Sarriá et al., 2014) and PC (Martínez-López et al., 2014b) on markers of cardiovascular health, two controlled, randomized, crossover studies were carried out in free-living healthy and moderately hypercholesterolemic (2000-2400 mg/L; 5.172-6.206 mmol/L). In both studies the inclusion criteria was: men and women (not including pregnant women), between 18-55 y old, with body mass index (BMI) < 30 kg/m², non-vegetarian, non-smoker, not suffering from any chronic pathology or gastrointestinal disorder. Briefly, each study consisted in a run-in (2 weeks), a control and cocoa intervention (4 weeks each). In Sarriá et al., (2014) volunteers consumed twice a day 15 g of DFC (30 g/day) and in Martínez-López et al., (2014b), two times a day 7.5 g per serving of PC (15 g/day). Along both studies other cocoa products, certain fruits and vegetables rich in polyphenols were restricted. Forty four subjects completed both studies, among these were the 13 subjects who carried out the bioavailability study.

Blood samples, blood pressure, heart rate and anthropometric measurements were taken at baseline and at the end of each intervention. In blood samples, biochemical, inflammatory, oxidation and antioxidant biomarkers were measured, for more details please see Sarriá et al., (2014) and Martínez-López et al., (2014b).

Both chronic and bioavailability studies were conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures were approved by the Clinical Research Ethics Committee of Hospital Universitario Puerta de Hierro Majadahonda in Madrid (Spain). Approvals were obtained by this Hospital because all human studies carried out in Consejo Superior de Investigaciones Científicas (CSIC) are submitted to the Ethics Committee of Hospital Puerta de Hierro, unless there are participants in the project who are affiliated to a hospital with an Ethics Committee, but in the present study this was not the case. Volunteer recruitment was carried out through placing advertisements in the Universidad Complutense campus as well as through giving short talks between lectures.

Results and Discussion

In the recent years, a Spanish cocoa manufacturer has been producing new soluble, powder, cocoa products with added value increasing the content in bioactive compounds, specifically flavanols, dietary fibre and methylxanthines, and lowering the sugar levels trying not to compromise the flavour and texture of the cocoa products but to keep or improve health effects.

Certainly, in cocoa and most cocoa products, the main bioactive components are phenolic compounds, methylxanthines and dietary fibre. Therefore, phenolic compound and methylxanthine bioavailability studies were carried out in humans, using realistic amounts of cocoa products from the same cocoa manufacturer with the purpose to better understand the metabolism of these components and to establish a relationship between the main cocoa phenolic and methylxanthine metabolites (Gómez-Juaristi et al., 2019; Martínez-López et al., 2014a) and the biological effects observed in chronic studies (Sarriá et al., 2014; Bravo & Saura-Calixto, 1998).

Characterization of cocoa products

The polyphenolic composition, according to the Folin-Ciocalteu assay and HPLC-DAD analysis, the methylxanthine and dietary fibre composition are shown in Table 1. Cocoa powder RC showed the highest total polyphenol content which may be explained because product RC is naturally rich, non-processed cocoa. Among the commercialized cocoa rich products, cocoa powder PC showed the highest level of total polyphenols followed by cocoa products MPC and CC, whereas product DFC, which was rich in DF but not in cocoa, showed the lowest polyphenol content. According to the results obtained in cocoa RC and DFC, it seems that the manufacturing process lowers the content of polyphenols approximately 62.6% and from this point cocoa enrichment has taken place to different extents to reach the polyphenol levels described in products PC, CC and MPC. Chromatographic analyses of the cocoa products studied showed that, among the monomeric flavanols, epicatechin (EC) was the most abundant, between 0.33-2.40 mg/g d.m., whereas catechin (CA) ranged between 0.26-0.83mg/g d.m. and, among the dimeric compounds the most abundant was procyanidin B2 (PB2), 0.39-2.04 mg/g d.m., whereas procyanidin B1 (PB1) was between 0-0.41 mg/g d.m. It is noteworthy that the “natural” content of total polyphenols obtained in product R is not reached by any of the cocoa enriched products. EC and CA contents observed in the natural cocoa powder were similar to those described by Miller et al., (2009) in other cocoa products.

Since many years ago, it has been established that food antioxidant capacity is strongly related to its polyphenol content (Benzie & Szeto, 1999). In accordance, in the present work the phenolic content of the studied cocoa powders was directly proportional to the antioxidant activity assessed using FRAP, ABTS and ORAC assays (Table 2), so that FRAP results were 5 times higher than the total polyphenol content of the cocoa products, whereas ABTS and ORAC results were 3-4 and 10-14 times higher, respectively. Taking the present results into account, consuming a dose of 15 g/d of cocoa PC as recommended by the manufacturer would yield 2626 (FRAP method), 1996 (ABTS method) or 3855 (ORAC method) μmol Trolox equivalents (TE) of antioxidant capacity, exceeding that provided by a 100 mL serving of coffee [2267

(FRAP) and 1328 (ABTS) μmol Trolox equivalents, respectively] or red wine [601 (FRAP) and 631 (ABTS) μmol Trolox equivalents, respectively], which are the beverages that have been described to have the highest antioxidant capacity within the Mediterranean diet by Saura-Calixto & Goñi (2006).

Furthermore, soluble cocoa products are also a relevant source of DF, in contrast to chocolate, due to DF being largely discarded in its production (Jenkins et al., 2000). Dietary guidelines recommend a minimum daily intake of DF of 25 g (<https://health.gov/dietaryguidelines/2015/>), which is considerably higher than the estimated intakes in Western countries (Jenkins et al., 2000). Bearing this in mind, the food industry prompted the production of dietary fibre rich foods, such as cocoa product DFC. In all products, insoluble DF (IDF) was the predominant fraction, varying between 73 and 92% of the total DF content, which is in agreement with previous results (Lecumberri et al., 2007) that showed that the main polysaccharides in cocoa IDF were cellulose, hemicellulose and some pectic substances, while cocoa soluble DF (SDF) was composed mainly of pectins and minor amounts of galactomannans.

As for the different cocoa products, it is noteworthy that cocoa product MPC, which is rich in polyphenols and methylxanthines, showed the highest content in total DF (30%), even higher than the fibre-rich cocoa product DFC, which contained 22% total DF. Both products DFC (92%) and MPC (90%) presented a similar proportion of IDF (IDF) with respect to total DF (Table 1). The contribution of SDF to the total dietary fibre was highest in product RC (27%) followed by product PC (20.7%). Interestingly, SDF content in the cocoa products followed the same order than the antioxidant capacity and their total flavanol content ($R > PC > MPC > CC > DFC$).

Regarding the methylxanthine contents in cocoa, theobromine was the primary methylxanthine in all the tested cocoa products (Table 1), followed by caffeine, in agreement with the literature (Carlin-Sinclair & Marc, 2009). Cocoa MPC was the product with the highest content in total methylxanthines, being particularly rich in caffeine, as this product was enriched in kola nut, which is a source of caffeine. Theophylline was only found in products MPC and R, and scarcely in PC, whereas in the other cocoa products the concentrations were under the limit of detection. Cocoa is

also a source of caffeine. A 15 g serving of cocoa MPC would provide 153.6 mg of methylxanthines (69% theobromine and 29.5% caffeine), which is similar to the amount of caffeine provided per serving in 330 mL of a cola drink (40 mg).

Bioavailability and metabolism

It is essential to better understand the bioavailability and metabolism of the main bioactive compounds in cocoa, i.e. polyphenols. However, another question that needs to be further clarified is what other compounds in cocoa, beyond flavanols, are responsible for the beneficial health effects. In this context, methylxanthines constitute another outstanding group of bioactive compounds in cocoa products, although less considered when biological effects of cocoa products on health are discussed. For this reason, it is relevant to explore the main metabolites formed from methylxanthines' bioavailability along with the metabolites derived from metabolism of flavanols, as it can help to better explain the compounds responsible of the health effects associated to cocoa consumption and the involved mechanisms.

In the bioavailability studies presented in this work, a main objective was, besides establishing flavanols and methylxanthines absorption and metabolism after intake of realistic doses of cocoa products, to assess if there was a dose-dependent response. Therefore, the product which showed the highest amount of methylxanthines of all the cocoa products and which was rich in polyphenols (MPC) was selected to compare with the conventional product (CC).

Methylxanthines

The absorption and metabolism of methylxanthines present in the two tested soluble cocoa products, CC corresponding to the conventional cocoa low in sugar and MPC enriched in methylxanthines and polyphenols, was evaluated in healthy subjects (Martínez-López et al., 2014a). The main methylxanthines and their plasma and urine metabolites were characterized after consuming realistic doses of cocoa products (15 g

of CC and 25 g of MPC, in 200 mL of milk) with different methylxanthine composition: CC provided 84.45 mg of theobromine (TB) and 9.9 mg of caffeine (CF) while MPC provided 177 mg TB, 75.75 mg CF and 3.25 mg theophylline (TP).

Metabolism of cocoa methylxanthines in humans has not been widely studied and some results obtained by our group were novel findings. Up to 12 different metabolites were identified and quantified in a single run in biological fluids by HPLC-DAD and LC-QTOF. On one hand, TB, CF, TP and paraxanthine (PX) together with two monomethylxanthines (3-methylxanthine (MX) and 7-methylxanthine) were identified and quantified in plasma by HPLC-DAD. TB the most abundant plasmatic compound of all the identified metabolites, followed by TP and PX with similar levels. These compounds were rapidly detected in plasma, showing their maximum concentration (C_{max}) between 1 to 4 hours after ingestion and maintaining similar levels throughout 8 h, not returning to basal levels in this period. The maximum concentration (C_{max}) and area under curve (AUC) values of all metabolites were significantly higher after the intake of MPC, showing a dose-dependent response.

These metabolites together with 1-methylxanthine and different mono-, di- and trimethyluric (MU) acids (1-MU, 1,3-MU, 1,7-MU, 3,7-MU and 1,3,7-MU) involved in the biotransformation of the ingested methylxanthines were detected by LC-QTOF and quantified by LC-DAD in urine, with 7-MX as the most abundant metabolite followed by TB and 3-MX (Figure 1). Most metabolites were extensively excreted in urine between 4 and 8 hours after CC and MPC intake; although similar to the behaviour described in plasma, higher levels of excreted metabolites were observed until 24 h post-intake. Results showed a strong dose–response effect after cocoa consumption at the doses tested, up to 256 mg of MX. It can be concluded that methylxanthines in cocoa are bioavailable, partially metabolized and rapidly eliminated, with a sustained urinary excretion for as long as 24 h after intake. The long permanence of methylxanthines and their metabolites in body fluids favours the involvement of these compounds on the health effects associated to cocoa products consumption.

Polyphenols

Although there are many bioavailability studies of cocoa flavanols, most of them have been carried out with unrealistic doses (Ottaviani et al., 2012a; Ottaviani et al., 2012b; among others). Addressing the need to carry out studies that follow realistic consumption guidelines to further understand the bioavailability and metabolism of real doses of flavanols, which may differ from higher doses used in previous articles, a recent clinical intervention study was performed in our research group (Gómez-Juaristi et al., 2019). This study was aimed at assessing in healthy humans the absorption and metabolism of polyphenols in a single, realistic dose of CC (15 g) and MPC (25 g), which provided 19.8 mg and 68.25 mg of flavanols, respectively.

Blood and urine samples were taken before and after CC and MPC consumption and analyzed by LC–MS–QToF. Up to 10 and 30 metabolites were identified in plasma and urine, respectively. Among the metabolites identified, phase II derivatives of epicatechin showed the maximum concentration in plasma from 1 to 2 h after ingestion and kinetics compatible with their absorption at small intestine. 3'-Methyl-epicatechin-5-sulfate and epicatechin-3'-sulfate were the predominant metabolites followed by 4'-methyl-epicatechin-5-sulfate, and epicatechin-3'-glucuronide among others. Phase II derivatives of epicatechin accounted for approximately 33% of total urinary metabolites. The most abundant group of metabolites were phase II derivatives of hydroxyphenyl- γ -valerolactones and valeric acid, formed at colonic level as 5C-ring fission metabolites by microbiota-mediated (Figure 1). These compounds showed a delayed kinetic due to their colonic origin and represented the most abundant group of metabolites (67% of total urinary metabolites) headed by 5-(4'-hydroxyphenyl)- γ -valerolactone-3'-sulfate (HPVL-3'-sulfate) followed by 4-hydroxy-5-(hydroxyphenyl)valeric acid sulfate (HHPHVA-sulfate), 5-(phenyl)- γ -valerolactone--sulfate (PVL--sulfate), 5-(4'-hydroxyphenyl)- γ -valerolactone-3'-glucuronide (HPVL-3'-glucuronide) and 5-(3',4'-dihydroxyphenyl)- γ -valerolactone (DHPVL), among others. According to these results, HPVL-3'-sulfate could be a sensitive biomarker of flavanol intake, considering its abundance in urine, amounting up to 40% of total urinary metabolites. Flavanols of CC and MPC showed a dose-dependent absorption with a recovery value of 35% of the ingested dose.

The relationship between the main cocoa metabolites and the cardiovascular health effects

To understand the health effects of cocoa or cocoa-based functional products, it is important to carry out randomized controlled human studies using normal, realistic doses of the food products. Such aspects were considered in two chronic studies carried out in our research group; in the first, the regular consumption of a cocoa product rich in dietary fibre (DFC) (Table 3) during 4 weeks led to a significant increase in HDL-cholesterol and a decrease in glucose and IL-1 β without producing changes in body weight neither other anthropometric parameters studied (Sarriá et al., 2014); in the second, the sustained consumption of the cocoa product rich in cocoa (PC) (Table 3) for the same length of time, 4 weeks, resulted in an increase in HDL-cholesterol without other changes in cardiovascular related biomarkers (Martínez-López et al., 2014b). When the health effects of both cocoa products were comparatively studied in relation to the amounts of bioactive compounds consumed, without considering the bioavailability of these compounds, on one hand, mainly the flavanol content provided by both cocoa products was related to the increase in HDL-cholesterol, and on the other hand, the IDF and theobromine were associated to the hypoglycemic and anti-inflammatory effects observed after consuming the fibre rich product (Sarriá et al., 2015). However, as aforementioned, the present study goes one step ahead and further looks into the relationship between the main phenolic and methylxanthine metabolites observed in bioavailability studies in humans, carried out using similar cocoa products to those used in the crossover, intervention studies.

Main phenolic metabolites after cocoa consumption in humans and their health effects

a) Epicatechin and phase II derivatives

Numerous human dietary intervention studies link the consumption of flavan-3-ols derived from cocoa to improved cardiovascular health (Heiss, Keen & Kelm, 2010; Rodriguez-Mateos et al., 2014), having epicatechin and its phase II derivatives, at least

partially, been linked with these beneficial effects (Borges et al., 2018). In contrast to the results of Dower et al., (2015), who investigated the effects of pure epicatechin supplementation and described no effects on cholesterol, many other studies have correlated the consumption of products rich in epicatechin with a decrease in serum levels of LDL-cholesterol in both hypercholesterolemic subjects (Grassi et al., 2005) and healthy subjects (Baba et al., 2007). The mechanisms responsible for the decrease in the levels of LDL include: inhibiting cholesterol absorption in the gut, reducing cholesterol synthesis by decreasing the activity and/or expression of hydroxymethylglutaryl coenzyme A (HMG-CoA) synthase, HMG-CoA reductase, sterol O-aciltransferase and microsomal triglyceride transport protein in the liver, suppressing hepatic secretion of apolipoprotein B100, increasing the expression of hepatic LDL receptors, and preventing oxidation of LDL (Baba et al., 2007). Separately, epicatechin and catechin, which have similar biological activity, have also shown to raise the concentration of HDL-cholesterol through the increased expression of scavenger receptor B type I (SR-BI), sterol regulatory element binding proteins (SREBPs), ATP binding cassette transporter A1 (ABCA1) or apolipoprotein A1, among others (Martínez-López et al., 2014b). Moreover, Rodriguez-Mateos et al., (2018) recently described that cocoa consumption in healthy humans induced effects on flow mediated dilatation, pulse wave velocity and blood pressure mediated by EC metabolites and not DHPVL metabolites.

In addition, it is important to emphasize epicatechin's antioxidant function, being able to adhere to LDL particles or apolipoprotein B and to recycle molecules of α -tocopherol donating a hydrogen atom, thus maintaining the concentration of endogenous antioxidants for longer time (Wan et al., 2001). This property, combined with the reduction of iron (non-heme) and malondialdehyde (MDA) concentrations and the suppression of pro-oxidant enzymes, have been described to be responsible for the increase in blood antioxidant activity after epicatechin consumption (Prakash, Basavaraj & Murthy 2019). However, not all phase II metabolites have the same antioxidant activity. Epicatechin and its 7-O-glucuronide derivative show a similar reduction in LDL oxidation, unlike 3'-O-glucuronide and 4'-O-methyl-3'-O-glucuronide derivatives, which show lower activity. Similarly, the O-methylated derivatives C-3' and C-4'-O-methyl ether

of epicatechin are less active than their base compound. Moreover, it should not be disregarded that the antioxidant activity of these metabolites is dependent on the pH of the medium, so that at 7.4 their activity is largely retained, which confirms their antioxidant action in physiological conditions (Monagas et al., 2010).

There is controversy regarding the hypoglycaemic activity of epicatechin and its derivatives. According to Rodriguez-Mateos et al., (2018), there were no effects of epicatechin or procyanidins on fasting glucose concentration. In contrast, a study conducted by Josic et al., (2010) in healthy subjects showed that consumption of green tea rich in epicatechin reduced glucose and insulin concentration. The main pathway by which epicatechin and its derivatives exert this effect is through increased sensitivity to insulin in hepatic or adipose cells, in which epicatechin inhibits the action of different signalling proteins such as PKC δ , IKK, JNK, NF- κ B and PTP-18, which leads to a decrease in insulin resistance (Cremonini et al., 2016). In addition, a recent study in mice has observed that the consumption of a food rich in epicatechin promotes the regeneration of cells β -pancreatic (Prakash, Basavaraj & Murthy 2019).

b) 5-(3',4'-dihydroxyphenyl)- γ -valerolactone (DHPV)

5-(3',4'-dihydroxyphenyl)- γ -valerolactone (DHPV) is a compound resulting from the degradation of procyanidins or monomers that may reach the colon and be cleaved by the colonic microbiota (Appeldoorn et al., 2009; Fogliano et al., 2011). Its antioxidant capacity has been tested *in vitro*, being superior to catechin, ascorbic acid and Trolox (vitamin E analogue; Appeldoorn et al., 2009).

Up to date, hardly any *in vivo* study has been conducted to demonstrate the effect of DHPVL on lipid and glycaemic metabolism. Only by extrapolating the results obtained in clinical studies with products rich in procyanidins, such as cocoa (Sarriá et al., 2014; Martínez-López et al., 2014b; Sarriá et al., 2015; Grassi et al., 2005; Baba et al., 2007; Khan et al., 2012), with its antioxidant capacity (Khan et al., 2012), and considering the fact that it is one of the main metabolites found in blood after the consumption of cocoa (Urpi-Sarda et al., 2009a; Urpi-Sarda et al., 2009b) it has been deduced that DHPVL has hypolipemic and hypoglycemic properties. This is in agreement with the results

obtained by Rodriguez-Mateos et al., (2018), who described that the consumption of cocoa procyanidins induced a health benefit in healthy humans related to the reduction of total cholesterol, having an impact on cholesterol absorption and faecal steroid excretion inside the gastrointestinal tract. This effect was particularly linked to procyanidins content, which are metabolized into DHPVL and its phase II derivatives.

Recently, a study carried out in rabbits with obesity and non-alcoholic fatty liver suggested that the effects of procyanidins, metabolized into DHPV derivatives, were due to changes in the intestinal microbiota. In fact, the supplementation with procyanidin B2 decreased the ratio Firmicutes/Bacteroidetes and increased the proportion of Akkermansia, which has the ability to maintain the thickness of intestinal mucus, reducing the permeability to lipopolysaccharides (LPS) and thus relieving inflammation. As a consequence of the maintenance of the intestinal barrier, a decrease in the concentration of LPS in serum was observed, which may explain the reduction in the accumulation of triglycerides in the liver and the protection against non-alcoholic fatty liver (NAFLD) (Xiang et al., 2019).

According to the results on phenolic metabolites (Table 3), it may be proposed that the main metabolites observed after intake of CC and MPC (HPVL-3'-sulfate, 3'-methyl-epicatechin-5-sulfate, HHPHVA-sulfate, PVL--sulfate and HPVL-3'-glucuronide), may be responsible for certain changes observed in the chronic studies with DFC and PC. Specifically, to the changes in cholesterol, and indirectly HDL-cholesterol, observed after regular intake of DFC or PC in both healthy and cardiovascular risk subjects.

Regarding the effects of the cocoa products on serum antioxidant capacity, along the bioavailability study, there were no significant differences between the two cocoa products according to the estimations carried out with FRAP method except at 2 h ($p=0.039$) and 6 h ($p=0.044$), and according to ABTS at 1 h ($p=0.004$) with MPC higher than CC (Table 4). In agreement, no significant differences in serum antioxidant capacity, measured by ORAC, FRAP and ABTS as well as levels of protein (carbonyl groups) and lipid (MDA), were observed in the chronic interventions (Sarriá et al., 2014; Martínez-López et al., 2014b).

475

476 ***Main methylxanthine metabolites after cocoa consumption in humans and their***
 477 ***health effects***

478 All the methylxanthine compounds have in common the capacity to act as
 479 phosphodiesterase inhibitors, modulators of GABA and adenosine receptors, as well as
 480 to regulate intracellular calcium levels (Williams et al., 1978; Monteiro et al., 2016).
 481 However, Williams et al., (1978) described many years ago that there are certain
 482 differences in the inhibitory and regulatory activity, with theophylline and 3-
 483 methylxanthine having greater activity than caffeine and 1-methylxanthine.

484 Traditionally, the HDL cholesterol raising effect of cocoa has been associated with
 485 cocoa flavonols (Baba et al., 2007; Mellor et al., 2010; Neufingerl et al., 2013).
 486 Nevertheless, methylxanthines, particularly theobromine, may increase HDL-
 487 concentration in blood (Neufingerl et al., 2013), having been postulated that the
 488 mechanism of action involves increased levels of apolipoprotein A-1 and is independent
 489 of its activity as an adenosine receptor inhibitor (Monteiro et al., 2016; Monteiro et al.,
 490 2019). It is also important to highlight the properties of methylxanthines in counteracting
 491 hyperglycaemia and insulin resistance. This hypoglycemic effect is explained by the
 492 ability of these compounds to regulate intracellular levels of cAMP, with the release of
 493 insulin by β -pancreatic cells and glucose by the liver being dependent on this second
 494 messenger (Monteiro et al., 2016).

495 To end, the anti-inflammatory properties of methylxanthines should also be
 496 emphasized. While caffeine is able to inhibit the expression of tumour necrosis factor
 497 (TNF)- α by suppressing the cyclic-adenosine monophosphate/protein kinase A
 498 (cAMP/PKA) pathway and inhibiting cAMP phosphodiesterase (Horrigan, Kelly &
 499 Connor, 2004) no variation in the production of interleukin (IL)-1 β , IL-12 and IL-10 has
 500 been observed in human studies (Horrigan, Kelly & Connor, 2004; Lofffield et al., 2015,
 501 Haskó & Cronstein, 2011). In the case of theobromine, a decrease in IL-1 β levels was
 502 reported in studies conducted *in vitro* (Fuggetta et al., 2019) and in mice (Camps-
 503 Bossacoma et al., 2019). In contrast, in humans, no studies with theobromine and

inflammatory markers other than C-reactive protein (CRP) were found. However, two studies, one carried out in moderate hypercholesterolemic and in healthy subjects (Sarriá et al., 2014), observed a decrease in the serum levels of IL-1 β and IL-10 after consumption of a cocoa product rich in fibre, so we cannot rule out a synergistic effect between the two bioactive compounds (Goya et al., 2016).

According to these results on methylxanthine metabolites, 7-methylxanthine (the main cocoa methylxanthine metabolite) and theobromine may be responsible for the antiinflammatory (decrease of IL-1 β) and hypoglycemic effects observed after the intervention with DFC.

Dietary fiber

The remaining bioactive compound in cocoa products, dietary fibre, has certainly played a role on the cardiovascular related effects observed. The different intake of dietary fibre, with DFC (over 10 g/d) and PC (less than 4 g/d) may be related to the decrease in glucose (p=0.029) levels observed only after regularly consuming DFC, but not PC (Heiss, Keen & Kelm, 2010). This result is in agreement with the significant decrease in plasma glucose concentration (p=0.019) observed in a previous study carried out with a dietary fibre-rich cocoa product (Sarriá et al., 2012), where the mechanisms responsible for this effect were suggested to be that dietary fibre rich foods delay glucose absorption from the small intestine (Glacco et al., 2000) and improve insulin sensitivity (Tokede, Gaziano & Djoussé, 2011). since it is well known that dietary fiber rich foods delay glucose absorption from the small intestine and improve insulin sensitivity (Sarriá et al, 2015). However, the higher dietary fibre intake with DFC did not increase HDL-cholesterol, in contrast to results described by Jenkins et al., (2000) who reported a significant increase in HDL-cholesterol after consumption a cocoa-bran (25 g DF/d) for two weeks in healthy subjects.

The effects of DFC on inflammation may be related to the down-regulation of cytokine gene expression and up-regulation of the expression of scgb1a1, gene that codifies a protein related to anti-inflammation, in the colon tissue (Massot-Cladera et al, 2017), as

observed in rats that consumed either a cocoa diet or a cocoa fiber diet. In addition to the positive effects on inflammation and glucose, DFC also produced positive gastrointestinal effects such as an increased number of daily bowel movements and reduced time to have a bowel movement, without inducing major adverse gastrointestinal symptoms (Sarriá et al, 2012).

This study presents the following limitations and strengths: the bioavailability and chronic studies were carried out separately, using different cocoa products although they were produced by the same manufacturer and the cocoa matrix was quite similar. However, it is relevant that health outcomes of chronic cocoa consumption in humans are related to the cocoa's composition taking into account the bioavailability of the main bioactive components, also evaluated in a human study. As aforementioned, no previous studies, to our knowledge, have considered this integrative and associative approach. Both the chronic and bioavailability studies are well designed and they have been carried out in an adequate number of subjects with similar characteristics using realistic amounts of cocoa. Moreover, the present work has been accomplished knowing that a dose-dependent response occurs with both polyphenols and methylxanthines, when the cocoa products were consumed at realistic doses, and the metabolites obtained are the same disregarding the different quantities of the bioactive components in cocoa products studied. It is pertinent to further investigate the results here presented, so that in future chronic cocoa studies in humans a bioavailability study will be performed in a subgroup of volunteers.

Conclusions

After the regular consumption of both DFC and PC, positive changes were observed in volunteer's lipid profile, which may be related to the long-lasting presence of colonic phenolic metabolites (headed by 5-(4'-hydroxyphenyl)- γ -valerolactone-3'-sulfate) in blood. In contrast, the anti-inflammatory and hypoglycemic effects were only observed with DFC, and these may be related to methylxanthine metabolites, particularly 7-methylxanthine which was the main cocoa methylxanthine metabolite, and theobromine, and it is likely that insoluble dietary fibre may have also played a role.

Acknowledgements

We are grateful to volunteers participating in the study. All authors revised and approved the final version of the manuscript.

References

- Appeldoorn M M, Vincken J P, Aura A M, Hollman P C, Gruppen H. 2009.** Procyanidin dimers are metabolized by human microbiota with 2-(3 4-dihydroxyphenyl) acetic acid and 5-(3 4-dihydroxyphenyl)-γ-valerolactone as the major metabolites. *Journal of Agricultural and Food Chemistry* **57**:1084-1092
- Arranz S, Valderas-Martinez P, Chiva-Blanch G, Casas R, Urpi-Sarda M, Lamuela-Raventós RM, Estruch R. 2013.** Cardioprotective effects of cocoa: clinical evidence from randomized clinical intervention trials in humans. *Molecular Nutrition and Food Research* **57**:936-947
- Baba S, Natsume M, Yasuda A, Nakamura Y, Tamura T, Osakabe N, Kanegae M, Kondo K. 2007.** Plasma LDL and HDL cholesterol and oxidized LDL concentrations are altered in normo- and hypercholesterolemic humans after intake of different levels of cocoa powder. *Journal of Nutrition* **137**:1436-1441
- Benzie IF, Szeto YT. 1999.** Total antioxidant capacity of teas by the ferric reducing/antioxidant power assay. *Journal of Agricultural and Food Chemistry* **47**:633-636
- Borges G, Ottaviani JI, Van der Hooft JJJ, Schroeter H, Crozier A. 2018.** Absorption metabolism distribution and excretion of (-)-epicatechin: A review of recent findings. *Molecular Aspects of Medicine* **61**:18-30
- Bravo L, Saura-Calixto F. 1998.** Characterization of the dietary fiber and the in vitro indigestible fraction of grape pomace. *American Journal of Enology and Viticulture* **49**:135-141

591 **Camps-Bossacoma M, Garcia-Aloy M, Saldaña-Ruiz S, Cambras T, González-**
 592 **Domínguez R, Franch A, Pérez-Cano FJ, Andres-Lacueva C, Castell M. 2019.** Role
 593 of theobromine in cocoa's metabolic properties in healthy rats. *Journal of Agriculture*
 594 *and Food Chemistry* **6**:3605-3614

595 **Carlin-Sinclair A, Marc I. 2009.** The determination of methylxanthines in chocolate and
 596 cocoa by different separation techniques: HPLC instrumental TLC and MECC. *Journal*
 597 *of Laboratory Chemical Education* **86**:1307-1310

598 **Cocoa - Food and Agriculture Organization of the United Nations**
 599 <http://www.fao.org/3/y5143e/y5143e0x.htm>

600 **Cremonini E, Bettaieb A, Haj F G, Fraga C G, Oteiza P I. 2016.** (-)-Epicatechin
 601 improves insulin sensitivity in high fat diet-fed mice. *Archives of Biochemistry and*
 602 *Biophysics* **599**:13-21

603 **Dower JI, Geleijnse JM, Gijsbers L, ZockPL, Kromhout D, Hollman PCH. 2015.**
 604 Effects of the pure flavonoids epicatechin and quercetin on vascular function and
 605 cardiometabolic health: a randomized double-blind placebo-controlled crossover trial.
 606 *American Journal of Clinical Nutrition* **101**:914-921

607 **Ellam S, Williamson G. 2013.** Cocoa and human health *Annual Review of Nutrition*
 608 **33**:105-128

609 **Fogliano V, Corollaro M L, Vitaglione P, Napolitano A, Ferracane R, Travaglia F,**
 610 **Arlorio M, Costabile A, Klinder A, Gibson G. 2011.** In vitro bioaccessibility and gut
 611 biotransformation of polyphenols present in the water-insoluble cocoa fraction.
 612 *Molecular Nutrition and Food Research* **55**:S44-55

613 **Fuggetta MP, Zonfrillo M Villivà, C Bonmassar, E, Ravagnan G. 2019.** Inflammatory
 614 Microenvironment and Adipogenic Differentiation in Obesity: The Inhibitory Effect of
 615 theobromine in a model of human obesity in vitro. *Mediators of Inflammation* **1515621**

616 **Giacco R, Parillo M, Rivellese A A, Lasorella G, Giaco A, D'Episcopo L, Ricardi J.**
 617 **2000.** Long-term dietary treatment with increased amounts of fiber-rich low-glycemic

- 618 index natural foods improves blood glucose control and reduces the number of
- 619 hypoglycaemic events in type 1 diabetic patients. *Diabetes Care* **23**:1461-1466
- 620 **Gómez-Juaristi M, González-Torres L, Bravo L, Vaquero MP, Bastida S, Sánchez-**
- 621 **Muniz FJ. 2011.** Beneficial effects of chocolate on cardiovascular health. *Nutrición*
- 622 *Hospitalaria* **26**:289-292
- 623 **Gómez-Juaristi M, Sarriá B, Martínez-López S, Bravo-Clemente L, Mateos R. 2019.**
- 624 Flavanol bioavailability in two cocoa products with different phenolic content. A
- 625 comparative study in humans. *Nutrients* **11**:1441
- 626 **Goya L, Martín M, Sarriá B, Ramos S, Mateos R, Bravo L. 2016.** Effect of cocoa and
- 627 its flavonoids on biomarkers of inflammation: studies of cell culture animals and
- 628 humans. *Nutrients* **8**:212
- 629 **Grassi D, Necozione S, Lippi C, Croce G Valeri L Pasqualetti P Desideri G**
- 630 **Blumberg JB Ferri C. 2005.** Cocoa reduces blood pressure and insulin resistance and
- 631 improves endothelium-dependent vasodilation in hypertensives. *Hypertension* **46**:398-
- 632 405
- 633 **Haskó G, Cronstein B. 2011.** Methylxanthines and inflammatory cells. *Handbook of*
- 634 *Experimental Pharmacology* **(200)**:457-468
- 635 **Heiss C, Keen CL, Kelm M. 2010.** Flavanols and cardiovascular disease prevention.
- 636 *European Heart Journal* **31**:2583e2592
- 637 **Hooper L, Kroon PA, Rimm EB, Cohn JS, Harvey I, Le Cornu KA, Ryder JJ, Hall**
- 638 **WL, Cassidy A. 2012.** Effects of chocolate, cocoa, and flavan-3-ols on cardiovascular
- 639 health: a systematic review and meta-analysis of randomized trials. *American Journal of*
- 640 *Clinical Nutrition* **95**:740-751
- 641 **Horrigan LA, Kelly JP, Connor TJ. 2004.** Caffeine suppresses TNF- α production via
- 642 activation of the cyclic AMP/protein kinase A pathway. *International Immunopharmacology*
- 643 **4**:1409-1417
- 644 **Huang D, Ou B, Hampsch-Woodill M, FlanaganJA, Prior RL. 2002.** High-throughput
- 645 assay of oxygen radical absorbance capacity (ORAC) using a multichannel liquid

646 handling system coupled with a microplate fluorescence reader in 96-well format.
 647 *Journal of Agricultural and Food Chemistry* **50**:4437-4444

648 **Jenkins DJA, Kendall CWC, Vuksan V, Vidgen E, Wong E, Augustin LS, Fulgoni V.**
 649 **2000.** Effect of cocoa bran on low-density lipoprotein oxidation and fecal bulking.
 650 *Archives of Internal Medicine* **160**:2374-2379

651 **Josic J, Olsson AT, Wickeberg J, Lindstedt S, Hlebowicz J. 2010** Does green tea
 652 affect postprandial glucose insulin and satiety in healthy subjects: A randomized
 653 controlled trial. *Nutrition Journal* **9**:63

654 **Khan N, Monagas M, Andres-Lacueva C, Casas R, Urpi-Sarda M, Lamuela-**
 655 **Raventos RM, Estruch R. 2012.** Regular consumption of cocoa powder with milk
 656 increases HDL cholesterol and reduces oxidized LDL levels in subjects at high-risk of
 657 cardiovascular disease. *Nutrition Metabolism and Cardiovascular Disease* **22**:1046-
 658 1053

659 **Lecumberri E, Mateos R, Izquierdo-Pulido M, Ruperez P, Goya L, Bravo L. 2007.**
 660 Dietary fibre composition, antioxidant capacity and physico-chemical properties of a
 661 fibre-rich product from cocoa (*Theobroma cacao* L.). *Food Chemistry* **104**:948-954
 662

663 **Lofffield E, Shiels MS, Graubard BI, Katki HA, Chaturvedi AK, Trabert B, Pinto LA,**
 664 **Kemp TJ, Shebl FM, Mayne ST, Wentzensen N, Purdue MP, Hildesheim A, Sinha**
 665 **R, Freedman ND. 2015.** Associations of coffee drinking with systemic immune and
 666 inflammatory markers. *Cancer Epidemiology Biomarkers Prevention* **24**:1052-1060

667 **Martínez-López S, Sarriá B, Gómez-Juaristi M, Goya L, Mateos R, Bravo-Clemente**
 668 **L. 2014a.** Theobromine, caffeine, and theophylline metabolites in human plasma and
 669 urine after consumption of soluble cocoa products with different methylxanthine
 670 contents. *Food Research International* **63**:446-455

671 **Martínez-López S, Sarriá B, Sierra-Cinos J L, Goya L, Mateos R, Bravo L. 2014b.**
 672 Realistic intake of a flavanol-rich soluble cocoa product increases HDL-cholesterol

673 without inducing anthropometric changes in healthy and moderately
 674 hypercholesterolemic subjects. *Food and Function* **5**:364-374

675 **Massot-Cladera M, Franch À, Castell M, Pérez-Cano FJ. 2017.** Cocoa polyphenols and
 676 fiber modify colonic gene expression in rats. *European Journal of Nutrition* **56**:1871-1885

677 **Mellor DD, Sathyapalan T, Kilpatrick E S, Beckett S, Atkin S L. 2010.** High-cocoa
 678 polyphenol-rich chocolate improves HDL cholesterol in Type 2 diabetes patients.
 679 *Diabetic Medicine* **27**:1318-1321

680 **Miller KB, Hurst WJ, Flannigan N, Ou B, Lee CY, Smith N, Stuart DA. 2009.** Survey
 681 of commercially available chocolate- and cocoa-containing products in the United
 682 States. 2. Comparison of flavan-3-ol content with nonfat cocoa solids total polyphenols
 683 and percent cacao. *Journal of Agricultural and Food Chemistry* **57**:9169-9180

684 **Monagas M, Urpi-Sarda M, Sánchez-Patán F, Llorach R, Garrido I, Gómez-**
 685 **Cordovés Andrés-Lacueva C, Bartolomé B. 2010.** Insights into the metabolism and
 686 microbial biotransformation of dietary flavan-3-ols and the bioactivity of their
 687 metabolites. *Food and Function* **1**:233-253

688 **Monteiro J, Alves M, Oliveira P, Silva B. 2016.** Structure-bioactivity relationships of
 689 methylxanthines: Trying to make sense of all the promises and the drawbacks.
 690 *Molecules* **21**:974

691 **Monteiro J, Alves MG, Oliveira PF, Silva BM. 2019.** Pharmacological potential of
 692 methylxanthines: Retrospective analysis and future expectations. *Critical Reviews in*
 693 *Food Science and Nutrition* **59**:2597-2625

694 **Neufingerl N, Zebregs YE, Schuring EA, Trautwein EA. 2013.** Effect of cocoa and
 695 theobromine consumption on serum HDL-cholesterol concentrations: a randomized
 696 controlled trial. *American Journal of Clinical Nutrition* **97**:1201-1209

697 **Ottaviani JI, Kwik-Urbe C, Keen CL, Schroeter H. 2012b.** Intake of dietary
 698 procyanidins does not contribute to the pool of circulating flavanols in humans.
 699 *American Journal of Clinical Nutrition* **95**:851-858

700 **Ottaviani JI, Momma TY, Kuhnle GK, Keen CL, Schroeter H. 2012a.** Structurally
701 related (–)-epicatechin metabolites in humans: assessment using de novo chemically
702 synthesized authentic metabolites. *Free Radical Biology and Medicine* **52**:1403-1412

703 **Prakash M, Basavaraj BV, Murthy KC. 2019.** Biological functions of epicatechin: Plant
704 cell to human cell health. *Journal of Functional Foods* **52**:14-24

705 **Pulido R, Bravo L, Saura-Calixto F. 2000.** Antioxidant activity of dietary polyphenols
706 as determined by a modified ferric reducing/antioxidant power assay. *Journal of*
707 *Agricultural and Food Chemistry* **48**:3396-3402

708 **Re N, Pellegrini A, Proteggente A, Pannala A, Yang M, Rice-Evans C. 1999.**
709 Antioxidant activity applying an improved ABTS radical cation decolorization assay.
710 *Free Radical Biology and Medicine* **26**:1231-1237

711 **Rodríguez-Mateos A, Weber T, Skene SS, Ottaviani JI, Crozier A, Kelm M,**
712 **Schroeter H , Heiss C. 2018.** Assessing the respective contribution of dietary flavanol
713 monomers and procyanidins in mediating cardiovascular effects in humans: randomized
714 controlled double-masked intervention trial. *American Journal of Clinical Nutrition*
715 **108**:1229-1237

716 **Rodriguez-Mateos AM, Vauzour D, Kreuger CG, Shanmuganayagam D, Reed D,**
717 **Canali L, Mena P, Del Rio D, Crozier A. 2014.** Flavonoids and related compounds
718 bioavailability bioactivity and impact on human health: an update. *Archives of*
719 *Toxicology* **88**:1803-1853

720 **Sarriá B, Martínez-López S, Fernández-Espinoza A, Gómez-Juaristi M, Goya L,**
721 **Mateos R, Bravo L. 2012.** Effects of regularly consuming dietary fibre rich soluble
722 cocoa products on bowel habits in healthy subjects: a free-living two-stage randomized
723 crossover single-blind intervention. *Nutrition & Metabolism* **18**:9-33

724 **Sarriá B, Martínez-López S, Sierra-Cinos J L, García-Diz L, Mateos R, Bravo L.**
725 **2014.** Regular consumption of a cocoa product improves the cardiometabolic profile in
726 healthy and moderately hypercholesterolaemic adults. *British Journal of Nutrition*
727 **111**:122-134

- 728 **Sarriá B, Martínez-López S, Sierra-Cinos JL, Garcia-Diz L, Goya L, Mateos R,**
729 **Bravo L. 2015.** Effects of bioactive constituents in functional cocoa products on
730 cardiovascular health in humans. *Food Chemistry* **174**:214-218
- 731 **Sarriá B, Mateos R, Sierra-Cinos JL, Goya L, García-Diz L, Bravo L. 2012.**
732 Hypotensive hypoglycaemic and antioxidant effects of consuming a cocoa product in
733 moderately hypercholesterolemic humans. *Food and Function* **3**:867-874
- 734 **Saura-Calixto F, García-Alonso A, Goñi I, Bravo L. 2000.** In vitro determination of the
735 indigestible fraction in foods: an alternative to dietary fiber analysis. *Journal of*
736 *Agricultural and Food Chemistry* **48**:3342-3347
- 737 **Saura-Calixto F, Goñi I. 2006.** Antioxidant capacity of the Spanish Mediterranean diet.
738 *Food Chemistry* **94**:442-447
- 739 **Tokede OA, Gaziano JM, Djoussé L. 2011.** Effects of cocoa products/dark chocolate
740 on serum lipids: a meta-analysis. *European Journal of Clinical Nutrition* **65**:879-886
- 741 **Urpi-Sarda M, Monagas M, Khan N, Lamuela-Raventos R M, Santos-Buelga C,**
742 **Sacanella E, Castell M, Permanyer J, Andres-Lacueva C. 2009a.** Epicatechin
743 procyanidins and phenolic microbial metabolites after cocoa intake in humans and rats.
744 *Analytical and Bioanalytical Chemistry* **394**:1545-1556
- 745 **Urpi-Sarda M, Monagas M, Khan N, Llorach R, Lamuela-Raventós R M, Jáuregui O**
746 **Andrés-Lacueva C. 2009b.** Targeted metabolic profiling of phenolics in urine and
747 plasma after regular consumption of cocoa by liquid chromatography–tandem mass
748 spectrometry. *Journal of Chromatography A* **1216**:7258-7267
- 749 **Wan Y, Vinson JA, Etherton TD, Proch J, Lazarus SA, Kris-Etherton PM. 2001.**
750 Effects of cocoa powder and dark chocolate on LDL oxidative susceptibility and
751 prostaglandin concentrations in humans. *American Journal of Clinical Nutrition* **74**:596-
752 602
- 753 **Williams J F, Lowitt S, Polson J B, Szentivanyi A. 1978.** Pharmacological and
754 biochemical activities of some monomethylxanthine and methyluric acid derivatives of
755 theophylline and caffeine. *Biochemical Pharmacology* **7**:1545-1550

756 **World Cocoa Conference, 2018.** <https://www.icco.org/>

757 **Xing Y W, Lei G T, Wu Q H, Jiang Y, Huang MX. 2019.** Procyanidin B2 protects
 758 against diet-induced obesity and non-alcoholic fatty liver disease via the modulation of
 759 the gut microbiota in rabbits. *World Journal of Gastroenterology* **25**:955

760

761

762

763

Figure 1

Bioactive compounds present in the cocoa products and the most abundant metabolites formed after flavanol and methylxanthine intake




Table 1(on next page)

Flavanols antioxidant activity dietary fibre and methylxanthines composition of the soluble cocoa products used in bioavailability and chronic interventions in humans: cocoa rich in cocoa (PC), cocoa rich in dietary fibre (DFC), conventional cocoa (CC), c

Table 1. Flavanols antioxidant activity dietary fibre and methylxanthines composition of the soluble cocoa products used in bioavailability and chronic interventions in humans: cocoa rich in cocoa (PC), cocoa rich in dietary fibre (DFC), conventional cocoa (CC), cocoa rich in methylxanthines and cocoa (MPC) and raw un-processed cocoa (RC).

	Product PC	Product DFC	Product CC	Product MPC	Product RC
Polyphenol composition					
Total Polyphenols (a) (μg equiv gallic acid /g product) n=6	34.04 \pm 2.28	15.75 \pm 0.67	21.70 \pm 1.40	25.63 \pm 1.00	42.11 \pm 2.50
EC (mg/g dry matter) (b)	1.26 \pm 0.18	0.33 \pm 0.09	0.57 \pm 0.07	1.15 \pm 0.06	2.40 \pm 0.50
CA (mg/g dry matter) (b)	0.47 \pm 0.03	0.26 \pm 0.12	0.32 \pm 0.03	0.53 \pm 0.04	0.83 \pm 0.17
PB1 (mg/g dry matter) (b)	0.20 \pm 0.04	n.d.	0.04 \pm 0.02	0.23 \pm 0.02	0.41 \pm 0.11
PB2 (mg/g dry matter) (b)	1.09 \pm 0.10	0.57 \pm 0.11	0.39 \pm 0.05	0.82 \pm 0.06	2.04 \pm 0.68
Total flavanols (b)	3.02 \pm 0.35	1.16 \pm 0.32	1.32 \pm 0.17	2.73 \pm 0.18	5.68 \pm 1.46
Methylxanthines Composition					
Theobromine (mg/g dry matter) (a)	6.43 \pm 0.84	5.11 \pm 0.14	5.63 \pm 0.06	7.08 \pm 0.22	6.33 \pm 0.17
Theophylline (mg/g dry matter) (a)	0.01 \pm 0.01	n.d.	n.d.	0.13 \pm 0.02	0.06 \pm 0.01
Caffeine (mg/g dry matter) (a)	0.88 \pm 0.08	0.51 \pm 0.05	0.66 \pm 0.06	3.03 \pm 0.28	1.46 \pm 0.21
Total methylxanthines	7.32 \pm 0.93	5.62 \pm 0.19	6.29 \pm 0.12	10.24 \pm 0.52	7.85 \pm 0.39
Dietary Fibre Composition					
Soluble dietary fibre (%)= NS + UA	3.13 \pm 0.59	1.68 \pm 0.13	2.69 \pm 0.70	3.00 \pm 0.86	4.27 \pm 0.27
Neutral sugars (%)	2.46 \pm 0.43	0.69 \pm 0.04	2.21 \pm 0.51	1.80 \pm 0.67	2.70 \pm 0.14
Uronic acid (%)	0.67 \pm 0.16	0.99 \pm 0.09	0.48 \pm 0.19	1.20 \pm 0.19	1.57 \pm 0.13
Insoluble dietary fibre (%)=NS + UA	11.96 \pm 1.05	20.32 \pm 1.67	14.31 \pm 0.51	26.63 \pm 2.54	11.58 \pm 1.30
Neutral sugars (%)	10.49 \pm 0.96	19.06 \pm 1.60	13.32 \pm 0.29	25.43 \pm 2.39	9.12 \pm 1.08
Uronic acid (%)	1.47 \pm 0.09	1.26 \pm 0.07	0.99 \pm 0.22	1.20 \pm 0.15	2.46 \pm 0.22
Total dietary fibre (%)	15.09 \pm 1.64	22.00 \pm 1.80	17.00 \pm 1.21	29.63 \pm 3.40	15.85 \pm 1.57

Analysis carried out using (a) Folin-Ciocalteu method (b) HPLC method (c) Bravo & Saura-Calixto, 1998. n.d.: not detected. EC: epicatechin; CA: catechin; PB1: procyanidin B1; PB2: procyanidin B2

10

11

12

Table 2(on next page)

Antioxidant activity of cocoa products rich in cocoa rich in cocoa (PC), cocoa rich in dietary fibre (DFC), conventional cocoa (CC), cocoa rich in methylxanthines and cocoa (MPC) and raw un-processed cocoa (RC).

Table 2. Antioxidant activity of cocoa products rich in cocoa (PC), cocoa rich in dietary fibre (DFC), conventional cocoa (CC), cocoa rich in methylxanthines and cocoa (MPC) and raw un-processed cocoa (RC).

	Product PC	Product DFC	Product CC	Product MPC	Product RC
Antioxidant activity					
FRAP (μmol TE/μg product) n=3	175.07 ± 1.52	75.88 ± 2.61	109.41 ± 2.33	120.12 ± 3.26	219.33 ± 6.90
ABTS (μmol TE/g product) n=3	133.09 ± 3.70	56.64 ± 3.26	66.64 ± 6.88	89.53 ± 2.99	173.42 ± 7.53
ORAC (μmol TE/g product) n=8	459.71 ± 9.40	234.51 ± 4.89	248.56 ± 4.32	257.39 ± 5.79	483.93 ± 19.88

Table 3(on next page)

Summary of the characteristics and outcomes of the cocoa bioavailability and chronic studies

Table 3. Summary of the characteristics and outcomes of the cocoa bioavailability and chronic studies

Reference	Characteristics of the study	Number of volunteers and criteria	Cocoa product and dose	Main cocoa metabolites*	Health effects
Gómez-Juaristi et al. 2019	<u>Polyphenol bioavailability study</u> Randomized and crossover, 8 hours long carried out on two separated days	13 Healthy subjects	Day 1: 15 g of conventional cocoa (CC) - 19.80 mg of flavanols/day Day 2: 25g of cocoa rich in methylxanthines and phenols (MPC) - 68.25 mg of flavanols/day	Plasma: Epicatechin and its phase II derivatives: 3'-Methyl-epicatechin-5-sulfate, epicatechin-3'-sulfate, 4'-methyl-epicatechin-5-sulfate, epicatechin-3'-glucuronide, among others. 5-(3',4'-Dihydroxyphenyl)-γ-valerolactone (DHPVL) and its phase II derivatives (HPVL-3'-glucuronide, HPVL-3'-sulfate, PVL-methyl-glucuronide). Urine: Phase II derivatives of epicatechin, approximately 33% of total. The majority were phase II derivatives of 5-(4'-hydroxyphenyl)-γ-valerolactone (HPVL) and 4-hydroxy-5-(4'-hydroxyphenyl)valeric acid (HHPVA): HPVL-3'-sulfate followed by HHPVA-sulfate, PVL-sulfate and HPVL-3'-glucuronide, among others Dose-dependent response	No changes in blood pressure along the 8 hours. No changes in antioxidant capacity in blood along the 8 hours, except 1h (ABTS) and 2 and 8 h (FRAP)
Martínez-López et al. 2014a	<u>Methylxanthine bioavailability study</u> Randomized and crossover, 8 hours long carried out on two separated days	13 Healthy subjects	Day 1: 15 g of conventional cocoa (CC) - 84.45 mg of theobromine (TB) and 9.9 mg of caffeine (CF)/day Day 2: 25g of cocoa rich in methylxanthines and phenols (MPC) - 177 mg TB, 75.75 mg CF and 3.25 mg theophylline (TP)/day	Plasma: TB, CF, TP and paraxanthine (PX) together with two monomethylxanthines (3-methylxanthine (MX) and 7-MX) Urine: TB, CF, TP and PX, along with monomethylxanthines (1-MX, 3-MX and 7-MX) and different mono-, di- and tri-methyluric (MU) acids (1-MU, 1,3-MU, 1,7-MU, 3,7-MU and 1,3,7-MU). 7-MX as the most abundant metabolite followed by TB and 3-MX Dose-dependent response	No changes in blood pressure along the 8 hours. No changes in antioxidant capacity in blood along the 8 hours, except 1h (ABTS) and 2 and 8 h (FRAP)

Sarriá et al. 2014.	Randomized, controlled and crossover chronic study. Cocoa and control interventions were 4 weeks long	44 subjects healthy (n 24) and moderately hypercholesterolemic (n 20)	<u>30 g/day (2 doses of 15 g of cocoa rich in dietary fiber (DFC))</u> 34.8 mg of flavanols/day 153.3 mg of TB and 15.3 mg of CF/day 6.6 g total dietary fiber (DF) 0.5 g soluble DF/day 6.1 g insoluble DF/day	Phenol metabolites according to Gómez-Juaristi et al. 2019: Epicatechin and phase II derivatives of epicatechin Phase II derivatives of hydroxyphenyl-γ-valerolactones and valeric acid Methylxanthine metabolites according to Martínez-López et al. 2014a: TB, CF, TP and paraxanthine (PX) together with monomethylxanthines (1-MX, 3-MX and 7-MX) and mono-, di- and tri-MU acids (1-MU, 1,3-MU, 1,7-MU, 3,7-MU and 1,3,7-MU) Dietary fiber according to Sarriá et al. 2014.	↑HDL-cholesterol ↓Blood glucose ↑IL-1β ↓IL-10
Martínez-López et al. 2014b.	Randomized, controlled and crossover-chronic study. Cocoa and control interventions were 4 weeks long	44 subjects healthy (n 24) and moderately hypercholesterolemic (n 20)	<u>15 g/day (2 doses of 7.5g) of cocoa rich in cocoa (PC)</u> 45.3 mg of flavanols/day 96.45 mg of TB and 13.2 mg of CF/day 2.3 g total dietary fiber (DF) 0.5 g soluble DF/day 1.8 g insoluble DF/day	Phenol metabolites according to Gómez-Juaristi et al. 2019: Epicatechin and phase II derivatives of epicatechin Phase II derivatives of hydroxyphenyl-γ-valerolactones and valeric acid Methylxanthine metabolites according to Martínez-López et al. 2014a: TB, CF, TP and PX together with monomethylxanthines and different mono-, di- and tri-MU acids. Dietary fiber	↑HDL-cholesterol

*5-(3',4'-dihydroxyphenyl)- γ -valerolactone (DHPVL); 5-(4'-hydroxyphenyl)- γ -valerolactone (HPVL); 5-phenyl- γ -valerolactone (PVL); 4-hydroxy-5-(3',4'-dihydroxyphenyl)valeric acid (HDHPVA) and 4-hydroxy-5-(hydroxyphenyl)valeric acid (HHPVA).

1

2

Table 4(on next page)

Antioxidant capacity in serum samples along the bioavailability study using ABTS and FRAP methods

1 Table 4. Antioxidant capacity in serum samples along the bioavailability study using ABTS and FRAP methods

ABTS (μM TE)								
Hours	0	0.5	1	2	3	4	6	8
CC	3830.59 ± 114.35	3682.88 ± 112.79	3605.06 ± 81.59 ^b	3658.54 ± 94.03	3696.70 ± 91.70	3665.45 ± 97.03	3687.98 ± 82.26	3797.14 ± 108.21
MPC	3944.78 ± 62.59	3913.31 ± 65.85	3953.47 ± 49.67 ^a	3860.47 ± 61.00	3831.18 ± 58.11	3873.56 ± 63.21	3828.77 ± 45.97	3815.37 ± 73.11
	N.S.	N.S.	p=0.004	N.S.	N.S.	N.S.	N.S.	N.S.
FRAP (μM TE)								
Hours	0	0.5	1	2	3	4	6	8
CC	592.82 ± 28.14	579.43 ± 32.27	551.75 ± 24.63	540.19 ± 26.67	550.39 ± 24.61	550.49 ± 27.44	556.48 ± 31.65	530.39 ± 24.16
MPC	596.70 ± 23.77	597.63 ± 22.09	580.28 ± 22.50	583.84 ± 23.20	578.61 ± 23.43	566.25 ± 26.51	583.85 ± 26.35	560.42 ± 28.83
	N.S.	N.S.	N.S.	p=0.039	N.S.	N.S.	N.S.	p=0.044

2 Conventional cocoa (CC), cocoa rich in methylxanthines and cocoa (MPC), Trolox Equivalents (TE)

3