

Role of reactive oxygen species and isoflavonoids in soybean resistance to the attack of the southern green stink bug

Ivana Sabljic^{1,2}, Jesica A Barneto¹, Karina B Balestrasse¹, Jorge A Zavala¹, Eduardo A Pagano^{Corresp. 1}

¹ Instituto de Investigaciones en Biociencias Agrícolas y Ambientales-INBA, Facultad de Agronomía, Universidad de Buenos Aires, Consejo Nacional de Investigaciones Científicas y Técnicas, Ciudad Autónoma de Buenos Aires, Argentina

² GDM, Chacabuco, Buenos Aires, Argentina

Corresponding Author: Eduardo A Pagano
Email address: epagano@agro.uba.ar

Southern green stink bugs (*Nezara viridula* L.) are one of the major pests in many soybean producing areas. They cause a decrease in yield and affect seed quality by reducing viability and vigor. Alterations have been reported in the oxidative response and in the secondary metabolites in different plant species due to insect damage. However, there is little information available on soybean-stink bug interactions. In this study we compare the response of undamaged and damaged seeds by *Nezara viridula* in two soybean cultivars, IAC-100 (resistant) and Davis (susceptible), grown under greenhouse conditions. Pod hardness, H₂O₂ generation, enzyme activities in guaiacol peroxidase (GPOX), catalase (CAT) and superoxide dismutase (SOD) as well as lipoxygenase expression and isoflavonoid production were quantified. Our results showed a greater resistance of IAC-100 to pod penetration, a decrease in peroxide content after stink bug attack, and higher GPOX, CAT and SOD activities in seeds due to the genotype and to the genotype-interaction with the herbivory treatment. Induction of *LOX* expression in both cultivars and higher production of isoflavonoids in IAC-100 were also detected. It was then concluded that the herbivory stink bug induces pathways related to oxidative stress and to the secondary metabolites in developing seeds of soybean and that differences between cultivars hold promise for a plant breeding program.

Role of reactive oxygen species and isoflavonoids in soybean resistance to the attack of the southern green stink bug

Ivana Sabljic^{1,2}, Jesica Alejandra Barneto¹, Karina Beatriz Balestrasse¹, Jorge Alberto Zavala¹, Eduardo Antonio Pagano¹

¹Universidad de Buenos Aires, Consejo Nacional de Investigaciones Científicas y Técnicas, Instituto de Investigaciones en Biociencias Agrícolas y Ambientales-INBA, Facultad de Agronomía, Buenos Aires, Argentina.

²GDM, Chacabuco, Provincia de Buenos Aires. Argentina.

Corresponding Author:

Eduardo Antonio Pagano¹

Av. San Martin 4453, Buenos Aires, C1417DSE, Argentina

Email address: epagano@agro.uba.ar

Abstract

Southern green stink bugs (*Nezara viridula* L.) are one of the major pests in many soybean producing areas. They cause a decrease in yield and affect seed quality by reducing viability and vigor. Alterations have been reported in the oxidative response and in the secondary metabolites in different plant species due to insect damage. However, there is little information available on soybean-stink bug interactions. In this study we compare the response of undamaged and damaged seeds by *Nezara viridula* in two soybean cultivars, IAC-100 (resistant) and Davis (susceptible), grown under greenhouse conditions. Pod hardness, H₂O₂ generation, enzyme activities in guaiacol peroxidase (GPOX), catalase (CAT) and superoxide dismutase (SOD) as well as lipoxygenase expression and isoflavonoid production were quantified. Our results showed a greater resistance of IAC-100 to pod penetration, a decrease in peroxide content after stink bug attack, and higher GPOX, CAT and SOD activities in seeds due to the genotype and to the genotype-interaction with the herbivory treatment. Induction of *LOX* expression in both cultivars and higher production of isoflavonoids in IAC-100 were also detected. It was then concluded that the herbivory stink bug induces pathways related to oxidative stress and to the secondary metabolites in developing seeds of soybean and that differences between cultivars hold promise for a plant breeding program.

Keywords: Plant–insect interaction – *Glycine max* - *Nezara viridula* – Oxidative stress – Herbivory – Lipoxygenase

Introduction

The stink bug complex is one of the most detrimental pests that affects soybean (*Glycine max* L.) yields in many countries. In South America, the main species that impair soybean yields are *Nezara viridula*, *Piezodorus guildinii* and *Euschistus heros*. They adversely affect seed vigor and may even kill the embryo and stop germination. Brazil and Argentina are two of the most important producers of this legume and damage caused by the mentioned species can result in significant economic losses (Schaefer & Panizzi, 2000). Soybean crops destined for seed production are less tolerant to stink bug damage than soybean crops used for animal feed (Jensen & Newsom, 1972). While stink bugs can be controlled by spraying pesticides, there are environmental and economic issues concerning the frequent use of these products. Biological control can be mentioned as an alternative method to fight against this complex of insects. For example, the well-studied egg-parasitoid *Trissolcus basalis* (Wollaston) has been used successfully by farmers in Brazil (Corrêa-Ferreira & Moscardi, 1996). However, soybean host plant resistance is an interesting strategy for managing plant breeding programs (Boerma & Walker, 2005), and it is necessary to first understand the mechanisms related to stink bug resistance in soybean breeding lines.

The soybean cultivar IAC-100 has been defined as resistant against stink bug attack (Campos *et al.*, 2010; McPherson, Buss & Roberts, 2007; Piubelli *et al.*, 2003a; de Souza *et al.*, 2014; Michereff *et al.*, 2011) and is used in breeding programs as a source of resistance (McPherson *et al.*, 2007). This variety was developed by the Agronomic Institute of Campinas in Sao Paulo (Brazil) by crossing breeding lines IAC 78-2318 and IAC-12 (Priolli *et al.*, 2002). Studies on this genotype showed that stink bug attack caused less seed damage and less seed weight loss than other genotypes (Campos *et al.* 2010). Previous studies found lower adult fresh weight of *Nezara viridula* when fed on 'IAC-100' and less lipids in females than in those fed on the other genotypes (Piubelli *et al.*, 2003b). However, the mechanisms through which soybean reacts to stink bug attack are still unknown.

Plants respond to herbivore injury through several direct and indirect morphological, biochemical and molecular mechanisms that help them avoid herbivore attack or affect its performance (Fürstenberg-Hägg, Zagrobelny & Bak, 2013). Direct defense mechanisms affect insect performance and feeding behavior, while indirect defense mechanisms can attract the natural enemies of the herbivores and thus reduce plant loss (War *et al.*, 2012). Insect herbivory induces early responses such as the oxidative burst (production of reactive oxygen species - ROS), the expression of defense-related genes, and late responses such as callose deposition and accumulation of proteinase inhibitors (Savatin *et al.*, 2014). Infestation of *Vicia faba* by *N. viridula* stimulates significantly the production of H₂O₂ (Ederli *et al.*, 2017). Moreover, in poplar the concentrations of H₂O₂ and malondialdehyde and the activities of ROS-scavenging enzymes, such as superoxide dismutase (SOD), catalase (CAT) and peroxidase (POD), were enhanced by herbivore wounding, suggesting that they are associated with insect resistance. The above-mentioned enzymes are related to plant signaling, synthesis of defense compounds, and to oxidative stress tolerance (Bi & Felton, 1995; Zhang, Hua & Zhang, 2008; Bednarski *et al.*, 2013; Zebelo & Maffei, 2014).

After insect attack the activation of defense genes that stimulate the production of the antiherbivory compounds is induced due to internal signals such as calcium ion flow, a phosphorylation cascade and responses mediated by hormones: jasmonic acid (JA) and ethylene (ET) (Waternack, 2007; Howe & Jander, 2008; Browse, 2009). JA and its methyl ester, methyl jasmonate (MJ), participate in the activation of plant defense mechanisms as signaling compounds in processes related to production of various secondary metabolites (i.e. terpenoids,

phenylpropanoids, and alkaloids) (Wang *et al.*, 2015; Misra *et al.*, 2014). In the case of soybean, isoflavones are the main chemical defense compounds against insects because they affect the performance and survival of the herbivores (Piubelli *et al.*, 2003a). It is well known that MJ and JA induce the accumulation of proteinase inhibitors as direct defense mechanisms against herbivorous insects (Farmer & Ryan 1990; Farmer, Johnson & Ryan, 1992). Studies on soybean analyzed the effects of jasmonic acid induction and showed that the soybean looper (*Chrysodeixis includens* Walker) selects plant leaves that have not been induced by JA (Accamando *et al.* (2012). In addition, soybean crops grown under conditions of high concentrations of ambient CO₂ were unable to express genes related to the JA synthesis, which made them highly susceptible to insect attack (Zavala *et al.*, 2008). The synthesis of jasmonates and many other oxilipins is initiated by lipoxygenases (LOXs), which catalyze dioxygenation of polyunsaturated fatty acids (reviewed by Blee, 2002, Feussner & Wasternack, 2002, Howe & Schilmiller, 2002, Wasternack, 2007). LOX activity was higher in soybean plants that were less vulnerable to the attack of *A. gemmatilis* when compared to controls (Fortunato *et al.* 2007). Since three lipoxygenases, Lox1, Lox2 and Lox3 were detected in soybean seeds (Axelrod, Cheesbrough & Laakso, 1981), it is possible that seed lipoxygenases regulate defenses against stink bug attack. As a first step to develop a resistant cultivar, it is necessary to identify the sources of resistance for any breeding program (de Moraes & Pinheiro, 2012). Therefore, it is important to determine the different characteristics of the diverse soybean cultivars as regards plant defense against insect attack.

The aim of this study is to identify and quantify physical and biochemical soybean defense mechanisms against the southern green stink bug, *Nezara viridula* attack. Our hypothesis is that herbivory by stink bugs lead to biochemical changes and that there are differences between cultivars at both biochemical and physical level. In this study we have compared soybean seed response to stink bug damage of two cultivars, IAC-100 (resistant) and Davis (susceptible; Orr, Boethel & Jones, 1985; Kester, Smith & Gilman 1984). Our results show that there are differences between genotypes as regards the oxidative stress response and isoflavonoid production in response to stink bugs attack. A better understanding of the resistance pathways can help find ways to improve varieties.

Materials & Methods

Insect and plant material

In order to determine soybean resistance to stink bug damage (*Nezara viridula* L.) in developing seeds, two soybean (*Glycine max* L.) cultivars, IAC-100 and Davis, were grown under greenhouse conditions and stink bug adults were placed and allowed to feed on pods. Plants were grown with 16:8 L:D light regime at 26±3°C, in pots with 3:1:1:1 soil:peat:sand:perlite mixture. The colony of stink bugs consisted of field-collected individuals from the city of Chacabuco (34°38'00"S 60°28'00"O) in the province of Buenos Aires (Argentina). The insects were reared at our laboratory (School of Agronomy, University of Buenos Aires) on a diet based on hydrated soybean seeds (cv. Williams) and peeled sunflower seeds, and were then provided water with ascorbic acid (0.5 w/v %) (Giacometti *et al.*, 2016).

Experimental design

Four experiments were carried out under the same greenhouse conditions in a completely randomized design. In the first experiment, seventeen replications of each cultivar were used, pod hardness was determined by collecting pods at the R6 soybean development stage (Fehr *et*

al., 1971) and by using a texturometer. In order to determine oxidative stress response of developing seeds to stink bug damage in the second experiment, a single adult stink bug was bagged in on one soybean pod of plants (n=5) at the R6 development stage (Fehr *et al.*, 1971). Pods of control plants were covered with tulle bags and collected 72 h after stink bugs started feeding. Soon after, they were fast frozen in liquid nitrogen and stored at -80°C until analysis. In addition, attacked and unattacked pods (one per plant) were collected for H_2O_2 detection in developing seeds. Five replications of each cultivar were used for stink bug infestation with an equal number of control plants. In the third experiment undamaged and damaged seeds by stink bugs were collected to determine isoflavonoid accumulation after 72-hr herbivory. Finally, a fourth experiment was performed to determine the LOX expression level in damaged and undamaged seeds, and pods were collected 24-hr after herbivory and were fast frozen in liquid nitrogen.

Hardness determination on pods

Hardness, the peak force required to penetrate a soybean pod, was measured with an INSTRON 4442 tensile tester calibrated to penetrate 12 mm at a rate of 2.0 mm sec^{-1} with a needle. The tests were performed on both soybean cultivars at the R6 phenological stage and the needle was inserted in pods at the second seed position. Results were expressed as gram-force per square centimeter (gf cm^{-2}) and represented the maximum force required to penetrate a pod with the tip needle in order to simulate the situation in which the sting bugs insert the stylus (de Santana Souza *et al.*, 2013).

Oxidative stress response

Extracts for determination of catalase (CAT), superoxide dismutase (SOD) and Guaiacol peroxidase (GPOX) activities were prepared from 0.3 g of soybean seeds homogenized under ice-cold conditions in 3mL of extraction buffer, containing 50mM phosphate buffer (pH 7.4), 1mMEDTA, 1 g PVP, and 0.5% (v/v) Triton X-100 at 4°C . The homogenates were centrifuged at $10\,000 \times g$ for 20 minutes and the supernatant fraction was used for the assays. Total protein content was determined by following the Bradford method for protein quantification (1976).

CAT activity was determined according to a modified protocol of Chance, Sies & Boveris (1979). The reaction was conducted in a quartz cuvette by measuring the decrease in absorption at 240 nm in a reaction medium containing 150 μL of enzyme extract, 50mM potassium phosphate buffer (pH 7.2) and 2mM H_2O_2 for 2 minutes. CAT activity was determined using the molar absorptivity of H_2O_2 at 240 nm and expressed as $\mu\text{moles mg protein}^{-1}\text{min}^{-1}$.

SOD activity was assayed by the inhibition of the photochemical reduction of NBT (Becana *et al.*, 1986). The reaction mixture consisted of 50, 100 and 200 μL of enzyme extract and 3.5mL $\text{O}_2^{\bullet-}$ generating solution which contained 14.3mM methionine, 82.5 μM NBT, and 2.2 μM riboflavin. Extracts were brought to a final volume of 0.3mL with 50mM K-phosphate (pH 7.8) and 0.1mM Na_2EDTA .

Sample tubes were shaken and placed in front of fluorescent lamps during 10 minutes. The reduction in NBT was followed by reading absorbance at 560 nm. Blanks and controls were run in the same way but without illumination and enzyme, respectively. One unit of SOD was defined as the amount of enzyme which produced a 50% inhibition of NBT reduction under the assay conditions.

GPOX activity was determined according to the protocol devised by Shannon *et al.* (1966) which consists in measuring the increase in absorption at 470 nm due to the formation of tetraguaiacol

at 30°C in a reaction extract, 50 mM buffer K-phosphate 50 mM, pH 7, 0.1 mM EDTA, 10 mM guaiacol and 10 mM H₂O₂. One unit of POD forms 1 µmol of guaiacol oxidised per min under assay conditions.

Detection of H₂O₂

To visually analyze H₂O₂ generation, stink bug damaged seeds and their respective controls were excised and immersed in a 1% solution of 3,3-Diaminobenzidine (DAB) for 24 h under light at 25 °C (Zilli *et al.*, 2008). DAB generates a reddish-brown compound that could be detected at the site of H₂O₂ formation in the presence of endogenous peroxidase activity. After staining, formation of brown precipitates was documented by photography.

Lipoxygenase expression

The expression level of Lipoxygenase 1 (*LOX1*) and Lipoxygenase 2 (*LOX2*) genes in soybean seeds was determined. Briefly, total RNA was isolated using the RNeasy Plant Mini kit (QIAGEN Inc., Valencia, CA, USA) according to the manufacturer's protocol. RNA concentration was determined with a fluorometer QubitTM (Invitrogen), and quality and quantity were assessed spectrophotometrically before the cDNA was synthesized using the Thermo Scientific RevertAid Reverse Transcriptase in a Bio-Rad My CyclerTM Thermal Cycler.

An ABI 7500 Fast Real-Time PCR system was used to perform the qRT-PCR (Applied Biosystems, Foster City, CA, USA) with the SYBR Green Real-time Master Mix. The *LOX1* and *LOX2* primer sequences in this study are detailed in Table 1. The housekeeping soybean elongation factor (*ELF1b*) was used as a reference gene for relative quantification and the target expression relative to the housekeeping was used for ANOVA.

Isoflavonoid determination

Isoflavonoid accumulation was determined after 72 h of herbivory treatment in seeds at R6 according to Fehr *et al* (1971), the lipid phase of seeds was removed with cyclohexane (Zavala *et al.*, 2015). 0.2 g of pulverized seeds were placed in a 50 ml tube and mixed with 10 mL of cyclohexane and incubated for 6 hours. Then, tubes were centrifuged at 1500 rpm for 10 minutes. Supernatant was discarded and the pellet was used for the extraction step. Isoflavonoids were extracted with methanol plus rutin that was used as an extraction internal control (0.1 g/ 5 ml MeOH). 10 mL of methanol were added to 50 mL tubes and incubated for 6 hours at room temperature. Then tubes were centrifuged at 1500 rpm for 10 minutes. Supernatant was saved. The solvent was then evaporated at 40 °C and samples were re-dissolved with 500 µL of MeOH. Samples were purified with a C18 Silica Cartridge (Agilent Technologies, Palo Alto, CA, USA) using different mixtures of MeOH–H₂O. Aliquots of 5 µL were subjected to high-performance liquid chromatography (HPLC; Agilent 1100 A series) using a reverse-phase octadecyl column (Eclipse XDB-C18 4.6 × 150 mm, 5 µm). Isoflavonoids were eluted using a mobile phase gradient of 15–60% acetonitrile in 0.1% acetic acid for 60 min at a flow rate of 1 mL min^{–1}. Compounds were measured with the detector set at λ 270 nm. Retention times and quantitative data for daidzin, daidzein, genistin and genistein were obtained by comparison to known standards (all from Sigma-Aldrich, St. Louis, MO, USA). External standard curves were performed for isoflavonoid quantification.

Statistical analysis

The experiments were set up as a completely randomized design with factorial treatments. One factor was the genotype (IAC-100 and Davis) and the second was insect damage (control and treated). Data were analyzed with the Infostat v. 2011 statistical package (Di Rienzo *et al.*, 2011). Hardness analysis of pods, and SOD, CAT and GPOX activities as well as seed isoflavonoid contents in each experiment were analyzed using analysis of variance (ANOVA). Data on isoflavonoid content were transformed as needed to meet the assumptions of ANOVA by square root transformation. We applied the Tukey pairwise multiple comparisons procedure for studying GENOTYPE and TREATMENT effects. Hierarchical clustering of oxidative stress enzymes was performed with the same data used for ANOVA using Euclidean distances in Infostat.

Results

Damage and hardness assessment of pods

Oxidative damage caused by stink bugs was observed by visual analyses of H₂O₂ generation. Davis seeds affected by *Nezara viridula* exhibited highly enhanced brownish staining compared to the IAC-100 genotype (Figure 1). Textural analysis of pods showed that IAC-100 has greater resistance to penetration than Davis ($p < 0.05$; Figure 2). This can be a physical barrier to the insect's stylet penetration.

Stress metabolism

According to our results, the cluster analysis of oxidative stress enzyme activities (CAT, SOD, GPOX) that was carried out using the statistical package Infostat v. 2011 (Di Rienzo *et al.*, 2011) showed two main clusters that grouped the genotypes with the treatments. Figure 3 A shows one cluster containing IAC-100 treated with stink bugs, and the second cluster containing IAC-100 control, Davis treated and Davis control. When exposed to insect treatment, the cluster distribution suggests a differential induction of enzyme activities between the resistant and susceptible genotypes. In addition, there is no Genotype x Treatment interaction in CAT activity and the differences are due to Genotype and Treatment ($p < 0.05$; Figure 3 B). Although the CAT enzyme is induced in both genotypes, its activity is always higher in IAC-100 than in Davis. SOD activity was significantly greater in IAC-100 ($p < 0.05$) when affected by stink bugs (Figure 3 C). When comparing cultivars, GPOX showed a higher activity in IAC-100 both with and without damage caused by the insect ($p < 0.05$, Figure 3 D).

Lipoxygenase expression quantification

Expression of *LOX1* and *LOX2* was induced by herbivory in both soybean cultivars when compared to their respective controls (Fig 4). It suggests that soybean plants respond to this type of stress by increasing *LOX* expression after 24 h following treatment ($p < 0.05$).

Isoflavonoids

Concentrations of isoflavonoids in soybean seeds showed differences between cultivars and types of compounds (Figure 5). After herbivory, the concentration of daidzin and genistin increased in both genotypes and constitutive concentrations were similar and very low ($p < 0.05$; Figure 5 A and C). Higher levels of daidzein were found in the IAC-100 genotype, regardless of the treatment, with a strong uprising trend under herbivory ($p < 0.05$; Figure 5 B). Except for one repeat of the cultivar Davis genistein was not detected in any treatment (supplementary information).

Discussion

Previous experiments with the IAC-100 soybean genotype reported low incidence of stink bugs (McPherson *et al.*, 2007). This cultivar has PI 229358 (Clement & Quisenberry, 1998) and PI 274454 (Souza *et al.*, 2015) in its genealogy (Carrao-Panizzi & Kitamura 1995), which exhibits a high degree of resistance to the stink bug complex. Some studies have shown antixenosis in IAC-100 against *N. viridula*, *P. guildinii* and *E. heros* (Campos *et al.*, 2010; Silva *et al.*, 2014 and de Souza *et al.*, 2014), and biological effects through isoflavonoids on *N. viridula* (Piubelli *et al.*, 2003a). Our study suggests that cultivar IAC-100 maybe more tolerant to the southern green stinkbug due to greater hardness in pods, which serves as the first line of defense. In addition, a higher antioxidative enzyme activity was involved in plant defense, and it showed high levels of isoflavonoids in response to insect attack which, according to some authors (Lane *et al.*, 1985; Murakami *et al.*, 2014; Simmonds & Stevenson, 2001) affect the performance and survival of the herbivores.

Antixenosis in resistant plants can be related to morphological factors such as thickened plant epidermal layers, waxy deposits on leaves, stems, or fruits, nutritional deficiency, and chemical compounds (Smith, 2005). The thickness of various plant tissues influences the degree of resistance in some crop cultivars. Stink bug (*N. viridula*) damage elicits activation of MAPK signal in soybean seeds and induced salicylic acid that induced genes related with cell wall restructuration and increased lignin content, which can increase resistance to new insect attack by hardening cell walls (Giacometti *et al.*, 2018). In this study, pods of IAC-100 cultivar showed greater resistance to penetration when compared to those of Davis, suggesting that hardening may be a physical barrier to insect feeding (Figure 2).

Under stress conditions in plants, such as insect feeding, many signaling pathways are activated by different types of ROS causing “oxidative burst” (Maffei, Mithöfer & Boland, 2007). After insect attack, ROS are accumulated acting as the first barrier against the attack of pathogens and herbivores. However, to avoid self-toxicity by ROS, plant cells have ROS scavenging systems that help to remove the excess concentration and maintain a relatively low and constant one (War *et al.*, 2012). Tolerance in plants is generally related to the control capacity of the cellular redox state resulting in lower oxidative damage (Slesak *et al.* 2007). ROS accumulation was detected after the feeding period of the chewing caterpillar *S. littoralis* or the piercing-sucking action of the spider mite *Tetranychus urticae* (Maffei *et al.*, 2006; Afify *et al.*, 2011). Infestation of lima beans by *S. littoralis* revealed, trough diaminobenzidine (DAB) staining, accumulation of H₂O₂ around the wounded area (Maffei *et al.*, 2006). In our study we found that IAC-100 had less accumulation of H₂O₂ after insect feeding when compared to the susceptible genotype, which suggests that it has a higher capacity to control the redox state. A previous study demonstrated that ROS detoxification enzymes were only induced in H₂O₂-accumulating resistant lines (Moloi & van der Westhuizen 2008). POD is one of such group of antioxidative enzymes, which scavenges the ROS besides having other defensive roles because they are also an important component of the immediate response of plants to insect damage (War *et al.*, 2012). Catalases remove the H₂O₂ and reduce H₂O₂ to 2H₂O. And SOD constitutes a frontline in the defense against ROS as they catalyze the dismutation of O₂⁻. (superoxide radical) to H₂O₂ (Caverzan, Casassola & Brammer, 2016). In wheat, an increase in *SOD* transcript in response to differential heat shock treatment was related to an enhanced tolerance to environmental stresses (Kumar *et al.*, 2013). The present study demonstrates that the resistant cultivar had more control of the

cellular redox state given its high level of enzyme activity after herbivory in SOD, GPOX y CAT (Figure 3).

Many studies have shown that LOXs are implicated in the defense systems of several plant species against pathogens and insects (Koch *et al.*, 1992; Melan *et al.*, 1993; Felton *et al.*, 1994; Christensen *et al.*, 2013). Induction of LOX activity after herbivory has previously been studied in tomato in response to *Helicoverpa armigera* (Yan *et al.*, 2013), in Arabidopsis following infestation by *Myzus persicae* and in lima bean in response to *Tetranychus urticae* (Moran & Thompson, 2001; Arimura *et al.*, 2000). In developing seeds of soybean, we found an induction of *LOX1* and *LOX2* after herbivory in both genotypes under study suggesting that this pathway is activated in response to the damage caused by *Nezara viridula*.

LOX activity regulates the production of the hormone JA, which modulates flavonoids and isoflavonoids that protect plants against insect pests and affect the behavior, growth and development of insects (Simmonds, 2003). In Arabidopsis, resistance against *Spodoptera frugiperda* is enhanced overexpressing a transcription factor that controls flavonoid production (Johnson & Dowd, 2004). In soybean leaves, flavonoids negatively affected the behavior of *Aphis glycines* and *Trichoplusia ni* (Meng *et al.*, 2011; Neupane & Norris 1990). Moreover, soybean seed damage caused by soybean pod borer (*Leguminivora glycinivorella*) was negatively correlated with higher levels of isoflavonoids, and showed positive correlations between isoflavonoid content in pods and undamaged seeds in treatments with stink bugs (*Nezara viridula* and *Piezodorus guildinii*) (Zhao *et al.*, 2015; Zavala *et al.*, 2015). Our results reveal that stink bug herbivory differentially promoted isoflavonoid production between genotypes with variations according to the type of compound. It can thus be concluded that not only total production of isoflavonoids but also the type of isoflavonoid produced can affect insect behavior.

Conclusions

Taken the results together, it can be inferred that stink bug herbivory injury generates cell wall changes, induces pathways related to oxidative stress and secondary metabolites in developing seeds of soybean. As a result of the resistance characteristics to insects, such as control of cellular redox state and production of secondary metabolites, IAC-100 is a promising cultivar for a breeding program. However, further studies are still required to understand the functions of genes and the regulatory factors involved in defense responses in order to develop molecular markers for a breeding program.

The study of host plant resistance is an important alternative for producers in Argentina, but even more in Brazil, not only because of the high insect pressure but also for the phytopathogenic fungi that lead to the frequent use of chemical control.

Acknowledgements

GDM Seeds for the availability for the use of greenhouses and other infrastructure. Verónica Feuring and Alina Crelier for technical assistance. Romina Giacometti and Natalia Ilina for the help in molecular biology methods.

References

Accamando AK, Cronin JT. (2012). Costs and benefits of jasmonic acid-induced responses in soybean. *Environmental entomology*, 41(3), 551-561

- 363 Afify AEMM, El-Beltagi HS, Fayed SA, Shalaby EA (2011). Acaricidal activity of different
364 extracts from *Syzygiumcumini* L. Skeels (Pomposia) against *Tetranychus urticae* Koch. *Asian*
365 *Pacific journal of tropical biomedicine*, 1(5), 359-364.
- 366
- 367 Arimura GI, Ozawa R, Shimoda T, Nishioka T, Boland W, Takabayashi, J (2000). Herbivory-
368 induced volatiles elicit defense genes in lima bean leaves. *Nature*, 406(6795), 512-515.
- 369
- 370 Axelrod B, Cheesbrough TM, Laakso S (1981). Lipoxygenase from soybeans: EC 1.13. 11.12
371 Linoleate: oxygen oxidoreductase. In *Methods in Enzymology* (Vol. 71, pp. 441-451). Academic
372 Press.
- 373
- 374 Becana M, Aparicio-Tejo P, Irigoyen JJ, Sanchez-Diaz M (1986). Some enzymes of hydrogen
375 peroxide metabolism in leaves and root nodules of *Medicago sativa*. *Plant Physiology*, 82(4),
376 1169-1171.
- 377
- 378 Blee E (2002). Impact of phyto-oxylipins in plant defense. *Trends Plant Sci* 7, 315–321.
- 379
- 380 Bi J.L., Felton G.W. (1995). Foliar oxidative stress and insect herbivory: primary compounds,
381 secondary metabolites, and reactive oxygen species as components of induced resistance.
382 *Journal of Chemical Ecology*, 21(10), 1511-1530.
- 383
- 384 Bradford MM (1976). A rapid and sensitive method for the quantitation of microgram quantities
385 of protein utilizing the principle of protein-dye binding. *Analytical biochemistry*, 72(1-2), 248-
386 254.
- 387
- 388 Bednarski W, Borowiak-Sobkowiak B, Wilkaniec B, Samardakiewicz S, Morkunas I (2013).
389 Oxidative stress in pea seedling leaves in response to *Acyrtosiphon pisum*
390 infestation. *Phytochemistry*, 93, 49-62.
- 391
- 392 Boerma HR, Walker DR (2005). Discovery and utilization of QTLs for insect resistance in
393 soybean. *Genetica*. 123, 181–189.
- 394
- 395 Browse J (2009). Jasmonate passes muster: a receptor and targets for the defense hormone. *Annu*
396 *Rev Plant Biol* 60, 183–205.
- 397
- 398 Campos M, Knutson A, Heitholt J, Campos C (2010). Resistance to seed feeding by southern
399 green stink bug, *Nezara viridula* (Linnaeus), in soybean, *Glycine max* (L.) Merrill. *Southwestern*
400 *Entomologist*, 35(3), 233-239.
- 401
- 402 Carrao-Panizzi MC, Kitamura K (1995). Isoflavone content in Brazilian soybean
403 cultivars. *Japanese Journal of Breeding*, 45(3), 295-300.
- 404
- 405 Caverzan A, Casassola A, Brammer SP (2016). Antioxidant responses of wheat plants under
406 stress. *Genetics and molecular biology*, 39(1), 1-6.
- 407

- Chance B, Sies H, Boveris A (1979). Hydroperoxide metabolism in mammalian organs. *Physiological reviews*, 59(3), 527-605.
- Chen H, Wang FW, Dong YY, Wang N, Sun YP, Li XY, Liu L, Fan XD, Yin HL, Jing YY, Zhang XY, Li YL, Chen G, Li HY (2012). Sequence mining and transcript profiling to explore differentially expressed genes associated with lipid biosynthesis during soybean seed development. *BMC plant biology*, 12(1), 122.
- Christensen SA, Nemchenko A, Borrego E, Murray I, Sobhy IS, Bosak L, De Blasio S, Erb M, Robert CAM, Vaughn K A, Herrfurth C, Tumlinson J, Feussner I, Jackson D, Turlings TCJ, Engelberth J, Nansen C, Meeley R, Kolomiets MV (2013). The maize lipoxygenase, ZmLOX10, mediates green leaf volatile, jasmonate and herbivore-induced plant volatile production for defense against insect attack. *The Plant Journal*, 74(1), 59-73.
- Clement SL, Quisenberry SS (1998). Global plant genetic resources for insect-resistant crops. CRC Press.
- Corrêa-Ferreira, B. S., Moscardi, F. (1996). Biological control of soybean stink bugs by inoculative releases of *Trissolcus basal*. *Entomologia experimentalis et applicata*, 79(1), 1-7.
- de Moraes AA, Pinheiro JB (2012). Breeding for Resistance to Insect Pests. In *Plant Breeding for Biotic Stress Resistance* (pp. 103-125). Springer Berlin Heidelberg.
- de Santana Souza E, Lopes Baldin EL, da Silva GF, Paulo J, Lourenção AL (2013). Feeding preference of *Nezara viridula* (Hemiptera: Pentatomidae) and attractiveness of soybean genotypes. *Chilean journal of agricultural research*, 73(4), 351-357.
- de Souza PV, Machado BR, da Silva DC, Menezes IPP, Araújo MS, de Jesus FG (2014). Effect of resistance and trichome inducers on attraction of *Euschistus heros* (Hemiptera: Pentatomidae) to soybeans. *African Journal of Agricultural Research*, 9(10), 889-894.
- Di Rienzo JA, Casanoves F, Balzarini MG, Gonzalez L, Tablada M, Robledo CW (2011) *InfoStat, Statistical Software version 2011*. Universidad Nacional de Córdoba Argentina.
- Ederli L, Brunetti C, Centritto M, Colazza S, Frati F, Loreto F, Marino G, Salerno G, Pasqualini S (2017). Infestation of broad bean (*Vicia faba*) by the green stink bug (*Nezara viridula*) decreases shoot abscisic acid contents under well-watered and drought conditions. *Frontiers in plant science*, 8, 959.
- Farmer EE, Johnson RR, Ryan CA (1992). Regulation of expression of proteinase inhibitor genes by methyl jasmonate and jasmonic acid. *Plant Physiology* 98, 995–1002.
- Farmer EE, Ryan CA (1990). Interplant communication: airborne methyl jasmonate induces synthesis of proteinase inhibitors in plant leaves. *Proc Natl Acad Sci*, 87, 7713-7716.

- 453 Fehr WR, Caviness CE, Burmood DT, Pennington JS (1971). Stage of development descriptions
454 for soybeans, *Glycine max* (L.) Merrill. *Crop Science* 11,929–931.
- 455
- 456 Felton GW, Bi JL, Summers CB, Mueller AJ, Duffey SS (1994). Potential role of lipoxygenases
457 in defense against insect herbivory. *Journal of Chemical Ecology*, 20(3), 651-666.
- 458 Feussner I., Wasternack, C. (2002). The lipoxygenase pathway. *Annu Rev Plant Biol* 53, 275–
459 297.
- 460
- 461 Freeman BC, Beattie GA (2008). An Overview of Plant Defenses against Pathogens and
462 Herbivores. *The Plant Health Instructor*. DOI: 10.1094/PHI-I-2008-0226-01
- 463
- 464 Fortunato FS, Oliveira MGA, Brumado MHN, Silva CHO, Guedes RNC, Moreira MA (2007).
465 Lipoxygenase-induced defense of soybean varieties to the attack of the velvetbean caterpillar
466 (*Anticarsia gemmatilis* Hubner). *Journal of Pest Science*, 80, 241–247.
- 467
- 468 Fürstenberg-Hägg J, Zagrobelny M, Bak S (2013). Plant defense against insect
469 herbivores. *International journal of molecular sciences*, 14(5), 10242-10297.
- 470
- 471 Giacometti R, Barneto J, Barriga LG, Sardoy PM, Balestrasse K, Andrade AM, Pagano EA,
472 Alemano SG, Zavala JA (2016). Early perception of stink bug damage in developing seeds of
473 field-grown soybean induces chemical defenses and reduces bug attack. *Pest management*
474 *science*. 72: 1585–1594
- 475
- 476 Giacometti R, Illina N, Pagano EA, Zavala JA (2018). Stink bug *nezara viridula* sustains late
477 MAPKs phosphorylation status and induces expression of genes related with cell wall
478 rearrangement in developing soybean seeds. *Arthropod-Plant Interactions*. 12: 531-541.
- 479
- 480 Howe G, Schilmiller AL (2002). Oxylin metabolism in response to stress. *Curr Opin Plant*
481 *Biol* 5, 230–236.
- 482
- 483 Howe G, Jander G (2008). Plant immunity to insect herbivores. *Annual Review of Plant Biology*,
484 59, 41-66.
- 485
- 486 Jensen RL, Newsom LD (1972). Effect of stink-bug-damaged soybean seeds on germination,
487 emergence and yield. *Journal of Economic Entomology*, 65, 261-264.
- 488
- 489 Jian B, Liu B, Bi YR, Hou WS, Wu CX, Han TF (2008) Validation of internal control for gene
490 expression study in soybean by quantitative real-time PCR. *BMC Mol Biol* 9:59. doi:
491 10.1186/1471-2199-9-59.
- 492
- 493 Johnson ET, Dowd PF (2004). Differentially enhanced insect resistance, at a cost, in *Arabidopsis*
494 *thaliana* constitutively expressing a transcription factor of defensive metabolites. *Journal of*
495 *agricultural and food chemistry*, 52(16), 5135-5138.
- 496

- 497 Kester K M, Smith CM, Gilman DF (1984). Mechanisms of resistance in soybean (*Glycine max*
498 [L.] Merrill) genotype PI171444 to the southern green stink bug, *Nezara viridula* (L.)
499 (Hemiptera: Pentatomidae). *Environmental entomology*, 13(5), 1208-1215.
- 500
- 501 Koch E, Meier BM, Eiben HG, Slusarenko A (1992). A lipoxygenase from leaves of tomato
502 (*Lycopersicon esculentum* Mill.) is induced in response to plant pathogenic pseudomonads. *Plant*
503 *Physiology*, 99(2), 571-576.
- 504
- 505 Kumar RR, Sharma SK, Goswami S, Singh K, Gadpayl KA, Singh GP, Pathak H, Rai RD
506 (2013). Transcript profiling and biochemical characterization of mitochondrial superoxide
507 dismutase (mtSOD) in wheat (*Triticum aestivum*) under different exogenous stresses. *Australian*
508 *Journal of Crop Science*, 7(3), 414.
- 509
- 510 Lane GA, Biggs DR, Russell GB, Sutherland OR, Williams EM, Maindonald JH, Donnell DJ
511 (1985). Isoflavonoid feeding deterrents for *Costelytra zealandica* Structure—Activity
512 relationships. *Journal of chemical ecology*, 11(12), 1713-1735.
- 513
- 514 Maffei ME, Mithöfer A, Boland W (2007). Insects feeding on plants: rapid signals and responses
515 preceding the induction of phytochemical release. *Phytochemistry*, 68(22), 2946-2959.
- 516
- 517 Maffei ME, Mithöfer A, Arimura GI, Uchtenhagen H, Bossi S, Berteau CM, Starvaggi Cucuzza
518 L, Novero M, Volpe, V, Quadro S, Boland W (2006). Effects of feeding *Spodoptera littoralis* on
519 lima bean leaves. III. Membrane depolarization and involvement of hydrogen peroxide. *Plant*
520 *Physiology*, 140(3), 1022-1035.
- 521
- 522 McPherson RM, Buss GR, Roberts PM (2007). Assessing stink bug resistance in soybean
523 breeding lines containing genes from germplasm IAC-100. *Journal of Economic Entomology*.
524 *Journal of economic entomology*, 100(4), 1456-1463.
- 525
- 526 Melan MA, Dong X, Endara ME, Davis KR, Ausubel FM, Peterman TK (1993). An Arabidopsis
527 thaliana lipoxygenase gene can be induced by pathogens, abscisic acid, and methyl
528 jasmonate. *Plant Physiology*, 101(2), 441-450.
- 529
- 530 Meng F, Han Y, Teng W, Li Y, Li W (2011). QTL underlying the resistance to soybean aphid
531 (*Aphis glycines* Matsumura) through isoflavone-mediated antibiosis in soybean cultivar
532 ‘Zhongdou 27’. *Theoretical and applied genetics*, 123(8), 1459-1465.
- 533
- 534 Michereff MFF, Laumann RA, Borges M, Michereff-Filho M, Diniz I R, Neto ALF, Moraes
535 MCB (2011). Volatiles mediating a plant-herbivore-natural enemy interaction in resistant and
536 susceptible soybean cultivars. *Journal of chemical ecology*, 37(3), 273-285.
- 537
- 538 Misra RC, Maiti P, Chanotiya CS, Shanker K, Ghosh S (2014). Methyl jasmonate-elicited
539 transcriptional responses and pentacyclic triterpene biosynthesis in sweet basil. *Plant*
540 *physiology*, 164(2), 1028-1044.
- 541

- Moloi MJ, van der Westhuizen AJ (2008). Antioxidant responses and the Russian wheat aphid (*Diuraphis noxia*) resistance responses in wheat (*Triticum aestivum*). *Plant Biology* 10, 403–407.
- Moran PJ, Thompson G A (2001). Molecular responses to aphid feeding in Arabidopsis in relation to plant defense pathways. *Plant Physiology*, 125(2), 1074-1085.
- Murakami S, Nakata R, Aboshi T, Yoshinaga N, Teraishi M, Okumoto Y, Ishihara A, Morisaka H, Huffaker A, Schmelz E A, Mori, N (2014). Insect-induced daidzein, formononetin and their conjugates in soybean leaves. *Metabolites*, 4(3), 532-546.
- Neupane FP, Norris DM (1990). Iodoacetic acid alteration of soybean resistance to the cabbage looper (Lepidoptera: Noctuidae). *Environmental entomology*, 19(2), 215-221.
- Orr DB, Boethel D J, Jones WA (1985). Biology of *Telenomus chloropus* (Hymenoptera: Scelionidae) from eggs of *Nezara viridula* (Hemiptera: Pentatomidae) reared on resistant and susceptible soybean genotypes. *The Canadian Entomologist*, 117(09), 1137-1142.
- Piubelli GC, Hoffmann-Campo CB, De Arruda IC, Franchini JC, Lara FM (2003a). Flavonoid increase in soybean as a response to *Nezara viridula* injury and its effect on insect-feeding preference. *Journal of chemical ecology*, 29(5), 1223-1233.
- Piubelli GC, Hoffmann-Campo CB, Arruda ICD, Lara FM (2003b). Nymphal development, lipid content, growth and weight gain of *Nezara viridula* (L.) (Heteroptera: Pentatomidae) fed on soybean genotypes. *Neotropical Entomology*, 32(1), 127-132.
- Priolli RHG, Mendes-Junior CT, Arantes NE, Contel EPB (2002). Characterization of Brazilian soybean cultivars using microsatellite markers. *Genetics and Molecular Biology*, 25(2), 185-193.
- Shannon LM, Kay E, Lew JY (1966). Peroxidase isozymes from horseradish roots I. Isolation and physical properties. *Journal of Biological Chemistry*, 241(9), 2166-2172.
- Savatin DV, Gramegna G, Modesti V, Cervone F (2014). Wounding in the plant tissue: the defense of a dangerous passage. *Frontiers in plant science*, 5(470):470.
- Schaefer, C. W., Panizzi, A. R. (Eds.). (2000). Heteroptera of economic importance. CRC press
- Silva JP, Baldin EL, Canassa VF, Souza ES, Lourenção AL (2014). Assessing antixenosis of soybean entries against *Piezodorus guildinii* (Hemiptera: Pentatomidae). *Arthropod-Plant Interactions*, 8(4), 349-359.
- Simmonds M S (2003). Flavonoid–insect interactions: recent advances in our knowledge. *Phytochemistry*, 64(1), 21-30.
- Simmonds M S, Stevenson P C (2001). Effects of isoflavonoids from Cicer on larvae of *Heliocoverpa armigera*. *Journal of chemical ecology*, 27(5), 965-977.

- Slesak I, Libik M, Karpinska B, Karpinski S, Miszalski Z (2007). The role of hydrogen peroxide in regulation of plant metabolism and cellular signaling in response to environmental stresses. *Acta Biochimica Polonica-English Edition*, 54(1), 39.
- Smith CM (2005). Plant resistance to arthropods: Molecular and conventional approaches, Dordrecht, The Netherlands: Springer.
- Souza ES, Silva JPGF, Baldin ELL, Pierozzi CG, Cunha LS, Canassa VF, Pannuti LER, Lourenção AL (2015). Response of soybean genotypes challenged by a stink bug complex (Hemiptera: Pentatomidae). *Journal of economic entomology*, 109(2), 898-906.
- Wang J, Qian J, Yao L, Lu Y (2015). Enhanced production of flavonoids by methyl jasmonate elicitation in cell suspension culture of *Hypericum perforatum*. *Bioresources and Bioprocessing*, 2(1), 5.
- War AR, Paulraj MG, Ahmad, T, Buhroo AA, Hussain B, Ignacimuthu S, Sharma H C (2012). Mechanisms of plant defense against insect herbivores. *Plant signaling and behavior*, 7(10), 1306-1320.
- Wasternack C (2007). Jasmonates: an update on biosynthesis, signal transduction and action in plant stress response, growth and development. *Ann Bot*. 100(4), 681-97.
- Yan L, Zhai Q, Wei J, Li S, Wang B, Huang T, Du M, Sun J, Kang L, Li CB, Li, C (2013). Role of tomato lipoxygenase D in wound-induced jasmonate biosynthesis and plant immunity to insect herbivores. *PLoS genetics*, 9(12), e1003964.
- Zavala JA, Mazza CA, Dillon FM, Chludil HD, Ballare CL (2015). Soybean resistance to stink bugs (*Nezara viridula* and *Piezodorus guildinii*) increases with exposure to solar UV-B radiation and correlates with isoflavonoid content in pods under field conditions. *Plant, cell and environment*, 38 (5), 920-928.
- Zavala J, Barneto J, Barriga L, Pagano E (2010). Plant-Insect Interactions in Soybean Crops: Plant Responses to Stink Bugs. 1st Latin American Meeting of Chemical Ecology (p. 82). Colonia del Sacramento: Latin American Association of Chemical Ecology.
- Zavala JA, Casteel CL, De Lucia EH, Berenbaum MR (2008). Anthropogenic increase in carbon dioxide compromises plant defense against invasive insects. *Proceedings of the national academy of sciences*, 105(13), 5129-5133.
- Zebelo SA, Maffei ME (2014). Role of early signalling events in plant-insect interactions. *Journal of Experimental Botany*, 66(2), 435-448.
- Zhang SZ, Hua BZ, Zhang F (2008). Induction of the activities of antioxidative enzymes and the levels of malondialdehyde in cucumber seedlings as a consequence of *Bemisia tabaci* (Hemiptera: Aleyrodidae) infestation. *Arthropod-Plant Interactions*, 2(4), 209-213.

634

635 Zhao G, Jiang Z, Li D, Han Y, Hu H, Wu L, Wang Y, Gao Y, Teng W, Li Y, Zeng G, Meng F,
636 Zeng, G Meng F, Li W (2015). Molecular loci associated with seed isoflavone content may
637 underlie resistance to soybean pod borer (*Leguminivora glycinivorella*). *Plant Breeding*, 134(1),
638 78-84.

639

640 Zilli CG, Balestrasse KB, Yannarelli GG, Polizio AH, Santa-Cruz, DM, Tomaro ML (2008).
641 Heme oxygenase up-regulation under salt stress protects nitrogen metabolism in nodules of
642 soybean plants. *Environmental and experimental botany*, 64(1), 83-89.

643

Figure 1

H₂O₂ generation in sites damaged by stink bug.

Brown color shows a more affected condition. Images were acquired with a digital camera Nikon Coolpix L110.




Figure 2

Mean hardness of soybean pods in R6

Means values (bars) and SEM (whiskers) are shown. Different letters are significantly different ($p < 0.05$ - $n = 17$). Data source in Supplemental Information.




Figure 3

Oxidative stress analysis in seeds of different soybean genotypes damaged by *Nezara viridula* and their respective controls without herbivory

(A) Cluster analysis of oxidative stress enzymes. **(B)** Catalase activity. **(C)** Superoxide dismutase activity. **(D)** Guaiacol peroxidase activity. Means values (dots and squares) and SEM (whiskers) are shown. Different letters are significantly different ($p < 0.05$). Data source in Supplemental Information.




Figure 4

Gene expression by Real-Time PCR of LOX1 and LOX2 in soybean seeds, either undamaged and damaged by *Nezara viridula*

(A) LOX1. (B) LOX2. Gene expression is relative to the soybean elongation factor (**ELF1b**). Means values (bars) and SEM (whiskers) are shown. Different letters are significantly different ($p < 0.05$). Data source in Supplemental Information.




Figure 5

Isoflavonoid concentration in seeds of cultivars Davis and IAC-100 damaged by stink bugs and their respective controls without herbivory

(A) Daidzin concentration. **(B)** Genistin concentration. **(C)** Daidzein concentration. Means values (bars) and SEM (whiskers) are shown. Different letters are significantly different ($p < 0.05$). Data source in Supplemental Information.




Table 1 (on next page)

LOX1, LOX2 and ELF1b primer sequences

1

<i>LOX1</i>	Forward: 5' CTGGTGTAATCCCTGCGTAA 3'	Chen <i>et al.</i> (2012)
	Reverse: 5' TACCAAGTGCCTCGTCCATT 3'	
<i>LOX2</i>	Forward: 5' AGATGGTTGCGGGTGTAAT 3'	Chen <i>et al.</i> (2012)
	Reverse: 5' GGGCATCTGCTGTTATCTTAC 3'	
<i>ELF1b</i>	Forward: 5'- GTTGAAAAGCCAGGGGACA -3'	Jian <i>et al.</i> (2008)
	Reverse: 5'- TCTTACCCCTTGAGCGTGG -3'	