

A peer-reviewed version of this preprint was published in PeerJ on 2 July 2015.

[View the peer-reviewed version](https://doi.org/10.7717/peerj.1068) (peerj.com/articles/1068), which is the preferred citable publication unless you specifically need to cite this preprint.

Mekov E, Slavova Y, Tsakova A, Genova M, Kostadinov D, Minchev D, Marinova D. 2015. Metabolic syndrome in hospitalized patients with chronic obstructive pulmonary disease. PeerJ 3:e1068
<https://doi.org/10.7717/peerj.1068>

Metabolic syndrome in hospitalized patients with chronic obstructive pulmonary disease

Evgeni Mekov, Yanina Slavova, Adelina Tsakova, Marianka Genova, Dimitar Kostadinov, Delcho Minchev, Dora Marinova

The metabolic syndrome (MS) affects 21-53% of patients with chronic obstructive pulmonary disease (COPD) with a higher prevalence in the early stages of COPD, with results being highly variable between studies. MS may also correlate with disease characteristics. The aim of the study is to examine the prevalence of MS and its correlation with comorbidities and COPD characteristics in patients with COPD admitted for exacerbation. 152 patients with COPD admitted for exacerbation were studied for presence of MS. All of them were also assessed for vitamin D status and diabetes mellitus type 2 (DM). Data were gathered for smoking status and exacerbations during the last year. All patients completed CAT (COPD assessment test) and mMRC (Modified Medical Research Council Dyspnea scale) questionnaires and underwent spirometry. Duration of current hospital stay was recorded. 25% of patients have MS. 23,1% of the male and 29,5% of the female patients have MS ($p>0.05$). The prevalence of MS in this study is significantly lower when compared to a national representative study (44,6% in subjects over 45 years). 69,1% of all patients and 97,4% from MS patients have arterial hypertension. The presence of MS is associated with significantly worse cough and sleep (1st and 7th CAT questions; $p=0.002$ and $p=0.001$ respectively) and higher total CAT score ($p=0.017$). Average BMI is 27,31. None of the patients have MS and BMI <25 . There is a correlation between the presence of MS and DM ($p=0.008$) and with the number of exacerbations in the last year ($p=0.015$). There is no correlation between the presence of MS and the pulmonary function. This study among hospitalized COPD patients finds comparable but relatively low prevalence of MS (25%) compared to previously published data (21-53%) and lower prevalence compared to general population (44,6%). MS may impact natural course and the number of exacerbations of COPD. Having in mind that MS is more common in the early stages and decreases with COPD progression, the COPD patients admitted for exacerbation may be considered as having advanced COPD.

Metabolic syndrome in hospitalized patients with chronic obstructive pulmonary disease

Evgeni Mekov¹, Yanina Slavova¹, Adelina Tsakova², Marianka Genova², Dimitar Kostadinov¹,
Delcho Minchev¹, Dora Marinova¹

1 - Clinical Center for Pulmonary Diseases, Medical University - Sofia, Sofia, Bulgaria

2- Central Clinical Laboratory, Medical University - Sofia, Sofia, Bulgaria

Abstract

Introduction: The metabolic syndrome (MS) affects 21-53% of patients with chronic obstructive pulmonary disease (COPD) with a higher prevalence in the early stages of COPD, with results being highly variable between studies. MS may also correlate with disease characteristics.

Aim: To examine the prevalence of MS and its correlation with comorbidities and COPD characteristics in patients with COPD admitted for exacerbation.

Material and methods: 152 patients with COPD admitted for exacerbation were studied for presence of MS. All of them were also assessed for vitamin D status and diabetes mellitus type 2 (DM). Data were gathered for smoking status and exacerbations during the last year. All patients completed CAT (COPD assessment test) and mMRC (Modified Medical Research Council Dyspnea scale) questionnaires and underwent spirometry. Duration of current hospital stay was recorded.

Results: 25% of patients have MS. 23,1% of the male and 29,5% of the female patients have MS ($p>0.05$). The prevalence of MS in this study is significantly lower when compared to a national representative study (44,6% in subjects over 45 years). 69,1% of all patients and 97,4% from MS patients have arterial hypertension. The presence of MS is associated with significantly worse cough and sleep (1st and 7th CAT questions; $p=0.002$ and $p=0.001$ respectively) and higher total CAT score ($p=0.017$). Average BMI is 27,31. None of the patients have MS and BMI <25 . There is a correlation between the presence of MS and DM ($p=0.008$) and with the number of exacerbations in the last year ($p=0.015$). There is no correlation between the presence of MS and the pulmonary function.

Conclusions: This study among hospitalized COPD patients finds comparable but relatively low prevalence of MS (25%) compared to previously published data (21-53%) and lower prevalence compared to general population (44,6%). MS may impact natural course and the number of exacerbations of COPD. Having in mind that MS is more common in the early stages and decreases with COPD progression, the COPD patients admitted for exacerbation may be considered as having advanced COPD.

Funding: This manuscript is part of a PhD project, which is partially funded by Medical University – Sofia, Sofia, Bulgaria (grant number 15-D/2014, project number 22-D/2014).

Introduction

Chronic Obstructive Pulmonary Disease (COPD) is a preventable and treatable disease with significant extrapulmonary effects that may contribute to the severity in individual patients. By 2030, COPD will be the fourth cause of mortality worldwide. The extrapulmonary comorbidities influence the prognosis of the patients with COPD (1).

Metabolic syndrome (MS) is common in patients with COPD. According to the available studies the prevalence of MS in COPD patients varies between 21-53% (2). The prevalence of MS in COPD patients is increased when compared to a control group (3-6).

Available studies suggest that MS may have impact on quality of life (7), lung function (8-12), natural course of COPD (number of exacerbations) (13,14) as well as to affect comorbidities in COPD patients (15).

Many studies examine prevalence of MS in COPD patients (3,4,6,13,16-21) with results being highly variable between studies. Prevalence of MS in an unselected Bulgarian population aged 20-80 years is 30,8%. The prevalence of MS for participants over 45 years (the most common age group for the COPD patients) is higher - 44,6% (22). An epidemiological study conducted in 2010 in Bulgaria in COPD patients (n=3598) showed that metabolic syndrome is found in 13.8% of the patients (23). These results differ significantly from the others probably because of the different criteria for metabolic syndrome. A more recent study indicates that the prevalence of metabolic syndrome is 41.8% in patients with COPD (n=141) compared to 39% in the control group (n=103). However this study was not conducted using a random sample (exclusion criteria was presence of DM, age 49-79 years for patients with COPD and 35-65 for controls) (24). The prevalence of MS in COPD patients, hospitalized for exacerbation is hard to predict because MS tends to be more prevalent in early stages of COPD while patients experiencing severe exacerbation often have advanced disease. On the other side MS may impact natural course of COPD and predispose to exacerbation which will lead to increased prevalence of MS in this group.

There is not enough data to determine whether the results from these studies are applicable to specific subgroups of patients such as COPD patients admitted for exacerbation. COPD is increasingly divided in subgroups or phenotypes based on specific features and association with prognosis or response to therapy, the most notable being the feature of frequent exacerbations (25). We hypothesize that the prevalence of MS in COPD patients admitted for exacerbation is low (because of the severity of COPD in this subgroup and the MS being more prevalent in the early stages). Presence of MS may also have distinctive characteristics for this subgroup ('severe' exacerbator phenotype). The aim of this study is to find out the prevalence of MS in patients with COPD admitted for exacerbation and the correlations of presence of MS with comorbidities and COPD characteristics.

Material and methods

A total of 152 COPD patients hospitalized for exacerbation were studied for the presence of MS, DM, and vitamin D deficiency and insufficiency using well-established criteria for:

- ❖ Presence of MS: at least 3 of the following: 1. Elevated waist circumference >102 cm in males, >88 cm in females; 2. Triglycerides >1.7 mmol/L (or on therapy); 3. HDL <1.0 mmol/L in males, <1.3 mmol/L in females (or on therapy); 4. Elevated blood pressure: systolic \geq 130 and/or diastolic \geq 85 mm Hg (or on therapy); 5. Fasting glucose >5.5 mmol/L (or on therapy) (26).

- ❖ Presence of DM: fasting plasma glucose ≥ 7.0 mmol/L OR 2-h plasma glucose ≥ 11.1 mmol/L during an oral glucose tolerance test (OGTT) OR HbA1c $\geq 6.5\%$ OR on therapy (27);
- ❖ Presence of prediabetes: fasting plasma glucose 5.6-6.9 mmol/L OR 2-h plasma glucose 7.8-11.0 mmol/L during an OGTT OR HbA1c 5.7-6.4% (27);
- ❖ Presence of vitamin D deficiency: 25(OH)D < 25 nmol/L; vitamin D insufficiency: 25(OH)D 25-50 nmol/L; vitamin D sufficiency: > 50 nmol/L (28,29).

The diagnosis of COPD was made according to GOLD (Global Initiative for Chronic Obstructive Lung Disease) criteria (DM1). Data were gathered for age, sex, smoking status and number of pack-years, number of bone fractures, therapy for arterial hypertension, therapy for DM, COPD therapy and number of exacerbations in the last year. The patients completed CAT and mMRC questionnaires and underwent pre- and post bronchodilatory spirometry. Blood pressure was obtained according to the American Heart Association Guidelines (30). A patient was considered as having arterial hypertension if taking antihypertensives.

The inclusion criteria were post bronchodilator spirometry obstruction defined as FEV1/FVC < 0.70 . All participants in this study signed informed consent.

The exclusion criteria were failure to comply with study procedures (no completed questionnaires, no medical and demographic information, no spirometry, no lab tests) or FEV1/FVC ratio > 0.70 after administration of bronchodilator.

Smoking status

Every participant was classified according to smoking status (31):

Never smoker – never smoked a cigarette or who smoked fewer than 100 cigarettes in their entire lifetime.

Former smoker – smoked at least 100 cigarettes in their entire life but were not currently smoking.

Current smoker – had smoked at least 100 cigarettes in their entire life and were still smoking.

Numbers of pack-years were calculated using the formula:

Number of pack-years = years of smoking X number of daily smoked cigarettes/20

Anthropometric indices

Body weight and height were measured and the body mass index (BMI) was calculated by dividing weight by height squared (kg/m^2). According to BMI all patients were classified as underweight (< 18.5), normal ($18.5 - 24.99$), overweight ($25-29.99$) and obese (> 30). Waist circumference was measured at the approximate midpoint between the lower margin of the last palpable rib and the top of the iliac crest according to the WHO STEPS protocol (32). Hip circumference was measured around the widest portion of the buttocks (32). Body adiposity index (BAI) was calculated as:

$\text{Hip circumference} / (\text{Height} \times \sqrt{\text{Height}}) - 18$

COPD exacerbations and duration of hospital stay

Data were gathered for number of severe exacerbations (hospitalizations) and moderate exacerbations (antibiotic or/and systemic steroid treatment without hospitalization due to worsening of pulmonary symptoms) (1) in the previous year. The duration of the current hospital stay was recorded.

127

128 **Quality of life**

129 Quality of life was assessed with the mMRC scale and CAT questionnaire. Patients were
130 instructed that there were no right or wrong answers. All patients' questions were answered. Patients
131 were classified according to GOLD as having less symptoms (CAT <10) and breathlessness (mMRC
132 grade 0-1) and more symptoms (CAT ≥10) and breathlessness (mMRC grade ≥2). Because all patients
133 were hospitalized due to exacerbation there were only group C (high risk, less symptoms) and group D
134 (high risk, more symptoms) patients according to GOLD (DM1).

135

136 **Pulmonary Function Testing**

137 The spirometry was performed using Minispir® New spirometer (MIR - Medical International
138 Research, Italy). Patients were instructed to withdraw using short-acting β₂-agonists at least 6 hours,
139 long-acting β₂-agonist at least 12 hours, long acting muscarinic antagonist 24 hours and short acting
140 muscarinic antagonist 12 hours before the spirometry (33). Post bronchodilator spirometry testing was
141 performed 15-30 min after inhalation of 400mcg Salbutamol according to ERS/ATS recommendations
142 (33). Pre- and post- values were obtained for: FVC, FEV₁, FEV₁/FVC, FEV₆, FEV₁/FEV₆, PEF,
143 FEF₂₅₇₅, FEV₃, FEV₃/FVC as well as the difference between post/pre values (delta values). GLI
144 (Global Lungs Initiative) predicted values were used (GLI-2012). Patients' obstruction were classified
145 according to the severity of airflow limitation based on post-bronchodilator FEV₁ as follows: mild
146 (≥80% predicted); moderate (80>FEV₁≥50% predicted); severe (50%>FEV₁≥30% predicted); very
147 severe (<30% predicted) (1).

148

149 **Blood samples and analyses**

150 A venous blood sample was collected from each subject after a 12-hour fasting. Blood samples
151 were taken as late as possible before discharging (usually on 6th or 7th day). Plasma glucose,
152 triglyceride (TG), high density lipoprotein (HDL), low density lipoprotein (LDL), and total cholesterol
153 (tChol) were measured with a Roche COBAS INTEGRA® 400 plus analyzer and an enzymatic
154 colorimetric assay and blood glucose was measured with an enzymatic reference method with
155 hexokinase. Vitamin D was measured with Elecsys 2010 (Roche) and Electro-chemiluminescence
156 immunoassay (ECLIA). Glycated hemoglobin (HbA_{1c}) was measured with a NycoCard device and
157 boronate affinity assay. For patients without established DM a 75g OGTT was performed with blood
158 samples for glucose taken on first and second hour.

159

160 **Statistical Analysis**

161 Statistical analysis was performed with the SPSS for Windows software, version 22.0 (SPSS
162 Inc., Chicago, IL, USA). Continuous variables were presented as mean ± standard deviation and 95
163 Confidence intervals (95%CI) and categorical variables - as percentages. Chi-square test was used to
164 determine the associations between categorical variables. Continuous variables were examined for
165 normality by Shapiro-Wilk test. For normally distributed variables, differences between the groups
166 were determined by independent-samples T test for two samples and analysis of variance (ANOVA)
167 for more than 2 samples. Mann-Whitney U test was used for abnormally distributed variables with 2
168 samples and Kruskal-Wallis test for variables with more than 2 samples. Regression analyses were

used to determine risk factors for presence of DM or the consequences of having DM. Significance value (p-value) was set at 0.05.

All patients signed informed consent. Medical University-Sofia Research Ethics Commission approved the study.

Results and discussion

Sample characteristics

A total of 152 COPD patients admitted for exacerbation were recruited from University Specialized Hospital for Active Treatment of Pulmonary Diseases 'Saint Sofia', Sofia, Bulgaria. Mean age of patients in this study was $65,1 \pm 9,9$ years. 71,1% (108/152) were males, 28,9% (44/152) were females; mean post-bronchodilator FEV₁ was $55,34 \pm 19,5\%$. 15,8% from the patients were never smokers, 57,9% - former smokers and 26,3% - current smokers.

Prevalence of MS

25,0% (38/152) of the patients have MS. 23,1% (25/108) of males have MS vs. 29,5% (13/44) of females but this difference is not statistically significant ($p=0.409$) (table 1). Mean age does not differ between the patients with and without MS ($64,5$ vs $65,3$ years, $p=0.688$).

Table 1. Prevalence of MS according to different factors

| | % MS | P value |
|------------------------------|------|----------|
| All | 25,0 | |
| Sex | | |
| Male | 23,1 | P=0.409 |
| Female | 29,5 | |
| Smoking status | | |
| Never | 25,0 | P=0.678 |
| Former | 27,3 | |
| Current | 20,0 | |
| Arterial hypertension | | |
| Yes | 35,2 | P<0.0005 |
| No | 2,1 | |
| Vitamin D status | | |
| >50 nmol/l | 24,2 | P=0.929 |
| 25-50 nmol/l | 24,6 | |
| <25 nmol/l | 28,0 | |
| DM | | |
| Yes | 37,7 | P=0.008 |
| No | 18,2 | |
| BMI | | |
| Underweight | 0 | P<0.0005 |

| | | |
|------------------------|------|----------------|
| Normal | 0 | |
| Overweight | 27,3 | |
| Obese | 54,8 | |
| BAI | | |
| Underweight | 0 | P=0.001 |
| Normal | 17,5 | |
| Overweight | 28,9 | |
| Obese | 50,0 | |
| Quality of life | | |
| CAT 0-9 | 16,0 | P=0.256 |
| CAT ≥10 | 26,8 | |
| mMRC 0 or 1 | 18,9 | P=0.201 |
| mMRC ≥2 | 28,3 | |
| FEV1 | | |
| FEV1≥50% | 21,3 | P=0.390 |
| FEV1<50% | 27,5 | |
| FEV1≥80% | 11,8 | P=0.852 |
| 80%>FEV1≥50% | 28,8 | |
| 50%>FEV1≥30% | 26,1 | |
| FEV1<30% | 25,0 | |

The prevalence of MS in COPD patients is summarized in table 2. It varies between 21-53% (table 2). This study found comparable but relatively low prevalence of MS compared to previous studies (fig. 1). The prevalence of MS in our study is significantly lower when compared to the general Bulgarian population (44,6% in subjects over 45 years) (22). The odds ratio for COPD patients admitted for exacerbation of having MS is 0.41 compared to general population (95% CI 0.28-0.61, $p<0.0005$).

Table 2. Prevalence of MS in patients with COPD

| Authors | N | Studied population | Prevalence of MS |
|-----------------------|------|---------------------------------|---|
| Akpınar et al. (16) | 133 | Patients with COPD and controls | 44.6% |
| Funakoshi et al. (3) | 7189 | Men aged 45-88 years | 16.8% , OR 0.72 (95% CI 0.51-1.02) in GOLD I; 28.7% , OR 1.33 (95% CI 1.01-1.76) in GOLD II-IV |
| Kupeli et al. (13) | 106 | Hospitalized patients with COPD | 27.3% |
| Lam et al. (4) | 7358 | General population >50 years | 22.6% ; OR 1.47 (95% CI 1.12-1.92) |
| Marquis et al. (17) | 72 | Patients with COPD and controls | 47% |
| Minas et al. (18) | 114 | Men with COPD | 21% |
| Ozgen Alpaydin et al. | 90 | Patients with COPD and controls | 43% |

| | | | |
|---------------------|------|---|--|
| (19) | | | |
| Park et al. (6) | 1215 | Patients with COPD and controls >40 years | 33% vs 22.2% for men; 48.5% vs 29.6% for women OR 2.03 (95% CI 1.08-3.80) |
| Poulain et al. (20) | 28 | Patients with COPD | Overweight – 50% ; Normal weight – 0% . |
| Watz et al. (21) | 200 | Patients with COPD and chronic bronchitis | GOLD I – 50% ; GOLD II – 53% ; GOLD III – 37% ; GOLD IV – 44% ; Chronic bronchitis – 53% |

197

Prevalence of MS in COPD patients




Fig. 1. Prevalence of MS in COPD patients

Two Bulgarian studies examined the prevalence of MS in COPD patients. An epidemiological study conducted in Bulgaria in COPD patients (n=3598) reported a prevalence of 13,8%. These results differ significantly from the literature data probably because of the different criteria for metabolic syndrome (presence of DM, BMI >30 and blood pressure >140/90 mmHg) which makes data comparing irrelevant (23). A more recent study indicates that the prevalence of metabolic syndrome is 41.8% in patients with COPD (n=141) compared to 39% in the control group (n=103). However this study was not conducted using a random sample (exclusion criteria was presence of DM, people were aged 49-79 years for patients with COPD and 35-65 years for controls) (24). However it uses similar criteria for MS and when comparing the results our study finds lower prevalence of MS.

Fulfilled criteria for MS (in all and in MS patients) are shown in table 3. Virtually all (37/38) patients with MS in this study have arterial hypertension, followed by elevated waist circumference (33/38). Arterial hypertension has greatest sensitivity but is not specific. Elevated waist circumference has greatest accuracy. Number of fulfilled criteria in all patients is given in table 4.

Table 3. Fulfilled criteria for MS in all patients and in patients with MS

| MS criteria | All patients (n=152) | MS only (n=38) | Without MS (n=114) | Accuracy |
|---|----------------------|----------------|--------------------|----------|
| Elevated blood pressure: systolic ≥ 130 and/or diastolic ≥ 85 mm Hg (or on therapy) | 69,1% (n=105) | 97,4% (n=37) | 59,6% (n=68) | 54,6% |

| | | | | |
|--|-----------------|-----------------|-----------------|-------|
| Elevated waist circumference >102 cm in males, >88 cm in females | 28,3% (n=43) | 86,8% (n=33) | 8,8% (n=10) | 90,1% |
| Triglycerides >1,7 mmol/L (or on therapy) | 29,6% (n=45) | 60,5% (n=23) | 19,3% (n=22) | 75,7% |
| Fasting glucose >5,5 mmol/L (or on therapy) | 34,2% (n=52) | 65,8% (n=25) | 23,7% (n=27) | 73,7% |
| HDL <1.0 mmol/L in males, <1.3 mmol/L in females (or on therapy) | 15,8% (n=24) | 39,5% (n=15) | 7,9% (n=9) | 78,9% |

Table 4. Number of fulfilled criteria for MS in all patients

| Number of fulfilled criteria | % | n |
|------------------------------|------|----|
| 0 | 13,8 | 21 |
| 1 | 33,6 | 51 |
| 2 | 27,6 | 42 |
| 3 | 13,2 | 20 |
| 4 | 10,5 | 16 |
| 5 | 1,3 | 2 |

The prevalence results could be explained with differences between the populations in different studies (physical activity, diet, lifestyle etc.). For example, Bulgaria is low-income country, which may impact diet preferences and treatment choices. Furthermore, patients in this study had been hospitalized due to exacerbation, which represents the most severe group of COPD patients. Having in mind that MS is more common in the early stages and decreases with COPD progression (21), COPD patients hospitalized for exacerbation may be considered as having advanced COPD.

MS have low prevalence in COPD patients hospitalized due to exacerbation suggesting advanced disease in these patients.

Lifestyle factors

According to NHANES III study smokers are more likely to develop MS than nonsmokers in general population and the risk increases with the number of pack-years even after adjusting for covariates (34). Our study did not find significant differences in prevalence of MS according to smoking status ($p=0.678$) and number of pack-years ($p=0.682$). These results could be explained with smoke being the biggest factor in developing COPD and effect of developing MS could be reduced. Moreover nicotine may be an appetite suppressant and lower the weight thus decreasing prevalence of metabolic syndrome (35). Third, hospitalized COPD patients are patients with predominantly advanced disease and prone to cachexia and wasting. Also lifestyle changes (quit smoking) in the presence of the two diseases should be considered which may change the prevalence of MS.

Treatment with inhalatory corticosteroids (ICS) is not risk factor for MS ($p=0.899$) similar to findings for DM (36). Fasting glucose level are not influenced by ICS ($p=0.837$).

MS is not associated with smoking status and treatment with ICS.

Comorbidity results

COPD is a disease that affects mainly the lungs, but is characterized by systemic inflammation and a number of extrapulmonary manifestations. Only 1/3 of patients with COPD die due to respiratory failure. Main cause of death is lung cancer and cardiovascular complications (37).

The vast majority of patients with COPD have a vitamin D deficiency (38). Aside from its role in the metabolism of calcium and phosphorus, vitamin D is involved in the pathogenesis of multiple diseases, including MS (39), mainly because it affects the secretion and the function of insulin (40). However in our study vitamin D levels do not significantly differ in relation to presence of MS (32,11 vs 31,92 nmol/l, $p=0.860$). Presence of MS is also not related to vitamin D status ($p=0.929$) (table 1).

There are no studies that examine the relationship between osteoporosis and MS in patients with COPD. However, both diseases share common risk factors such as smoking, lack of physical activity, and treatment with corticosteroids. Some of the components of the metabolic syndrome (arterial hypertension, elevated triglycerides, reduced HDL cholesterol) are risk factors for low bone density. Systemic inflammation in MS plays a role in the pathogenesis of osteoporosis (41). On the other hand, studies examining the relationship between MS and osteoporosis showed inconsistent results, probably due to the protective effect of obesity (42,43). However number of fractures in our study does not significantly differ regarding the presence of MS ($p=0.443$). Presence of at least one fracture also does not differ significantly in relation to presence of MS ($p=0.511$).

Most patients with DM have MS, but the opposite is not necessarily true (44). In our study there is a correlation between the presence of MS and DM ($p=0.008$). 52,6% (20/38) from patients with MS have DM. BMI and BAI differ significantly according to presence of MS – 32,51 vs 25,58, $p<0,0005$ for BMI and 31,01 vs 25,58, $p<0,0005$ for BAI. There is also significant difference in prevalence of MS in BMI and BAI groups (table 1). It is notable that none of the patients have MS and BMI <25 (table 1).

Linear regression showed presence of MS as risk factor for higher BMI ($R=0.542$, $r^2=0.293$, $p<0.0005$, $B=6.928$, 95% CI 5.193-8.662) and BAI ($R=0.406$, $r^2=0.165$, $p<0.0005$, $B=5.423$, 95% CI 3.455-7.392).

A logistic regression analysis was conducted to predict presence of MS in a relation to presence of other comorbidities. Presence of DM slightly improves the model (chi square = 6.818, $p=0.009$ with $df=1$). Nagelkerke's R^2 of 0.065 indicates a weak relationship. Odds ratio was 2,73. Vitamin D status does not improve the model ($p>0.05$).

The presence of MS is associated with presence of DM, higher BMI and BAI, but is not associated with vitamin D levels, vitamin D status and number of fractures.

Exacerbations results and duration of hospital stay

Hyperglycemia is associated with elevated glucose concentrations in tissues and bronchial aspirates (45) where it may stimulate infection by enhancing bacterial growth (46) and by promoting bacterial interaction with the airway epithelium (47). Hyperglycemia also impairs both innate and adaptive immunity, suppressing the host response to infection.

The presence of MS in patients with COPD increases the frequency of exacerbations (2.4 vs. 0.7) and their duration – (7.5 vs. 5.0 days) according to Kupeli et al. (13), and 8 versus 5.5 days, according to Abdelghaffar et al. (14).

Our study found a significant difference between the number of total exacerbations according to the presence of MS (table 5, $p=0.015$). The number of severe exacerbation, moderate exacerbation and duration of hospital stay did not reach significance ($p=0.130$, $p=0.188$ and $p=0.553$ respectively).

Table 5. Number of severe, moderate and total exacerbations in previous year and duration of hospital stay according to the presence of MS

| | No MS | MS |
|--|-------|----|
|--|-------|----|

| | | |
|-------------------------|-------------------------|-------------------------|
| Moderate exacerbations | 0,61 (0,49-0,76) | 0,92 (0,59-1,34) |
| Severe exacerbations | 1,79 (1,61-1,97) | 2,08 (1,71-2,50) |
| All exacerbations | 2,40 (2,19-2,61) | 3,00 (2,56-3,52) |
| Hospital stay (in days) | 7,47 (7,24-7,70) | 7,63 (7,28-8,05) |

Triglycerides and blood glucose levels in our study did not correlate with number of exacerbations as reported by other authors (13) (all $p > 0.05$).

Linear regression showed presence of MS as risk factor for higher number of exacerbations ($R=0.207$, $r^2=0.043$, $p=0.010$, $B=0.596$, 95% CI 0.143-1.050). From the MS components presence of arterial hypertension is strongest risk factor for exacerbation ($R=0.228$, $r^2=0.052$, $p=0.005$, $B=0.615$, 95% CI 0.192-1.038).

MS is associated with increased number of exacerbations, but not with the duration of hospital stay.

Quality of life results

The presence of MS is associated with significantly worse cough and sleep (1st and 7th CAT questions; $p=0.002$ and $p=0.001$ respectively) and higher total CAT score ($p=0.017$) (table 6). This confirms the data about reduced quality of life in patients with MS (7). However prevalence of MS is not significantly different between patients with less symptoms (CAT 0-9) and breathlessness (mMRC 0 or 1) compared to patients with more symptoms (CAT ≥ 10) and breathlessness (mMRC ≥ 2) (all $p > 0.05$) (table 1). These mixed results may be explained with COPD having higher negative impact on quality of life than MS (physical limitation due to shortness of breath) like suggested for DM (48) and ameliorating the effect in patients having both diseases.

Table 6. Mean CAT score on every question and in total according to presence of MS

| MS | Mean CAT score | N | P value |
|----------|-----------------|-----|----------------|
| MS – no | CAT1 1,95 | 114 | P=0.002 |
| MS - yes | CAT1 2,63 | 38 | |
| MS – no | CAT2 1,92 | 114 | P=0.063 |
| MS - yes | CAT2 2,34 | 38 | |
| MS – no | CAT3 2,54 | 114 | P=0.092 |
| MS - yes | CAT3 2,97 | 38 | |
| MS – no | CAT4 3,52 | 114 | P=0.361 |
| MS - yes | CAT4 3,74 | 38 | |
| MS – no | CAT5 1,23 | 114 | P=0.198 |
| MS - yes | CAT5 1,66 | 38 | |
| MS – no | CAT6 1,54 | 114 | P=0.695 |
| MS - yes | CAT6 1,68 | 38 | |
| MS – no | CAT7 1,28 | 114 | P=0.001 |
| MS - yes | CAT7 2,21 | 38 | |
| MS – no | CAT8 2,62 | 114 | P=0.068 |
| MS - yes | CAT8 3,08 | 38 | |
| MS – no | Total CAT 16,61 | 114 | P=0.017 |
| MS - yes | Total CAT 20,32 | 38 | |

Regression analyses also showed that MS is a risk factor for reduced quality of life, measured with total CAT score ($R=0.205$, $r^2=0.042$, $p=0.011$, $B=3.711$, 95% CI 0.859-6,562). Presence of MS also impairs cough and sleep – first ($R=0.285$, $r^2=0.081$, $p<0.0005$, $B=0.684$, 95% CI 0.313-1,055) and seventh ($R=0.268$, $r^2=0.072$, $p=0.001$, $B=0.930$, 95% CI 0.390-1,470) CAT questions.

In this study the presence of MS is associated with reduced quality of life, measured with CAT questionnaire.

Pulmonary function test (PFT) results

COPD is characterized by airflow obstruction, which is not fully reversible. MS is associated with a reduction of lung volumes (8-12). It should be noted that some studies found no association between lung function and the presence of MS (49).

Our study did not find differences in FVC, FEV1, FEV1/FVC, FEV6, FEV1/FEV6, PEF, FEF2575 and FEV3 according to the presence of MS. It should be noted that there is tendency for FVC and FEV1/FVC ratio ($p=0.094$ and $p=0.091$ respectively). However, because of this there is significant difference in FEV3/FVC ratio ($p=0.033$) (table 7).

Table 7. Mean PFT values according to the presence of MS

| MS | Mean PFT value | N | P value |
|-----|----------------|-----|----------------|
| No | FEV1 55,56% | 114 | P=0.811 |
| Yes | FEV1 54,68% | 38 | |
| No | FVC 80,46% | 114 | P=0.094 |
| Yes | FVC 72,45% | 38 | |
| No | FEV1/FVC 0,53 | 114 | P=0.091 |
| Yes | FEV1/FVC 0,57 | 38 | |
| No | FEV6 73,89% | 114 | P=0.277 |
| Yes | FEV6 68,63% | 38 | |
| No | FEV1/FEV6 0,57 | 114 | P=0.107 |
| Yes | FEV1/FEV6 0,61 | 38 | |
| No | PEF 55,62% | 114 | P=0.735 |
| Yes | PEF 56,66% | 38 | |
| No | FEF2575 38,89% | 114 | P=0.316 |
| Yes | FEF2575 40,95% | 38 | |
| No | FEV3 66,62% | 114 | P=0.601 |
| Yes | FEV3 63,89% | 38 | |
| No | FEV3/FVC 0,81 | 114 | P=0.033 |
| Yes | FEV3/FVC 0,85 | 38 | |

Regression analyses also showed that MS is not a risk factor for reduced pulmonary function (all $p>0.05$). However some of the components of MS are associated with reduced pulmonary function with highest impact of HDL on FVC ($R=0.183$, $r^2=0.033$, $p=0.024$, $B=-8.517$, 95% CI -15.904;-1,130) and FEV1 ($R=0.251$, $r^2=0.063$, $p=0.001$, $B=-10.391$, 95% CI -16.865;-3,918). Fasting glucose is associated with increased FEV1/FVC ratio ($R=0.186$, $r^2=0.035$, $p=0.022$, $B=1.238$, 95% CI 0.183; 2.294) probably because of lowering FVC.

There is no difference in prevalence of MS in patients with FEV1 <50%, when compared to patients with FEV1 >50% ($p=0.390$) or regarding GOLD stage ($p=0.852$) (table 1).

MS in our study is not associated with worsen pulmonary function. Prevalence of MS is not different regarding GOLD stage.

337 **Conclusions**

338 This study finds 25% prevalence of MS in COPD patients admitted for exacerbation which is
339 significantly lower than general population. MS is more prevalent in females, but the gender difference
340 is not statistically significant. In this study most of the patients are former smokers and prevalence of
341 MS does not differ regarding smoking status and treatment with ICS.

342 The presence of MS is associated with presence of DM, higher BMI and BAI, more
343 exacerbations during the previous year and lower quality of life. MS is not associated with increased
344 hospital stay and lower pulmonary function.

345 This study finds comparable but relatively low prevalence of MS compared to previously
346 published data (21-53%). MS may impact natural course of COPD. Having in mind that MS is more
347 common in the early stages and decreases with COPD progression, the COPD patients admitted for
348 exacerbation may be considered as having advanced COPD.
349

350 References:

- 351 1. From the Global Strategy for the Diagnosis, Management and Prevention of COPD, Global Initiative for
352 Chronic Obstructive Lung Disease (GOLD) 2014. Available from: <http://www.goldcopd.org/>.
- 353 2. Mekov E, Slavova Y. Diabetes mellitus and metabolic syndrome in COPD – part 1: introduction and
354 epidemiology. *Thoracic medicine* 2013;5(4):6-18.
- 355 3. Funakoshi Y, Omori H, Mihara S, et al. Association between airflow obstruction and the metabolic
356 syndrome or its components in Japanese men. *Intern Med* 2010;49:2093–9.
- 357 4. Lam KB, Jordan RE, Jiang CQ, et al. Airflow obstruction and metabolic syndrome: the Guangzhou Biobank
358 Cohort Study. *Eur Respir J* 2010;35:317–23.
- 359 5. Marquis K, Debigare R, Lacasse Y, et al. Midthigh muscle cross-sectional area is a better predictor of
360 mortality than body mass index in patients with chronic obstructive pulmonary disease. *Am J Respir Crit*
361 *Care Med* 2002;166:809–13.
- 362 6. Park BH, Park MS, Chang J, et al. Chronic obstructive pulmonary disease and metabolic syndrome: a
363 nationwide survey in Korea. *Int J Tuberc Lung Dis* 2012;16:694–700.
- 364 7. Ford E, Li C. Metabolic Syndrome and Health-Related Quality of Life among U.S. Adults. *Annals of*
365 *epidemiology* 2008;18(3):165-71
- 366 8. Fimognari FL, Pasqualetti P, Moro L, et al. The association between metabolic syndrome and restrictive
367 ventilatory dysfunction in older persons. *J Gerontol A Biol Sci Med Sci* 2007;62:760–5.
- 368 9. Leone N, Courbon D, Thomas F, et al. Lung function impairment and metabolic syndrome: the critical role
369 of abdominal obesity. *Am J Respir Crit Care Med* 2009;179:509–16.
- 370 10. Lin WY, Yao CA, Wang HC, Huang KC. Impaired lung function is associated with obesity and metabolic
371 syndrome in adults. *Obesity (Silver Spring)* 2006;14:1654–61.
- 372 11. Nakajima K, Kubouchi Y, Muneyuki T, Ebata M, Eguchi S, Munakata H. A possible association between
373 suspected restrictive pattern as assessed by ordinary pulmonary function test and the metabolic syndrome.
374 *Chest* 2008;134:712–8.
- 375 12. Yeh F, Dixon AE, Marion S, et al. Obesity in adults is associated with reduced lung function in metabolic
376 syndrome and diabetes: the Strong Heart Study. *Diabetes Care* 2011;34:2306–13.
- 377 13. Kupeli E, Ulubay G, Ulasli SS, et al. Metabolic syndrome is associated with increased risk of acute
378 exacerbation of COPD: a preliminary study. *Endocrine* 2010;38:76–82.
- 379 14. Abdelghaffar H, Tangour E, Fenniche S et al. Relation between metabolic syndrome and acute exacerbation
380 of COPD. *Eur Respir J* 2012;40(Suppl. 56):886s.
- 381 15. Mekov E, Slavova Y. Diabetes mellitus and metabolic syndrome in COPD – part 3: consequences. *Thoracic*
382 *medicine* 2014;6(4):23-36.
- 383 16. Borissova A-M, Kovatcheva R, Shinkov A et al. Prevalence and features of the metabolic syndrome in
384 unselected Bulgarian population. *Endocrinologia* 2007;12(2):68-77.
- 385 17. Pavlov P, Ivanov Y, Glogovska P, Popova T, Borissova E, Hristova P. Metabolic syndrome and COPD. *Eur*
386 *Respir J* 2010;36:Suppl. 54,66s.
- 387 18. Stratev V, Petev J, Galcheva S, Peneva M. Chronic inflammation and metabolic syndrome (MS) in patients
388 with chronic obstructive pulmonary disease (COPD). *Thoracic Medicine* 2012;4(3):50-7.
- 389 19. Akpinar EE, Akpinar S, Ertek S, Sayin E, Gulhan M. Systemic inflammation and metabolic syndrome in
390 stable COPD patients. *Tuberk Toraks* 2012;60(3):230-7.
- 391 20. Marquis K, Maltais F, Duguay V, et al. The metabolic syndrome in patients with chronic obstructive
392 pulmonary disease. *J Cardiopulm Rehabil* 2005;25:226-32.
- 393 21. Minas M, Kostikas K, Papaioannou AI, et al. The association of metabolic syndrome with adipose tissue
394 hormones and insulin resistance in patients with COPD without co-morbidities. *COPD* 2011;8:414–20.
- 395 22. Ozgen Alpaydin A, Konyar Arslan I, Serter S, et al. Metabolic syndrome and carotid intima-media thickness
396 in chronic obstructive pulmonary disease. *Multidiscip Respir Med* 2013;8(1):61.
- 397 23. Poulain M, Doucet M, Drapeau V, et al. Metabolic and inflammatory profile in obese patients with chronic
398 obstructive pulmonary disease. *Chron Respir Dis* 2008;5:35–41.

- 399 24. Watz H, Waschki B, Kirsten A, et al. The metabolic syndrome in patients with chronic bronchitis and
400 COPD: frequency and associated consequences for systemic inflammation and physical inactivity. *Chest*
401 2009;136:1039–46.
- 402 25. Vestbo J. COPD: definition and phenotypes. *Clin Chest Med* 2014;35(1):1-6.
- 403 26. Alberti KGMM, Eckel RH, Grundy SM, et al. Harmonizing the metabolic syndrome: a joint interim
404 statement of the International Diabetes Federation Task Force on Epidemiology and Prevention; National
405 Heart, Lung, and Blood Institute; American Heart Association; World Heart Federation; International
406 Atherosclerosis Society; and International Association for the Study of Obesity. *Circulation* 2009;120:1640–
407 5.
- 408 27. American Diabetes Association. Diagnosis and Classification of Diabetes Mellitus. *Diabetes Care*
409 2012;35:S64-S71
- 410 28. Borissova A-M, Shinkov A, Vlahov J et al. Determination of the optimal level of 25(OH)D in the Bulgarian
411 population. *Endocrinologia* 2012;17(3):135-42.
- 412 29. Institute of Medicine. 2011. Dietary reference intakes for calcium and vitamin D. Washington DC: The
413 National Academies Press.
- 414 30. Pickering TG, Hall JE, Appel LJ, Falkner BE, Graves J, Hill MN, et al.; Subcommittee of Professional and
415 Public Education of the American Heart Association Council on High Blood Pressure Research.
416 Recommendations for blood pressure measurement in humans and experimental animals. Part 1: blood
417 pressure measurement in humans. *Hypertension* 2005;45:142–61.
- 418 31. US Centers for Disease Control and Prevention (2010). Health behaviors of adults: United States, 2005-
419 2007. Vital and Health Statistics, Series 10, Number 245, Appendix II, p. 80.
- 420 32. WHO. WHO STEPwise approach to surveillance (STEPS). Geneva, World Health Organization (WHO),
421 2008b.
- 422 33. Miller M.R, Hankinson J, Brusasco V. et al. Standardisation of spirometry. *Eur Respir J* 2005; 26: 319–338.
- 423 34. Park Y-W, Zhu S, Palaniappan L, Heshka S, Carnethon M, Heymsfield S. The Metabolic Syndrome
424 Prevalence and Associated Risk Factor Findings in the US Population From the Third National Health and
425 Nutrition Examination Survey, 1988-1994 *Arch Intern Med* 2003;163(4):427-436.
- 426 35. Chiolerio, A; Faeh, D; Paccaud, F; Cornuz, J. Consequences of smoking for body weight, body fat
427 distribution, and insulin resistance. *The American journal of clinical nutrition* 2008;87(4):801–9.
- 428 36. O'Byrne PM, Rennard S, Gerstein H et al. Risk of new onset diabetes mellitus in patients with asthma or
429 COPD taking inhaled corticosteroids. *Respir Med*. 2012;106(11):1487-93.
- 430 37. Calverley PM, Anderson JA, Celli B, et al. Salmeterol and fluticasone propionate and survival in chronic
431 obstructive pulmonary disease. *N Engl J Med* 2007;356:775–89.
- 432 38. Romme EA, Rutten EP, Smeenk FW, et al. Vitamin D status is associated with bone mineral density and
433 functional exercise capacity in patients with chronic obstructive pulmonary disease. *Ann Med*
434 2013;45(1):91-6.
- 435 39. Ju S, Jeong H, Kim H. Blood vitamin D status and metabolic syndrome in the general adult population: a
436 dose-response meta-analysis. *J Clin Endocrinol Metab* 2014;99(3):1053-63.
- 437 40. Mitri J, Muraru MD, Pittas AG. Vitamin D and type 2 diabetes: a systematic review. *Eur J Clin Nutr*
438 2011;65:1005-15.
- 439 41. McFarlane SI. Bone metabolism and the cardiometabolic syndrome: pathophysiologic insights. *J*
440 *Cardiometab Syndr* 2006;1:53–7.
- 441 42. Muhlen D, Safii S, Jassal S, Svarthberg J, Barret-Connor E. Associations between the Metabolic Syndrome
442 and Bone Health in Older Men and Women: The Rancho Bernardo Study. *Osteoporosis International*
443 2007;18(10):1337-44.
- 444 43. Zhou J, Zhang Q, Yuan X, et al. Association between metabolic syndrome and osteoporosis: a meta-
445 analysis. *Bone* 2013;57(1):30-5.
- 446 44. Ginsberg HN, Stalenhoef AF. The metabolic syndrome: targeting dyslipidaemia to reduce coronary risk. *J*
447 *Cardiovasc Risk*. 2003;10:121-128.
- 448 45. Philips BJ, Redman J, Brennan A, et al. Glucose in bronchial aspirates increases the risk of respiratory
449 MRSA in intubated patients. *Thorax* 2005;60:761–4.

- 450 46. Brennan AL, Gyi KM, Wood DM, et al. Airway glucose concentrations and effect on growth of respiratory
451 pathogens in cystic fibrosis. *J Cyst Fibros* 2007;6:101–9.
- 452 47. Brennan AL, Baines DL, Woollhead A, et al. Development of an in vitro model to investigate the effect of
453 glucose on the interaction between respiratory epithelia and bacterial pathogens. *Thorax* 2006;61(Suppl
454 II):ii101.
- 455 48. Arne M, Janson C, Janson S. Physical activity and quality of life in subjects with chronic disease: chronic
456 obstructive pulmonary disease compared with rheumatoid arthritis and diabetes mellitus. *Scand J Prim*
457 *Health Care*. 2009;27(3):141-7.
- 458 49. Yamamoto Y, Oya J, Nakagami T, Uchigata Y. Association Between Lung Function and Metabolic
459 Syndrome Independent of Insulin in Japanese Men and Women. *Jpn Clin Med*.2014; 5:1–8.