

A peer-reviewed version of this preprint was published in PeerJ on 1 October 2015.

[View the peer-reviewed version](https://doi.org/10.7717/peerj.1264) (peerj.com/articles/1264), which is the preferred citable publication unless you specifically need to cite this preprint.

Waqas A, Ahmad W, Haddad M, Taggart FM, Muhammad Z, Bukhari MH, Sami SA, Batool SM, Najeeb F, Hanif A, Rizvi ZA, Ejaz S. 2015. Measuring the well-being of health care professionals in the Punjab: a psychometric evaluation of the Warwick-Edinburgh Mental Well-being Scale in a Pakistani population. PeerJ 3:e1264
<https://doi.org/10.7717/peerj.1264>

Measuring the well-being of health care professionals in the Punjab: a psychometric evaluation of the Warwick-Edinburgh Mental Well-being Scale in a Pakistani population

Background: There is growing awareness of the public health importance of the mental well-being of population in general. This study was carried out to examine the wellbeing of Pakistani healthcare professionals, and to evaluate the psychometric properties of Warwick-Edinburgh Mental Well-being Scale in in this population. The well-being of health professionals is likely to influence their care delivery. **Methods:** A cross-sectional survey was carried out from June, 2013 to December, 2014 among 1271 Pakistani health personnel (HCPs) belonging to seven different cities of Punjab province, Pakistan, to examine the acceptability, internal consistency, test-retest reliability and the content and construct validity of the English version of the Warwick-Edinburgh Mental Well-being Scale in a Pakistani population sample. All data were analyzed in SPSS v.21. **Results:** Our analysis demonstrated a unidimensional construct, a high internal consistency (0.89) and test-retest reliability, good convergent validity and easy readability of WEMBS among Pakistani HCPs. The mean on WEMWBS score was 48.1 (9.4) which is lower than general population sample in other nations. Male healthcare professionals scored significantly higher on WEMWBS than their female counterparts ($P < .05$), and older respondents had higher scores. **Conclusion:** The WEMBS appears feasible and acceptable for use in the Pakistani healthcare professionals, and findings from this study indicate its validity and internal consistency. The well-being scores of the health professional respondents were lower than those identified in general population surveys in the UK.

Measuring the well-being of health care professionals in the Punjab: a psychometric evaluation of the Warwick-Edinburgh Mental Well-being Scale in a Pakistani population

Authors: Ahmed Waqas¹, Waqas Ahmad¹, Mark Haddad², Frances Taggart³, Zerwah Muhammad¹, Muhammad Hamza Bukhari¹, Shahzad Ahmed Sami¹, Sayyeda Mehak Batool¹, Fiza Najeeb¹, Ayesha Hanif¹, Zehra Ali Rizvi¹, Sumbul Ejaz¹

Affiliation: ¹CMH Lahore Medical College and Institute of Dentistry, Lahore Cantt. Pakistan

²Centre for Mental Health Research, School of Health Sciences, City University London, Northampton Square, London EC1V 0HB, United Kingdom

³Statistics and Epidemiology Unit, Division of Health Sciences, University of Warwick Medical School, Coventry, United Kingdom

Corresponding author: Ahmed Waqas

Funding support: none

Conflict of interest: none

Manuscript word count: 3490 (Excluding references)

Abstract word count:

Keywords: mental fatigue, burnout, Warwick-Edinburgh Mental Well-being Scale, WEMWBS, Health personnel, Mental health

Abstract

Background:

There is growing awareness of the public health importance of the mental well-being of population in general. This study was carried out to examine the wellbeing of Pakistani healthcare professionals, and to evaluate the psychometric properties of Warwick-Edinburgh Mental Well-being Scale in in this population. The well-being of health professionals is likely to influence their care delivery.

Methods:

A cross-sectional survey was carried out from June, 2013 to December, 2014 among 1271 Pakistani health personnel (HCPs) belonging to seven different cities of Punjab province, Pakistan, to examine the acceptability, internal consistency, test-retest reliability and the content and construct validity of the English version of the Warwick-Edinburgh Mental Well-being Scale in a Pakistani population sample. All data were analyzed in SPSS v.21.

Results:

Our analysis demonstrated a unidimensional construct, a high internal consistency (0.89) and test-retest reliability, good convergent validity and easy readability of WEMBS among Pakistani HCPs. The mean on WEMWBS score was 48.1 (9.4) which is lower than general population sample in other nations. Male healthcare professionals scored significantly higher on WEMWBS than their female counterparts ($P < .05$), and older respondents had higher scores.

Conclusion:

The WEMBS appears feasible and acceptable for use in the Pakistani healthcare professionals, and findings from this study indicate its validity and internal consistency. The well-being scores of the health professional respondents were lower than those identified in general population surveys in the UK.

Introduction:

Subjective or psychological well-being is increasingly recognized as a crucial aspect of overall health that highlights the importance of *positive* mental health. It relates to the classic World Health Organization (WHO) definition of health as more than just the absence of disease; “Health is a state of complete physical, mental and social well-being and not merely the absence of disease or infirmity” [1]. The importance of mental health within overall well-being is emphasized in international policy as the 'foundation for well-being and effective functioning for both the individual and the community' (which) allows individuals to realise their abilities, cope with the normal stresses of life, work productively and fruitfully, and make a contribution to their community.'[2,3]

Subjective well-being has a long tradition as a key part of the conceptualisation of quality of life [3] and its measurement has been embedded within this construct, with measures of quality of life and of health related quality of life commonly incorporating questions concerning positive

and negative affect [4,5] as well as life satisfaction and considerations of fulfilment and purpose, as for instance within the triennial European Quality of Life Survey [6].

However, over the past two decades, interest in measuring subjective well-being has expanded, with the emergence of evidence that it can be measured in ways that are valid and reliable [3], and here is now a widespread acknowledgement that focusing on the measurement of subjective well-being is an essential part of measuring quality of life that merits particular measures and approaches [7].

This focus on the positive elements of mental health and functioning is augmenting a previous emphasis on deficits, problems and symptoms. This is evident within psychiatry research, mental health policy and clinical practice [8] where the concepts of recovery and the increasing use of wellbeing measures are increasingly commonly used [9]. Within broader public health and policy it relates to an acknowledgment that promoting wellbeing and independence are essential to preventing ill health, which is of particular relevance in light of the growing burden of long-term conditions [10].

Within health services, there is a recognition that staff wellbeing is relevant to the quality and safety of healthcare delivery [11] as well as to efforts and expenditure to recruit and retain clinical staff [12]

The healthcare sector rates highly on important potential causes of work stress [13] with health professionals (in particular nurses) among the occupations reporting the highest rates of work-related stress, depression or anxiety (three-year averages) [14]. An extensive literature has

explored stress among health professionals, using measures of psychological distress as well as more conceptually developed and specific tools such as burnout instruments [15]. Stressors that may be evident within health professionals' work include working long and unsocial hours, prolonged and cumulative involvement with seriously ill patients and their relatives, and exposure to organisational conflicts. Even during their training, HCPs are exposed to a plethora of academic, psychosocial and health-related stressors [16]. Medical students and junior doctors have been reported to be at particular risk, but estimates of prevalence vary widely, from 18% to 82% [17].

Studies among healthcare professionals in Pakistan have reported a high prevalence of anxiety, depression, high stress levels and burn-out, highlighting the need to address the issues of poor job satisfaction, sleep deprivation due to shift patterns, patient overload, authoritarian work relationships and limited peer and social support [18–20].

In line with the growing emphasis on measures of positive health rather than a focus on deficits and problems, it was considered useful to measure the subjective mental well-being of health care professionals in Pakistan. The most widely known measure of mental wellbeing in the UK is the Warwick Edinburgh Mental Well-being Scale (WEMWBS). WEMWBS is a measure of mental wellbeing, developed by the Universities of Warwick and Edinburgh, and funded by NHS Scotland.

The current study was devised to examine the psychometric characteristics of the WEMWBS measure among English-speaking Pakistani health professionals. This is useful because there is a

paucity of psychological and psychiatric rating scales that have been cross-culturally validated for use within Pakistan population [21], and the mental health scales that have been evaluated for use in this population are largely measures of symptoms of mental disorder rather than of positive mental health.

.

Materials and methods:

Questionnaire and Instrument:

The questionnaire consisted of two sections. The first section recorded information related to the demographics and profession of the respondents, and the second section assessed mental well-being levels with WEMWBS.

The Warwick-Edinburgh Mental Well-being Scale (WEMWBS) was developed and extensively validated in the UK general population [22]. It is a self-administered questionnaire and comprises 14 positively worded items assessing eudaimonic (focused on psychological functioning and a sense of meaning and purpose in life), and hedonic (focused on happiness, contentment and life satisfaction) constructs of mental wellbeing. Responses are recorded on a 5 point Likert scale ranging from “none of the time” to “all the time”. For analysis, scores ranging from 14-70 are obtained by summing all of the items. There is no reverse coding. The overall score is the sum of all items, with higher scores indicating greater well-being, varying from 14 (low well-being) to 70 (high well-being).

Use in the UK, and in other nations [23] reveal the scale has good psychometric properties, and is feasible, reliable, and sensitive to change. Chronbach’s α for the scale lies between 0.87 and

0.91 and the one-week test-retest reliability at $r = 0.83$. Validation studies have been undertaken in different populations in the UK such as Scottish undergraduate and postgraduate students and general population samples [22], English and Scottish teenagers [24], as well as Pakistani and Chinese ethnic minorities in the UK [25]. It has also demonstrated high internal consistency, good content, face validity, normal distribution and no floor and ceiling effects in previous validation studies [22,24,25].

138

139 **Pilot Survey:**

A pilot study was conducted in which a convenience sample of 50 healthcare professionals including doctors, nurses and dentists in the Combined Military Hospitals, Lahore, completed and commented on the WEMWBS to assess if the original English language version WEMWBS could be easily understood and completed. We received positive comments from the participants that WEMBS was very easy to understand. Therefore, we did not feel the need to translate this instrument to Urdu language.

146 **Sample size calculation:**

In general terms sample size calculation is based on variability in the sample and the size of effects expected. Usually both of these are unknown before starting the study and figures are estimated from previous studies or sometimes “rules of thumb” based on previous experience are used. Most of the studies on validation of questionnaires in social sciences use 5-10 respondents per questionnaire item for factor analyses. Factor analyses are subject to the strength of communalities and factor loading and these statistical tests should instead be based on large samples for reliable results [26]. Comrey and Lee recommend at least 500 cases and according to

154 their rating scale a sample size of 1000 or more is considered as excellent for factor analytic
155 studies [27]. Therefore, we aimed for a sample size of at least 1000 HCPs.

Procedure:

157 This cross-sectional survey was carried out from June, 2013 to December, 2014. The study
158 sample consisted of practicing doctors, nurses, physiotherapists, pharmacists and dentists
159 practicing either in hospital based or private settings in various cities of the province of Punjab,
160 Pakistan. Seven districts; Lahore, Faisalabad, Gujrat, Multan, Rawalpindi, Islamabad and
161 Sheikhupura were included in the survey for a study sample representative of the Punjabi
162 population. However, due to resources limitations, we could not conduct a systematic random
163 sample. Data were collected by convenience sampling approach. Sixteen medical students
164 (currently final year MBBS) approached HCPs in all available clinical settings in the Punjab
165 province. An independent sample of 50 HCPs from CMH Lahore participated in test-retest
166 assessment of WEMWBS and completed the questionnaire twice with a test-retest interval of 7
167 days.

168 Ethical approval was obtained from the Ethical Review committee of CMH Lahore Medical
169 College, Lahore Cantt, Pakistan. Written informed consent was provided by all the study
170 participants. The respondents were ensured anonymity and informed that only group level
171 findings would be reported.

Statistical analysis:

173 All data were analyzed in SPSS v. 21 (IBM Chicago, IL, USA). Frequencies and descriptive
174 statistics were calculated for demographic variables and total scores on WEMWBS. Histograms
175 and QQ-plots were visualized to assess the assumption of normality for WEMWBS scores and

176 floor and ceiling effects in the response distribution of WEMWBS. Factor analysis with principal
177 component analysis (PCA) and Quartimax rotation method was done to evaluate the construct of
178 WEMWBS. Prior to performing principal component analysis (PCA), its suitability was assessed
179 with following criteria: correlation coefficient > 0.3 for all variables, an overall Kaiser-Meyer-
180 Olkin (KMO) greater than 0.6 and a statistically significant Barlett's Test of Sphericity test ($P <$
181 $.05$). The number of components to retain were assessed with Cartell's scree plot, eigen values $>$
182 1, interpretability criterion, the amount of variance explained and reliability analysis.

183 Only those statements were included that had a factor loading > 0.3 . Cronbach's alpha
184 coefficient was calculated to assess the internal consistency of WEMWBS in our study sample
185 and a value of 0.70 - 0.9 was considered acceptable [28]. Item-total correlations were used to
186 assess the convergent validity of the questionnaire. Correlations were calculated using Pearson's
187 product moment correlation coefficient, and values that were substantial but not excessive
188 (greater than or equal to 0.2 and less than 0.8) were sought [28]. Test-retest reliability was
189 evaluated by the intra-class correlation coefficient using a two factor mixed effects model and
190 type consistency.

191 Floor and ceiling effects were assessed by examining response patterns for each of the factors
192 derived from factor analysis. Scores were graphed as a histogram and the distribution of scores
193 inspected; the percentage of individuals with the lowest and highest possible score in each of the
194 factors was recorded, and values greater than 20% were considered as floor and ceiling effects.

195 Association between participant characteristics and WEMWBS scores were examined with
196 independent sample t-tests. Although focus groups have been used to evaluate understanding and
197 meaning to participants [25] but in previous studies on WEMBWS, its readability had not been

assessed. Therefore, we used the Flesch Reading Ease score and Flesch-Kincaid Grade level functions to assess the readability of the questionnaire [29].

Results:

Response rate:

Total response rate was 90.7% (1271/1400).

Demographics:

1271 individuals responded to the questionnaire. As noted in Table 1, participants' ages ranged between 20 and 68 years, with mean age (SD) 31.8 (9.4). Male participants were older, 34.1 (10.3) than females, 30.0 (8.1). Most of the respondents were doctors (n=840, 66%) or nurses (n=218, 17.2%), female (n=720, 57%), and practicing in Lahore (n= 70%).

There were some differences in respondent age and gender in relation to profession: nurses were almost all (97%) female, whereas among doctors and dentists there was near equal gender representation. 42 of the pharmacists were male (79%), as were 64% of the physiotherapists. There were no significant age differences between the professional groups, except for the dentists in the sample, who were younger than the other professions - mean age 28.5 (6.6); $p < 0.001$)

Respondent characteristics are presented in Table 1.

Descriptive Statistics and floor or ceiling effects:

Mean (SD) and median scores on WEMBWS were 48.1 (9.4) and 48 respectively. The assessment of item response frequencies from respondents showed little evidence of highly skewed distributions: scores exhibited mild skewness of -.31 (.07) and kurtosis of .42 (.14). Visual inspection of histogram and QQ plots did not reveal any significant deviation of response distribution from normality. All response categories were used by at least one person for all the scale items. Mean values for individual items ranged from 2.88 (1.1) for “I’ve had energy to spare” to 3.74 (.99) for “I’ve been feeling confident” (Table 2). A histogram representing distribution of mean scores of participants on WEMWBS is given in Figure 1. Based on these results, it can be concluded that WEMWBS validated for Pakistani population is free from any floor or ceiling effects.

As in the UK validation study (Tenant et al, 2007), mean WEMBWS scores were significantly higher for men (49.16) than for women (47.30) ($t = 3.486$, $df = 1269$, $p = 0.001$) (See Table 1). There were differences in well-being scores between the professional groups, with doctors’ and nurses’ WEMWBS mean scores lower than those for the other professional groups. The difference between WEMWBS for doctors and nurses combined, compared to the other professional groups combined was statistically significant: $p = 0.002$ (mean difference = 2.22; $t = -3.14$, $df = 1269$). A difference in wellbeing score was evident across age group, with participants aged 30 years and older noting higher WEMWBS than the younger participants: ($p = 0.013$, mean difference = 1.33, $t = 2.48$, $df = 1269$).

Items 2, 10, 11 and 12 were the most positively endorsed items by both male and female respondents; whereas, items 3, 4 and 5 were least endorsed by the HCPs (both males and females).

241

242

243 **Factor analysis with Principal Component Analysis (PCA):**

244 A principal components analysis (PCA) was run on WEBWS responses. Inspection of the
 245 correlation matrix showed that all variables had at least one correlation coefficient greater than
 246 0.3. The overall Kaiser-Meyer-Olkin (KMO) measure was 0.83 with individual KMO measures
 247 all greater than 0.7, classifications of 'middling' to 'meritorious' according to Kaiser (1974).
 248 Bartlett's Test of Sphericity was statistically significant ($p < .0005$), indicating that the data was
 249 suitable for factor analysis. PCA revealed two components that had eigenvalues greater than one
 250 and which explained 42.11% and 8.5% of the total variance, respectively. Eigen values, visual
 251 inspection of the Catell's scree plot, interpretability criterion and Cronbach's alpha value,
 252 indicated that one component should be retained. A single-component solution explained 42.1%
 253 of the total variance. Total score was obtained by summing all of the items. Factor loadings of
 254 the rotated component solution for 14-item WEMWBS based on eigen values greater than 1 are
 255 presented in Table 3. The Scree plot is given as Figure 2.

256 For interest of readers, exploratory factor analysis with principal axis factoring method have
 257 been provided as a supplementary file Table S1.

258 **Internal Consistency and content validity:**

259 The WEMBS consists of 14 items. The Cronbach's alpha value obtained for one factor structure
 260 of WEMBS was found to be .89. Corrected item correlations of all the items were greater than
 261 0.3, thus, exhibiting the same construct of all the statements of WEMWBS. Item-total statistics
 262 for all items of WEMWBS are given in Table 4. For further assessment of internal consistency,

item-total score correlations, adjusted for overlap, were calculated for each item and ranged between 0.46 and 0.68, that is substantial but not excessive values. The item-total statistics for all items of WEMWBS are given in Table 4.

Test-retest reliability:

The test-retest reliability evaluated by intraclass correlation coefficient was .921 with a 95 % confidence interval from .861-.955, thus, indicating a very high degree of reliability.

Readability:

The values for Flesch reading ease and FK grade level score were calculated on Microsoft Word 2013. Flesch reading ease score for 14-item WEMWBS and FK grade level score 73.5 and 4.4 respectively which indicates the easy readability and understanding level of WEMWBS.

Discussion:

The present study confirms one factor construct, a high internal consistency, convergent validity, high readability of WEMWBS in the Pakistani population.

It also indicates no floor and ceiling effects exist in response distribution of WEMWBS for Pakistani population. Our study sample was relatively large and consisted of a diverse range of healthcare professionals -physicians, surgeons, general practitioners, nurses, pharmacists, dentists and physiotherapists - which is an important strength of this study.

The mean (SD) well-being score of HCP respondents in this study was 48.1 (9.4), which is lower than values obtained in general population surveys in the UK. The Health Survey for England identified a mean WEMWBS score for adults of 52.3 [30]; whilst the mean score found in a

284 recent population survey for Scotland was 50.0 [31]. The median score was 48, compared to 51 in
285 the initial validation study [22] and 50 in a large-scale population sample in Northern Ireland
286 [32].

287 We also identified gender differences in well-being scores, as well as some differences between
288 professions and age groups. In this study, male HCPs scored higher on WEMWBS than females,
289 which is consistent with the findings of UK population studies [30,31]

290 Items 3, 4 and 5 were least endorsed by the HCPs in our study. According to Taggart et al
291 (2013), item 4 (I've been feeling interested in other people) was misinterpreted by Chinese and
292 Pakistani groups in the UK who took it in a sexual context [25], which might be a potential
293 reason for low positive response to this item in our study.

294 Similar to other studies conducted in the general population of the UK, our analysis has
295 confirmed the unidimensional construct of WEMWBS and a high internal consistency [22,24].
296 However, the factor structure was not as clear cut as in the original validation study [22] and
297 other analyses, with some suggestion of two factors; but the extent of cross-loading and the
298 modest additional variance explained by 2 factors/components, together with the lack of
299 explanatory or theoretical underpinning for this structure indicated that a unidimensional model
300 be retained.

301 A study conducted with Pakistani people in the UK also showed a high internal consistency, but,
302 as in the current study, the factor structure was less clear, with three factors evident, but the
303 limited variance explained by the second and third factors supporting a one-factor model. [25]

304
305 There was some correspondence between the item-total correlations identified in our study and
306 in the Taggart et al study of UK English-speaking Pakistanis. The three items with lowest item

total correlations in our study (item 1: ‘I’ve been feeling optimistic about the future’; item 4: ‘I’ve been feeling interested in other people’; and item 5: ‘I’ve had energy to spare’) had lowest correlations in the UK-Pakistani study, though the markedly low value for item found in the UK sample (0.28) was not evident in our sample of Pakistani HCPs (.0.488). In focus groups conducted as part of the UK study it was indicated that a quarter of Pakistanis completing the questionnaire were Pashtun speaking and born abroad and there is no translation for “optimistic” in the Pashto language [25]. However, our sample consisted of Punjabi and Urdu speaking healthcare professionals.

The histogram was slightly negatively skewed and kurtotic, however, no floor and ceiling effects were observed in WEMWBS in consonance with other validation studies. WEMWBS can also prove useful for assessing mental wellbeing of undergraduates due to its easy readability and lucid content as demonstrated by its values for Flesch reading ease and FK grade level score of 4.4. This fact was also supported by favorable comments from the participants in the pilot survey. Based on this data, we can speculate that this scale can be easily used in teenagers, college and university students as well.

Limitations:

This cross-sectional study used a convenience sample of HCPs and so, unlike a systematically derived probability sample, the extent to which the respondents are representative of the target population is uncertain. This is likely to be of lesser importance to the psychometric evaluation than the descriptive results, and is partially compensated for by the relatively large sample size and high response rate; 91% of those requested to complete the measure did so, which limits participation and selection bias.

329 Comparator instruments were not used alongside WEMWBS to enable criterion validity
330 evaluation.

331 Translation and back-translation to develop an Urdu language version (the national language of
332 Pakistan) was not conducted as part of this study, because our sample consisted of HCPs who
333 were all capable of reading and writing in English language. So although the WEMWBS has
334 been shown suitable for assessing mental wellbeing in a diverse group of HCPs, it cannot yet be
335 used for studies within the general population of Pakistan, where the extent of fluency in English
336 is considerably less.

337 WEMWBS assesses hedonic and eudaimonic constructs of mental wellbeing but does not have
338 any items assessing spiritual and religious wellbeing of an individual which predominate in
339 Eastern traditions and culture of Pakistan where religious tendencies have emerged as an
340 important coping mechanism [33,34], emphasized in a study conducted in ethnic minority
341 groups of Malay, Chinese and Indian Muslims and Hindus [35]. Therefore, it is suggested that
342 new generation of psychometric instruments should include items assessing these constructs as
343 well.

344 **Recommendations:**

345 This scale is particularly useful in monitoring mental wellbeing levels of HCPs and medical
346 students working in a highly demanding roles and environments. Further studies are required to
347 develop and validate an Urdu version of this scale in the general community. Alongside general
348 investigations of the range of factors that are associated with mental health and well-being, the
349 WEMWBS may also have value in evaluating efficacy of stress reduction programs such as

mindfulness programs, relaxation exercises, cognitive therapies, and social support and group therapies that have proven useful in academic and hospital based settings [36].

Conclusion:

According to our knowledge, this is the first study validating a scale for assessing mental wellbeing in Pakistan and thus, it is an important addition to the limited arsenal of validated psychometric instruments that exist for the Pakistani population. The WEMWBS has demonstrated excellent psychometric properties in Pakistani HCPs and it appears a valid and reliable tool for use among English-speaking Pakistani people.

Acknowledgement:

The authors thank Professor Dr. Atif Rahman, Professor of Child Psychiatry at University of Liverpool, United Kingdom, for his valuable suggestions to improve this manuscript.

References:

1. Official Records of the World Health Organization. Preamble to the Constitution of the World Health Organization as adopted by the International Health Conference, New York, 19-22 June, 1946; signed on 22 July 1946 by the representatives of 61 States and entered into force on 7 April 1948. 1946.
2. World Health Organisation. WHO: Promoting Mental Health: Concepts, Emerging Evidence, Practice. Geneva. 2004.
3. OECD. OECD Guidelines on Measuring Subjective Well-being. 2013. doi:<http://dx.doi.org/10.1787/9789264191655-en>
4. World Health Organization, Regional Office for Europe, Psychiatric Research Unit. WHO-5 well-being index (WHO, 1998 World Health Organization info package: Mastering depression in primary care. Frederiksborg. 1998.

- 375 5. McHorney C, Ware JJ, Lu J, Sherbourne C. The MOS 36-item Short-Form Health Survey
376 (SF-36): III. Tests of data quality, scaling assumptions, and reliability across diverse
377 patient groups. *Med care.* 1994;32: 40–66.
- 378 6. Eurofound. European quality of life survey [Internet]. 2012 [cited 12 Jun 2015].
379 Available: [http://www.eurofound.europa.eu/surveys/european-quality-of-life-surveys-](http://www.eurofound.europa.eu/surveys/european-quality-of-life-surveys-eqls/european-quality-of-life-survey-2012)
380 [eqls/european-quality-of-life-survey-2012](http://www.eurofound.europa.eu/surveys/european-quality-of-life-surveys-eqls/european-quality-of-life-survey-2012)
- 381 7. Stiglitz J, Sen A, Fitoussi J. Report by the Commission on the Measurement of Economic
382 Performance and Social Progress. 2009.
- 383 8. Slade M. Mental illness and well-being: the central importance of positive psychology and
384 recovery approaches. *BMC Heal Serv Res.* 2010;10: 26.
- 385 9. Warner R. Recovery from schizophrenia and the recovery model. *Curr Opin Psychiatry.*
386 2009; 374–80. doi:10.1097/YCO.0b013e32832c920b
- 387 10. Nolte E, McKee M. Caring for people with chronic conditions. A health system
388 perspective. World Health Organization 2008 on behalf of the European Observatory on
389 Health Systems and Policies. Maidenhead: OUP [Internet]. 2008. Available:
390 http://www.euro.who.int/__data/assets/pdf_file/0006/96468/E91878.pdf
- 391 11. Royal College of Physicians. Work and wellbeing in the NHS: why staff health matters to
392 patient care [Internet]. 2015. Available:
393 https://www.rcplondon.ac.uk/sites/default/files/comms_wwnhs_a4_12pp_final.pdf
- 394 12. Boorman S. NHS health and wellbeing review: interim report. London: Department of
395 Health. 2009.
- 396 13. Eurofound. Fifth European Working Conditions Survey. [Internet]. 2012. Available:
397 www.eurofound.europa.eu/publications/htmlfiles/ef1182.htm. Luxembourg
- 398 14. Health and safety executive. Stress-related and psychological disorders in Great Britain
399 2014 [Internet]. 2015. Available: <http://www.hse.gov.uk/statistics/causdis/stress/>
- 400 15. McManus C. Stress in Health Professionals. [Internet]. Ayers et al., editor. Cambridge
401 Handbook of Psychology, Health and Medicine. Cambridge: CUP; Available:
402 [http://www.ucl.ac.uk/medical-education/reprints/2007-StressInHealthProfessionals-](http://www.ucl.ac.uk/medical-education/reprints/2007-StressInHealthProfessionals-CambridgeHandbook.pdf)
403 [CambridgeHandbook.pdf](http://www.ucl.ac.uk/medical-education/reprints/2007-StressInHealthProfessionals-CambridgeHandbook.pdf)
- 404 16. Waqas A, Khan S, Sharif W, Khalid U, Ali A. Association of academic stress with
405 sleeping difficulties in medical students of a Pakistani medical school: a cross sectional
406 survey. *PeerJ.* 2015;3: e840. doi:10.7717/peerj.840
- 407 17. Prins J, Gazendam-Donofrio S, Tubben B, van der Heijden F, van de Wiel H, Hoekstra-
408 Weebers J. Burnout in medical residents: a review. *Med Ed.* 2007;41: 788–800.

- 409 18. Aslam HD, Mansoor N, Suleman Q. Analysis of Level of Stress among Doctors in Public
410 and Private Hospitals of Pakistan. *Int J Learn Dev*. 2013;3: 109–135.
411 doi:10.5296/ijld.v3i2.6247
- 412 19. Usman R, Nafees F, Farooqui SI, Faruq NM. Level of satisfaction of physiotherapists with
413 regards to their profession. *Pakistan J Rehabil*. 2013;2: 41–47.
- 414 20. Fradelos E, Mpelegrinos S, Mparo C, Vassilopoulou C, Argyrou P, Tsironi M, Zyga S,
415 Theofilou P. Burnout syndrome impacts on quality of life in nursing professionals : The
416 contribution of perceived social support. *Prog Heal Sci*. 2014;4: 102–109.
- 417 21. Ahmer S, Faruqui RA, Aijaz A. Psychiatric rating scales in Urdu: a systematic review.
418 *BMC Psychiatry*. 2007;7: 59. doi:10.1186/1471-244X-7-59
- 419 22. Tennant R, Hiller L, Fishwick R, Platt S, Joseph S, Weich S, Parkinson J, Secker J,
420 Stewart-Brown S. et al. The Warwick-Edinburgh Mental Well-being Scale (WEMWBS):
421 development and UK validation. *Health Qual Life Outcomes*. 2007;5: 63.
422 doi:10.1186/1477-7525-5-63
- 423 23. Castellví P, Forero C, Codony M, Vilagut G, Brugulat P, Medina A, Gabilondo A,
424 Mompart A, Colom J, Tresserras R et al. The Spanish version of the Warwick-Edinburgh
425 mental well-being scale (WEMWBS) is valid for use in the general population. *Qual Life*
426 *Res*. 2014;23: 857–68.
- 427 24. Clarke A, Friede T, Putz R, Ashdown J, Martin S, Blake A, Adi Y, Parkinson J, Flynn P,
428 Platt S, et al. Warwick-Edinburgh Mental Well-being Scale (WEMWBS): validated for
429 teenage school students in England and Scotland. A mixed methods assessment. *BMC*
430 *Public Health*. BioMed Central Ltd; 2011;11: 487. doi:10.1186/1471-2458-11-487
- 431 25. Taggart F, Friede T, Weich S, Clarke A, Johnson M, Stewart-Brown S. Cross cultural
432 evaluation of the Warwick-Edinburgh mental well-being scale (WEMWBS) -a mixed
433 methods study. *Health Qual Life Outcomes*. *Health and Quality of Life Outcomes*;
434 2013;11: 27. doi:10.1186/1477-7525-11-27
- 435 26. MacCallum R, Widaman K, Zhang S, Hong S. Sample size in factor analysis. *Psychol*
436 *Methods*. 1999;4.
- 437 27. Comrey AL, Lee H. A first course in factor analysis. Hillsdale, NJ: Erlbaum. 1992.
- 438 28. Streiner D. Health measurement scales: a practical guide to their development and use. 4th
439 ed. Oxford University Press; 2008.
- 440 29. Flesch R. A new readability yardstick. *J Appl Psychol*. 1948;32: 221–33.

30. Bridges S. Health Survey for England, 2012. The Health and Social Care Information Centre. [Internet]. 2013. Available: <http://www.hscic.gov.uk/catalogue/PUB13218/HSE2012-Ch5-Wellbeing.pdf>
31. Wilson, Kellock C, Adams D, Landsberg J. The Scottish Health Survey: Topic Report – Mental Health and Wellbeing. Published by The Scottish Government. [Internet]. 2015. Available: <http://www.gov.scot/Resource/0046/00469088.pdf>
32. Lloyed K, Devine P. Psychometric properties of the Warwick-Edinburgh mental well-being scale (WEMWBS) in Northern Ireland. *J Ment Heal*. 2012;21: 257–263.
33. Waqas A, Zubair M, Ghulam H, Wajih Ullah M, Zubair Tariq M. Public stigma associated with mental illnesses in Pakistani university students: a cross sectional survey. *PeerJ*. 2014;2: e698. doi:10.7717/peerj.698
34. Voll JO. Conservative and Traditional Brotherhoods. *Ann Am Acad Polit Soc Sci*. 1992;524: 66–78.
35. Vaingankar J, Subramaniam M, Chong S, Abdin E, Orlando E, Picco L, Lim YW, Phua MY, Chua BY, Tee JY, et al. The positive mental health instrument: development and validation of a culturally relevant scale in a multi-ethnic Asian population. *Health Qual Life Outcomes*. 2011;9: 1–18.
36. Shapiro SL, Shapiro DE, Schwartz GE. Stress management in medical education: a review of the literature. *Acad Med*. 2000;75: 748–59. Available: <http://www.ncbi.nlm.nih.gov/pubmed/10926029>

473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493

494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512

Table 1 (on next page)

Demographic characteristics of HCPs participating in the survey (n=1271)

1 Table 1: Demographic characteristics of HCPs participating in the survey (n=1271)

Variable		Frequency (n)	WEMWBS Mean (SD)
Gender	Male	551 (43.4%)	49.16 (9.16)
	Female	720 (56.6%)	47.30 (9.59)
Healthcare Profession	Doctor	840 (66.1%)	47.74 (9.57)
	Nurse	218 (17.2%)	47.74 (9.45)
	Pharmacist	53 (4.2%)	50.28 (8.35)
	Physiotherapist	25 (2%)	49.44 (6.90)
	Dentist	135 (10.6%)	49.93 (9.26)
City	Lahore	893 (70.3%)	
	Multan	86 (6.8%)	
	Sheikhupura	54 (4.2%)	
	Gujrat	58 (4.6%)	
	Rawalpindi	101 (7.9%)	
	Faisalabad	54 (4.2%)	
	Islamabad	25 (2%)	
Age	>= 30	542 (42.6%)	48.87 (9.11)
	< 30	729 (57.4%)	47.54 (9.65)

2

Table 2 (on next page)

Item level statistics for WEMWBS for Pakistani HCPs (n = 1271)

2 **Table 2. Item level statistics for WEMWBS for Pakistani HCPs (n = 1271)**

Statement	Mean	Std. Deviation
I've been feeling optimistic about the future	3.36	1.109
I've been feeling useful	3.74	.991
I've been feeling relaxed	3.03	1.085
I've been feeling interested in other people	2.97	1.140
I've had energy to spare	2.88	1.113
I've been dealing with problems well	3.53	.959
I've been thinking clearly	3.59	1.034
I've been feeling good about myself	3.58	1.052
I've been feeling close to other people	3.33	1.053
I've been feeling confident	3.74	.999
I've been able to make up my own mind about things	3.72	.992
I've been feeling loved	3.64	1.038
I've been interested in new things	3.58	1.111
I've been feeling cheerful	3.42	1.022

3

Table 3 (on next page)

Factor loadings for 14 item WEMWBS in Pakistani health personnel (n = 1271)

Table 3: Factor loadings for 14 item WEMWBS in Pakistani health personnel (n = 1271)

Statements	Factor Loading	
	I	II
I've been feeling confident (Item 10)	.789	
I've been feeling good about myself (Item 8)	.776	
I've been thinking clearly (Item 7)	.744	
I've been able to make up my own mind about things (Item 11)	.722	
I've been dealing with problems well (Item 6)	.703	
I've been feeling cheerful (Item 14)	.686	
I've been feeling close to other people (Item 9)	.670	
I've been feeling useful (Item 2)	.637	
I've been feeling loved (Item 12)	.625	
I've been interested in new things (Item 13)	.578	
I've been feeling optimistic about the future (Item 1)	.541	
I've been feeling interested in other people (Item 4)	.390	.663
I've had energy to spare (Item 5)	.378	.646
I've been feeling relaxed (Item 3)	.508	.528

Extraction Method: Principal Component Analysis.

Rotation Method: Quartimax with Kaiser Normalization.

a. Rotation converged in 3 iterations.

Table 4(on next page)


Item total statistics for 14-item WEMWBS (n = 1271)

Table 4: Item total statistics for 14-item WEMWBS (n = 1271)

	Scale Mean if Item Deleted	Scale Variance if Item Deleted	Corrected Item-Total Correlation	Squared Multiple Correlation	Cronbach's Alpha if Item Deleted
I've been feeling optimistic about the future	44.75	78.422	.488	.293	.887
I've been feeling useful	44.37	78.627	.548	.350	.884
I've been feeling relaxed	45.08	77.650	.544	.355	.884
I've been feeling interested in other people	45.13	78.590	.462	.292	.888
I've had energy to spare	45.23	79.177	.445	.289	.889
I've been dealing with problems well	44.58	77.894	.616	.447	.881
I've been thinking clearly	44.52	76.722	.631	.490	.880
I've been feeling good about myself	44.53	75.597	.684	.525	.878
I've been feeling close to other people	44.78	76.788	.614	.424	.881
I've been feeling confident	44.37	76.938	.644	.518	.880
I've been able to make up my own mind about things	44.39	77.959	.587	.421	.882
I've been feeling loved	44.47	78.118	.547	.348	.884
I've been interested in new things	44.52	77.251	.550	.352	.884
I've been feeling cheerful	44.69	76.463	.656	.492	.879

1

Frequency distribution of total scores of Pakistani HCPs on WEMWBS (n = 1271)



2

Scree plot for 14-item WEMWBS

