

An unusual archosauromorph tooth increases known tetrapod diversity in the lower Chinle Formation (Late Triassic) of southeastern Utah

Lopez, Andres; St. Aude, Isabella; Alderete, David; Alvarez, David; Aultman, Hannah; Busch, Dominique; Bustamante, Rogelio; Cirks, Leah; Lopez, Martin; Moncada, Adriana; Ortega, Elizabeth; Verdugo, Carlos; Gay, Robert J *.

Mission Heights Preparatory High School, 1376 E. Cottonwood Ln., Casa Grande, Arizona 85122

[*rob.gay@leonagroup.com](mailto:rob.gay@leonagroup.com) 520-836-9383

Abstract:

An unusual tetrapod tooth was discovered in the Late Triassic Chinle Formation of southeastern Utah. The tooth was originally thought to belong to *Revueltosaurus* but further investigations have rejected that hypothesis. In this paper we compare MNA V10668 to other known fossil teeth found in the Chinle Formation and identify the least inclusive clade it may belongs to. Using data found in other publications and pictures of other teeth, we compare this specimen to other Triassic dental taxa. MNA V10668 shares some similarities with *Crosbysaurus*, *Tecovasaurus*, and several other named taxa but possesses unique characteristics not found in other diapsid teeth. We conclude that it is most likely an archosauromorph and probably an archosauriform. This increases the known diversity of tetrapods from the Chinle Formation and represents the first tooth morphotype completely unique to Utah in the Late Triassic Period.

1 Introduction:

2 The recovery of vertebrate life from the Permian-Triassic transition resulted in an
3 amazing array of new body forms as life filled ecological voids. This is especially
4 noticeable in the archosaur-line diapsids. Many archosauromorph, archosauriform, and
5 archosaurian reptiles adapted and radiated across the globe, filling or creating
6 numerous niches with novel body forms (Nesbitt et al., 2010) and dietary specializations
7 (Heckert, 2004; Parker et al., 2005). The ecological revolution of the Triassic Period laid
8 the groundwork for dinosaurs (including modern birds), crocodiles, and mammals to
9 dominate terrestrial vertebrate assemblages for the next 200 million years.

10 It is perhaps somewhat surprising then that the terrestrial record of the Late
11 Triassic Period from Utah has not reflected the global diversification of tetrapod clades.
12 Some of this may be attributed to the greater attention that Late Triassic deposits in
13 neighboring Arizona and New Mexico have received (Long and Murray, 1995; Parker et
14 al., 2006). Until recently (Heckert et al., 2006; Gibson, 2013; Martz et al., 2014) the
15 Triassic tetrapod record published from Utah has mainly consisted of the ubiquitous
16 phytosaurs (Morales and Ash, 1993). This is especially true when looking at body fossils
17 only. Even with this recent work, Utah's Triassic tetrapod record is low in diversity
18 compared to adjoining states, with the majority of specimens being identified as either
19 phytosaurs or aetosaurs (Martz et al., 2014).

20 In May of 2014 a paleontological expedition was conducted by Mission Heights
21 Preparatory High School to Comb Ridge in southeastern Utah. During the expedition
22 two of the authors (AM and IS) discovered a new, very rich microsite they dubbed The

Hills Have Teeth (Museum of Northern Arizona Locality 1724), near a locality that was previously discovered by the senior author (RG). Both at The Hills Have Teeth and the alluvial fan immediately adjacent to the hill a dozen partial and complete tetrapod teeth were collected. Most of these teeth belonged to phytosaurs and temnospondyl amphibians. Two teeth were notably different from the dominant taxa. One, discovered by IS, is described elsewhere (Gay and St. Aude, 2015). The other was collected by one of the authors (AM) and defied classification at the time of discovery. Since then we have had the opportunity to compare this new specimen to other identified teeth from across the Chinle and Dockum Formations. That specimen, MNA V10668, is compared here to many Triassic diapsids to help classify it. We compare it to the non-archosauriform archosauromorphs *Azendohsaurus* (Flynn et al., 2010), *Mesosuchus browni* (Dilkes, 1998), and *Terraterpeton hrynewichorum* (Sues, 2003), several non-archosaurian archosauriforms including *Crosbysaurus harrisae* (Heckert, 2004), *Crosbysaurus sp.* (Gay and St. Aude, 2015), *Krzyzanowskisaurus hunti* (Heckert, 2005), *Lucianosaurus wildi* (Hunt and Lucas, 1995), *Protecovasaurus lucasi* (Heckert, 2004), *Revueltosaurus callendari* (Hunt, 1989), *Tecovasaurus murrayi* (Hunt and Lucas, 1994), unidentified or unnamed archosauriform teeth (Heckert, 2004), and several archosaurs (Colbert, 1989; Dalla Vecchia, 2009; Heckert, 2004).

Materials and Methods:

Standard paleontological field materials and methods were used to collect all specimens from MNA locality 1725, as described in Gay and St. Aude (2015). MNA V10668 was recorded using Backcountry Navigator Pro running on an Android OS smartphone. It was collected in a zip-seal collection bag after being removed from the

surface exposure by a hand. Measurements of MNA V10668 were obtained using a set of Craftsman metal calipers (model 40257) with 0.05mm precision. Figures were created using GIMP 2.8.4. Photos were captured taken with an Olympus E-500 DSLR and PC USB digital microscope. MNA V10668 was collected under Bureau of Land Management permit UT14-001S and is permanently housed at the Museum of Northern Arizona (MNA).

Geologic Setting:

MNA V10668 was found on the surface of Lower Member of the Chinle Formation at Comb Ridge, Utah (Figure 1), roughly 6 meters from the base of the Lower Member along with teeth of phytosaurs, temnospondyls, and *Crosbysaurus*. (Gay & St. Aude 2015) at MNA Locality 1725. As with earlier work, we hold that material from locality 1725 has washed down slope from The Hills Have Teeth outcrop, MNA locality 1724. In May of 2015 the precise fossil-bearing horizon was located at The Hills Have Teeth. The horizon is a light grey mudstone with interspersed carbonaceous clasts and numerous teeth (Figure 2). This mudstone is 13 cm below the red brown mudstone-grading-to-shale, 8.75 meters above the base of the Chinle Formation (Gay and St. Aude, 2015; figure 4). The fossil-bearing Hills Have Teeth bed is exposed locally for about half a kilometer in the Rainbow Garden area and appears be present where the base of the Chinle Formation is exposed all along the western face of Comb Ridge.

Description:

MNA V10668 is a single tooth crown that is flattened labiolingually and concial in profile. It measures 5 mm apicobasally and 3mm mesiodistally. The distal side of the tooth has a continuous serrated edge from the base to the apex. These distal serrations

are 0.1 mm in length. There are eight serrations per millimeter with an estimated thirty serrations along the entirety of the distal keel. The serrations show increasing wear apically with the apex itself completely worn away. These serrations are stacked and do not stagger as they progress to the apex of the specimen. The mesial side is missing most of its enamel so identification of features is difficult. None the less the dentine does preserve the traces of several apical serrations but there is no evidence of a pronounced keel mesially. There is no root preserved and a small resorption pit is present on the base, suggesting this is a shed tooth crown. The tooth has a small chip on its base, distal to the midline (Figure 3, 4).

Differential Diagnosis:

MNA V10668 differs from most described Triassic teeth with serrations on only one side. Because this morphology may be due to taphonomic processes, we compare MNA V10668 to other diapsids with thecodont or sub-thecodont dentition with both mesial and distal serrations as well as those only possessing distal serrations.

Azendohsaurus is an archosauromorph reptile from Madagascar known from reasonably complete remains (Flynn et al., 2010). Its dentition is well documented and illustrated, allowing comparisons to be made easily. *Azendohsaurus* teeth are slightly recurved with a basal constriction while MNA V10668 appears to be conical with no mesiodistal constriction apical to the base. The teeth of *Azendohsaurus* do not possess significant wear facets or worn denticles, as MNA V10668 does. The denticles that exist on the teeth of *Azendohsaurus* are apically directed. In MNA V10668 the preserved distal denticles appear perpendicular to the long axis of the tooth. The denticles of

Azendohsaurus are also much larger and fewer in number than those of MNA V10668. MNA V10668 clearly does not represent a specimen of *Azendohsaurus*.

Mesosuchus browni is a basal rhynchosaur, deeply nested within archosauromorpha, (Dilkes, 1998) known from multiple specimens. The dentition of *Meosuchus* is rounded in cross-section and conical in profile. The tooth-jaw junction is not well preserved enough to say whether the teeth had thecodont implantation. Dilkes (1998) noted an unusual wear facet on the teeth of *Mesosuchus*, which is why it is included here. Despite MNA V10668 and *Mesosuchus* both having erosional surfaces, those on *Mesosuchus* are mesiolabially directed while in MNA V10668 the wear is mesiobasal. Coupled with the differences in cross-sectional profile, MNA V10668 does not represent a specimen of *Mesosuchus* or any rhynchosaur by extension.

The unusual archosauromorph *Terraterpeton hrynewichorum* from the Triassic of Nova Scotia was first described by Sues (2003). The teeth of *Terraterpeton* are as odd as the rest of its skull. The teeth are round to oval in cross-section, with the posterior-most teeth being much broader labiolingually than mesiodistally. The teeth have a distal triangular cusp and a flattened area mesially on each occlusal surface. The narrow, conical profile and labiolingually compressed cross-section of MNA V10668 strongly differs from the teeth of *Terraterpeton* in all these aspects, excluding it as the animal that possessed MNA V10668 during the Triassic.

Crosbysaurus (Heckert, 2004) is an archosauriform that has serrations on both mesial and distal sides of the tooth, with the distal serrations being much larger than those on the mesial keel. These denticles are subdivided and on the distal keel they point apically. *Crosbysaurus harrisae* and MNA V10668 have a similar shape and size.

Both MNA V10668 and *Crosbysaurus* teeth are similar in size apicobasally and have the same triangular shape in labial and lingual views. *Crosbysaurus* teeth are distally curved on the apicomesimal keel, a condition not seen in MNA V10668.

MNA V10668 and MNA V10666, referred to *Crosbysaurus* sp. by Gay and St. Aude (2015), were both found at the same locality. Because of the close association between these two specimens we paid special attention to MNA V10666 when considering the affinities of this new specimen. MNA V10666 does lack serrations on the mesiobasal keel, as does MNA V10668. That is where the similarities end. The tooth referred to as *Crosbysaurus* sp. by Gay and St. Aude (2015) has clear mesial denticles towards the apex. The distal denticles are much larger and subdivided, as in all other *Crosbysaurus* teeth. While MNA V10668 is labiolingually compressed like MNA V10666 and other known *Crosbysaurus* teeth, it is not as mesiodistally narrow. Considering that *Crosbysaurus* serrations are larger, present on the mesial side, apically directed, and the teeth tend to be mesiodistally narrower it is doubtful that MNA V10668 is a *Crosbysaurus* tooth.

Krzyzanowskisaurus hunti (Heckert 2005) is a (presumably) small herbivorous pseudosuchian known only from dental remains. It superficially resembles *Revueltosaurus* but can be diagnosed by the presence of a cingulum on the base of the tooth. Since MNA V10668 does not have a cingulum it is obvious that it cannot be a specimen of *Krzyzanowskisaurus*.

Lucianosaurus wildi (Hunt and Lucas, 1995) is similar to other isolated Triassic teeth described in the literature by having enlarged denticles and a squat shape with convex mesial and distal edges, being mesiodistally broad while apicobasally short.

138 MNA V10668 is taller than it is long and has relatively small denticles. MNA V10668
139 does not represent *Lucianosaurus*.

140 *Protecovasaurus lucasi* (Heckert, 2004) is diagnosed by having a recurved
141 mesial surface where the apex is even with or overhangs the distal margin. The
142 denticles on both the mesial and distal keels are apically directed. In all these features
143 the teeth of *Protecovasaurus* do not match the features seen in MNA V10668.

144 *Revueltosaurus* (Hunt, 1989; Heckert, 2002; Parker et al., 2005) has serrations
145 on both the mesial and labial sides. Its serrations are proportionally larger and closer
146 together. The teeth of *Revueltosaurus* are broader mesiodistally compared to their
147 apicobasal height. In general *Revueltosaurus* teeth have more serrations on the distal
148 keel of the tooth than at the mesial side of the tooth. Furthermore, *Revueltosaurus* has
149 been distinguished by more than it's teeth (Parker et al., 2005). MNA V10668 is
150 labiolingually narrower than the teeth of *Revueltosaurus*. These differences rule out the
151 possibility that MNA V10668 is *Revueltosaurus*.

152 Heckert (2004) described some tetrapod teeth found from other localities across
153 the Chinle Formation. Some of these teeth are from phytosaurs (Heckert, 2004, figure
154 43). NMMNH P-30806 for example is roughly conical in outline and somewhat
155 labiolingually compressed. The serrations are orientied perpendicular to the long axis of
156 the tooth. In these regards young phytosaur teeth are similar to MNA V10668. Unlike
157 MNA V10668, however, these teeth are moderately curved and have serrations on their
158 mesial surface. In addition the serrations on phytosaur teeth, like the ones figured in
159 Heckert (2004), are more dense per millimeter compared to MNA V10668. Phytosaur
160 teeth in general, especially the teeth from segments of the jaw posterior to the

premaxillary rosette, tend to be more robust than MNA V10668. Although phytosaurs are the most common taxa represented at The Hills Have Teeth it not likely MNA V106668 is a phytosaur tooth.

Heckert described another specimen, NMMNH P-34013 (Heckert, 2004, figure 20), that is roughly the same size as MNA V10668. Both have a resorption pit at the base. However the serrations on NMMNH P-34013 are smaller than MNA V10668, and has a slight curve unlike MNA V10668. Heckert described this tooth as belonging to an indeterminate archosauriformes. Despite their differences this tooth, NMMNH P-34013, is the closest tooth to MNA V10668 yet identified.

Based on the examination of a skull cast of *Coelophysis bauri* at Mission Heights Preparatory High School and from the literature (Colbert, 1989), it can be seen that *Coelophysis* and MNA V10668 have a similar tooth shape and size. This is especially true for teeth from the mid-posterior region of the maxilla of *Coelophysis*. Both teeth are 5mm tall from the apex to the base. When they are looked at closely many things stand out as to why they are different. *Coelophysis* teeth are naturally recurved, at least slightly, whereas MNA V10668 does not have a noticeable curve to it. *Coelophysis* teeth have small serrations along the mesial and distal sides. *Coelophysis* teeth tend to be even more mesiodistally compressed and the serrations at the distal side are completely different. *Coelophysis* tooth serrations are smaller and are closer together to each other. We can conclude that MNA V10668 cannot be a *Coelophysis* tooth.

Austriadactylus teeth (Dalla Vecchia, 2009) and MNA V10668 are completely different in shape and size. *Austriadactylus* teeth are smaller and sharper; also they have serrations at the mesial and labial sides of the tooth. The serrations are completely

different because they are larger and possess more distinct tips. *Austriadactylus* has a few different types of teeth. Most teeth are small, have three cusps, and a slight curve to them. Other teeth have only one distinct cusp and have a slight curve to them. They have very few and large serrations. MNA V10668 differs from all of the *Austriadactylus* teeth as it has no visible curve, and serrations along the mesial side. Seeing this, MNA V10668 does not represent *Austriadactylus*.

Reported Chinle prosauropod teeth, such as those figured in Heckert (2004, figures 45, 83, 84) are extremely mesiolaterally compressed. They also exhibit serrations on the mesial and labial sides of the tooth. Its serrations are relatively larger, closer together, and are apically directed. Also prosauropod teeth have a distinctly “pointy” apex with no wear facets. Its shape is completely different because this MNA V10668 is relatively wider labiolingually and apicobasally smaller than the reported prosauropod specimens. There is no possibility that the specimen is a prosauropod. It should also be noted that the extreme convergence seen in *Azhendousaurus* (Flynn et al., 2010) makes the identification of prosauropods from the Chinle Formation tentative at best.

The most common vertebrate remains from the Chinle Formation are phytosaur teeth. Despite the small size of MNA V10668 it is possible that this specimen pertains to a juvenile phytosaur. To test this hypothesis two juvenile phytosaur snouts were examined at the Museum of Northern Arizona. One of these, PEFO 13890/MNA V1789 was collected by Billingsly in 1979 from the Upper Petrified Forest Member of the Chinle Formation in Petrified Forest National Park (PEFO). It represents articulated paired premaxillae with 15 preserved alveoli on the right and 14 on the left, all of which save

one are empty. The total preserved length of this specimen is 9.3 cm. While identified in collections as *Pseudopalatus zunii* there are no preserved autapomorphies to support this assignment.

The second specimen, MNA V3601, is a partial right dentary from the Blue Mesa Member of the Chinle Formation (Ramezani et al., 2014) Placerias Quarry, near St. Johns, Arizona identified as *Leptosuchus* sp. (Long and Murray, 1995). MNA V3601 is 4.95 cm in length, preserving the anterior tip and eight alveoli. In this specimen several of the tooth crowns are present and show wear while others are broken off at the gum line or inside the alveolus.

In both specimens the juvenile phytosaurs exhibit remarkable homodonty, especially considering the heterodonty seen in more mature phytosaurs (Heckert, 2004). While MNA V10668 is roughly the right size of tooth to have come from a juvenile phytosaur similar in age to PEFO13890/MNA V1789 or MNA V3601, the basal structure of the tooth is unlike any preserved juvenile phytosaur teeth or alveoli. Both undisputed juvenile phytosaur specimens have round alveoli with serrated or unserrated conical teeth preserved (Figure 5). In addition, all preserved teeth in MNA V3601 do not show any lingual curvature as seen in MNA V10668. While adult phytosaurs have triangular, lingually curved teeth in their dentition, especially as one moves posteriorly (Long and Murray, 1995; Heckert, 2004), these seem to be absent in juveniles from the specimens we have on hand. The lingually curved teeth of adult phytosaurs are also much more robust, with labiolingually wide basal and mid-crown section, unlike the laterally compressed and teardrop-shaped base of MNA V10668. It may be that phytosaur dentition changed during ontogeny to adapt to a changing diet. Even

considering this we do not think that MNA V10668 can be assigned to the phytosauria due to the marked differences between it and all other known phytosaur teeth.

Conclusions:

MNA V10668 cannot be identified as any previously described Triassic taxon as it does not have any distinguishing autapomorphies preserved. However, this tooth can be identified at least as archosauriformes *incertae sedis*. MNA V10668 has many characteristics that match up with other archosauriformes. Another indeterminate tooth, NMMNH P-34013, is the closest tooth to MNA V10668. Despite their similarities it is obvious that MNA V10668 is morphologically distinct from NMMNH P-34013. Although isolated teeth have been described before from Utah (Heckert et al., 2006; Gay and St. Aude, 2015) this is the first tooth morphotype described from Utah to not be assigned to an existing genus of Triassic tetrapod. As such it may represent an animal endemic to what is now Utah.

These findings are important since they demonstrate the existence of a previously unrecognized clade of diapsids from the Chinle Formation in Utah. In addition, most of the tetrapod record from Utah's Chinle Formation has come from the Church Rock Member (Martz et al., 2014; RG pers. obs.) This specimen, coming from the Lower Member of the Chinle Formation, demonstrates increased diversity in an older part of the formation that has not been studied until recently (Gay and St. Aude, 2015).

Work is ongoing at Comb Ridge by crews from Mission Heights Preparatory High School. The tetrapod diversity of Chinle Formation at Comb Ridge will continue to

increase as new discoveries come to light. It is hoped that additional taxa can be added to the growing faunal list with additional fieldwork in the near future.

Acknowledgements:

The authors would like to thank Jason Durivage, Gary Shepler, Steven Hall, and Deborah Avey for their assistance with fieldwork while MNA V10668 was collected. We would also like to thank David and Janet Gillette for their assistance with collections and access to specimens. We would also like to thank ReBecca Hunt-Foster for her assistance with our permit.

Works Cited

Colbert EH. 1989. The Triassic dinosaur *Coelophysis*. Flagstaff, AZ: Museum of Northern Arizona Press.

Dalla Vecchia, FM. 2009. The first Italian specimen of *Austriadactylus cristatus* (Diapsida, Pterosauria) from the Norian (Upper Triassic) of the Carnic Prealps. *Rivista Italiana di Paleontologia e Stratigrafia*, **115**(3):291-304.

Dilkes DW. 1998. The early Triassic rhynchosaur *Mesosuchus browni* and the interrelationships of basal archosauromorph reptiles. *Philosophical Transactions of the Royal Society B: Biological Sciences* **353**:501–541.

Flynn JJ., Nesbitt SJ., Parrish JM., Ranivoharimanana L., Wyss AR. 2010. A new species of *Azendohsaurus* (Diapsida: Archosauromorpha) from the Triassic Isalo Group of southwestern Madagascar: cranium and mandible. *Palaeontology* **53**:669–688.

Gay RJ., Aude IS. 2015. The first occurrence of the enigmatic archosauriform *Crosbysaurus* Heckert 2004 from the Chinle Formation of southern Utah. *PeerJ* **3**.

- 276 **Gibson SZ.** 2013. A new hump-backed ginglymodian fish (Neopterygii,
277 Semionotiformes) from the Upper Triassic Chinle Formation of southeastern Utah.
278 Journal of Vertebrate Paleontology **33**:1037–1050.
- 279 **Heckert AB.** 2002. Heckert AB, Lucas SG, eds. *A revision of the Upper Triassic*
280 *ornithischian dinosaur Revueltosaurus, with a description of a new species*, Upper
281 Triassic stratigraphy and paleontology. Albuquerque: New Mexico Museum of Natural
282 History and Science. vol. **21**:253-268
- 283 **Heckert AB.** 2004. Late Triassic microvertebrates from the lower Chinle (Otischalkian-
284 Adamanian:Carnian), southwestern USA. Albuquerque: New Mexico Museum of Natural
285 History and Science. **Vol. 27**.
- 286 **Heckert AB.** 2005. Heckert AB, Lucas SG, eds. *Krzyzanowskisaurus*, a new name for a
287 probable ornithischian dinosaur from the Upper Triassic Chinle Group, Arizona and New
288 Mexico, USA, Vertebrate paleontology in Arizona. Albuquerque: New Mexico Museum
289 of Natural History and Science. vol. **29**:77-83
- 290 **Heckert, AB., Lucas, SG., DeBlieux, DD., Kirkland, JI.** Harris JD, Lucas SG,
291 Spielmann JA, Lockley MG, Milner ARC, Kirkland JI, eds. *A revueltosaurus-like tooth from*
292 *the Petrified Forest Formation (Upper Triassic: Revueltian) from Zion National Park*, The
293 Triassic-Jurassic Terrestrial Transition. Albuquerque: New Mexico Museum of Natural
294 History and Science. vol. **37**:588-591
- 295 **Hunt AP.** 1989. A new ?ornithischian dinosaur from the Bull Canyon Formation (Upper
296 Triassic) of east-central New Mexico. In: Lucas SG, Hunt AP, eds. *Dawn of the age of*
297 *dinosaurs in the American Southwest*. Albuquerque: New Mexico Museum of Nature
298 and Science. 355-358

- 299 **Hunt AP, Lucas SG.** 1994. Ornithischian dinosaurs from the Upper Triassic of the
300 United States. In: Sues H-D, Fraser NC, eds. *In the shadow of the dinosaurs: early*
301 *Mesozoic tetrapods*. Cambridge: Cambridge University Press. 227-241
- 302 **Long RA., Murry PA.** 1995. Late Triassic (Carnian and Norian) tetrapods from the
303 southwestern United States. Albuquerque: New Mexico Museum of Natural History and
304 Science.
- 305 **Martz JW, Irmis RG, Milner ARC.** 2014. Lithostratigraphy and biostratigraphy of the
306 Chinle Formation (Upper Triassic) in southern Lisbon Valley, southeastern Utah. *UGA*
307 *Publication* **43**(2014):397-446.
- 308 **Morales, M., Ash, SR.** 1993. The last phytosaurs. *The Nonmarine Triassic*. New
309 Mexico Museum of Natural History and Science. vol. **3**:357-358
- 310 **Nesbitt SJ., Sidor CA., Irmis RB., Angielczyk KD., Smith RMH., Tsuji LA.** 2010.
311 Ecologically distinct dinosaurian sister group shows early diversification of Ornithodira.
312 *Nature* **464**:95–98.
- 313 **Parker WG., Ash SR., Irmis RB.** 2006. A century of research at Petrified Forest
314 National Park: geology and paleontology. Flagstaff: Museum of Northern Arizona.
- 315 **Ramezani J., Fastovsky DE., Bowring SA.** 2014. Revised chronostratigraphy of the
316 Lower Chinle Formation strata in Arizona and New Mexico (USA): high-precision U-Pb
317 geochronological constraints on the Late Triassic evolution of dinosaurs. *American*
318 *Journal of Science* **314**: 981-1008
- 319 **Sues H-D.** 2003. An unusual new archosauromorph reptile from the Upper Triassic
320 Wolfville Formation of Nova Scotia. *Can. J. Earth Sci. Canadian Journal of Earth*
321 *Sciences* **40**:635–649.




Figure 1: The location of Comb Ridge in the state of Utah. Modified from Gay and St. Aude (2015)




Figure 2: Tetrapod tooth *in situ* at The Hills Have Teeth (MNA locality 1724), showing the lithology of the fossil-bearing layer.




Figure 3: MNA V10668 in A) lingual, B) labial, C) distal, D) mesial, E) apical, and F) basal views. Scale bar = 1 mm.




Figure 4: Interpretive line drawing of MNA V10668 in A) lingual, B) labial, C) distal, D) mesial, E) apical, and F) basal views. In A-D, dark gray represents missing enamel, light gray represents a resorption pit. In E and F no missing enamel is figured; dark gray represents worn apex, light gray represents resorption pit. Scale bar = 1 mm.



Figure 5: Juvenile phytosaur jaws. Top: PEFO 13890/MNA V1789, *Macheroprosopus zunii* premaxillae in A) ventral view. Bottom: MNA V3601 right dentary in B) lateral C) dorsal views. Scale bar = 1 cm.